

REVIEW ARTICLE

A REVIEW ON THE MEDICINAL PLANT *PSIDIUM GUAJAVA* LINN. (MYRTACEAE)*Shirur Dakappa Shruthi^{1,2}, Adhikari Roshan¹ Sanjay Sharma Timilsina¹ and Sajjekhan Sunita²¹P.G. Department of Biotechnology, The Oxford College of Science, Bangalore 560102, Karnataka, India.²P.G. Department of Studies and Research in Biotechnology & Bioinformatics, Kuvempu University, Shankaraghatta – 577451, Karnataka, India*Corresponding author's Email: sdshruthi@gmail.com

ABSTRACT

Psidium guajava is an important food crop and medicinal plant available in tropical and subtropical countries, widely used in food and folk medicines around the world. It contains important phytoconstituents such as tannins, triterpenes, flavonoid: quercetin, pentacyclic triterpenoid: guajanoic acid, saponins, carotenoids, lectins, leucocyanidin, ellagic acid, amritoside, beta-sitosterol, uvaol, oleanolic acid and ursolic acid. In view of the immense medicinal importance of the plant, this review is an effort to compile all the information reported on its ethanobotanical, phytochemical and pharmacological activities. The present work attempts to generate interest among the masses regarding its potential in preventing and treating several common diseases. Many pharmacological studies have demonstrated the ability of this plant to exhibit antioxidant, hepatoprotective, anti-allergy, antimicrobial, antigenotoxic, antiplasmodial, cytotoxic, antispasmodic, cardioactive, anticough, antidiabetic, antiinflammatory and antinociceptive activities, supporting its traditional uses. Suggesting a wide range of clinical applications for the treatment of infantile rotaviral enteritis, diarrhoea and diabetes.

Key words: ethanobotany, myrtaceae, pharmacology, physicochemical, phytochemical, *Psidium guajava*

INTRODUCTION

Psidium guajava L. known as Guava is a medicinal plant belonging to the family Myrtaceae. *P. guajava* is a well known traditional medicinal plant used in various indigenous systems of medicine. It is widely distributed throughout India¹. The leaves and bark of *P. guajava* tree have long history of medicinal uses, that is still employed today. It is a native of Central America but is now widely cultivated, distributed and the fruits enrich the diets of millions of people in the tropics of the world^{2,3}. It is a genus of about 133 genera and more than 3,800 species of tropical shrubs and a small tree of about 10m high with spreading branches that thrives on all kinds of soils. It is one of the most gregarious fruit trees and is widely known by its common English name (guava). In Nigeria, it is called guava (Hausa), gurfa (Yoruba) and Gwaibwa (Igbo)³. *P. guajava* also known as the 'poor man's apple' of the tropics has a long history of traditional use, and a good proportion of which have been validated by scientific research⁴.

Nutritional value of guavas are often included among superfruits, being rich in dietary fiber, vitamins A and C, folic acid; and the dietary minerals, potassium, copper and manganese. Having a generally broad, low calorie profile of essential nutrients, a single common guava fruit contains about four times the amount of vitamin C as an orange. The food value and contents of guava fruit is listed in the **Table 1**. However, nutrient content varies across guava cultivars.

Distribution

It is now cultivated in Southern Florida, Bermuda, and throughout the West Indies from the Bahamas and Cuba to Trinidad, and south to Brazil⁶.

Table 1: Nutrient value of guava fruit⁵

| Nutrients | Content |
|----------------------|--------------|
| Moisture | 2.8-5.5g |
| Crude fiber | 0.9-1.0g |
| Protein | 0.1-0.5mg |
| Fat | 0.43-0.7mg |
| Ash | 9.5-10mg |
| Carbohydrate | 9.1-17mg |
| Calcium | 17.8-30mg |
| Phosphorous | 0.30-0.70mg |
| Iron | 200-400 I.U. |
| Carotene (Vitamin A) | 0.046mg |
| Thiamin | 0.03-0.04mg |
| Riboflavin | 0.6-1.068mg |
| Niacin | 40 I.U. |
| Vitamin | 36-50mg |

Taxonomy

Kingdom : Plantae
 Order : Myrtales
 Family : Myrtaceae
 Subfamily : Myrtoideae
 Genus : *Psidium*
 Species : *Guajava*
 Binomial name : *Psidium guajava* Linn.

P. guajava has a long history of traditional use, a good proportion of which have been validated by scientific research⁴. The ethno-medicinal uses include the crushing of the leaves and the application of the extract on wounds, boils, skin and soft tissue infectious site⁷. Stem, bark and root-bark are astringent. Unripe fruit is indigestible, causes vomiting and feverish. Fruit is laxative, leaves are astringent⁸. Locally, decoction of the leaves is with much

benefit to the prolapsus ani of children⁸; ache upsets and for vertigo⁶. *P. guajava* leaf is a phytotherapeutic used to treat gastrointestinal and respiratory disturbances and is used as anti-inflammatory medicine. Its anti-amoebic and antimalarial effects have also been documented^{9,10}. Guava fruit paste and cheese are popular dishes in Florida, the West Indies and parts of South America¹¹. The plant *P. guajava* Linn. has an ethnomedical history as it has various activities especially functionally against the hyperglycemia.

Table 2: Showing worldwide ethnomedical uses of Guava⁵

| | |
|--------------------|--|
| Amazonia | for diarrhea, dysentery, menstrual disorders, stomach ache, vertigo |
| Brazil | for anorexia, cholera, diarrhea, digestive problems, dysentery, gastric insufficiency, inflamed mucous membranes, laryngitis, mouth(swelling), skin problems, sore throat, ulcers, vaginal discharge |
| Cuba | for cold, dysentery, dyspepsia |
| Ghana | coughs, diarrhea, dysentery, toothache |
| Haiti | for dysentery, diarrhea, epilepsy, itch, piles, scabies, skin sores, sore throat, stomachache, wounds and as an antiseptic and astringent |
| India | for anorexia, cerebral ailments, childbirth, chorea, convulsions, epilepsy, nephritis |
| Malaya | for dermatosis, diarrhea, epilepsy, hysteria, menstrual disorders |
| Mexico | for deafness, diarrhea, itch, scabies, stomachache, swelling, ulcer, worms, wounds |
| Peru | for conjunctivitis, cough, diarrhea, digestive problems, dysentery, edema, gout, hemorrhages, gastroenteritis, gastritis, lung problems, PMS, shock, vaginal discharge, vertigo, vomiting, worms |
| Philippines | for sores, wounds and as an astringent |
| Trinidad | bacterial infections, blood cleansing, diarrhea, dysentery |
| Elsewhere | for anorexia, aches, bacterial infections, boils, bowel disorders, bronchitis, catarrh, cholera, chorea, colds, colic, convulsions, coughs, diarrhea, dysentery, dyspepsia, edema, epilepsy, fever, gingivitis, hemorrhoids, itch, jaundice, menstrual problems, nausea, nephritis, respiratory problems, rheumatism, scabies, sore throat, spasms, sprains, stomach problems, swelling, tonic, toothache, ulcers, worms, wounds and as an antiseptic and astringent |

Various parts of the plant has been used in traditional medicine.

Leaves: The decoction or infusion of the leaves is used as febrifuge, antispasmodic and for rheumatism in India¹². The leaves are used in USA as an antibiotic in the form of poultice or decoction for wounds, ulcers and tooth ache^{13,14}. Bronchitis, asthma attacks, cough, pulmonary diseases could be also treated with guava teas^{15,16}.

Bark: The bark in the form of decoction and poultice is used as an astringent in the treatment of ulcers wounds and diarrhea in Philippines while in Panama, Bolivia and Venezuela, the bark is used in treatment of dysentery and skin ailments. In the form of decoction and poultice, it is used to expel the placenta after childbirth and in infections

of the skin, vaginal hemorrhage wounds, fever, dehydration and respiratory disturbances¹⁷.

Root: The root is used in West Africa as a decoction to relieve diarrhea, coughs, stomach ache, dysentery, toothaches, indigestion and constipation; while in Philippines, Fiji and South Africa, the roots are used in the form of decoction and poultice as an astringent in ulcers wounds and in treatment of diarrhea¹⁷.

Whole plant: In general, the whole plant or it shoots are used in the form of infusion, decoction and paste as skin tonic in Tahiti and Samoa and as analgesia in painful menstruation, miscarriages, uterine bleeding, premature labor and wounds¹⁷.



Figure 1: Whole plant of *Psidium guajava* L.(A) along with leaves (B); fruit (C) and flower (D) Pharmacognosy

Macroscopy

P. guajava is a large dicotyledonous shrub, or small evergreen tree, generally 3-10 m high with many branches¹⁸. The stems are crooked and the bark is light to reddish brown, thin, smooth and continuously flaking. Root system is generally superficial and very extensive, frequently extending well beyond the canopy. Each has some deep roots but no distinct taproot¹⁷. The leaves are opposite and simple; stipules are absent, petiole short, 3-10 mm long; blade oblong to elliptic, veins prominent, gland dotted¹⁸. The flowers are white, incurved petals, 2 or 3 in the leaf axils; they are fragrant, with four to six petals and yellow anthers. The fruit is small, 3 to 6 cm long, pear-shaped, reddish-yellow when ripe⁶. The fruit contains several small seeds and consists of a fleshy pericarp and seed cavity with pulp^{19,20,21}.

Microscopy

The microscopy reveals the presence of abundant unicellular trichome, paracytic stomata, xylem vessels, calcium crystals and few crystal sheath^{22,23,24}. The average stomatal index on upper epidermis is more than on lower epidermis for mature leaf and found opposite in case of young leaf²³. Young leaves contain more moisture, moreover, water and alcohol soluble compounds are present nearly in same amount in both young and mature leaves. The transverse section of midribs show gutter shaped xylem and phloem and beneath the phloem pericycle present, which contain collenchymatous cells²⁴.

Physico-chemical studies

The proximate analysis of powdered sample of *P. guajava* leaves showed low moisture content of 1.0% and low ash value of 2.80%. The ash value indicates the quantity of inorganic components of the plant; hence a low value indicates that the powdered leaves of *P. guajava* contain more of organic components. There is also low quantity of

protein and fat but an abundance of carbohydrate which makes it a source of energy. The metal analysis of powdered sample of *P. guajava* showed the presence of all the metals screened for which include; magnesium, manganese, zinc, calcium, iron, sodium and potassium. The quantity of these metals revealed that they were well below tolerable upper intake level and within the recommended daily intake in healthy individuals established by the Dietary Reference Intakes¹⁸.

The plant extracts were also screened for the presence of bio-active components in the leaves and roots^{25,26,27}. The results of phytochemical analysis showed that the crude hydro-ethanolic leaf extract of *P. guajava* have different classes of bioactive constituents such as saponins, alkaloids, tannins, cardiac-glycosides, terpenes, flavonoids and sterols. The results showed that saponins, tannins and alkaloids are present in high concentrations, followed by terpenes, flavonoids and sterols while small concentration of cardiac-glycosides²⁸.

Phytochemistry

Guava contains broad spectrum of phytochemicals including minerals, enzymes, proteins²⁹, sesquiterpenoid alcohols and triterpenoid acids^{30,31,32}, alkaloids, glycosides, steroids, flavanoids, tannins, saponins^{33,34}. Guava is very rich in antioxidants and vitamins and also high in lutein, zeaxanthine and lycopene^{35,36}. The guava leaves contain several chemical constituents such as α -pinene, β -pinene, limonene, menthol, terpenyl acetate, isopropyl alcohol, longicyclene, caryophyllene, β -bisabolene, caryophyllene oxide, β -copanene, farnesene, humulene, selinene, cardinene and curcumene, mallic acids, nerolidiol, β -sitosterol, ursolic, crategolic, and guayavolic acids, cineol, quercetin, 3-L-4-4-arabinofuranoside (avicularin) and its 3-L-4-pyranoside (essential oil), resin, tannin, eugenol,

caryophyllene (1a α -, 4a α -, 7 α -, 7a β -, 7b α -)-decahydro-1H-cycloprop[e] azulene, Guajavolide (2 α -,3 β -,6 β -,23-tetrahydroxyurs-12-en-28,20 β -olide; 1) and guavenoic acid (2 α -,3 β -,6 β -,23-tetrahydroxyurs-12,20(30)-dien-28-oic acid, triterpene oleanolic acid, triterpenoids, flavinone-2 2'-ene, prenol, dihydrobenzophenanthridine and cryptonine^{37,8,38,39,32,40}. Guavas contain carotenoids and polyphenols, the major classes of antioxidant pigments giving them relatively high potential antioxidant value among plant foods¹⁹. As these pigments produce the fruit skin and flesh color, guavas that are red-orange have more pigment content as polyphenol, carotenoid and pro-vitamin A, retinoid sources than yellow-green ones.

Guavas contain both carotenoids and polyphenols like (+)-galocatechin, guajaverin, leucocyanidin and amritoside⁴¹. It was reported that the leaves of *P. guajava* contain an essential oil rich in cineol, tannins and triterpenes. In addition three flavonoids (quercetin, avicularin, and guajaverin) have been isolated from the leaves⁴². The leaves of guava are rich in flavonoids, particularly quercetin. The bark of guava tree contains considerable amounts of tannins (11-27%), and hence is used for tanning and dyeing purposes.

Leucocyanidin, luectic acid, ellagic acid and amritoside have been isolated from the stem bark. Five constituents, including one new pentacyclic triterpenoid: guajanoic acid and four known compounds beta-sitosterol, uvaol, oleanolic acid and ursolic acid, have been recently isolated from the leaves of *P. guajava* by Begum *et al.*³². The essential oil contains alpha pinene, caryophyllene, cineol, D-limonene, eugenol, and myrcene. The major constituents of the volatile acids include (E)-cinnamic acid and (Z)-3-hexenoic acid⁴³. The guava fruit has high water content with lesser amounts of carbohydrates, proteins and fats. The fruit also contains iron, vitamins A and C, thiamine, riboflavin, niacin and manganese. The characteristic fruit odor is attributed to carbonyl compounds. Unripe fruits are high in tannins. The major constituent of the fruit skin is ascorbic acid, largely destroyed by canning and processing¹⁷.

Active factors of *P. guajava* fruits involve ursolic acid, oleanolic acid, arjunolic acid and glucuronic acid⁴⁴. In comparison, huge amounts of β -sitosterol glucoside, brahmic acid, and polyphenolics including gallic acid, ferulic acid, and quercetin⁴⁵ and triterpenoids⁴⁶, exist in guava leaves⁴⁵. Thus, it is clear that *P. guajava* contains many components reported to display efficacy against various diseases.

Pharmacology

Pharmacology of extracts:

The aqueous extract of *P. guajava* leaves exhibited good antibacterial activity against various test cultures. Report says that flavanoids extracted from guava leaves believed to be responsible for antibacterial activity⁴⁷. The microbicidal activity of *P. guajava* is also attributable to guajaverine and to psydiolic acid⁴⁸. Joseph *et al.*⁴⁰ reported that guava leaf essential oil contains more terpenoids and that can strongly inhibit human cervical cancer cells. The leaf extract was found to possess anticestodal⁴⁹, analgesic, anti-inflammatory properties⁵⁰, antimicrobial⁵¹,

hepatoprotective and antioxidant activities⁵². In addition, the leaf extract is used in many pharmaceutical preparations as a cough sedative⁵³. It has demonstrated antibacterial and anti-diarrheal effects and is able to relax the intestinal smooth muscle and inhibit bowel contractions. Guava has antioxidant properties attributed to polyphenols found in its leaves. The presence of the metabolites such as cardiac glycosides, saponins, tannins, alkaloids in *P. guajava* may be responsible for its potential use as a drug against pathogenic bacteria^{54,55}. Alkaloids, flavonoids are phenolics structure containing one carbonyl group complexes with extracellular and soluble protein and with bacterial cell wall⁵⁶, thus exhibits antibacterial activity through these complexes.

In the pharmacological action guava leaf extracts have also been indicated to inhibit disturbances of the central nervous system: insomnia, convulsions and epilepsy^{57,58}. In addition, anti-rotavirus activity has also been reported to exist in these extracts⁵⁹. Bark and leaf extracts were shown to have *in vitro* toxic action against numerous bacteria. Water and chloroform extracts of guava were effective in activating the mutagenicity of *Salmonella typhimurium*⁶⁰. It was shown that *P. guajava* leaf extracts might be beneficial in treating acne especially those that have anti-inflammatory activities⁶¹.

In several studies, guava showed significant antibacterial activity against common diarrhea causing bacteria such as *Staphylococcus*, *Shigella*, *Salmonella*, *Bacillus*, *E. coli*, *Clostridium* and *Pseudomonas*, which was concluded that guava has good curative effect on infantile rotaviral enteritis⁶². In a study carried out with leaf extract of the plant, inhibition of gastrointestinal release of acetylcholine by quercetin present in extract was suggested as a possible mode of action in the treatment of acute diarrheal disease⁵⁷. Guava fruit and leaf showed antioxidant and free radical scavenging capacity⁵². Its leaf extract possess anticough activity by reducing the frequency of cough induced by capsaicin aerosol⁶³. Leaf extract of guava had inotropic effect on guinea pig atrium⁶⁴. Another study investigated that the hypoglycemic and hypotensive effects of *P. guajava* leaf aqueous extract in rats showed hypoglycemic activity. The hypoglycemic effect of plant extract was examined in normal and diabetic rats, using streptozotocin (STZ) induced diabetes mellitus model⁶⁵. In the study, i.p. treatment with 1 g/kg guava juice produced a markable hypoglycemic action in normal and alloxan-treated diabetic mice⁶⁶. The anti-stress and adaptogenic activity exhibited by ethanol extract of *P. guajava* possess anti-stress property. It may be useful in the treatment of several disorders caused by stress by its immunostimulating, immunomodulating properties and also by enhancing the homeostatic mechanisms. Aqueous extract of *P. guajava* budding leaves has been shown to possess anti-prostate cancer activity in a cell line model. Guava leaf essential oil has been shown to possess cytotoxic effect on human cervical cancer cell lines^{46,67}.

The leaf extract has an excellent capacity to form coloured complex with iron. Iron chelation therapy has been shown to be anti-parasitic especially in African trypanosomiasis. Furthermore, decreasing of free-radicals has antioxidizing effect in the body, meaning that polyphenols can prevent arterial sclerosis, thrombosis, cataract and inhibit senescence of the body and skin⁶⁸. In other animal studies,

guava leaf extracts have shown central nervous system (CNS) depressant activity⁶⁹. It can also be used as antihypertensive and antidiarrhoeal agents in traditional medicine, by inhibiting intracellular calcium release⁷⁰. *P. guajava* is reputed for its medicinal use in hyperactive gut disorders⁷¹. In a recent study with guinea pigs Brazilian researchers reported that guava leaf extracts have numerous effects on the cardiovascular system which might be beneficial in treating irregular beat (arrhythmia). Previous research indicated guava leaf provided antioxidant effects beneficial to the heart, heart protective properties, and improved myocardial function. Guava leaf extracts decreased spasms associated with induced diarrhea in rodents.

The *P. guajava* infusion at the higher concentration caused a statistically significant inhibition of cellular division in the onion root-tip cells⁷². During various episodes of screening of medicinal plants, extract from *P. guajava* leaves exhibited significant inhibitory effect on the protein tyrosine phosphatase1B (PTP1B). In a study including 17 Thai medicinal plants on anti-proliferative effects on human mouth epidermal carcinoma and murine leukemia cells using MIT assay, guava leaf showed anti-proliferative activity, which was 4.37 times more than vincristine⁷³.

Pharmacology of pure compounds:

Gallocatechin isolated from the methanol extract of guava leaf showed antimutagenic activity against *E. coli*⁷⁴. The active flavonoid compound quercetin-3-O-alpha-l-arabinopyranoside (guaijaverin) extracted from leaves has high potential antiplaque activity by inhibiting the growth of *Streptococcus mutans*⁷⁵. Lectin chemicals in guava were shown to bind to *E. coli*, preventing its adhesion to the intestinal wall and thus preventing infection and resulting diarrhea⁷⁶. Quercetin has several pharmacologic actions; it inhibits the intestinal movement, reduces capillary permeability in the abdominal cavity⁷⁷ and possesses dose-dependent antioxidant properties⁷⁸.

Clinical trials

In two randomized human studies, the consumption of guava fruit for 12 weeks was shown to reduce blood pressure by an average 8%, decrease total cholesterol level by 9%, decrease triglycerides by almost 8% and increase HDL cholesterol by 8%. The effects were attributed to the

high potassium and soluble fiber content of the fruit. A randomized, single-blind, controlled trial was conducted to examine the effects of guava fruit intake on blood pressure and blood lipids in patients with essential hypertension. It is possible that an increased consumption of guava fruit can cause a substantial reduction in blood pressure and blood lipids without decreasing HDL-cholesterol level^{79,80}. A double-blind clinical study of the effects of a Phytodrug (QG- 5) developed from guava leaf showed a decrease in duration of abdominal pain, which is attributed to antispasmodic effect of quercetin present in leaf extract²⁰. The fruit or fruit juice has been documented to lower blood sugar levels in normal and diabetic animals and humans. Most of these studies confirm the plant's many uses in tropical herbal medicine systems⁶⁴. In a clinical study with 62 infants with infantile rotaviral enteritis, the recovery rate was 3 days in those treated with guava, and diarrhea ceased in a shorter period than controls. It was concluded in the study that guava has 'good curative effect on infantile rotaviral enteritis'⁶².

CONCLUSION

Psidium guajava (Linn.) is popularly known as 'poor man's apple of the tropics', has a long history of traditional use for a wide range of ailments. The fruit as well as its juice is freely consumed for its great taste and nutritional benefits. Much of the traditional uses have been validated by scientific research. Toxicity studies in mice and other animal models as well as controlled human studies show both leaf and fruit are safe without any side effects. The plant has been extensively studied in terms of pharmacological activity of its major components, and the results indicate potent anti-diarrheal, antihypertensive, hepatoprotective, antioxidant, antimicrobial, hypoglycemic and anti-mutagenic activities. A number of chemicals isolated from plants like quercetin, guaijaverin, flavonoids and galactose-specific lecithins have shown promising activity in many human trials. In recent years, emphasis of research has been on utilizing traditional medicines that have a long and proven history of treating various ailments. Quite a significant amount of work has been done on the pharmacological and biological activity and possible application of chemical compounds from whole part of the plant. Hence extensive investigation on its pharmacodynamics, kinetics, and proper standardization and clinical trials is needed to exploit their therapeutic utility to combat various diseases.

REFERENCES

- Mital Kaneria, Sumitra chanda, Phytochemical and Pharmacognostic Evaluation of leaves of *Psidium Guajava* L. (Myrtaceae), *Pharmacognosy Journal*, 2011, 3(23), 41-45.
- Rathish N, Sumitra C, *In-vitro* antimicrobial activity of *Psidium guajava* l. leaf extracts against clinically important pathogenic microbial strains, *Brazilian Journal of Microbiology*, 2007, 38, 452-458.
- El-Mahmood MA, The use of *Psidium guajava* Linn. in treating wound, skin and soft tissue infections, *Scientific Research and Essay*, 2009, 4(6), 605-611.
- Burkil HM. The useful plants of west Tropical Africa. Royal Botanical Gardens, Kew; 1994. P. 21-150.
- Kamath JV, Nair Rahul, Ashok Kumar CK, Mohana Lakshmi S, *Psidium guajava* L: A review, *International Journal of Green Pharmacy*, 2008, 2(1), 9-12.
- Little EL, Wadsworth FL. Common trees of Puerto Rico and the Virgin Islands. *Agriculture Handbook 249*. U.S. Department of Agriculture, Washington, DC; 1964. P. 548.
- Bala SA. *Psidium guajava*. In: some ethno-medicinal plants of the savannah regions of West Africa: Description and phytochemicals. Triumph publishing company limited, Kano, Nigeria. (Vol. II); 2006. P. 21-56.
- Nadkarni KM, Nadkarni AK. *Indian Materia Medica - with Ayurvedic, Unani- Tibbi, Siddha, Allopathic, Homeopathic, Naturopathic and Home remedies*. Popular Prakashan Private Ltd., Bombay, India; 1991. P. 142-49.
- Morton JF. *Atlas of medicinal plants of Middle America*. Springfield, Charles C; 1981.
- Tona L, Kambu K, Mesia K, Cimanga K, Apers S, Biological screening of traditional preparations from some medicinal plants used as antidiarrhoeal in Kinshasa, Congo. *Phytomedicine*, 1999, 6, 59-66.
- Datta SC. *Systemic Botany*. (4th Edn). Willey Eastern Ltd, New Delhi, India; 1988. P. 406.
- Hernandez DF. *Plants of the Philippines*. 2nd Edn., M&L Licudine Enterprises, Philippines; 1971.
- Heinrich M. *Plants as antidiarrhoeals in medicine and diet*. Proceedings from a Joint Meeting of the Society for Economic Botany and the International Society London, Royal Botanic Gardens, Kew, UK; 1998. P. 17-30.
- Leonti M, Vibrans H, Stiche O, Heinrich M, *Ethnopharmacology of the Popoluca, Mexico: An evaluation*. *J. Pharmacy Pharmacol*, 2001, 53, 1653-1669.
- Batick MJ. *Ethnobotany of Palms in the Neotropics*. In: Prance GT, Kallunki JA, editors. *Advances in Economic Botany: Ethnobotany in the Neotropics*. New York, USA: New York Botanical Garden; 1984. P. 9-23.
- Khan MLH, Ahmad J, A pharmacognostic study of *Psidium guajava* L., *International Journal of Crude Drug Research*, 1985, 23, 95-103.
- Gutierrez RM, Mitchell S, Solis RV, *Psidium guajava*: a review of its traditional uses, phytochemistry and pharmacology, *J Ethnopharmacol*, 2008, 117(1), 1-27.
- Lucky O Okunrobo, Kate E Imafidon, Adeyemi A Alabi, *Phytochemical, Proximate and Metal Content Analysis of the Leaves of Psidium guajava* Linn (Myrtaceae), *International Journal of Health Research*, 2010, 3(4), 217-221.
- Jimenez-Escrig M, Rincon M, Pulido R, Saura-Calixto F, Guava fruit (*Psidium guajava* L.) as a new source of antioxidant dietary fiber, *Journal of Agricultural and Food Chemistry*, 2001, 49(11), 5489-5493.
- Lozoya X, Reyes-Morales H, Chavez-Soto MA, Martinez-Garcia MC, Soto-Gonzalez Y, Doubova SV, Intestinal antispasmodic effect of a phytodrug of *Psidium guajava* folia in the treatment of acute diarrhoeal diseases. *Journal of Ethnopharmacology*, 2002, 83(1-2), 19-24.
- Lapik O, Klejdus B, Kokoska L, Identification of isoflavones in *Acca sellowiana* and two *Psidium* species (Myrtaceae), *Biochem Syst Ecol*, 2005, 33, 983-992.
- Laddha KS. *Practical Pharmacognosy*. 1st ed., New Vrinda Publishing House; 1992. P. 110.
- Kokate CK, Gokhle AP, Khandelwal P. *Practical Pharmacognosy, Techniques and Experiment*, 3rd ed., Nirali Prakashan; 1994. P. 115-121.
- Rahul Mishra, Kalyan K Sethi, Manesh Kumar, S Jha, Harshita Jain, A Pharmacognostical Approach for Study of *Psidium Guajava* Linn, *International Journal of Current Pharmaceutical Research*, 2011, 3(4).
- Evans WC. *Trease and Evans Pharmacognosy*. (13th Edn), Bailere Traiadal, London; 1989. P. 101-104.
- Sofowora A. *Medicinal Plants and Traditional Medicine in Africa*. (2nd Edn). Spectrum Books Limited, Ibadan, Nigeria; 1993. P. 1-153.
- Akinjogunla OJ, Adegoke AA, Udokang IP, Adebayo-Tayo BC, Antimicrobial potential of *Nymphaea lotus* (Nymphaeaceae) against wound pathogens. *Journal of Medicinal Plants Research*, 2009, 3(3), 138-141.
- Akinjogunla OJ, Etok CA, Oshoma CE, Preliminary phytochemistry and *in-vitro* antibacterial efficacy of Hydro-Ethanollic leaf extracts of *Psidium guajava* on common urinary tract Bacterial Pathogens, *Bioresearch Bulletin*, 2011, 5, 329-336.
- Deo A, Satri NV, Purification and characterization of polygalacturonase inhibitory proteins from *Psidium guajava* Linn. (Guava) fruit, *Plant science*, 2003, 164, 147-56.
- Smith RM, Siwatibau S, Sesquiterpene hydrocarbons of Fijian guavas, *Phytochemistry*, 1975, 14, 2013-2015.
- Wilson CW, Shaw PE, Terpene hydrocarbons from *Psidium guajava*, *Phytochemistry*, 1978, 17, 1435-6.
- Begum S, Hassan SI, Ali SN, Siddiqui BS, Chemical constituents from the leaves of *Psidium guajava*, *Nat Prod. Res*, 2004, 18(2), 135-140.
- Cho EJ, Yokozanawa T Yokozawa, Rhyu DY, Kim SC, Shibahara N, Park JC, Study on the inhibitory effects of Korean medicinal plants and their main compounds on the 1,1-diphenyl-2-picrylhydrazyl radical, *Phytomed*, 2003, 10, 544-551.
- Narayana KR, Reddy MS, Chaluvadi MR, Krishna DR, *Bioflavonoids: classification, pharmacology, biochemical effects and therapeutic potential*, *Indian Journal of Pharmacol*, 2001, 33, 2-16.
- Tee ES, Mohd Ismail N, Mohd Nasir A, Khatijah I. *Nutrient Composition of Malaysian Foods*, Institute for Medical Research, Kuala Lumpur; 1997.
- Hobert I, Tietze HW. *Guava as Medicine: A Safe and Cheap Form of Food Therapy*. Pelanduk Publications, Kelana Jaya, Selangor, Malaysia; 1998.
- Oliver-Bever. *Bep, Medicinal Plants in tropical West Africa*. Cambridge University Press, Cambridge; 1986. ISBN No. 0-521-26815.
- Zakaria M, Mohd MA. *Traditional Malay Medicinal Plants*. Fajar Bakti Sdn. Bhd., Kuala Lumpur; 1994. ISBN: 967-65-2476.
- Wyk, Ben-Erik van, Oudtshoorn, Bosch van, Gericke, Nige. *Medicinal Plants of South Africa*, Briza Publications, Pretoria, South Africa. First edition; 1997. ISBN No. 1-875093-09-5.
- Joseph B, Priya RM, *Phytochemical and biopharmaceutical aspects of Psidium guajava* (L.) essential oil: A review, *Res J Med Plant*, 2011, 5, 432-442.
- Seshadri T, Vasishta K, Polyphenols of the leaves of *Psidium guajava*; quercetin, guaijaverin, leucocyanidin, amritoside, *Phytochemistry*, 1965, 4, 989-92.
- El Khadem H, Mohamed YS, *Constituents of the leaves of Psidium guajava* L: Part II: Quercetin, avicularin, and guaijaverin, *J Chem Soc*, 1958, 32, 3320-3.
- Latza S, Ganber D, Berger RG, *Carbohydrate esters of cinnamic acid from fruits of Physalis peruviana, Psidium guajava and Vaccinium vitis-idaea*, *Phytochemistry*, 1996, 43, 481-485.
- Chang WS. *Studies on Active Principles of Hypoglycemic Effect from Psidium Guajava* (L). Master Thesis, The Graduate Institute of Pharmacy, Taipei Medical College; 1982.

45. Peng RY, Hsieh CL, Review on the medicinal uses of *Psidium guajava* L., Recent Progress in Medicinal Plants, 2006, 20, 215–248.
46. Chen KC, Peng CC, Chiu WT, Cheng YT, Huang GT, Hsieh CL, Action mechanism and signal pathways of *Psidium guajava* L. aqueous extract in killing prostate cancer LNCaP cells, Nutr Cancer, 2010, 62(2), 260–270.
47. Buvaneswari S, Raadha CK, Krishnaveni N, Jayashree S, *In vitro* Antimicrobial activity of *Psidium guajava* against clinically important strains, EJLS, 2011, 1(1), 14-22.
48. Berdy J, Aszalos A, Bostian M, Mcnitt KL. CRC Handbook of antibiotic compounds. Boca Raton, CRC Press, v. 8, Part 1; 1981.
49. Tangpu TV, Yadav AK, Anticestodal efficacy of *Psidium guajava* against experimental *Hymenolepis diminuta* infection in rats, Indian J Pharmacol, 2006, 38, 29–32.
50. Nundkumar N, Ojewole JA, Studies on the antiplasmodial properties of some South African medicinal plants used as antimalarial remedies in Zulu folk medicine, Methods Find Exp Clin Pharmacol, 2002, 24, 397-401.
51. Nair R, Chanda S, *In vitro* antimicrobial activity of *Psidium guajava* L. leaf extracts against clinically important pathogenic microbial strains, Braz J Microbiol, 2007, 38, 452-458.
52. Hui-Yin Chen, Gow-Chin Yen, Antioxidant activity and free radical-scavenging capacity of extracts from guava (*Psidium guajava* L.) leaves, Food Chemistry, 2007, 101(2), 686-694.
53. Metwally AM, Omar AA, Harraz FM, El Sohafy SM, Phytochemical investigation and antimicrobial activity of *Psidium guajava* L. leaves, Phcog Mag, 2010, 6, 212-8.
54. Sofowora A. Medicinal plant and traditional medicine in Africa II, John Wiley Chichester; 1986. P. 178.
55. Akinjogunla OJ, Yah SC, Eghafona NO, Ogbemudia FO. Antibacterial activity of leaf extracts of *Nymphaea lotus* (*Nymphaeaceae*) on Methicillin Resistant *Staphylococcus aureus* (MRSA) and Vancomycin Resistant *Staphylococcus aureus* (VRSa) Isolated from clinical samples, Annals of Biological Research, 2010, 1(2), 174-184.
56. Cowan MM, Plant products as Antimicrobial agents, Clinical Microbiological Review, 1999, 12, 564-583.
57. Lutterodt GD, Maleque A. Effects on mice locomotor activity of a narcotic-like principle from *Psidium guajava* leaves, Journal of Ethnopharmacology, 1988, 24(2-3), 219–231.
58. Meckes M, Calzada F, Tortoriello J, Gonzalez JL, Martinez M, Terpenoids isolated from *Psidium guajava* with depressant activity on central nervous system, Phytotherapy Research, 1996, 10(7), 600–603.
59. Goncalves JLS, Lopes RC, Oliveira DB, Costa SS, Miranda MMFS, Romanos MTV, Santos NSO, Wigg MD, *In vitro* antirotavirus activity of some medicinal plants used in Brazil against Diarrhea, J Ethnopharmacol, 2005, 99, 403.
60. Grover IS, Bala S, Study on anti mutagenic effects of guava in *S. typhimurium*, Mutation Research/Genetic Toxicology, 1993, 300(1), 1-3.
61. Qadan F, Thewaini AJ, Ali DA, Afifi R, Elkhawad A, Matalka KZ, The antimicrobial activities of *Psidium guajava* and *Juglans regia* leaf extracts to acne developing organisms, Am Chinese Med, 2005, 33, 197-204.
62. Wei L, Li Z, Chen B, Clinical study on treatment of infantile rotaviral enteritis with *Psidium guajava* L., Zhongguo Zhong Xi Yi Jie He Za Zhi, 2000, 20(12), 893-895.
63. Jaiarj P, Khoohaswan P, Wongkrajang Y, Peungvicha P, Suriyawong P, Saraya MLS, Ruangsomboon O, Anticough and antimicrobial activities of *Psidium guajava* Linn. Leaf extract, Journal of Ethnopharmacology, 1999, 67(2), 203–212.
64. Abreu PRC, Almeida MC, Bernardo RM, Bernardo LC, Brito LC, Garcia EAC, Fonseca AS, Bernardo-Filho M, Guava extract (*Psidium guajava*) alters the labelling of blood constituents with technetium-99m, Journal of Zhejiang University Science, 2006, 7(6), 429–435.
65. Chiwororo WD, Ojewole JA, Biphasic effect of *Psidium guajava* Linn.(Myrtaceae) leaf aqueous extract on rat isolated vascular smooth muscles, J Smooth Muscle Res, 2008, 44(6), 217-229.
66. Cheng JT, Yang RS, Hypoglycaemic effect of guava juice in mice and human subject, Am J Chinese Med, 1983, 11, 74-76.
67. Joseph B, Priya RM, Review on nutritional, medicinal and pharmacological properties of Guava (*Psidium guajava* Linn), Int J Pharma Bio Sci, 2011, 2, 53-69.
68. Okuda T, Yoshida T, Hatano T, Yakazi K, Ashida M, Ellagitannins of the casuarinaceae, stachyuraceae and myrtaceae, Phytochemistry, 1982, 21(12), 2871-2874.
69. Shaheen HM, Effect of *Psidium guajava* leaves on some aspects of central nervous system in mice, Phytother Res, 2000, 14(2), 107-111.
70. Belemtougri RG, Constantin B, Cognard C, Raymond G, Sawadogo L, Effects of *Sclerocarya birrea* (A. rich) hochst (anacardiaceae) leaf extracts on calcium signalling in cultured rat skeletal muscle cells, J Ethnopharmacology, 2001, 76, 247–252.
71. Abdul Jabbar Shah, Sabira Begum, Syed Imran Hassan, Syed Nawazish Ali, Bina Shaheen Siddiqui, Anwarul-Hassan Gilani, Pharmacological basis for the medicinal use of *Psidium guajava* leave in hyperactive gut disorders, Bangladesh Journal of Pharmacology, 2011, 6(2).
72. Rosângela de Oliveira Teixeira, Marjori Leiva Camparoto, Mário Sérgio Mantovani, Veronica Elisa Pimenta Vicentini, Assessment of two medicinal plants, *Psidium guajava* L. and *Achillea millefolium* L., in *in vitro* and *in vivo* assays, Genet Mol Biol, 2003, 26.
73. Manosroi J, Dhumtanom P, Manosroi A, Antiproliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines, Cancer Lett, 2006, 235, 114-120.
74. Matsuo N Hanamura, Koyoko SY Nakamura, Tomita I, Identification of (+) galocatechin as a bio-antimutagenic compound in *Psidium guajava* leaves, Phytochemistry, 1994, 36(4), 1027-1029.
75. Limsong J, Benjavong kulchai E, Kuvataanasuchati J, Inhibitory effects of some herbal extracts on adherence of *S. Mutans*, J Ethnopharmacol, 2004, 92(2-3), 281-289.
76. Rodriguez RC, Cruz PH, Rios HG, Lectins in fruits having gastrointestinal activity their participation in hemagglutinating property of *Escherichia coli* O157, Arch Med Res, 2001, 32(4), 251-257.
77. Zhang WJ, Chen BT, Wang CY, Zhu QH, Mo ZX, Mechanism of quercetin as an anti-diarrheal agent, Di Yi Jim Yi Da Xue Xue Bao, 2003, 23, 1029-1031.
78. Nakamura Y, Ishimitsu S, Tonogai Y, Effects of quercetin and rutin on serum and hepatic lipid concentrations, fecal steroid excretion and serum antioxidant properties, J Health Sci, 2000, 46, 229–40.
79. Singh RB, Rastogi SS, Singh NK, Ghosh S, Niaz MA, Effects of guava intake on serum total and high-density lipoprotein cholesterol levels and on systemic blood pressure, Am J Cardiol, 1992, 70(15), 1287-1291.
80. Singh RB, Rastogi SS, Singh NK, Ghosh S, Gupta S, Niaz MA, Can guava fruit intake decrease blood pressure and blood lipids, J Hum Hypertens, 1993, 7(1), 33-38.