



Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open  Access

Research Article

Physicochemical, Qualitative and Quantitative Determination of Secondary Metabolites and Antioxidant Potential of *Kalanchoe Pinnata* (Lam.) Pers. Leaf Extracts

Santosh Kumar Singh^{1*}, Jay Ram Patel², Arvind Dangi³

¹ Ph.D. Scholar, Pacific college of Pharmacy, Pacific University, Pacific Hills, Airport Road, Pratap Nagar Extension, Debari, Udaipur-313024, Rajasthan, India.

² Principal, Rewa College of Pharmacy, NH-7, Allahabad Road, Ratahra, Rewa- 486001, Madhya Pradesh, India.

³ Truba Institute of Pharmacy, Karond, Bhopal-462038, Madhya Pradesh, India.

ABSTRACT

Kalanchoe pinnata (Lam.) Pers. has many pharmacological properties such as anticancer, antidiabetic, insecticidal, antimicrobial, anti-urolithiatic, etc. The plant leaves are used as ethno medicine traditionally. Many diseases are associated with oxidative stress caused by free radicals. The aim of the present study was to evaluate physicochemical, qualitative and quantitative phytochemical analysis and in vitro antioxidant activities of leaf of *Kalanchoe pinnata* collected from Bhopal region of Madhya Pradesh. The physicochemical evaluations carried out in terms of loss on drying, ash value, extractive values and acid insoluble ash value ect. Qualitative analysis of various phytochemical constituents and quantitative analysis of total phenolics and flavonoids were determined by the well-known test protocol available in the literature. Quantitative analysis of phenolic and flavonoids was carried out by Folin Ciocalteu reagent method and Aluminium chloride method respectively. The In vitro antioxidant activity of ethanolic extract of the leaf was assessed against DPPH assay method using standard protocols. Phytochemical analysis revealed the presence of phenols, flavonoids, tannins, saponins, alkaloids. The total phenolics content of leaves ethanolic extract was (4.63mg/100mg), followed by flavonoids (2.03mg/100mg). The activities of ethanolic leaves extract against DPPH assay method were concentration dependent with IC 50 values of ascorbic acid and extracts 24.44 and 49.81µg/ml respectively. These studies provided information for standardization and correct identification of this plant material. The diverse array of phytochemicals present in the plant thus suggests its therapeutic potentials which may be explored in drug manufacturing industry as well as in traditional medicine.

Keywords: *Kalanchoe pinnata*, Physicochemical, Qualitative, Quantitative phytochemical, Antioxidant

Article Info: Received 03 Dec 2018; Review Completed 09 Jan 2019; Accepted 11 Jan 2019; Available online 15 Jan 2019



Cite this article as:

Singh SK, Patel JR, Dangi A, Physicochemical, Qualitative and Quantitative Determination of Secondary Metabolites and Antioxidant Potential of *Kalanchoe Pinnata* (Lam.) Pers. Leaf Extracts Journal of Drug Delivery and Therapeutics. 2019; 9(1):220-224 DOI: <http://dx.doi.org/10.22270/jddt.v9i1.2225>

*Address for Correspondence:

Santosh Kumar Singh, Pacific college of Pharmacy, Pacific University, P.B.-12 Pacific Hills, Airport Road, Pratap Nagar Extension, Debari, Udaipur-313024, Rajasthan, India.

INTRODUCTION

Drugs from the plants are easily available, less expensive, safe and efficient and rarely have side effects¹. According to World Health Organization (WHO), medicinal plants would be the best source to obtain variety of drugs. Large sections of the population in developing countries still rely on traditional practitioners and herbal medicines for their primary care². Medicinal plants are plants in which one or more of their organs contain substances that can be used for therapeutic purposes or which are precursors for the synthesis of useful drugs. WHO consultative group that formulated this definition stated also that, such a description makes it possible to distinguish between medicinal plants whose therapeutic properties and constituents have been established scientifically and plants that are regarded as medicinal but which have not yet been subjected to a

thorough scientific study³. Such plants should be investigated to better understand their properties, safety and efficacy. The medicinal properties of plants are due to some chemical constituents that produce certain pharmacological action on the humans. The qualitative analysis of phytochemicals of a medicinal plant is reported as vital step in any kind of medicinal plant research. Screening of plant constituents accurately can be done by employing chromatographic techniques⁴. Quantification usually employs the use of gravimetric and spectroscopic methods with several advanced approaches now available⁵. Crude extract acquired from the *Kalanchoe pinnata* was evaluated for its radical scavenging properties and assessed that it could be a rich source of natural oxidants with potential applications⁶. *Bryophyllumpinnatum* (Lam.) Synonym *Kalanchoe pinnata* (Lam.) is a perennial herb growing widely and used in folkloric medicine in tropical Africa, India, China,

Australia and tropical America⁷⁻¹⁰. Classified as a weed, the plant flourishes throughout the Southern part of Nigeria. A number of active compounds, including flavonoids, glycosides, steroids, bufadienolides and organic acids, have been identified in *Kalanchoe pinnata*⁹. Its bufadienolides are structurally similar to cardiac glycosides and have demonstrated in clinical research to possess antimicrobial, antifungal, anticancer, anti tumour, insecticidal actions. It also possess other activities like anti-ulcer, anti-inflammatory and analgesic, antihypertensive, hepatoprotective, Nephroprotective, diuretic, antidiabetic, anticonvulsion, antioxidant, uterine relaxant, muscle relaxant and neurosedative activity and tocolysis activity^{11,12}. *Kalanchoe pinnata* is used in ethno medicine for treatment of earache, burns, abscesses, ulcer, insect bites, diarrhea and Lithiasis⁷. In traditional medicine, its stem, leaves and the whole plant have been extensively worked on while there is limited work done on the leaf of *Kalanchoe pinnata*. The aim of this work was to determine the quality (types), quantity (amount) of bioactive compounds and in vitro antioxidant activity of leaf of *Kalanchoe pinnata* in Bhopal region of Madhya Pradesh.

MATERIALS AND METHODS

Plant material

The leaves of plant *Kalanchoe pinnata* was collected from rural area of Bhopal (M.P), India. **Chemical reagents**

All the chemicals used in this study were obtained from HiMedia Laboratories Pvt. Ltd. (Mumbai, India), Sigma Aldrich Chemical Co. (Milwaukee, WI, USA), SD Fine-Chem Chem. Ltd. (Mumbai, India) and SRL Pvt. Ltd. (Mumbai, India). All the chemicals used in this study were of analytical grade.

Organoleptic characters

Plant material (leaves) selected for the study were washed thoroughly under running tap water and then were rinsed in distilled water; they were allowed to dry for some time at room temperature. Then the plant material was shade dried without any contamination for about 3 to 4 weeks. Dried plant material was grinded using electronic grinder to obtain a powdered form and then subsequently used for organoleptic characterization. A small amount of powdered plant part was spread on a white tile and physically examined for general appearance i.e. color, taste, texture etc. Dried plant material was packed in air tight container and stored for Phytochemical and biological studies¹³.

Physicochemical study¹⁴⁻¹⁶

Determination of loss on drying

Two grams of crude powder was taken in an evaporating dish and then dried in an oven at 105°C till constant weight was obtained. The weight after drying was noted and loss on drying was calculated. The percentage was calculated on the basis of sample taken initially.

Total ash

3 g of powdered drug was accurately weighed and taken in a tarred silica crucible which was previously ignited and weighed. The powdered drug was spread as a fine even layer on the bottom of the crucible. The crucible was incinerated gradually by increasing the temperature to make it dull red hot until free from carbon. The crucible was cooled and weighed. The ash was weighed and the total ash content was calculated with reference to the air dried drug.

Acid insoluble ash

The ash obtained as described in total ash was boiled with 25ml of dilute hydrochloric acid for 5 minutes. The insoluble ash was collected on an ashless filter paper and washed with hot water. This insoluble ash was transferred into a silica crucible and it was ignited, cooled and weighed. The process was repeated to get constant weight. The percentage of acid insoluble ash was calculated with reference to the quantity of air dried crude drug.

Water soluble ash

Total ash obtained was boiled for 5 minutes with 25ml of water. The insoluble matter was collected in ashless filter paper and washed with hot water. The insoluble ash was transferred into silica crucible and was ignited, cooled and weighed. The process was repeated to get constant weight. The weight of insoluble matter was subtracted from the weight of the total ash. The difference of the weight was considered as the water soluble ash. The percentage of water soluble ash was calculated with reference to the air dried drug.

Solvent ether soluble extractive values

Accurately weighed 5gm of powdered air dried leaves was taken with 100 ml of solvent ether in a stopper flask and kept for 24 hours. The flask was shaken frequently (Maceration) then the solvent ether extract was filtered rapidly through filter paper to prevent excessive loss of solvent ether. 25 ml of solvent ether extract was evaporated to dryness on a water bath and complete the drying in an oven at 100°C. Then the residue was cooled weighed and kept in desiccators. Then the percentage w/w of solvent ether soluble extractive with reference to the air-dried drug was calculated.

Alcohol soluble extractive values

Accurately weighed 5 gm of powdered air dried leaves was taken with 100ml of alcohol (90 % v/v) in a stopper flask and kept for 24 hours. The flask was shaken frequently (Maceration). Then the alcohol extract was filtered rapidly through filter paper to prevent excessive loss of alcohol. 25ml of alcoholic extract was evaporated to dryness on a water bath and complete the drying in an oven 100°C. Then the residue was cooled, weighed and kept in desiccators. Then the percentage w/w of alcohol soluble extractive with reference to the air-dried drug was calculated.

Water soluble extractive values

Accurately weighed 5gm of powdered air-dried leaves was taken with 100 ml of water in a stopper flask and kept for 24 hours. The flask was shaken frequently (Maceration). Then the aqueous extract was filtered rapidly through filter paper. 25 ml of aqueous extract was evaporated to dryness on a water bath and complete the drying in an oven at 100°C. Then the residue was cooled, weighed and kept in desiccators. Then percentage w/w of soluble extractive with reference to the air dried drug was calculated.

Extraction procedure

Defatting of plant material

Powdered plant material (leaves) of *Kalanchoe pinnata* was shade dried at room temperature. The shade dried flower was coarsely powdered and subjected to extraction with petroleum ether (60-80°C) in a soxhlet apparatus. The extraction was continued till the defatting of the material had taken place.

Extraction

50 g. of *Kalanchoe pinnata* dried leaves were successive extracted with various solvent (chloroform, ethyl acetate,

ethanol and aqueous) and using different drug: solvent ratios using hot continuous percolation for different time (soxhlet apparatus). The extracts were evaporated above their boiling points and stored in an air tight container free from any contamination until it was used. Finally the percentage yields were calculated of the dried extracts¹⁷.

Qualitative phytochemical analysis of plant extract

The *Kalanchoe pinnata* leaves extract obtained was subjected to the preliminary phytochemical analysis following standard methods by Khandelwal and Kokate^{18,19}. The extract was screened to identify the presence or absence of various active principles like phenolic compounds, carbohydrates, flavonoids, glycosides, saponins, alkaloids, fats or fixed oils, protein and amino acid and tannins.

Quantification of secondary metabolites

Quantitative analysis is an important tool for the determination of quantity of phytoconstituents present in plant extracts. For this TPC and TFC are determined. Extracts obtained from leaves of *Kalanchoe pinnata* plant material of subjected to estimate the presence of TPC and TFC by standard procedure.

Total phenol determination

The total phenolic content was determined using the method of Olufunmiso et al²⁰. A volume of 2 ml of *Kalanchoe pinnata* leaves extracts or standard was mixed with 5 ml of Folin Ciocalteu reagent (previously diluted with distilled water 1:10 v/v) and 4 ml (75g/l) of sodium carbonate. The mixture was allowed to stand for 15 min under room temperature. The blue colour developed was read at 765 nm using UV/visible spectrophotometer. The total phenolic content was calculated from the standard graph of gallic acid and the results were expressed as gallic acid equivalent (mg/g).

Total flavonoids determination

The total flavonoid content was determined using the method of Olufunmiso et al²⁰. 1 ml of 2% AlCl₃ methanolic solution was added to 1 ml of extract or standard and allowed to stand for 60 min at room temperature; the absorbance of the reaction mixture was measured at 420 nm using UV/visible spectrophotometer. The content of flavonoids was calculated using standard graph of quercetin and the results were expressed as quercetin equivalent (mg/g).

DPPH free radical scavenging assay

DPPH scavenging activity was measured by modified method²⁰. DPPH scavenging activity was measured by the spectrophotometer. Stock solution (6 mg in 100ml methanol) was prepared such that 1.5 ml of it in 1.5 ml of methanol gave an initial absorbance. Decrease in the absorbance in presence of sample extract at different concentration (10-100 µg/ml) was noted after 15 minutes. 1.5 ml of DPPH solution was taken and volume made till 3 ml with methanol, absorbance was taken immediately at 517 nm for control reading. 1.5 ml of DPPH and 1.5 ml of the test sample of different concentration were put in a series of

volumetric flasks and final volume was adjusted to 3 ml with methanol. Three test samples were taken and each processed similarly. Finally the mean was taken. Absorbance at zero time was taken for each concentration. Final decrease in absorbance was noted of DPPH with the sample at different concentration after 15 minutes at 517 nm. The percentage inhibition of free radical DPPH was calculated from the following equation: % inhibition = [(absorbance of control - absorbance of sample)/absorbance of control] × 100%. Though the activity is expressed as 50% inhibitory concentration (IC₅₀), IC₅₀ was calculated based on the percentage of DPPH radicals scavenged. The lower the IC₅₀ value, the higher is the antioxidant activity.

RESULTS AND DISCUSSIONS

Organoleptic evaluation represents those properties of materials that can be done by using the sense organs. It thereby defines some specific characteristics of the material which can be considered as a first step towards establishing the identity and degree of purity of the material. The organoleptic parameters were evaluated represent in Table 1. The crude extracts so obtained after each of the successive soxhlet extraction process were concentrated on water bath by evaporation the solvents completely to obtain the actual yield of extraction. The percentage yield of extraction is very important in phytochemical extraction in order to evaluate the standard extraction efficiency for a particular plant, different parts of same plant or different solvents used. The yield of extracts obtained from the leaves of the plants using petroleum ether, chloroform, ethyl acetate, ethanol and water as solvents are depicted in the Table 2. The physical constituent's estimation of the drugs is an essential parameter to determine adulteration or inappropriate handling of drugs. The physicochemical characters of powder drug of leaves of *Kalanchoe pinnata* such as total alcohol soluble extractive, water soluble extractive, ash value, acid insoluble ash, and water soluble ash, loss after drying and foreign substances are given in Table 3. The leaves showed less moisture content; it was 2.35%. Moisture content of drugs could be at minimal level to discourage the growth of bacteria, yeast or fungi during storage. These can serve as a valuable basis of information and provide suitable standards to establish the quality of this plant material as future prospects. An ash values are used to decide quality and purity of crude drug, it indicates presence of various impurities like, silicate, oxalate and carbonate. The water soluble ash is used to determine the quantity of inorganic compounds present in drugs. The acid insoluble ash helps to estimate the amount of silica present in the material. The total water soluble portion of the ash is considered as water soluble ash. Less amount of these three parameters indicate that the inorganic matter and silica were less in *Kalanchoe pinnata* leaf. The results of qualitative phytochemical analysis of the crude powder of leaves of *Kalanchoe pinnata* are shown in Table 4. Ethanolic and aqueous extracts of leaves sample of *Kalanchoe pinnata* showed the presence of flavonoids, phenols, tannins, carbohydrate, glycosides and proteins but in chloroform and ethyl acetate extracts all phytoconstituents was absents.

Table 1 Organoleptic characters of *Kalanchoe pinnata*

Plant parts	Color	Odour	Taste	Texture
Leaves	Green color	Characteristic	Slightly Bitter	Decussate and foliolate

Table 2 Results of percentage yield of leaf extracts

Plant Name	Percentage yield (%)				
	Pet. Ether	Chloroform	Ethyl acetate	Ethanol	Water
<i>Kalanchoe pinnata</i>	1.8	1.23	2.03	4.43	6.9

Table 3 Results of physicochemical parameters of leaf

S. No.	Parameters (% w/w)	<i>Kalanchoe pinnata</i>
1.	Total ash	10.2%
2.	Acid Insoluble Ash	6.8%
3.	Water soluble Ash	2.3%
4.	Alcohol soluble extractive value	15.1%
5.	Water soluble extractive value	9.8%
6.	Ether soluble extractive value	11.1%
7.	Moisture content	2.35%

Table 4 Phytochemical Evaluation of *Kalanchoe pinnata* leaves

Chemical Tests	Pet. ether	Chloroform	Ethyl acetate	Ethanol	Aqueous
Alkaloids					
Mayer's reagent	-	-	-	-	+
Hager's reagent	-	-	-	-	+
Wagner's reagent	-	-	-	-	+
Dragendorff's reagent	-	-	-	-	+
Glycosides (+Ve)					
Baljet test	-	-	-	-	-
Legal's test	-	-	-	-	-
Keller-Kiliani	-	-	-	-	-
Phenols/Tannins					
Ferric chloride	-	-	+	+	+
Gelatin Solution	-	-	+	+	+
Lead acetate test	-	-	+	+	+
Flavonoids					
FeCl ₃ test	-	-	+	+	+
Alkaline reagent test	-	-	+	+	+
Shinoda test	-	-	+	+	+
Saponins					
Foam test	-	-	-	-	+
Hemolytic test	-	-	-	-	+
Lead acetate	-	-	-	-	+
Fixed oil/Fats					
Spot	+	-	-	-	-
Saponification	+	-	-	-	-
Gums & Mucilage					
Water	-	-	-	-	-
Carbohydrates					
Molish test	-	-	-	-	-
Fehling's solution	-	-	-	-	-
Benedict's test	-	-	-	-	-
Amino acids					
Ninhydrin Test	-	-	-	-	+
Millons Test	-	-	-	-	+
Xantoprotein Test	-	-	-	-	+
Terpenoids					
L B Test	-	-	-	-	-
Salkowski test	-	-	-	-	-
Steroids					
Lieberman Test	-	-	-	-	-
Protein					
Biuret test	-	-	-	+	-

(+) Indicates 'Presence'; (-) Indicates 'Absence'

The determination of the total phenolic content, expressed as mg gallic acid equivalents and per 100 mg dry weight of sample. TPC of ethanolic, aqueous and ethyl acetate extract of *Kalanchoe pinnata* leaves showed the content values of 4.6363, 2.8181 and 2.0363 respectively. But chloroform extracts of *Kalanchoe pinnata* leaves have no phenolic content. The total flavonoids content of the extracts was expressed as percentage of quercetin equivalent per 100 mg dry weight of sample. The total flavonoids estimation of ethanolic, aqueous and ethyl acetate extracts of leaves of *Kalanchoe pinnata* showed the content values of 2.0300,

1.2475 and 1.0075 respectively. The above results showed that aqueous extract contain less phenolic and flavonoids content than the alcoholic extract. It may be due to the solubility of principle contents presence be higher in case of alcoholic solvent, thus it has been accepted that it is a universal solvent for the extraction of plant constituents. Results are provided in (Table 5 and Fig. 1, 2).

Table 5: Estimation of total phenolics and total flavonoids content in *Kalanchoe pinnata* leaves

S. No	<i>Kalanchoe pinnata</i> Extracts	Total phenolic content (mg/100mg of dried extract)	Total flavonoids content (mg/ 100 mg of dried extract)
1.	Chloroform	-	-
2.	Ethyl acetate	2.0363	1.0075
3.	Ethanol	4.6363	2.0300
4.	Aqueous	2.8181	1.2475




Figure 1: Graph of estimation of total phenolic content




Figure 2 Graph of estimation of total flavonoids content

Antioxidant activity of the samples was calculated through DPPH assay. % inhibition was calculated as an indicative of antioxidant potency. The higher the % inhibition the better the activity. Ascorbic acid was taken as standard and the values were comparable with concentration ranging from 20 µg/ml to 100 µg/ml. A dose dependent activity with respect to concentration was observed Table 6.

Table 6 DPPH assay of ascorbic acid and ethanolic extract

S. No.	Conc. (µg/ml)	% Inhibition	
		Ascorbic acid	Ethanolic extract
1	20	47.66	19.80
2	40	58.93	30.43
3	60	68.11	41.22
4	80	81.48	58.77
5	100	90.98	79.87
IC 50		24.44	49.81

CONCLUSION

It can be concluded that from present investigation the Physicochemical and preliminary phytochemical investigation study of *Kalanchoe pinnata* leaves yielded a set of standards that can serve as an essential basis of evidence to determine the identity and to determine the quality and purity of the plant material as per its future perspectives. The phytochemical investigation gave valuable information about the different phytoconstituents present in the plant, which helps the future investigators concerning the selection of the particular extract for further investigation of isolating the active principle and also gave idea about different phytochemical have been found to possess a wide range of activities. The total phenolic and flavonoid content in ethanolic leaves extract was found to be higher than all the

extracts which is further proved by in vitro antioxidant studies. Potential antioxidant activity has good correlations with the therapeutic use in the treatment of cardiovascular disorders. Further research to isolate individual compounds, their in-vivo antioxidant activities with different mechanism is needed.

REFERENCES

1. Yadav RNS, Agarwala M. Phytochemical analysis of some medicinal plants. Journal of Phytology 2011; 3(12):10-14.
2. WHO. Traditional medicine-growing needs and potentials. WHO Policy Perspectives Med, 2002.
3. WHO. Traditional Medicine and Modern Health Care Progress Report by the Direct General World Health Organization, Document No A-144110, 1991.
4. Sneader W, The discovery of aspirin: A reappraisal, BMJ (Clinical research ed, 2000; 321(7276):1591-1594.
5. Aguru CU, Ameh SJ, Olasan O. Comparative phytochemical studies on the presence and quantification of various bioactive compounds in the three major organs of okoko plant (*Cissampelos guineensis*) in Benue State, North Central Nigeria, West Africa, European Journal of Advanced Research in Biological and Life Sciences, 2014; 2(2):16-22.
6. Sindhu S, Manorama S. Exploration of Antioxidant Properties in Various Extracts of *Bryophyllum pinnatum* (Lam.). Research in Pharmacy 3(4): 01-08, 2013.
7. Nwali BU, Okaka ANC, Ibiam UA, Aja PM. Phytochemical Composition of *Bryophyllum pinnatum* leaves. International Journal of Advanced Biological Research 2012; 2: 614-616.
8. Umebese CE, Falana FD. Growth, phytochemicals and antifungal activity of *Bryophyllum pinnatum* L. subjected to water deficit stress: African Journal of Biotechnology 2013; 12: 6599-6604.
9. Dassharma K, Bhatkar A, Pandey P, Shaikh N. *Bryophyllum pinnatum* (Lam.): A potential source of antioxidants. Bionano Frontier 2013; 6:140-144.
10. Kamboj A, Saluja AK. *Bryophyllum pinnatum* (Lam.) Kurz.: Phytochemical and pharmacological profile. A review Phcog Rev 2009; 3: 364-374.
11. Mensor LL, Menezes FS, Leitão GG, Reis AS, dos Santos TC, et al. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. Phytother Res 2001; 15: 127-130.
12. Ruqaiyah Khan, Afzal M, Kazmi I, Chauhan M, Bisht T. Chemical composition of *bryophyllum*. International Journal of Research in Biological Science 2012; 2:143-149.
13. Quality standards of Indian medicinal plants. Medicinal Plants Unit, ICMR, New Delhi, 2013, 11.
14. WHO Quality control methods for medicinal plants. Geneva, 2002, 28-31.
15. Government of India-Ministry of Health and Family Welfare. Indian Pharmacopoeia, Edn. 1, India Controller of Publications, New Delhi, 1996, II
16. Harborne JB. Phytochemical Methods: A Guide to Modern techniques of plant analysis, Edn 2, Chapman and Hall, London, 1984, 123-75.
17. Mukherjee PK. Quality control of herbal drugs. 2nd Ed. Business Horizons; 2007.
18. Khandelwal KR. Practical pharmacognosy technique and experiments. 23rd Ed. NiraliPrakashan; 2005.
19. Kokate CK. Practical pharmacognosy. 4th Ed. Vallabh Prakashan; 1994.
20. Olufunmiso OO, Afolayan AJ. Phenolic content and antioxidant property of the bark extract of *Ziziphus mucronata* Willd. Subsp. *mucronata* Willd, BMC Complement Alternative Medicine, 2011; 11:130.