

Available online on 15.03.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Evaluation of Anticonvulsant and Antioxidant Activity of *Senna occidentalis* Seeds Extracts

Vijay Vikram Singh^{1*}, Jainendra Jain¹, Arun Kumar Mishra²¹ Department of Pharmacy, Ram-Eesh Institute of Vocational and Technical Education, Greater Noida, India² Phytochemistry Laboratory, Faculty of Pharmacy, IFTM University, Moradabad, India

ABSTRACT

Aim: The aim of present work was to determine the anticonvulsant and antioxidant activity of *Senna occidentalis* L. ethanolic seed extract by different models. **Methods:** For evaluation of anticonvulsant activity, Pentylentetrazole (PTZ) seizure model and Maximal electroshock (MES) seizure model were used. For antioxidant activity, (1, 1-diphenyl - 2-picryl hydrazine (DPPH) and hydrogen peroxide (H₂O₂) method were used. **Results:** The finding suggested that the ethanolic extract (EAE) of *Senna occidentalis* in the dose 400 mg/kg body weight posses potent anticonvulsant activity. The EAE showed anticonvulsant action in dose dependent fashion. It was observed that upon increasing the concentration of extract, it showed reduced absorbance and increased free radical inhibition, and when comparison was made with Ascorbic acid, it showed marked antioxidant property in DPPH as well as H₂O₂ method. The IC₅₀ of Ascorbic acid and EAE by DPPH method were found to be 14.56 and 14.8 respectively whereas the IC₅₀ of Ascorbic acid and EAE by H₂O₂ method were found that 14.3 and 14.8 respectively. **Conclusion:** The results of the present study concluded that the EAE of *Senna occidentalis* L. possesses significant antioxidant and anticonvulsant activity. The activity was in dose dependent fashion. This study will assist in future research associated with formulation development of seeds of *Senna occidentalis* L.

Keyword: *Senna occidentalis* L., Anticonvulsant, Antioxidant, DPPH model

Article Info: Received 28 Jan 2019; Review Completed 27 Feb 2019; Accepted 06 March 2019; Available online 15 March 2019



Cite this article as:

Singh VV, Jain J, Mishra AK, Evaluation of Anticonvulsant and Antioxidant Activity of *Senna occidentalis* Seeds Extracts, Journal of Drug Delivery and Therapeutics. 2019; 9(2):183-187 <http://dx.doi.org/10.22270/jddt.v9i2.2400>

*Address for Correspondence:

Vijay Vikram Singh, Department of Pharmacy, Ram-Eesh Institute of Vocational and Technical Education, Gr Noida, India

INTRODUCTION

Senna occidentalis, growing mainly in lower region, is an erect tropical annual herb. The seed are dark brown and curved with slightly upward, the seeds are brown and flattened on both ends. The seeds, in the long pods, can be roasted and made into a coffee like drink¹. *Senna occidentalis* is known by various names, e.g. Coffee Senna, Fedegoso and Negro coffee. It is common weed scattered from Himalayas to the Western Bengal, South India, Burma and Ceylon. The main phytoconstituents present in *Senna occidentalis* L. includes aloe emodin, anthraquinones, anthrones, apigenin, aurantiobtusin, campesterol, cassiollin, chrysophanol, chrysoeriol, cmodin, physicon quarcetin, rhamnosides, rhein, sitosterols, and xanthorine etc². The plant is bitter in taste, thermogenic, purgative, expectorant, antipyretic and anticonvulsants as it is used by tribals to treat such problems.

Senna occidentalis is reported to have number of medicinal properties in *Ayurveda*. *Senna occidentalis* is used in traditional system of medicine as antipyretic and to treat convulsions in Andhra Pradesh, India with very popularity

but still its pharmacological potential for antioxidant activity and other activities including anticonvulsant has not been fully explored. In present work, with scientific model are used to prove the claim of this drug. The traditional treatise of Ayurveda states the claim of this important drug as antiepileptic (Aakeshpa treatment) and health supplement. Since thousands of year ago, it is used to treat pyrexia and inflammation by tribals.

The literature review revealed that the potential of *Senna occidentalis* as anticonvulsant and antioxidant drug still to be experimentally proven. The present investigation includes pharmacological evaluation of *Senna occidentalis* ethanolic seed extract.

MATERIALS AND METHODS

Collection and identification of plant material

The seeds of *Senna occidentalis* belonging to the family Fabaceae were cultivated and collected from the Herbal garden area of Ram-Eesh Institute of Vocational and Technical Education, Greater Noida, District Gautam Budhdha Nagar, U.P., India.

Identification

The plant and the seeds of *Senna occidentalis* were identified and authenticated by Dr. Sunita Garg, Chief Scientist, National Institute of Science Communication and Information Resources (NISCAIR), New Delhi.

Preparation of the extract

The seeds of *Senna occidentalis* were shade dried and reduced to the coarse powder. The coarse powder (1 Kg) was evenly packed in the Soxhlet apparatus, was subjected to defatting³. The powdered seeds were defatted by petroleum ether (60-80°C) until the color has been changed from dark yellow to colorless. The marc was subjected to the extraction in the presence of ethanol as extraction solvent. The ethanolic extract (EAE) was filtered and filtrate was concentrated by rotary evaporator. The EAE in different concentration was subjected to evaluation of anticonvulsant activity by Pentylentetrazole (PTZ) seizure model and Maximal electroshock (MES) seizure model. For antioxidant activity evaluation of EAE, (1, 1-diphenyl - 2-picryl hydrazine (DPPH) and hydrogen peroxide (H₂O₂) method were used

Experimental animals

For anticonvulsant activity using PTZ model, Swiss albino mice (approximately weighing 30-40 g) and for MES model, Swiss Albino rats of either sex (approximately weighing 200-250 g) were used. The animals for the activity were procured from Institutional Animal House, Department of Pharmacy, Ram-Eesh Institute of Vocational and Technical Education, Greater Noida. All the laboratory animals were kept in temperature controlled room conditions, with 12 h alternating light and dark cycle. The animals were given adequate nutrition and water *ad libitum*. The protocols were followed as per "Guidelines for the Care and Use of Laboratory Animals" and approved by the Institutional Animal Ethics Committee (IEAC) (Ref. Number, RGI/RIT/01/2016; dated 16.01.2016)

Experimental design

Anticonvulsant Activity

Pentylentetrazole (PTZ) induced seizure test^{4,5}

The swiss albino mice were randomly divided into 05 groups having six (n=6) animals in each groups. The animals were divided into all five groups as follows-

Group I : (Negative Control): Treated with normal saline (10 ml/kg) b.w

Group II : (Positive Control): Treated with Valproic acid (300 mg/kg) b.w

Group III : (Test Group): Treated with EAE (100 mg/kg) b.w

Group IV : (Test Group): Treated with EAE (200 mg/kg) b.w

Group V : (Test Group): Treated with EAE (400 mg/kg) b.w

The mice were given the treatment of extract and normal saline to control groups (i.p.), 30 m prior to the administration of 25 mg/kg b.w (PTZ). The animals were placed individually in plastic boxes and observed immediately after PTZ injection for a period of 30 m and after 24 h. The onset time of hind limb tonic extensions (HLTEs) and the ratio of convulsion survivors to total animals tested (mortality protection) were recorded (Table 1; Figure 1, 2)

Maximal Electroshock Induced Seizures (MES) Model^{6,7}

In order to procure tonic convulsion in Albino rats, crocodile ear clip were used and through this, 150 mA current maximum for 0.2 S was discharged through electroconvulsimeter (Inco, Ambala). The electric shock was given just after administration of either vehicle (normal saline 10 ml/kg b.w) or test drug (EAE in dose of 100, 200 and 400 mg/kg b.w) and after 90 m of standard drug i.e. Valproic acid 300 mg/kg p.o.)

The number of animals protected from tonic hind limb extension (HLTE) indicates the abolition of tonic hind limb extension within 10 S after delivery of the electroshock and the duration of observed, HLTE was recorded for each group. For measurement of this, rats were placed in clear rectangular plastic cages with an open top, permitting full view of the animal's motor responses to seizure. The parameters selected for present study were tonic flexion, extension, clonus, stupor and mortality. The findings are presented in table 2 and Figure 3,4. The albino rats were randomly divided into 5 groups of n=6 as follows-

Group I: (Negative Control): Treated with normal saline (10 ml/kg) b.w

Group II: (Positive Control): Treated with Valproic acid (300 mg/kg) b.w

Group III: (Test Group): Treated with EAE (100 mg/kg) b.w

Group IV: (Test Group): Treated with EAE (200 mg/kg) b.w

Group V: (Test Group): Treated with EAE (400 mg/kg) b.w

Antioxidant Activity

Free radical scavenging activity of Senna occidentalis L by DPPH Method^{8,9}

Different conc. (10 µl - 50 µl) of EAE sample and standard sample were prepared. To this, 3 ml of a 0.004% (w/v) solution of DPPH in methanol was added in each test tube. The reaction mixtures were shaken and then incubated at room temperature for period of 30 m. A blank was prepared in similar way, without DPPH and absorbance was measured at 517 nm using double beam UV-Visible spectrophotometer (Shimadzu 1800). Free radical scavenging activity was expressed as the percentage inhibition calculated using formula -

$$\text{Percentage Inhibition DPPH scavenging effect (\%)} =$$

$$\frac{\text{Absorbance of control sample (A}_0\text{)} - \text{Absorbance of test sample (A}_1\text{)}}{\text{Absorbance of control sample (A}_0\text{)}} \times 100$$

A₀ was absorption of control reaction and A₁ was absorption in presence of test or standard sample. Ascorbic acid was used as a positive control. IC₅₀ value was calculated from % inhibition. The IC₅₀ value of the sample, which is the concentration of sample required to inhibit 50% of the DPPH free radical, was calculated using inhibition curve. Lower absorbance of reaction mixture indicates higher free radical activity (Table 3,4).

Free radical scavenging activity of Senna occidentalis L by Hydrogen peroxide method^{10,11}

1 ml of 10% (v/v) sample / standard solution Ascorbic Acid (10-50 µg/ml) was added to the 0.6 ml of the hydrogen peroxide solution which is prepared in phosphate buffer (pH - 7.4). The sample solutions were incubated for 10 m at 37°C. Hydrogen peroxide in phosphate buffer solution it was used as control. Absorbance was measured at 230 nm (Ebrahizadeh et al., 2010). The formula employed for % inhibition was as follows-

Percentage Inhibition (%) =

$$\frac{\text{Absorbance of control sample (A}_0\text{)} - \text{Absorbance of test sample (A}_1\text{)}}{\text{Absorbance of control sample (A}_0\text{)}} \times 100$$

A₀ was absorption of control reaction and A₁ was absorption in presence of test or standard sample (Table 5,6)

RESULTS AND DISCUSSION

Anticonvulsant Activity

Pentylentetrazole -induced seizure test

Table 1: Effect of *Senna occidentalis* on Anticonvulsant effect (PTZ model)

Treatment	Dose (mg/kg)	Onset of Tonic Convulsions	Onset of Clonic Convulsions	Survived/Used
Group I	Control	75.24±1.62	150.27±2.45	5/6
Group II	Valproic Acid 300	935.0±2.31***	1556±2.23***	6/6
Group III	EAE 100	507.21±2.14**	1254.0±3.64*	6/6
Group IV	EAE 200	753.32±2.54***	1398.21±2.31**	6/6
Group V	EAE 400	912.12±1.4***	1498.5±2.15***	6/6

Values are expressed in mean±SEM are represent various phases of convulsion in seconds. Significant at P<0.001*** and P<0.01**, compared with the control group.




Figure 1: Effect of *Senna occidentalis* on onset of tonic convulsions




Figure 2: Effect of *Senna occidentalis* on onset of clonic convulsions

Maximal Electro Shock Induced seizures (MES)

Table 2: Effect of *Senna occidentalis* on Maximum electroshock induced seizures

Treatment	Dose (mg/kg)	Flexon	Extensor	Clonus	Stupor	survived/used
Group I	Control	15.98±0.16	13.72±0.94	14.67±0.91	9.27±0.94	0/6
Group II	Val Acid 300	3.18±0.51***	1.12±0.01***	3.21±0.07***	1.19±0.01***	6/6
Group III	EAE 100	8.14±0.09	4.08±0.11**	10.14±0.51**	6.27±0.05	6/6
Group IV	EAE 200	6.26±0.03**	2.92±0.03***	6.24±0.65***	3.42±0.07***	6/6
Group V	EAE 400	4.18±0.06***	2.02±0.06***	3.92±0.98***	1.84±0.04***	6/6

Values are expressed in mean±SEM are represent various phases of convulsion in seconds. Significant at P<0.001*** and P<0.01**, compared with the control group.




Figure 3: Effect of *Senna occidentalis* on MES induced seizures

The findings suggested that the EAE was effective against seizures. Upon comparison of all dose level, it was observed that EAE (400 mg/kg) b.w was most effective among all the dose level. The activity was observed in dose dependent manner. The results of this study demonstrated that the ethanolic extract have anticonvulsant activity. Data showed that the ethanolic extract displayed anticonvulsant effect in the PTZ induced seizure model. In the MES test, the ethanolic extract reduced the duration of HLTE. According to the data, the extract exhibited protective effects against mortality in MES experiments.

Antioxidant Activity**Free radical scavenging activity of *Senna occidentalis* L by DPPH.**

Polyphenolic compounds present in plant contribute significantly to the total antioxidant capacity of the seeds. Flavonoids play some important pharmacological roles against diseases, such as cardiovascular diseases, cancer, inflammation and allergy. In the present study, reduction of the DPPH radicals was found in concentration dependent

manner. The *Senna occidentalis* ethanolic extract reduced the stable DPPH radical to yellow colored unstable compound. However, ascorbic acid displays significant scavenging activity over the *Senna occidentalis* ethanolic extract. This might be due to the presence of flavanoid content which increases the quenching of the free radical. The absorbance was measured using spectrophotometer and the same is presented in Table 3. The findings of % inhibitions of free radicals are presented in table 4.

Table 3: *In-vitro* Antioxidant effect of extracts of *Senna occidentalis* L. seeds (by DPPH method)

Sl. No	Concentration $\mu\text{g/ml}$	Absorbance	
		Ascorbic Acid	EAE
1	10	0.3254 \pm 0.001	0.2645 \pm 0.004
2	20	0.1917 \pm 0.002	0.2314 \pm 0.005
3	30	0.1012 \pm 0.001	0.1874 \pm 0.003
4	40	0.0742 \pm 0.001	0.1542 \pm 0.012
5	50	0.0654 \pm 0.002	0.0874 \pm 0.011

Table 4: Percentage of Inhibition of EAE with Ascorbic acid

Concentration ($\mu\text{g/ml}$)	Ascorbic Acid (%)	EAE (%)
10	22.76	41.82
20	54.49	51.2
30	75.97	63.39
40	82.38	68.02
50	84.8	79.25

Free radical scavenging activity of *Senna occidentalis* by H_2O_2 method

H_2O_2 is highly important because of its ability to penetrate biological membranes. H_2O_2 itself is not very reactive, but it can sometimes be toxic to cell as it may give rise to hydroxyl radical in the cells. The different fractions of ethanol extract are tested for hydrogen peroxide scavenging activity. The results showed that extracts of *Senna occidentalis* had an effective H_2O_2 scavenging activity. The highest inhibition of free radical was recorded for 50 $\mu\text{g/ml}$ concentration of EAE where as least free radical scavenging was observed for 10 $\mu\text{g/ml}$ concentration of EAE. The inhibition of free radical was recorded in dose dependent manner. The absorbance was measured using spectrophotometer and the same is presented in Table 5.

Table 5: *In-vitro* Antioxidant effect of extract of *Senna occidentalis* seeds (by H_2O_2 method)

Concentration $\mu\text{g/ml}$	Absorbance	
	Ascorbic Acid	EAE
10	0.2145 \pm 0.005	0.2421 \pm 0.002
20	0.1984 \pm 0.004	0.2032 \pm 0.004
30	0.1645 \pm 0.008	0.1874 \pm 0.003
40	0.1121 \pm 0.010	0.1745 \pm 0.011
50	0.0765 \pm 0.007	0.1654 \pm 0.004

Table 6: Percentage of Inhibition of PEE, EAE and AEE with Ascorbic acid

Concentration $\mu\text{g/ml}$	Ascorbic Acid	EAE (%)
10	18.21	24.23
20	52.01	54.28
30	73.21	66.21
40	81.20	69.02
50	84.22	74.25

The % inhibition was recorded for dilution of extracts (10-50 $\mu\text{g/ml}$), using Ascorbic acid as standard and the findings are presented in table 6.

The finding suggested that the EAE of *Senna occidentalis* possesses potent activity of anticonvulsant. The EAE showed anticonvulsant action in dose dependent fashion. The EAE showed less anticonvulsant activity when it was given 100 mg/kg body weight but the effect was increased when dose increased. The maximum anticonvulsant effect was found when the dose was given in 400 mg/kg body weight in PTZ induced seizure. The effect was increased when dose was high. When all the three dose levels of EAE were compared with positive control, it was seen that the EAE (400mg/kg bw.) was most effective extract.

The EAE showed anticonvulsant property in MES induced model in dose dependent fashion. The effect was highest when the dose was given 400 mg/kg and the effect was lowest when the dose was 100 mg/kg b.w. When all the three dose levels of EAE extracts were compared with control group, it was found that the EAE (400mg/kg bw) is having significant anticonvulsant property.

The findings suggested that the EAE possessed a potent antioxidant property. It was seen that as the concentration of the extract increases, the extract showed decrease of absorbance and in comparison with Ascorbic acid, it showed marked antioxidant property in DPPH as well as Hydrogen peroxide model. The IC_{50} is the half maximal inhibitory concentration^{12,13}. The IC_{50} of Ascorbic acid and EAE in DPPH method were found that 14.56 and 14.8 respectively. The IC_{50} of Ascorbic acid and EAE in Hydrogen peroxide model were found that 14.3 and 14.8 respectively. It was seen that, as the extract concentration increases, the absorbance decreases. The result of antioxidant activity using both models suggested that 50 $\mu\text{g/ml}$ concentration of EAE was most potent in exhibiting the antioxidant response when comparison was made with Ascorbic acid.

CONCLUSION

The research conducted on assessment of anticonvulsant and antioxidant profile of *Senna occidentalis* extracts revealed that EAE in 400mg/kg b.w. exhibits most potent anticonvulsant response when evaluation was done on MES and PTZ induced model. Similarly, the 50 µg/ml dilution of EAE showed maximum response in Antioxidant activity in both the models viz DPPH model and Hydrogen peroxide model.

ACKNOWLEDGEMENT

The authors are thankful to Management, Ram-Eesh Institute of Vocational and Technical Education, Greater Noida (U.P.) for assistance towards this research work. The present work is a part of Ph.D from IFTM University, Moradabad.

CONFLICT OF INTEREST

The authors have no conflict of interest.

REFERENCES

1. The Wealth of India. A dictionary of Indian Raw Material and Industrial Products. New Delhi: Council of Scientific and Industrial Research, 1998; pp. 350.
2. Sini KR, Sinha BN, Karpakavalli M, Sangeetha PT. Analgesic and antipyretic activity of *Senna occidentalis* Linn. Annals of Biological Research. 2011; 2 (1):195-200.
3. Hong N, Van TN, Quan VV. Effect of Extraction Solvents and Drying Methods on the Physicochemical and Antioxidant Properties of *Helicteres hirsuta* Lour. Leaves. 2015; 3:285-301
4. Shyamjith M, Deepa B, Anu E Joy and Rao S N. Anticonvulsant Activity of *Phyllanthus Amarusin* Experimental animal Models. International Journal of Applied Biology and Pharmaceutical Technology. 2011; 2(4):144-149.
5. Joshi R, Reeta KH, Sharma SK, Tripathi M, Gupta YK. Pharmacodynamic and pharmacokinetic interaction of Panchagavya Ghrita with phenytoin and carbamazepine in maximal electroshock induced seizures in rats. Ayu. 2015; 36: 196-202.
6. Chauhan AK, Dobhal M.P, Joshi B.C., A review of medicinal plants showing anticonvulsant activity, Journal of Ethnopharmacology, 1988, 22:11-23.
7. McNamara JO., Drugs effective in the therapy of the epilepsies: In: Goodman and Gillman's "The pharmacological basis of therapeutics", McGraw-Hill, New York, 2001, 521-539.
8. Pandey MM, Vijayakumar M, Rastogi S, Rawat AKS. Phenolic content and antioxidant properties of selected Indian spices of Apiaceae. J Herb, Spice Med Plant 2012; 18:246-56.
9. Ebrahimzadeh AM, Nabavi SM, Nabavi SF, Bahramiyan F, Bekhradni AR. Antioxidant and Free Radical Scavenging Activity of *H. Officinalis* L. *angustifolius*, *V. odorata*, *B. hyrcana* and *C. speciosum*. Pakistan Journal of Pharmaceutical Science. 2010; 23(1):29-34.
10. Furuta, S., I. Suda, Y. Nishiba and O. Yamakaw. High tertbutylperoxy radical scavenging activities of sweet potato cultivars with purple flesh. Food Sci., Technol. 1998; 4:33-35.
11. Zahin M, Aqil F, Ahmad I. Broad spectrum antimutagenic activity of antioxidant active fraction of *Punica granatum* L. peel extracts. Mutat Res. 2010; 703:99-107.
12. Zidane H, Elmiz M, Aouinti F, Tahani A, Wathélet J, Sindic M, Elbachiri A. Chemical composition and antioxidant activity of essential oil, various organic extracts of *Cistus ladanifer* and *Cistus libanotis* growing in Eastern Morocco. African J. of Biotechnology, 2013; 34(12): 5314-5320.
13. Tepe B, Sokmen M., Akpulat H.A., Sokmen A. Screening of the antioxidant potentials of six *Salvia* species from Turkey. Food Chem. 2006; 95:200-204
14. Jain S. The role of epilepsy management guidelines in a developing country. Neurology Asia 2011; 16:57-8

