

Available online on 20.04.2019 at <http://jddtonline.info>

# Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Review Article

## Free Radical Pharmacology and its role in various diseases

Priya Gupta \*, RB Jyoti, Saibya Gupta

Department of Pharmaceutical Science, Banasthali Vidyapith, Jaipur, Rajasthan. (India)

### ABSTRACT

Free Radicals are molecules with an unpaired electron and are important intermediates in natural processes. Free radicals are very unstable and react quickly with other compounds, and try to capture the needed electron to gain stability. Once the process is started, it can cascade, and finally results in the disruption of a living cell. Generally, harmful effects of reactive oxygen species on the cell are most often like damage of DNA, oxidations of polydesaturated fatty acids in lipids, oxidations of amino acids in proteins, oxidatively inactivate specific enzymes by oxidation of co-factors. Free radicals cause many human diseases like cancer Alzheimer's disease, cardiac reperfusion abnormalities, kidney disease, fibrosis, etc. The free radicals formed in our body are combated by antioxidants that safely interact with free radicals and terminate the chain reaction before vital molecules are damaged. Excessive exercise has been found to increase the free radical level in the body and causes intense damage to the Regular physical exercise enhances the antioxidant defense system and protects against exercise induced free radical damage.

**Keywords:** Radicals, Free Radicals, Reactive oxygen species, Anti-oxidant, Redox signaling.

**Article Info:** Received 15 March 2019; Review Completed 16 April 2019; Accepted 18 April 2019; Available online 20 April 2019



### Cite this article as:

Gupta P, Jyoti RB, Gupta S, Free Radical Pharmacology and its role in various diseases, Journal of Drug Delivery and Therapeutics. 2019; 9(2-s):690-694 <http://dx.doi.org/10.22270/jddt.v9i2-s.2593>

### \*Address for Correspondence:

Priya Gupta, Department of Pharmaceutical Science, Banasthali Vidyapith, Jaipur, Rajasthan. (India)

### Introduction

Free Radicals are molecules with an unpaired electron. Due to the presence of a free electron, these molecules are highly reactive. They are important intermediates in natural processes involved in cytotoxicity, control of vascular tone, and neurotransmission. Radiolysis is a powerful method to generate specific free radicals and measure their reactivity.<sup>1</sup>

### Types of free radicals:

1. Super oxide
2. Hydrogen oxide
3. Hydroxyl radical
4. Singlet Oxygen
5. Hydroperoxy Radical.
6. Lipid peroxide Radical.
7. Nitric Oxide.

### Generation of Free Radical:

The generation can be endogenous (within the body) or exogenous (outside the body). The exogenous source of free radical is from the environment. These include oxidizing agent i.e. Ionizing radiations (from industry, sun exposure,

cosmic rays, medical x-ray); Ozone and nitrous oxide (primarily from automobile exhaust); Heavy metals (such as mercury, cadmium and lead); Cigarette smoking (both active and passive); Alcohol; Unsaturated fat (this may create a strain on the natural antioxidants of the body) and other chemicals (pesticides) and compounds from food, water, drugs and air. Endogenous free radical formation occurs continuously in the cells as a consequence of both enzymatic and non-enzymatic reactions. Endogenous free radicals are produced in the body by four different mechanisms:

- The normal metabolism of oxygen requiring nutrients.
  - White blood cells destroy parasites, bacteria and viruses by using oxidants (free radicals) such as nitric oxide, super oxide and hydrogen peroxide.
1. Other cellular components called peroxisomes produce hydrogen peroxide as a by-product of the degradation of fatty acids and other molecules.
  2. An enzyme in the cells called cytochrome P450 is one of the body's primary defenses against toxic chemicals ingested with food.<sup>2,3,4</sup>

## Sources of Free Radical:

### 1. Endogenous:

- Mitochondria
- Xanthine oxidase
- Phagocytes
- Peroxisomes
- Arachidonate Pathways
- Ischemia/Reperfusion
- Inflammation

### 2. Exogenous:


- Cigarette Smoking
- Environmental Pollutants
- Radiation
- Ozone
- Industrial Solvents

### 3. Physiological Factor:

- Mental Status
- Disease Condition

## Free Radical Scavengers (Antioxidant)

**Antioxidant:** Those substances when present at low concentration, compared to those of an oxidizable substrate, will significantly delay or inhibit oxidation of that substrate.



## Sources of Antioxidant:


**Endogenous:** Superoxide dismutase, Catalase, Glutathione Peroxidase

**Food:** Vitamin A, C, E, beta, Carotene, Selenium, Lycopene.

### Targets:

- 1. Lipids:** Peroxidation of lipids in cell membrane can damage cell membrane by disrupting fluidity and permeability. Lipid peroxidation can also adversely affect the function of membrane bound protein such as enzyme and receptor.
- 2. Protein:** Direct damage to protein can be caused by free radical; it affects many proteins interfering with enzyme activity and that of structural proteins.
- 3. DNA:** Fragmentation of DNA is caused by free radical attack which causes activation of **polysynthetase** enzyme; it splits NAD<sup>+</sup> to aid the repair of DNA.<sup>5</sup>

## Role of free radical in various diseases:



### Diabetes Mellitus:

Diabetes mellitus is heterogeneous group of chronic disorders characterized by enhanced blood glucose levels (hyperglycemia) resulting from defective insulin secretion (in type I diabetes), resistance to insulin action (in type II diabetes) or both. The major symptoms are thirst, hunger, emaciation, and weakness, eventually lead to coma. DM is associated with the increased production of free radicals or decreased activity of the antioxidant systems, which leads to development of oxidative stress.

### The hyperglycemic condition induces increased free radical production via four different routes namely:

1. Increased glycolysis, results in increased ratio between the rate of oxidation of G3P to 1, 3-DPG, following increased NADH/NAD<sup>+</sup> ratio (redox imbalance).
2. Activated sorbitol (or polyol) pathway, causes the accumulation of both sorbitol and fructose, results in decreased reduced GSH and increased NADH/NAD<sup>+</sup> ratio.
3. Autoxidation of glucose, results in the generation of different free radicals such as H<sub>2</sub>O<sub>2</sub>, OH<sup>•</sup>, O<sub>2</sub><sup>•-</sup> and keto aldehydes.
4. Non enzymatic protein glycation, results in the formation of AGEs which upon interacting with RAGEs generate oxidative stress.<sup>6,7,8,9.</sup>

### Neurodegenerative Diseases

The central nervous system (CNS) is mainly susceptible to the oxidants due to the presence of high lipid content, high consumption of oxygen, and low levels of antioxidant enzymes, for example, SOD is localized primarily in neurons, and GSH and GPx are restricted in astrocytes. The lipid peroxidation by ROS leads to progressive loss of membrane fluidity, decreases membrane potential, and increases permeability to ions such as Ca<sup>2+</sup>. The region of the brain such as hippocampus nigr and the striatum are particularly susceptible to attack by free radicle.<sup>10,11,12</sup>

### Alzheimer's Disease:

AD is characterized by the accumulation of amyloid protein plaques (formed from the improper folding and processing of amyloid  $\beta$  pr.....P-O-P-ecursor protein and intracellular neurofibrillary tangles made up of abnormal and hyper phosphorylated tau protein. The hyper phosphorylated tau protein aggregates binds to Fe<sup>3+</sup>, results in the production of neurofibrillary tangles. The Amyloid- $\beta$  peptide (A $\beta$ ) can chelate with transition metal ions (Cu<sup>2+</sup>, Zn<sup>2+</sup> and Fe<sup>3+</sup>). The lipid peroxidation is also extensive in AD patients, which can induce neuronal death by multiple mechanisms such as impairment of function of ion pumps (both Na<sup>+</sup>/K<sup>+</sup>-ATPase

and Ca<sup>2+</sup>-ATPase), glucose transporters and glutamate transporters. The other oxidative markers of protein damage such as protein carbonyls and 3-nitrotyrosine have been also observed in AD patients.<sup>17,18,19</sup>

### Cancer

It is one of the leading causes of death in humans. Free radicals cause different types of chemical changes in DNA. Cancer cells in particular, in comparison to normal cells, have higher levels of ROS and are more susceptible to mitochondrial dysfunction due to their higher metabolic rate. Cancer cells display elevated levels of oxidative stress due to activation of oncogenes and loss of tumor suppressors. ROS by altering the growth signals and gene expression cause continuous proliferation of cancer cells. ROS can ...damage DNA by inducing base modifications, deletions, strand breakage, chromosomal rearrangements and hyper- and hypo-methylation of DNA.<sup>20,21,22,23,24,25</sup>

### Breast Cancer

In majority of breast carcinomas the oxidative stress can be induced by the over expression of thymidine phosphorylase enzyme which catabolizes thymidine to thymine and 2-deoxy-D-ribose-1-phosphate; the latter is a powerful reducing sugar that rapidly glycosylates proteins, generating oxygen radicals within the carcinoma cell. Another breast specific mechanism of oxidative stress induction involves a mammary gland specific lactoperoxidase enzyme catalyzed one electron oxidation of 17- $\beta$ -oestradiol to a reactive phenoxyl radical.<sup>26,27</sup>

### Prostate Cancer

The ROS produced are responsible for the cellular proliferation of prostate cancer cells. Overexpression of NADPH oxidase 1 (Nox1) protein is an early event in the development of prostate cancer. The superoxide produced (by NOX) in prostate cancer cells facilitates cellular immortality through resistance to programmed cell death which results in cancer-promoting effect.<sup>28,29,30</sup>

### Lung Cancer

Lung cancer has been the most commonly diagnosed cancer and is the central cause of cancer death in men worldwide. Lung cancer mortality account 30 % of all cancer related deaths. Cigarette Smoking is the most crucial environmental risk factor in lung cancer etiology. It is estimated that smoking accounts for ~80 % of global lung cancer burden in males and 50 % in females. Cigarette smoke particulate matter contains complex mixture of numerous carcinogens and stable ROS with very long half-lives. These ROS can damage the tissues resulting in progressive transformation of cells into the malignant form, which leads to increased frequency of mutations by the oxidative damage to DNA and, eventually leading to lung cancer. Smokers develop lung cancer a 10-fold higher than non-smokers. Lung cancer (LC) and chronic obstructive pulmonary disease (COPD) commonly coexist in smokers, and the presence of COPD increases the risk of developing LC.<sup>31,32,33,34,35</sup>

### Bladder Cancer

The most common risk factors for bladder cancer are cigarette smoking, exposure to industrial carcinogens (aromatic amines), high levels of arsenic intake and diet. Oxidative stress critically contributes to the development of bladder cancer. Various lines of evidence reported an increased oxidative stress in patients with breast cancer. Increased NO levels have been reported in bladder cancer patients. This NO stimulates matrix metalloproteinase (MMPs), especially prolydase activity, which is involved in

the terminal step of collagen degradation. Significantly higher serum prolydase activities were reported in patients with bladder cancer than healthy controls.<sup>36,37,38,39,40,41</sup>

### Cardiovascular Diseases (CVDs)

Cardiovascular diseases are a class of pathologies involving the heart and blood vessels (arteries, capillaries, and veins). They include cardiac diseases, vascular diseases of the brain and kidney, and peripheral arterial disease. Most of the people are dying due to CVDs compared to other diseases<sup>42</sup>.

### Hypertension (HT)

Hypertension (HT) is a major health problem. Persons with hypertension are at an increased risk for stroke, heart disease, kidney failure, and premature mortality. Increased ROS generation eliminates NO<sup>•</sup> by forming ONOO<sup>-</sup>; thus reducing NO<sup>•</sup> bioavailability which leads to decreased endothelium-dependent vasodilation subsequent in hypertension. A decrease in NO bioavailability and an increase in oxidative stress are present in human hypertension. Oxidation-induced impairment of NO also results in reduced opposition to the vasoconstrictive and hypertensive effects of angiotensin II. Angiotensin II decreases NO bioavailability by promoting oxidative stress.<sup>43,44,45,46</sup>

### Rheumatoid Arthritis (RA)

RA is a chronic disease. The disease is characterized by synovial and systemic inflammation with joint swelling, morning stiffness, destruction of articular tissues, joint deformities, fatigue, loss of appetite and weakness. It is believed to be a T-lymphocyte. This cytokine release and subsequent migration is thought to be responsible for the chronic inflammation and characteristic changes in RA.

Both ROS and RNS damage cartilage. Tissue injury in inflammation results in NO<sup>•</sup> production by articular chondrocytes and synovial fibroblasts and elevated levels of NO<sup>•</sup> are observed in the serum and synovial fluid of RA patients. The free radicals, particularly NO<sup>•</sup> and O<sub>2</sub><sup>-•</sup>, inhibit the synthesis of matrix components like proteoglycans by chondrocytes and also damage the extracellular matrix through activation and up regulation of matrix metalloproteinase. The HOCl, produced by myeloperoxidase (MPO) in neutrophils, chlorinate the tyrosine residues to form 3-chlorotyrosine and damage the collagen, thus implicated in arthritogenesis. RA patients have increased plasma MPO concentrations.<sup>47,48</sup>

### Asthma

Asthma is the most common disorders of the airways of the lungs and is one of the major global health problems. It is characterized by chronic inflammation of the airways involving variable and recurrent airflow obstruction and bronchial hyper reactivity associated with airway remodeling. Airway remodeling is a dynamic process involving mucous hyper secretion, collagen deposition, wall thickening, myocyte hypertrophy and hyperplasia, myofibroblast hyperplasia, vascular proliferation and alterations in airway elastic fibers, all of which culminate in persistent structural alterations of the airway. NO is endogenously produced in mammalian airways by NOS and is known to regulate many aspects of human asthma, including modulation of airway and vascular smooth muscle tone and the inflammation.

## The Benefits of Free Radicals

- In body they control the flow of blood through our arteries, to fight infection, to keep our brain alert and focus.
- Similar to antioxidants, some free radicals are involved as signaling molecules, i.e, they are responsible for turning on and off genes.
- Some free radicals like NO and superoxide produced in a very high amount by immune cells to poison viruses and bacteria.
- Some free radicals kill cancer cells, in fact certain cancer drugs aim in increasing the free radical amount in the body.<sup>49</sup>

## References

- [1] R. T. Oakley, Prog. Inorg. Chem., 1998, 36,
- [2] Lobo V, et al. "Free radicals, antioxidants and functional foods: Impact on human health". *Pharmacognosy Reviews* 4.8 (2010): 118-126.
- [3] Dhalla NS, et al. "Role of Oxidation Stress Cardiovascular Diseases". *Hypertension* 18.6 (2000): 655.
- [4] Antioxidants and Free Radical (2005).
- [5] <https://www.slideshare.net/Aejaz/mechanisim-and-generation-of-free-radicals>
- [6] Gavin JR, Alberti KGMM, Davidson MB, DeFronzo RA, Drash A, Gabbe SG, et al. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care*. 1997; 20:1183-1197
- [7] Oberley LW. Free radicals and diabetes. *Free Radic Biol Med*. 1988; 5(2):113-124. [PubMed]
- [8] Bashan N, Kovsan J, Kachko I, Ovadia H, Rudich A. Positive and negative regulation of insulin signaling by reactive oxygen and nitrogen species. *Physiol Rev*. 2009; 89(1):27-71.
- [9] Ahmed RG. The physiological and biochemical Effects of diabetes on the balance between oxidative stress and Antioxidant defense system. *Med J Islam World Acad Sci*. 2005; 15(1):31-42.
- [10] Pollack M, Leeuwenburgh C. Molecular mechanisms of oxidative stress in aging: free radicals, aging, antioxidants and disease. Elsevier Science B.V. Handbook of Oxidants and Antioxidants in Exercise. 1999; 881-923.
- [11] Rivas-Arancibia S, Guevara-Guzmán R, López-Vidal Y, Rodríguez-Martínez E, Zanardo-Gomes M, Angoa-Pérez M, et al. Oxidative stress caused by ozone exposure induces loss of brain repair in the hippocampus of adult rats. *Toxicol Sci*. 2010; 113(1):187-197. [PubMed]
- [12] Santiago-López JA, Bautista-Martínez CI, Reyes-Hernandez M, Aguilar-Martínez S, Rivas- Arancibia S. Oxidative stress, progressive damage in the substantianigra and plasma dopamine oxidation, in rats chronically exposed to ozone. *ToxicolLett*. 2010; 197(3):193-200. [PubMed]
- [13] Pan XD, Zhu YG, Lin N, Zhang J, Ye QY, Huang HP, Chen XC. Microglial phagocytosis induced by fibrillar  $\beta$ -amyloid is attenuated by oligomeric  $\beta$ -amyloid: implications for Alzheimer's disease. *MolNeurodegener*. 2011; 6(45):1-17
- [14] Sevcsik E, Trexler AJ, Dunn JM, Rhoades E. Allosteric in a disordered protein: oxidative modifications to  $\alpha$ -synuclein act distally to regulate membrane binding. *J Am Chem Soc*. 2011; 133(18):7152-7158
- [15] Zhao W, Varghese M, Yemul S, Pan Y, Cheng A, Marano P, et al. Peroxisome proliferator activator receptor gamma coactivator-1alpha (PGC-1 $\alpha$ ) improves motor performance and survival in a mouse model of amyotrophic lateral sclerosis. *MolNeurodegener*. 2011; 6(1):1-8
- [16] Witherick J, Wilkins A, Scolding N, Kemp K. Mechanisms of oxidative damage in multiple sclerosis and a cell therapy approach to treatment. *Autoimmune Dis*. 2010; 1-11. [PMC free article] [PubMed]
- [17] Butterfield DA, Perluigi M, Sultana R. Oxidative stress in Alzheimer's disease brain: new insights from redox proteomics. *Eur J Pharmacol*. 2006; 545(1):39-50.
- [18] Chang Y, Kong Q, Shan X, Tian G, Ilieva H, Cleveland DW, et al. Messenger RNA oxidation occurs early in disease pathogenesis and promotes motor neuron degeneration in ALS. *PLoS ONE*. 2008; 3(8):1-19
- [19] Uttara B, Singh AV, Zamboni P, Mahajan RT. Oxidative stress and neurodegenerative diseases. A Review of upstream and downstream antioxidant therapeutic options. *CurrNeuropharmacol*. 2009; 7(1):65-74
- [20] Goldstein BD, Witz G. Free radicals and carcinogenesis. *Free Radic Res Comm*. 1990; 11(1-3):3-10. [PubMed]
- [21] Dreher D, Junod AF. Role of oxygen free radicals in cancer development. *Eur J Cancer*. 1996; 32A(1):30-38. [PubMed]
- [22] Acuna UM, Wittwer J, Ayers S, Pearce CJ, Oberlies NH, De Blanco EJ. Effects of (5Z)-7-Oxozeaenol on the Oxidative pathway of cancer cells. *Anticancer Res*. 2012; 32(7):2665-2671.
- [23] Cairns RA, Harris I, McCracken S, Mak TW. Cancer cell metabolism. *Cold Spring Harb Symp Quant Biol*. 2011; 76:299-311.
- [24] Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal hitological functions and human disease. *Int J Biochem Cell Biol*. 2007; 39(1):44-84.
- [25] Valko M, Izakovic M, Mazur M, Rhodes CJ, Telser J. Role of oxygen radicals in DNA damage and cancer incidence. *Mol Cell Biochem*. 2004; 266(1-2):37-56. [PubMed]
- [26] Brown NS, Jones A, Fujiyama C, Harris AL, Bicknell R. Thymidine phosphorylase induces carcinoma cell oxidative stress and promotes secretion of angiogenic factors. *Cancer Res*. 2000; 60(22):6298-6302.
- [27] Sipe HJ, Jr, Jordan SJ, Hanna PM, Mason RP. The metabolism of 17 beta-estradiol by lactoperoxidase: a possible source of oxidative stress breast cancer. *Carcinogenesis*. 1994; 15(11):2637-2643.
- [28] Arnold RS, He J, Remo A, Ritsick D, Yin-Goen Q, Lambeth JD, et al. Nox1 expression determines cellular reactive oxygen and modulates c-fos-induced growth factor, interleukin-8, and Cav-1. *Am J Pathol*. 2007; 171(6):2021-2032. [PMC free article] [PubMed]
- [29] Lim SD, Sun C, Lambeth JD, Marshall F, Amin M, Chung L, et al. Increased Nox1 and hydrogen peroxide in prostate cancer. *Prostate*. 2005; 62(2):200-207.
- [30] Veeramani S, Yuan TC, Lin FF, Lin MF. Mitochondrial redox signaling by p66Shc in involved in regulating androgenic growth stimulation of human prostate cancer cell. *Oncogene*. 2008; 27(37):5057-5068.
- [31] WHO World cancer Report 2008. In: Boyle P, Levin B, editors. Lung cancer, 12.Chapter 5.10.
- [32] Azad N, Rojanasakul Y, Vallyathan V. Inflammation and lung cancer: roles of reactive oxygen/nitrogen species. *J Toxicol Environ Health*. 2008; 11(1):1-15.
- [33] Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin*. 2011; 61(2):69-90.
- [34] Hoagland LF 4th, Campa MJ, Gottlin EB, Herndon 2nd JE, Patz Jr EF. Haptoglobin and posttranslational glycan-modified derivatives as serum biomarkers for the diagnosis of nonsmall cell lung cancer. *Cancer*. 2007; 110(10):2260-2268.
- [35] Pastora MD, Nogala A, Molina-Pinelo S, Meléndez R, Salinas A, González De la Penaa M, et al. Identification of proteomic signatures associated with lung cancer and COPD. *J Proteomics*. 2013; 89:227-237.
- [36] Wynder EL, Goldsmith R. The epidemiology of bladder cancer: a second look. *Cancer*. 1977; 40:1246-1268. [PubMed]



- [37]. Opanuraks J, Boonla C, Saelim C, Kittikowit W, Sumpatanukul P, Honglertsakula C, et al. Elevated urinary total sialic acid and increased oxidative stress in patients with bladder cancer. *Asian Biomedicine*. 2010; 4(5):703–710.
- [38]. Opanuraks J, Boonla C, Saelim C, Kittikowit W, Sumpatanukul P, Honglertsakula C, Tosukhowong P. Elevated urinary total sialic acid and increased oxidative stress in patients with bladder cancer. *Asian Biomedicine*. 2010; 4(5):703–710.
- [39]. Soini Y, Haapasaari KM, Vaarala MH, Turpeenniemi-Hujanen T, Karja V, Karihtala P. 8-hydroxydeguanosine and nitrotyrosine are prognostic factors in urinary bladder carcinoma. *Int J ClinExpPathol*. 2011; 4(3):267–275.
- [40]. Eijan AM, Piccardo I, Riveros MD, Sandes EO, Porcella H, Jasnis MA, et al. Nitric oxide in patients with transitional bladder cancer. *J SurgOncol*. 2002;81:203–208.
- [41]. Gecit I, Aslan M, Gunes M, Pirincci N, Esen R, Demir H, et al. Serum prolidase activity, oxidative stress, and nitric oxide levels in patients with bladder cancer. *J Cancer Res ClinOncol*. 2012; 138(5):739–743.
- [42]. Ellidag HY, Eren E, Aydın O, Akgol E, Yalcınkaya S, Sezer C, et al. Ischemia Modified Albumin Levels and Oxidative Stress in Patients with Bladder Cancer. *Asian Pacific J Cancer Prev*. 2013; 14(5):2759–2763.
- [43]. A global brief on hypertension. World health day 2013. WHO
- [44]. Dzau VJ. Tissue angiotensin and pathobiology of vascular disease: a unifying hypothesis. *Hypertension*. 2001; 37:1047–1052.
- [45]. Touyz RM. Reactive oxygen species, vascular oxidative stress, and redox signaling in hypertension: what is the clinical significance? *Hypertension*. 2004; 44(3):248–252.
- [46]. Touyz RM. Reactive oxygen species and angiotensin II signaling in vascular cells: implications in cardiovascular disease. *Braz J Med Biol Res*. 2004; 37(8):1263–1273.
- [47]. Stamp LK, Khalilova I, Tarr JM, Senthilmohan R, Turner R, Haigh RC, et al. Myeloperoxidase and oxidative stress in rheumatoid arthritis. *Rheumatology (Oxford)* 2012; 51(10):1796–1803.
- [48]. Desai PB, Manjunath S, Kadi S, Chetana K, Vanishree J. Oxidative stress and enzymatic antioxidant status in rheumatoid arthritis: a case control study. *Eur Rev Med Pharmacol Sci*. 2010; 14(11):959–967.
- [49]. Xiao M, Zhu T, Wang T, Wen FQ. Hydrogen-rich saline reduces airway remodeling via inactivation of NF- $\kappa$ B in a murine model of asthma. *Eur Rev Med Pharmacol Sci*. 2013; 17(8):1033–1043.
- [50]. <https://www.floressencetea.com/The-Benefits-and-Risks-of-Free-Radicals-a/257.htm>

