

Available online on 15.05.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Review Article

Varicose Vein of Bilateral Lower Limb: A Review

Arsheed Iqbal*¹, Afroza Jan², Haider Ali Quraishi³, Arjumand Shah⁴, Huma⁴, Md. Sheeraz⁴

- 1- Research officer Unani (Scientist-III), Regional Research Institute of Unani Medicine, University of Kashmir, Srinagar, J & K., India.
- 2- Assistant Professor, Institute of Asian Medical Science, Srinagar.
- 3- M.D. Scholar Unani Medicine, Regional Research Institute of Unani Medicine, University of Kashmir, Srinagar, J & K., India.
- 4- Research officer Unani, Regional Research Institute of Unani Medicine, University of Kashmir, Srinagar, J & K., India.

ABSTRACT

Varicose veins are tortuous, winded veins in the subcutaneous tissues of legs. These veins are visible and can be easily diagnosed clinically. The varicose veins are usually incompetent and may result in the reflux of blood and ultimately venous hypertension. Varicose veins are usually found asymptomatic however it may result in local pain, heaviness in the limb, change in the color, itching, dryness and may develop ankle swelling including cramps. Conditions that increase pressure on leg veins such as obesity, sedentary life style, lack of activity or exercise, hormonal fluctuations during pregnancy, menopause, smoking, heavy weight lifting may lead to varicose veins.

Keywords: - Varicose, Reflux, Vein, Incompetent.

Article Info: Received 24 March 2019; Review Completed 30 April 2019; Accepted 02 May 2019; Available online 15 May 2019



Cite this article as:

Iqbal A, Jan A, Quraishi HA, Shah A, Huma, Sheeraz M, Varicose Vein of Bilateral Lower Limb: A Review, Journal of Drug Delivery and Therapeutics. 2019; 9(3):584-587 <http://dx.doi.org/10.22270/jddt.v9i3.2634>

*Address for Correspondence:

Dr. Arshid Iqbal, Research Officer Unani (Scientist -III & M.D. Scholar), Regional Research Institute of Unani Medicine, Kashmir University, Srinagar. J & K.190006

INTRODUCTION

The great saphenous vein is a large, subcutaneous superficial vein of the leg. It is the largest vein in the body running along the length of the lower limb, returning blood from the foot, leg and thigh to the deep femoral vein at the femoral triangle. Veins have one-way valves that prevent blood from flowing backwards. When the walls of a vein become weak especially near a valve, the valve becomes leaky and malfunctioning. The blood then begins to flow backwards instead of towards the heart. Once this happens, it puts a lot of pressure on the vein which can cause the vein to widen and the occurrence of more leaky valves. Varicose veins often affect the legs. Varicose veins are more common in women than in men and are linked with heredity,¹ other related factors are pregnancy, obesity, menopause, aging, prolonged standing, leg injury and abdominal straining. Varicose veins are unlikely to be caused by crossing the legs or ankles.² More recent research has shown the importance of pelvic vein reflux (PVR) in the development of varicose veins. Hobbs showed varicose veins in the legs could be due to ovarian vein reflux⁴. Whiteley and his team reported that both ovarian and internal iliac vein reflux causes leg varicose veins and that this condition affects 14% of women with varicose veins or 20% of women who have had vaginal

delivery and have leg varicose veins⁶. Varicose veins could also be caused by hyperhomocysteinemia in the body, which can degrade and inhibit the formation of the three main structural components of the artery. Homocysteine permanently degrades cysteine disulfide bridges and lysine amino acid residues in proteins, gradually affecting function and structure. Simply put, homocysteine is a 'corrosive' of longliving proteins, i.e. collagen or elastin, or lifelong proteins, i.e. fibrillin. These longterm effects are difficult to establish in clinical trials focusing on groups with existing artery decline.¹⁰

MATERIAL AND METHODS

The databases used to get information from journals and articles are Google, Pub Med, Science Direct, Scopus and Google Scholar. For the search of primordial and current Unani Classical literature author visited Library of Regional Research Institute of Unani Medicine (RRIUM), Srinagar, J & K, India.

Incidence

A study in UK population shows that it affects 40 % in men and 32% in woman, however women are found more in clinics for treatment.¹¹ Varicose veins often appear first in

pregnancies can make them more worse. Varicose veins can also be found in teen age and also shows positive family history in its occurrence.¹² The risk of varicose vein increase with aging. The vein becomes weak with age and eventually wear and tear can occur which many cause the valvular malformation to malformation.

Precaution

- To avoid the prolonged standing.
- Maintain the body weight as per medical recommendation.
- Try to check and regulate the hormone levels.
- Post-menopausal care.
- Management of life style during pregnancy.
- Avoid sedentary work.
- Avoid heavy weight lifting.
- Precaution and prophylactic care for genetically predisposed family.
- Elevating the leg.
- Wearing the intermittent pneumatic compression devices have been shown to reduce swelling and regulate circulation.

Treatment

The immediate treatment for varicose veins is compression stockings which will squeeze the legs and ultimately squeezing the veins to let the blood flow easily towards the heart. Certain exercises may be good for varicose veins such as swimming, Yoga, and Pilates. Regimental therapy like massage, steaming, cupping and leeching also helps in the management of the varicose veins. The other modalities of treatment are

- Laser surgeries for varicose veins.
- Catheter assisted procedures using radiofrequency or laser energy.
- High ligation and vein stripping.
- Endoscopic vein surgery in severe cases.

Some other particular methods of treatment

I. Stripping

II. Chiva

III. Sclerotherapy

IV. Endovenous Thermal Ablation

I-Stripping

Stripping consists of removal of all or part of the saphenous vein. The complications included deep vein thrombosis (5.3%)¹³, pulmonary embolism (0.06%), and wound complications including infection (2.2%). There is evidence for the great saphenous vein regrowing after stripping.¹⁴ In addition, since stripping removes the saphenous main trunks, they are no longer available for use as venous bypass grafts in the future.¹⁵

II- Chiva

Experimental evidence shows that the ambulatory conservative hemodynamic correction of venous insufficiency method (CHIVA), which intends saving the veins, decreases varicose veins and is safer than vein

stripping in those with chronic venous insufficiency.¹⁶ CHIVA comes from the French translation for the technique, *conservatrice et hémodynamique de l'insuffisance veineuse en ambulatoire*. The operation involves performing one to four small incisions on average, under local anesthesia, to remove by precise ligatures abnormal blood flow due to valvular incontinence and responsible for varicose vein dilation. The patient returns home the same day. This method tends to correct the venous functions in order to cure the symptoms of venous insufficiency such as varicose veins, edema, and ulcers.¹⁷ The superiority of the CHIVA treatment in recurrences has been demonstrated in several Randomized Controlled Trials judged to be of average quality.¹⁸⁻²⁶

III- Sclerotherapy

A commonly performed non-surgical treatment for varicose and "spider" leg veins is sclerotherapy, in which medicine (sclerosant) is injected into the veins to make them shrink. The medicines that are commonly used as sclerosants are polidocanol, sodium tetradecyl sulphate (STS), Sclerodex (Canada), Hypertonic Saline, Glycerin and Chromated Glycerin. Foams may allow more veins to be treated per session with comparable efficacy. Sclerotherapy has been used in the treatment of varicose veins for over 150 years. Sclerotherapy is often used for telangiectasias (spider veins) and varicose veins that persist or recur after vein stripping.^{28,29} Sclerotherapy can also be performed using foamed sclerosants under ultrasound guidance to treat larger varicose veins, including the great saphenous and small saphenous veins.^{30,31} Complications of sclerotherapy are rare but can include blood clots and ulceration. Anaphylactic reactions are "extraordinarily rare but can be life-threatening," and doctors should have resuscitation equipment ready.^{32,33}

IV- Endovenous thermal ablation

There are three kinds of endovenous thermal ablation treatment possible: laser, radiofrequency, and steam.³⁴ The Australian Medical Services Advisory Committee (MSAC) in 2008 determined that endovenous laser treatment/ablation (ELA) for varicose veins "appears to be more effective."³⁵ Complications for ELA include minor skin burns (0.4%) and temporary paresthesia (2.1%).³⁶ Complications for ERA include burns, paraesthesia, clinical phlebitis and slightly higher rates of deep vein thrombosis (0.57%) and pulmonary embolism (0.17%). Steam treatment consists of injection of pulses of steam into the sick vein. This treatment which works with a natural agent (water) has similar results than laser or radiofrequency.³⁷ The steam presents a lot of post-operative advantages for the patient.³⁸

V- Other

Other surgical treatments are: Ambulatory phlebectomy Vein ligation is done at the saphenofemoral junction after ligating the tributaries at the saphenofemoral junction without stripping the long saphenous vein provided the perforator veins are competent and absent DVT in the deep veins. With this method, the long saphenous vein is preserved. Cryosurgery- A cryoprobe is passed down the long saphenous vein following saphenofemoral ligation. Then the probe is cooled with NO₂ or CO₂ to -85 °F. The vein freezes to the probe and can be retrogradely stripped after 5 seconds of freezing. It is a variant of Stripping. The only point of this technique is to avoid a distal incision to remove the stripper.²⁷

CONCLUSION

Varicose vein is a commonest manifestation due to change in life style and work culture. Various pharmacological

therapies where used like anticoagulant, Subcutaneous LDUH, LMWH and phlethysmography were later replaced by surgical intervention and superseded by venous sclerotherapy, endovenous laser ablation, radiofrequency ablation, ambulatory phlebectomy, foam sclerotherapy and cutaneous laser therapy. In Unani Medicine the intervention of Hirudotherapy proven highly efficacious in varicose veins. Leeches not only suck blood over the varicose veins but also inject the bioactive substances like Hirudin, Hyluridase, Acetylcholine, Hirudin, Decorcin, Calin, Typtase inhibitors and natural steroids, which is observed to regulate the venous flow, improves the valvular functioning and thus preventing the back flow of venous blood. Hirudotherapy also balances deranged humoral level of the body which may help in improving the general body vascular tone and venous valvular physiology which prevents the predisposition of varicose vein in the subjects with history of recurrence or with positive family history.

Acknowledgement

Authors duly acknowledge the Assistant Director Incharge, clinical and library staff of Regional Research Institute of Unani medicine, Srinagar, J and K, India.

REFERENCES

- Ng M, Andrew T, Spector T, Jeffery S, "Linkage to the FOXC2 region of chromosome 16 for varicose veins in otherwise healthy, unselected sibling pairs" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1736007>). *Journal of Medical Genetics.*, 2005; 42 (3): 235-9.
- Kate Griesmann (March 16, 2011). "Myth or Fact: Crossing Your Legs Causes Varicose Veins" (https://web.archive.org/web/20140305174414/http://www.dukehealth.org/health_library/health_articles/myth-or-fact-crossing-your-legs-causes-varicose-veins). Duke University Health System. Archived from the original (http://www.dukehealth.org/health_library/health_articles/myth-or-fact-crossing-your-legs-causes-varicose-vein) on 2014-03-05. Retrieved March 1, 2014.
- Franceschi, Claude, "Physiopathologie Hémodynamique de l'Insuffisance veineuse", p. 49 in *Chirurgie des veines des Membres Inférieurs*, AERCV éditions 23 rue Royale 75008 Paris France, 1996.
- Hobbs JT "Varicose veins arising from the pelvis due to ovarian vein incompetence" *Int J Clin Pract.*, October 2005; 59: 1195-203.
- Giannoukas AD, Dacie JE, Lumley JS., "Recurrent varicose veins of both lower limbs due to bilateral ovarian vein incompetence". *Ann Vasc Surg.*, July 2000; 14: 397-400.
- Marsh P, Holdstock J, Harrison C, Smith C, Price BA, Whiteley MS., "Pelvic vein reflux in female patients with varicose veins: comparison of incidence between a specialist private vein clinic and the vascular department of a National Health Service District General Hospital". *P. hlebology.*, June 2009; 24: 108-13.
- Whiteley A.M.; Taylor D.C.; Dos Santos S.J.; Whiteley M.S., "Pelvic Venous Reflux is a Major Contributory Cause of Recurrent Varicose Veins in More Than a Quarter of Women". *Journal of Vascular Surgery: Venous and Lymphatic Disorders.*, 2014; 2: 390-396.
- Whiteley MS., "Part One: For the Motion. Venous Perforator Surgery is Proven and Does Reduce Recurrences". *European Journal of Vascular and Endovascular Surgery.*, September 2014; 48 (3):239-42.
- Rutherford EE, Kianifard B, Cook SJ, Holdstock JM, Whiteley MS., "Incompetent perforating veins are associated with recurrent varicose veins". *European Journal of Vascular and Endovascular Surgery.*, May 2001; 21 (5):458-60.
- Pathophysiology for the Boards and Wards, Fourth Edition
- Lee AJ, Evans CJ, Allan PL, Ruckley CV, Fowkes GFR. Lifestyle factors and the risk of varicose veins. *J Clin Epidemiol* 2003; 56:171-9.
- Campbell WB, Decaluwe H, MacIntyre J, Thompson JF, Cowan AR. Most patients with varicose veins have fears or concerns about the future, in addition to their presenting symptoms. *Eur J Vasc Endovasc Surg* 2006; 31:332-4.
- Van Rij AM, Chai J, Hill GB, Christie RA., "Incidence of deep vein thrombosis after varicose vein surgery". *J Surg.*, 2004; 91 (12):1582-5.
- Munasinghe A, Smith C, Kianifard B, Price BA, Holdstock JM, Whiteley MS., "Strip-tract revascularization after stripping of the great saphenous vein". *Br J Surg.*, 2007; 94 (7): 840-3.
- Hammarsten J, Pedersen P, Cederlund CG, Campanello M., "Long saphenous vein saving surgery for varicose veins. A long-term follow-up". *Eur J Vasc Surg.*, 1990; 4 (4): 361-4.
- Bellmunt-Montoya S; Escribano JM; Dilme J; Martinez-Zapata MJ., "CHIVA method for the treatment of chronic venous insufficiency". *The Cochrane Database of Systematic Reviews* (6): 29 June 2015; CD009648.
- Milone M, Salvatore G, Maietta P, Sosa Fernandez LM, Milone *Recurrent varicose veins of the lower limbs after surgery. Role of surgical technique (stripping vs. CHIVA) and surgeon's experience.* *F. G Chir.* 2011 Nov-Dec; 32(11-12):460-3.
- Carandina S, Mari C, De Palma M, Marcellino MG, Cisno C, Legnaro A, et al. *aricose Vein Stripping vs Haemodynamic Correction (CHIVA): a long term randomised trial.* *European Journal of Vascular and Endovascular Surgery.* 2008; 35(2):230-7
- Parés JO, Juan J, Tellez R, Mata A, Moreno C, Quer FX et l. *Varicose vein surgery: stripping versus the CHIVA Method: a randomized controlled trial.* *Annals of Surgery* 2010; 251(4):624-31.
- Iborra-Ortega E, Barjau-Urrea E, Vila-Coll R, Ballon-Carazas H, Cairols-Castellote MA. *Comparative study of two surgical techniques in the treatment of varicose veins of the lower extremities: results after five years of followup. Estudio comparativo de dos técnicas quirúrgicas en el tratamiento de las varices de las extremidades inferiores: resultados tras cinco años de seguimiento.* *Angiología* 2006; 58(6):459-68.
- P.Zamboni et al: *Minimally Invasive Surgical management of primary venous Ulcer vs. Compression* *Eur J vasc Endovasc Surg* 00,16 (2003)
- Bellmunt-Montoya S, Escribano JM, Dilme J, Martinez-Zapata MJ. *CHIVA Method for the treatment of chronic venous insufficiency.* *Cochrane Database of Systematic Reviews* 2012, Issue 2. Art. No.: CD009648.
- Chan, C.-Y.a, Chen, T.-C.b, Hsieh, Y.-K.a, Huang, J.-H.c. *Retrospective comparison of clinical outcomes between endovenous laser and saphenous vein-sparing surgery for treatment of varicose vein, World Journal of Surgery,* 2011; 35(7):1679-1686
- Hemodynamic classification and CHIVA treatment of varicose veins in lower extremities (VVLE) Department of Vascular Surgery, Xinhua Hospital, China (<http://www.ijcem.com/files/ijcem0016552.pdf>)
- Marta Zmudzinski, BSc, Pierre Malo, MD, FCSP, Christine Hall, MD, FRCPC, Allen Hayashi, MD, FRCSC *HIVA, A prospective study of a vein sparing technique for the management of varicose vein diseases* ([http://www.americanjournalofsurgery.com/article/S0002-9610\(16\)31046-7/fulltext](http://www.americanjournalofsurgery.com/article/S0002-9610(16)31046-7/fulltext)) *The American Journal of Surgery* - 21 Mars 2017.
- Mark D. Iafrati And Thomas F. O'donnell Jr. *Rutherford's Vascular Surgery And Endovascular Therapy, Ninth Edition - Saphenous-Sparing Operations 2030; Section 23, Chapter 154* 2018 ISBN=978-0-323-42791-3
- Shouten R, Mollen RM, Kuijpers HC., "A comparison between cryosurgery and conventional stripping in varicose vein surgery: perioperative features and complications" *Annals of Vascular Surgery.*, 2006; 20 (3): 306-11.
- Pak, L. K. et al. "Veins & Lymphatics," in *Lange's Current Surgical Diagnosis & Treatment*, 11th ed., McGraw-Hill.
- Tisi, Paul V; Beverley, Catherine; Rees, Angie., Tisi, Paul V, ed. "Cochrane Database of Systematic Reviews, Chapter: Injection sclerotherapy for varicose veins" *Cochrane Database Syst Rev* (4): 2006; CD001732. (<https://www.ncbi.nlm.nih.gov/pubmed/17054141>).
- Thibault, Paul., *Sclerotherapy and Ultrasound-Guided Sclerotherapy, The Vein Book*, John J. Bergan (ed.), 2007.
- Padbury A, Benveniste GL., "Foam echosclerotherapy of the small saphenous vein". *Australian and New Zealand Journal of Phlebology.*, December 2004; 8 (1).
- Finkelmeier, William R., "Sclerotherapy", Ch. 12 in *ACS Surgery: Principles & Practice*, WebMD, ISBN 0-9748327-4-X., 2004.

33. Scurr JR, Fisher RK, Wallace SB, "Anaphylaxis Following Foam Sclerotherapy: A Life Threatening Complication of Non Invasive Treatment For Varicose Veins". *EJVES Extra.*, 2007; 13 (6):87-89.
34. Malskat WS, Stokbroekx MA, van der Geld CW, Nijsten TE, van den Bos RR, "Temperature Profiles of 980- and 1,470-nm Endovenous Laser Ablation, Endovenous Radiofrequency Ablation and Endovenous Steam Ablation". *Lasers Med Sci.*, March 29, 2014; 29: 423-9.
35. Medical Services Advisory Committee, E LA for varicose veins ([http://www.msac.gov.au/internet/msac/publishing.nsf/Content/2E0BACBB8704139ACA25745E001C2F21/\\$File/1113report.pdf](http://www.msac.gov.au/internet/msac/publishing.nsf/Content/2E0BACBB8704139ACA25745E001C2F21/$File/1113report.pdf))MSAC application 1113, Dept of Health and Ageing, Commonwealth of Australia, 2008.
36. Elmore FA, Lackey D., "Effectiveness of ELA in eliminating superficial venous reflux" *P.hlebology.*, 2008; 23 (1):21-23.
37. Van den Bos, RR; Malskat, WS; De Maeseneer, MG; De Roos, KP; Groeneweg, DA; Kockaert, MA; Neumann, HA; Nijsten, T (2014-06-30). "Randomized clinical trial of endovenous laser ablation versus steam ablation (LAST trial) for great saphenous veins". *Br J Surg.* 101: 1077-1083.
38. Milleret, René., "Obliteration of varicose veins with superheated steam" *P. hlebology.*, 2011; 19 (4):174-181.

Journal of Drug Delivery & Therapeutics



JDDT