

Available online on 15.07.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Evaluation of Hepatoprotective Activity of Hydroalcoholic Extract of Onion Peels Containing Protocatechuic Acid

Mrinalini C. Damle¹, Sachin V. Tembhurne², Swapnil R. Todkar²¹ Department of Pharmaceutical Quality Assurance, All India Shri Shivaji Memorial Society's College of Pharmacy, Kennedy Road, Near RTO, Pune-411001.² Department of Pharmacology, All India Shri Shivaji Memorial Society's College of Pharmacy, Kennedy Road, Near RTO, Pune-411001.

ABSTRACT

India is the second largest country for cultivation of onion. 19.90% of onion is cultivated in India. Onion that is *Allium cepa* family Amaryllidaceae used in daily diet for taste. Onions also have number of medicinal properties, such as hepatoprotection, anticancer, antifungal, antioxidant, antiulcer, anti-aging, anti-inflammatory, anticancer etc.^[1-2] The dried scales or peels of onion also have the same medicinal constituents with same activity as raw onions. The onion peels contains flavonoids, such as anthocyanins, flavones (quercetin and its derivatives), ferulic, Gallic, protocatechuic acids, sulphur, vitamins etc. In the present study determination of protocatechuic acid in onion peels extract has been performed using HPTLC. HPTLC separation was carried out on Merck TLC aluminium sheets pre-coated with silica gel 60F₂₅₄ using Toluene: Ethyl acetate: Formic acid (6: 6: 1.2 v/v/v) as mobile phase.^[3-11] Quantitative analysis was carried out in the absorbance mode at 258 nm. Hydroalcoholic extract was tested for hepatoprotective activity in wistar rats (either sex) by using CCl₄ as hepatotoxicity inducing agent and Silamyrin as standard. Hydroalcoholic extract shows hepatoprotective activity as indicated by decrease in the level of SGOT, SGPT, total protein, bilirubin in which hepatotoxicity was induced by CCl₄ intraperitoneal injection route to animal, from the result it may be concluded that onion peels extract may be used for hepatoprotective activity.

Keywords: Onion peel extract, Protocatechuic acid, HPTLC, hepatoprotection.

Article Info: Received 14 May 2019; Review Completed 12 June 2019; Accepted 19 June 2019; Available online 15 July 2019



Cite this article as:

Damle MC Tembhurne SV, Todkar SR, Evaluation of Hepatoprotective Activity of Hydroalcoholic Extract of Onion Peels Containing Protocatechuic Acid, Journal of Drug Delivery and Therapeutics. 2019; 9(4):82-88
<http://dx.doi.org/10.22270/jddt.v9i4.2982>

*Address for Correspondence:

Dr. Mrinalini C. Damle, Professor, Department of Quality Assurance, AISSMS College of Pharmacy, (Affiliated to Savitribai Phule Pune University), Maharashtra, India- 411001.

INTRODUCTION

There is an increasing interest in herbal remedies because of their effectiveness, less side effects in clinical experiments and relatively low cost. Protocatechuic acid (PCA) is widely distributed and present in most edible plants used in medicine. It is also a very common compound present in human diet, present in bran and brown rice (*Oryza sativa* L.) and onion (*Allium cepa* L.). Onion peels are the external part of onion bulb and having several folds, it is thin, light weight, strong often translucent paper. Onion peel contains numbers of phytoconstituent such as sulphur, quercetin, protocatechuic acid, calcium, flavonoids, phenolic acid, etc.^{1,5}.

PCA has been reported to possess antioxidant, antibacterial, anticancer, antiulcer, antidiabetic, anti-ageing, antifibrotic, antiviral, anti-inflammatory, analgesic, antiatherosclerotic, cardio protective, hepatoprotective, neurological and nephroprotective activities. As per literature search there is

no method for determination of protocatechuic acid in onion peel extract and evaluation of hepatoprotective activity in rats⁶⁻⁹.

Liver is one of the largest organs in the human body and chief site for intense metabolism and excretion. Liver diseases are one of the major health problems in the world. These are caused by toxic chemicals, autoimmune disorders, infections and excess consumption of alcohol. The hepatotoxic chemicals can induce lipid peroxidation and oxidative damages. It is involved in almost all the biochemical pathways to growth, fight against the disease, nutrient supply, energy provision and reproduction.¹⁰⁻¹⁴

The hydroalcoholic extract of onion peels contains PCA as active constituent. There are many literature records indicating hepatoprotective activity of PCA. Despite of there is no report available on action of onion peel extract. For that reason the present study changed into undertaken to

evaluate the hepatoprotective activity of hydroalcoholic extract of onion peels in CCl₄ induced hepatotoxicity in wistar rats.

MATERIAL AND METHODS:

Procurement of drug: PCA was procured from Sisco Research Laboratories. Pvt Ltd.

Collection and Identification of plant Material:

Onion peels were collected from Talegaon Dhamdhare, Pune and authenticated by botanical survey of India, Pune (sample no. ST: 01)

Extraction method ¹²⁻¹³

Peels were separated from onion bulb. Onion peels was sundried and crushed to obtain powder. This powder was

weighed, added 70% ethanol with same quantity of 0.1M NaOH solution and kept into glass bottle for cold maceration. This mixture was placed for 72 hrs with intermittent shaking. After 72hrs the macerates were removed from bottle, filtered through whattman filter paper and filtrate was concentrated on waterbath.

Selection of Analytical Wavelength

For the selection of the analytical wavelength, the Protocatechuic acid solution was prepared by dissolving 10 mg in 10 ml (1000µg/ml) of methanol and further diluted with methanol to produce (10 µg/ ml) its absorbance was taken on UV visible spectrophotometer in range of 200-400 nm. The compound absorb at 258nm which is selected as an analytical wavelength.




Figure 1: UV Spectrum of protocatechuic acid(10µg/ml)

HPTLC analysis¹³⁻¹⁶

The chromatographic separation was carried on aluminium plates precoated with silica gel 60F₂₅₄ in (10 cm × 10 cm with 250 µm layer thickness). Sample was applied on the plate with band width of 6 mm width using Camag 100 µL sample syringe (Hamilton, Switzerland) with Linomat 5 applicator (Camag, Switzerland). A solution of 100µg/ ml in methanol of PCA was prepared and then spotted using Camag Linomat V sample applicator. The optimised mobile phase comprises of Toluene: Ethyl acetate: Formic acid (6:6:1.2, v/v/v). CAMAG twin trough glass chamber was used for linear ascending development of TLC plate with 15 min saturation time for

mobile phase, migration distance was 90 mm. Densitometric scanning was performed using Camag TLC scanner 3 at 258 nm, operated by win CATS software (version 1.4.3, Camag), slit dimensions were 4.00 × 0.45 mm and deuterium lamp was used as a radiation source.

The identification of the protocatechuic acid in sample densitogram was confirmed by comparing the R_f value with that obtained from pure marker. The identity of protocatechuic acid in the hydro alcoholic extract was confirmed by comparing the UV absorption spectra with that of standard using a CAMAG TLC scanner 3.




Figure 2: HPTLC densitogram of protocatechuic acid (R_f=0.52)




Figure3: Typical HPTLC densitogram obtained from extract showing peak at Rf = 0.52




Fig.4. UV spectrum match of protocatechuic acid marker and onion peel extract at Rf. 0.52

Regression data obtained from calibration curve demonstrated excellent linear relationship over 100-500ng/ band concentration range. The linear regression equation was found to be

$$y = 12.436x + 176.82 \text{ having correlation coefficient } 0.9949.$$




Fig.5. 3D densitogram of protocatechuic acid (track 1 methanol, track 2 to 6 std. protocatechuic acid 100-500ng/band, track 7-8 and 9-10 onion peel extract 2000 and 4000ng/band)




Fig.no. 6: Calibration curve of PCA (100-500ng/band)

With the help of above statistical data, the content of protocatechuic acid was determined in the hydroalcoholic onion peel plant extract which was found to be 6.5%. The HPTLC data obtained for the quantification of protocatechuic acid is summarized in Table 1. HPTLC analysis was carried out to determine the content and to quantify the protocatechuic acid present in onion peel extract.

Table no.1: HPTLC data for Protocatechuic acid

Sr. No	Validation Parameters	Protocatechuic acid
1	Linearity Equation (r^2) Range	$y = 12.43x + 176.8$ $R^2 = 0.994$ 100-500 ng/ band
2	Precision a) Intra day b) Interday	(% RSD) 1.44 1.11 0.85 1.13
3	Accuracy	% Recovery
	80%	99.84
	100	100.22
	120%	99.80
4	Limit of Detection	3.74 ng/band
5	Limit of Quantitation	11.35ng/band
6	Specificity	Specific
7	Robustness	Robust
8	Solution stability	Stable

Hepatoprotective activity^[15-24]

Wister rats of either sex weighing around 150 to 200 gm housed in standard conditions of temperature ($22 \pm 2^\circ\text{C}$), relative humidity ($55 \pm 5\%$), and light (12hrs light/dark cycles) were used. They were feed with standard pellet diet and water ad libitum. The experimental protocol was approved by Institutional Animal Ethical Committee as per the guidelines of CPCSEA, Government of India (protocol no. CPCSEA/IAEC/QA-02/01-2K18)

Experimental design

A total of 30 rats were divided into 5 groups of 6 rats each. Group I served as normal control and received only the vehicle, Group II received CCl_4 1 mL/kg (1:1 of CCl_4 in olive oil) i.p. once daily for 7 days. Group III received CCl_4 1 mL/kg (1:1 of CCl_4 in olive oil) i.p. and silymarin as standard drug 100 mg/kg orally (p.o.) for 7 days. Groups IV, V, were administered hydro alcoholic onion peels extract 300 and 600 mg/kg body weight p.o.^[15], as low and high dosage respectively and dose of 1 mL/kg i.p. of CCl_4 (1:1 of CCl_4 in olive oil) for 7 days.

All rats were sacrificed by cervical dislocation 24 hrs after the last treatment. Just before sacrifice, blood was collected from the retro-orbital sinus plexus under mild ether anaesthesia. Collected blood was allowed to clot and serum was separated at 3500 rpm for 15 min to carry out further biochemical investigations such as SGPT, SGOT, total protein and billirubin. Liver was dissected out and used for histopathological examination.

Histopathological examination of Liver

After decapitation, liver from each rat was collected, rinsed in 0.9% saline and fixed in 10% formalin. After fixation, these tissues were trimmed and processed. Tissue processing was done to dehydrate in ascending grade of alcohol, clearing in xylene and embedded in paraffin wax. Paraffin wax embedded tissue were sections at 3-5 μm thickness with rotary microtome. Slides were stained with hematoxylin and eosin stain. The prepared slides were examined under microscope.

Statistical analysis:

Results were expressed as Mean \pm S.E.M. The data was analysed by using one way ANOVA (analysis of variance) followed by Tukey-Kramer's test and $p < 0.05$ was considered as statistically significant.

RESULTS

In the present investigations rats treated with chronic dose of CCl_4 developed significant hepatic damage which was observed through a significant increase in the concentration of SGOT, SGPT and total billirubin. While there was substantial reduction in the content of the total protein. The treatment with hydro alcoholic extract of onion peels in CCl_4 treated group result in decreased in hepatic injury to a considerable extent which was reflected by the ability of the extract to lower the elevated serum enzyme levels. The increased level of SGOT, SGPT, and total billirubin in serum is indicative of cellular leakage and loss of functional integrity of cell membrane in liver. With respect to that extract mediated reduction in the level of SGOT, SGPT and total billirubin towards the respective normal values is an indication of stabilization of plasma membrane as well as repair of hepatic tissue damage caused by CCl_4 . This effect is in agreement with commonly accepted view that serum levels of transaminase return to normal level with healing of hepatic parenchyma and the regeneration of hepatocyte.






Fig.no.7: Effect of hydroalcoholic extract of *onion peels* on SGOT, SGPT, Toal protein and Total Bilirubin. Values are considered as Mean \pm S.E.M. (n = 6), one way ANOVA followed by the Tukey- Kramer's Test.

Histopathological Examination


Microscopic examination of liver treated with CCL₄ showed mild hepatocellular necrosis with infiltration of inflammatory cells when compared with control group.

Severity of hepatocellular necrosis with infiltration of inflammatory cells was decreased in animals treated with Standard and Extract given at 300 mg/kg body weight and animals treated with Extract at 600 mg/kg did not show any hepatic changes when compared with control group.







CCl₄; Liver; Male: Showing hepatocellular necrosis with infiltration of inflammatory cells (Mild). 40 X, H & E




CCl₄ ; Liver; Female: Showing hepatocellular necrosis with infiltration of inflammatory cells (Mild). 40 X, H & E




Standard (Silymarin); Liver; Male: Showing hepatocellular necrosis with infiltration of inflammatory cells (Minimal). 40 X, H & E




Standard (Silymarin); Liver; Female: Showing hepatocellular necrosis with infiltration of inflammatory cells (Minimal). 40 X, H & E




Low Dose; Liver; Male: Showing hepatocellular necrosis with infiltration of inflammatory cells (Minimal). 40 X, H & E



Low Dose; Liver; Female: Showing hepatocellular necrosis with infiltration of inflammatory cells (Minimal). 40 X, H & E



High Dose; Liver; Male: Showing normal hepatic parenchyma. 40 X, H & E



High Dose; Liver; Female: Showing normal hepatic parenchyma. 40 X, H & E

DISCUSSION:

CCl₄ is a hepatotoxin used extensively for inducing liver injury in various experimental models to elucidate the mechanisms underlying hepatotoxicity. CCl₄ mediated

hepatotoxicity is developed from the biotransformation of CCl₄ by cytochrome P450 2E1 to the trichloromethyl free radical (•CCl₃). Hepatocyte damage is characterised by different hepatic marker enzymes (SGPT, SGOT) and the levels of total bilirubin and total protein. When liver cells

are damaged, these enzymes leak into the bloodstream from liver tissue and produce markedly elevated serum levels. Both SGOT and SGPT are associated with liver parenchymal cells. SGPT is found predominantly in the liver with negligible quantities found in heart, kidneys and skeletal muscles, whereas SGOT is found in liver, cardiac muscles, skeletal muscles, brain, kidney and red blood cells. Thus SGPT is a more specific indicator of liver intoxication as levels of SGOT may also be increased in diseases affecting other organs.

On the other hand, bilirubin levels are related to the functions of hepatic cell. Elevation in level is due to increased synthesis, in presence of increased biliary pressure. Our experiment showed that rats intoxicated with CCl₄ develop a significant liver necrosis which was evidenced by increased activities of hepatic marker enzymes (SGPT, SGOT) and the levels of total bilirubin, whereas levels of total protein were decreased due to liver injury. The result of this study showed that after administration with *onion peels extract* the activities of the serum marker enzymes SGPT, SGOT and the levels of total bilirubin and total protein were restored to normal level, thus indicating *onion peels extract* preserved the structural integrity of hepatocellular components and protected the liver from the harmful effect of this hepatotoxin.

The above inferences were further confirmed by histopathological studies. The result of the histopathology showed that CCl₄ administration caused severe acute liver damage in rats which is characterised by hepatic cell necrosis, ballooning degeneration, fatty changes or inflammatory cell infiltration and other histological manifestations, which were consistent with previous findings. After treatment with *onion peels extract* hepatic injury caused by CCl₄ administration was significantly prevented and the structure of hepatocytes was almost restored to normal. The results of the different biochemical parameters and histopathological findings are co-related well with each other, which indicated that *onion peels extract* possesses significant hepatoprotective effect in CCl₄-intoxicated liver injury in rats.

CONCLUSION:

The present study demonstrated that the hydroalcoholic extract obtained from onion peels contains Protocatechuic acid as active phytoconstituent and has significant hepatoprotective activities against CCl₄-induced hepatotoxicity in Wistar albino rats. The hepatoprotective activity of onion peels extract due to the presence of protocatechuic acid in the extract which was determined by HPTLC method. Therefore, the present study shows that there is a prospective future in the use of Onion peel as a source of natural medicine for curing hepatic diseases.

ACKNOWLEDGEMENT

Authors are thankful to the Principal and the management of AISSMS College of Pharmacy Pune, for providing the necessary facilities for research work.

REFERENCES

1. https://en.wikipedia.org/wiki/Protocatechuic_acid (Accessed on 12-07-2018).
2. <https://www.drugbank.ca/drugs/DB03946> (Accessed on 13-07-2018).
3. Sahil Kakkar and Souravh Bais, A Review on Protocatechuic Acid and Its Pharmacological Potential, International scholarly Research Notices Pharmacology journal, Volume 2014, page no. 1-9.
4. Manek Ravi, Patel N. M.2, Determination of Protocatechuic Acid by HPTLC Method in Amomum subulatum Roxb. Fruit Extracts PHCOG Journal, August 2009, page no. 130 to 137.
5. Abida kalsoom khan1, pharmacological activities of protocatechuic acid, Acta Poloniae Pharmaceutica n Drug Research, 2015, Vol. 72 No. 4 pp. 643-650.
6. Pandey Shivanand, R. Mahalaxmi, Identification and determination of protocatechuic acid present in greater cardamom fruit extracts by HPTLC technique, International Journal of Pharmaceutical Sciences Review and Research Volume 1, issue 1, page no. 27 to 32.
7. Rana Arslan, The possible mechanisms of protocatechuic acid induced central analgesia, Saudi pharmaceutical journal 2018, page no. 541-545.
8. Brahma N. Singh polyphenolics from various extract/fraction of red onion (Allium Cepa) peel with potent antioxidant and antimutagenic activity, journal of Food and Chemical Toxicology 47 (2009) 1161-1167.
9. Jungmi Kirn cytotoxic and anti-inflammatory effect of onion peel extract on lipopolysaccharide stimulated human colon carcinoma cells, journal of food and chemical toxicology 2003, page no. 199-204.
10. Srinivasan, K. Plant foods in the management of diabetes mellitus: spices as beneficial antidiabetic food adjuncts. *Int. J. Food Sci. Nutr.* 2005;56(6):399-414. 16361181
11. Ozougwu, Jervas C, Physiology of the liver, International Journal of Research in Pharmacy and Biosciences Volume 4, Issue 8, 2017, PP 13-24
12. <http://www.berkem.com/en/expertise-en/plant-extraction> (Accessed on 22 April 19)
13. http://agritech.tnau.ac.in/horticulture/extraction_techniques%20_medicinal_plants.pdf (Accessed on 22 April 19)
14. <https://www.webmd.com/digestive-disorders/picture-of-the-liver#1> (Accessed on 22 April 19)
15. Salami H.A, John A.I. et al., The effect of aqueous preparation of Allium Cepa (onion) and Allium Sativa (garlic) on erythrocyte osmotic fragility in wistar rats: in vivo and in vitro studies, Nigerian journal of Physiological Science, volume 27. June 2012, page no. 029-034.
16. Butura A. Drug and Alcohol Induced Hepatotoxicity. 2008 Ph. D Thesis Department of Physiology and Pharmacology Karolinska Institutet, Stockholm, Sweden. 55 pp.
17. Kmiec Z. Co-operation of liver cells in health and diseases. *Advances in Anatomy, Embryology and Cell Biology* 2001, 161: 3 - 12.
18. Rang and dales book of pharmacology, 8th edition 2016, page no. 456-460
19. Kavita Gulati, Mohd Rafi Reshi et al., Hepatotoxicity: Its Mechanisms, Experimental Evaluation and Protective Strategies, *American Journal of Pharmacology*, 2018; 1(1): 1004(article no)
20. Rakesh Sagar, Amrita Bhajji, et al., A comprehensive review on herbal drugs for hepatoprotection of 21st Century, *International Journal of Nutrition, Pharmacology, Neurological Diseases* | October-December 2014 | Vol 4| Issue 4, pg no 191-197
21. G. Dineshkumar hepatoprotective activity of leaves extract of eichhornia crassipes against ccl4 induced hepatotoxicity albino rats, *International Journal of Pure and Applied Zoology*, Vol. 1, Issue 3, pp: 209-212, 2013.
22. Prashant Tiwari Evaluation of hepatoprotective activity of alcoholic and aqueous extracts of Selaginella lepidophylla, *Asian Pacific Journal of Tropical Disease*, page no. S81 to S86.
23. Sumanta Mondala, Hepatoprotective activity of Macrothelypteris torresiana (Gaudich.) aerial parts against CCl₄-induced hepatotoxicity in rodents and analysis of polyphenolic compounds by HPTLC, *Journal of Pharmaceutical Analysis*, page no. 181-189.
24. ZHANG Zhi-Feng, Hepatoprotective activity of Gentiana veitchiorum Hemsl. against carbon tetrachloride-induced hepatotoxicity in mice, *Chinese Journal of Natural Medicines* 2014, section 12(7) page no.0488-0494.
25. Kalpana Patil, Hepatoprotective activity of Mentha arvensis Linn. leaves against CCl₄ induced liver damage in rats, *Asian Pacific Journal of Tropical Disease*, page no. 5223-5226