

Available online on 15.07.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Assessment of the In-vitro antioxidant activity of the alcoholic extract/fractions of *Bombax malabaricum* DC.

Singh Charanjeet ^{1*}, Ahuja Dharmendra², Mehta SC²¹ Research Scholar, Faculty of Pharmaceutical Science, Jayoti Vidyapeeth Women's University, Jaipur, Rajasthan, India.² Professor, Faculty of Pharmaceutical Science, Jayoti Vidyapeeth Women's University, Jaipur, Rajasthan, India.

ABSTRACT

Various parts of *Bombax malabaricum* are used in traditional Indian medicine for the treatment of inflammatory diseases, diarrhea, fever, acute swelling and diuretic. Current research on *Bombax malabaricum* DC is aimed at finding of natural plants derived from antioxidants that provide sustenance through additional components or synergistic training as the plant's cellular reinforcements are necessary to prevent the movement of interfaced free radical problems. The antioxidant activity in *Bombax malabaricum* DC can be detected by alcoholic extract fraction using unique laboratory models. These include free radical activity of 1,1-Diphenyl-2-picryl-hydrazil (DPPH) assay, ABTS assay, total antioxidant capacity, and o-phenanthroline assay/Iron chelating assay. The plant contains a large number of phenolic and flavonoid compounds. The plant shows great antioxidant activity. The thought was taught to investigate the use of the *Bombax malabaricum*

Keywords: *Bombax malabaricum*, Antioxidant, DPPH, ABTS, O-phenanthroline.**Article Info:** Received 13 May 2019; Review Completed 16 June 2019; Accepted 23 June 2019; Available online 15 July 2019**Cite this article as:**Singh CJ, Ahuja D, Mehta SC, Assessment of the In-vitro antioxidant activity of the alcoholic extract/fractions of *Bombax malabaricum* DC, Journal of Drug Delivery and Therapeutics. 2019; 9(4):135-140
<http://dx.doi.org/10.22270/jddt.v9i4.2989>***Address for Correspondence:**

Charanjeet Singh, Research Scholar, Faculty of Pharmaceutical Science, Jayoti Vidyapeeth Women's University, Jaipur

INTRODUCTION:

Responsive oxygen species (ROS) such as Superoxide (O_2^-), hydrogenradicals (OH^\cdot), and hydrogen peroxide (H_2O_2) are considered as significant elements causing numerous disorders like cardiovascular, diabetes, swelling, malignant growth and neurogenerative diseases^{1,2}. ROS are corrupted to non-receptive structures by Enzymatic and Non-Enzymatic resistance components. Free radicals respond with known biological molecule and harm protein, breakdown of DNA strands and starts peroxidation of different particles. Antioxidant agents go about as a noteworthy guard against radical-interceded poisonous quality, by securing the harms brought about by free radicals³. Antioxidative segments of common source have pulled in unique intrigue since they can shield human body from free radicals. Enzymatic cancer prevention agents incorporate essential catalysts like Superoxide Dismutase, Catalase, Glutathione peroxidase and auxiliary proteins incorporate Glutathione reductase etc⁴.

Non-Enzymatic antioxidant agents incorporate either water-solvent (Vitamin C and phenolic agents) or lipid-dissolvable (vitamin E and carotenoids) agents which go about as protection against oxidative stress^{5,6}. Anyway antioxidant

agent like Phenols, Flavonoids scavenge these free radicals and shield the framework from the Oxidative mechanism.

The plants of antiquity are used for traditional treatment therapies and the literature explains the use of different parts of the plant in different dosage forms for diseases of different origins, including the infectious nature^{7,8}. Products of natural plant origin are known for their strong medicinal activities. These plants are the sources of medicines from the many years. All plants contain active chemical constituents and groups of secondary plant metabolites such as alkaloids, steroids, glycosides etc.^{9,10}. The Indian subcontinent is blessed by the nature and strength of traditional local healers, helping society maximize the impact of nature⁷. Many antioxidant compounds from plant sources that occur naturally as free radicals or oxygen and active nitrogen. Flavonoids are the most important group of natural antioxidants. Its biological effects are mainly related to the cleaning of free radicals or ionic minerals. Metal ions participate directly in the formation of interactive species¹¹.

Bombax malabaricum DC belongs from family Bombacaceae, commonly known as salmali. It is widely distributed throughout India, in the forest up to about 1500 meters high, since it was raised on the farm and found as rice in Malaysia.

In India, it is distributed from Rajasthan, the southern wing of sarakallu and in the adjacent area of chittoor district, Andhra Pradesh. The leaves are large, scattered, saturated, digitates, leaflets, insoles, 3-7 entire. The flowers are red, many, appear, when the tree is stripped of the stems arranged largely in beautiful bundles of 9-12 each and an inner package of 15. Fruit capsule, exposing 5 leather or woody valves. The seeds are smooth, black or gray, set in a long white wool. Gray or brown bark covered with hardness, sharpness and conical prickles. It is clear that the gums are brownish in color and gradually become dark brown. Various parts of the *Bombax malabaricum* such as roots, leaves, seed, stem bark, flowers, fruits and gums are documented to have healing properties in ethnographic investigations conducted by ethnologists and in traditional medicine systems such as Ayurveda^{12,13}.

This plant is widely used in traditional Chinese and Indian medicine for the treatment of inflammatory diseases, diarrhea, fever, chronic swelling, acute illness and as diuretic^{14,15}. Earlier phytochemical researches of this plant have led to in the separation of naphthol and naphthoquinone derivatives, anthocyanins¹⁶, flavonoids¹⁵, several sesquiterpenoids¹⁷, and polysaccharide¹⁸. Though, the viable activities or pharmacological properties of *B. malabaricum* explained, activity information antioxidant fraction of the extract is still limited. Bark utilized for astringent, diuretics, agonists, diuretics, treatment of cysts, wounds and other skin disorders. The leaves are anti-inflammatory, the roots are aphrodisiac, anti-diarrhea cure. The flowers are diuretics and laxatives and the gums are used in hemodialysis. seeds are used in gonorrhoea¹⁹. decoction bark traditionally used abroad in the inflammatory process, and seal secondary infections, the wounded skin healing in paste form and the leaves of this plant is grounded and the mixed milk is provided for choking and inflammation. Despite the traditional use of this plant, no scientific report has focused on the biological activity of the *Bombax malabaricum*. The chemical prevention of cancer is one of the main areas investigated in the last years²⁰.

The consumption of many vegetables is associated with a reduced risk of degenerative diseases, which depend on oxidative stress such as atherosclerosis, tumors, diabetes and Alzheimer's disease^{21,22}. Good attention is given to naturally occurring antioxidants, which can play important functions that inhibit both free radicals and oxidative chain reactions within tissues and membranes²³. Consequently, the screening of plant materials based on potential antioxidants can have a central task to identify potential extracts or fractions to remove all the roots and initiate a chain reaction or seal catalysts for oxidizing reactions, such as some minerals²⁴. From the point of view of its high antioxidant strength, it is recommended to consume high levels of many plants, so analyzing antioxidant activity in extracts and fractions is an important step before detoxifying the antioxidant and it²⁵. In the present study an attempt was made to estimate the impact of antioxidants effect of extract fraction of the *Bombax malabaricum* against by various in vitro models.

MATERIALS AND METHODS

Collection of Plant material

The plant material (*Bombax malabaricum*) was collected from smriti van, Jaipur (Jaipur, Rajasthan, India) in the month of September and authenticated by Herbarium department, University of Rajasthan, Jaipur, Rajasthan, India. The plant was deposited in the herbarium at Department of Botany (University of Rajasthan, Jaipur, Rajasthan, India).

Preparation of extract

Alcoholic extract: - The coarsely powdered (1000 g) oven-dried *Bombax malabaricum* was extracted with alcohol by using soxhlet apparatus for 20 h. After completion of extraction, the solvent was removed by distillation and concentrated. The yield obtained was 20.54 % w/v.

Fractionation of crud extract

Fractionation of alcoholic extract Completely dried ethanolic extract was suspended in distilled water and extracted successively and exhaustively with solvents of increasing polarity like petroleum ether, dichloromethane, n-butanol, ethyl acetate. Each fraction was concentrated using rotary evaporator (Rotavapor, R-210, BUCHI Laborte, Switzerland) and stored in vacuum desiccator. The percentage yield of various extracts/fractions was ethanol (BME) [26.31% w/v], petroleum ether (BMP) [2.51 % w/v], dichloromethane (BMD) [8.14 % w/v] and n-butanol (BMB) [22.81 % w/v]

Aqueous extract

1000 g of the coarsely powdered root of *Bombax malabaricum* was extracted by chloroform water (1:99) by cold maceration process for 7 days. After completion of extraction, the marc was filtered through muslin cloth and concentrated. The yield of BMA was obtained 21.56 %.

Qualitative determination of the chemical constituents

Presence of alkaloids, anthraquinones, cardiac glycosides, coumarins, flavonoids, saponins, phlobatannins, tannins and terpenoids in the extract and various fractions was confirmed individually by following standard procedures.

Test for alkaloids

Mixture of methanol extract of *Bombax malabaricum* extract/fraction and its various derived fractions (0.4 g) in 8 ml of 1% HCl was warmed on water bath. After filtration 2 ml filtrate from the extract and each fraction was allowed to react with few drops of potassium mercuric iodide and with potassium bismuth, separately. Turbidity or precipitation formation was considered as a confirmation for presence of alkaloids.

Test for saponins

The criterion of oil emulsion formation of saponins was used for the screening of saponins. Briefly, extract and various fractions (20 mg) suspended in 20 ml of distilled water and boiled for 5 min. In 10 ml of the above filtrate 5 ml of distilled water was added and mixed well to develop the froth. Development of emulsion after mixing the froth with olive oil confirmed the existence of saponins.

Test for terpenoids

Briefly, 2 ml of chloroform was mixed with 5 ml (1 mg/ml) of each sample in a test tube then 3 ml of concentrated H₂SO₄ was added to develop the color. Exhibition of reddish-brown coloration at the interface confirmed the presence of terpenoids.

Test for anthraquinones

To a volume of 6 ml of 1% HCl, 200 mg of each sample was added separately and boiled. Benzene (5 ml) was mixed with the filtrate and after separation of benzene layer 2 ml of 10% ammonia solution was lowered. Development of pink, violet or red color in the ammoniacal phase indicated the existence of anthraquinones²⁶.

Cardiac glycosides determination

An aliquot of 5 ml of ethanol extract of *Bombax malabaricum* and its various fractions (10 mg/ml in methanol) were added in the sequence of glacial acetic acid (2 ml) and FeCl₃ solution (one drop). Concentrated H₂SO₄ (1 ml) was added and the formation of brown ring at the interface confirmed the presence of cardiac glycosides.

Test for coumarins

In a vial having 300 mg/ml of the extract and each fraction was plugged with filter paper dipped in 1 N NaOH and boiled in a boiling water bath for few minutes. Yellow fluorescence of filter paper under UV light confirmed the presence of coumarins.

Test for phlobatannins

An amount of 80 mg of the extract and various fractions was boiled in 1% HCl. Development of red precipitate indicated the existence of phlobatannins²⁷.

Test for flavonoids

Mixture of methanol extract and various fractions of *Bombax malabaricum* were prepared by adding 50 mg of each sample to 100 ml of distilled water and filtered. An aliquot of 5 ml of dilute ammonia solution was mixed with 10 ml of the filtrate. Appearance of yellow coloration by addition of few drops of concentrated sulfuric acid indicated the presence of flavonoid.

Test for tannins

A mixture was prepared by mixing 50 mg of methanol extract and each fraction in 20 ml of distilled water and boiled. Appearance of brownish green or blue-black coloration after mixing few drops of 0.1% FeCl₃ confirmed the existence of tannins²⁸.

Determination of in-vitro scavenging activities³⁴:

DPPH radical scavenging activity:

The free radical scavenging activity of the extract and fractions was evaluated using the stable DPPH free radical. One mL of 0.1 mM DPPH solution in methanol was added to 1.0 mL of standard and extract/fractions solution at different concentrations. The mixture was incubated for 20 min and the absorbance recorded at 517 nm. Ascorbic acid was used as positive control. DPPH radical scavenging activity was calculated using the formula:

Percent scavenging = $\frac{(A_0 - A_t)}{A_0} \times 100$; where A₀ = Absorbance of control (without extract) and A_t = Absorbance of sample. All determinations were carried out in triplicate²⁹.

ABTS radical scavenging activity:

ABTS free radical was generated by reacting 7 mM ABTS solution with 2.45 mM potassium per sulphate. The mixture was allowed to stand for 15 h in dark at room temperature. ABTS solution was diluted with methanol to obtain the absorbance of 0.7 ± 0.2 units at 750 nm. The standard/extract solutions were prepared at different concentrations in methanol and 20 µL of test solutions were added to 180 µL of ABTS free radical solution. The absorbance was measured after 20 minutes incubation at 750 nm. Ascorbic acid was used as positive control. The ABTS free radical scavenging activity was calculated using the formula:

$$\text{Percent scavenging} = \frac{(A_0 - A_t)}{A_0} \times 100;$$

where A₀ = Absorbance of control (without extract) and A_t = Absorbance of sample³⁰. All the tests were performed in triplicate.

Total antioxidant capacity:

The total antioxidant capacities of the extract and fractions of *Bombax malabaricum* were determined using phosphomolybdenum method. Briefly, 0.1 mL of standard/extract solution was mixed with 0.3 mL of reagent solution (0.6 M sulfuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate) and incubated at 95°C for 90 min. The mixture was cooled down to room temperature and absorbance recorded at 695 nm. The blank solution contained all the reagents except the test sample. Ascorbic acid was used to plot the standard curve. The results were expressed as ascorbic acid equivalents³¹. All the tests were performed in triplicate.

o-Phenanthroline assay/Iron chelating activity:

The 1, 10-Phenanthroline-iron (III) reagent was prepared by mixing 0.198 g of 1, 10-phenanthroline monohydrate, 2 mL of 1 M hydrochloric acid and 0.16 g of ferric ammonium sulphate in 100 mL water. Briefly, 0.2 mL standard/extracts were mixed with 0.2 mL 1, 10-phenanthroline-iron (III) reagent, 0.6 mL methanol and 4 mL water. The solutions were incubated at 50°C for 30 min and absorbance read at 510 nm. Ascorbic acid was used as positive control. A higher absorbance indicated higher iron chelating activity

Percentage scavenging was calculated by using the following formula:

Percent scavenging = $\frac{(A_t - A_0)}{A_t} \times 100$; where A₀ = Absorbance of control (without extract) and A_t = Absorbance of sample^{32,33}. All the tests were performed in triplicate.

RESULTS AND DISCUSSION

Different concentrations of 3.125, 6.25, 12.5, 25, 50, 100 and 200 µg / ml of alcoholic extracts were compared with their antioxidant activity in different in vitro models. Free radicals are removed from all selected extract fractions in a concentration-based manner until concentration is given in all models. On a comparative basis, ethanol and butanol fraction showed good antioxidant activity compared to remaining fraction these are ether and DCM part (Table 2 and 3).

DPPH is a stable free radical. The in vitro study carried out on this radical is based on the measurement of the scavenging ability of antioxidants towards the stable radical DPPH. This radical reacts with suitable reducing agents, the electrons become paired off and the solution loses color stoichiometrically depending on the number of electrons taken up³⁵. From the present results, it may be concluded that the extracts reduce the radical to the corresponding hydrazine when they react with the hydrogen donors in the antioxidant principles. The result show that ethanol and butanol fraction compared to pet. Ether and DCM fraction % SC₅₀ found both potent fractions 6.98% and 7.19% respectively by DPPH assay method (Table no. 2 and fig. 1).

In the present study, examined by ABTS radical scavenging assay method the observation shows ethanol and butanol fraction compared to pet. Ether and DCM fraction % SC₅₀ found both potent fractions 4.69% and 4.06% respectively (Table no. 2 and fig. 2). This may be due to the antioxidant principles in the ABTS radical cation was produced in the stable form using potassium per sulphate. After getting the stable absorbance, the antioxidant plant extract is added to the reaction medium and the antioxidant power was measured by studying decolorization³⁶. ABTS assay is an

excellent tool for determining the antioxidant activity of hydrogen-donating antioxidants and of chain-breaking antioxidants³⁷. Different artificial free radical species, such as ABTS + radical cation, has been used to assess radical scavenging ability and antioxidant activity.

Ortho-substituted phenolic compounds may exert pro-oxidant effects by interacting with iron. O-phenanthroline quantitatively forms complexes with ferric ion which get disrupted in the presence of chelating agents. The extract fractions interfered with the formation of ferrous-O-phenanthroline complex, thereby suggesting that the extract fractions have metal chelating activity and show in table no. 2 and fig. 3. The result of this assay method shows pet. Ether and DCM fraction compared to ethanol and butanol fractions % SC₅₀ found both potent fractions 9.63% and 101.9% respectively.

The total antioxidant activity of the extract fractions was calculated based on them formation of phosphomolybdenum complex which was measured spectrophotometrically³⁷ the obtained result show in table no. 2. Free radicals induce lipid peroxidation in polyunsaturated lipid rich areas like liver and brain. In this study, lipid peroxidation was induced in vitro and the extracts showed concentration dependent prevention towards generation of lipid peroxides.

Preliminary phytochemical screening revealed the presence of alkaloids, carbohydrates, phenolic compounds, tannins, saponins and flavonoids (show in Table no. 1) in the selected plant³⁹. Phenolics, flavonoids, saponins, and tannins have been proved to be responsible for the antioxidant activity of various medicinal plants reported earlier⁴⁰. Hence, these may be responsible for the observed activity in the plant. The present study proved the antioxidant potential of *Bombax malabaricum*.

Table 1: Estimation of phytochemicals of extract and its fractions of *Bombax malabaricum*.

Phytochemical tests	Ethanol	Pet. Ether	Butanol	Dichloromethane	Aqueous
Alkaloids	+	+	-	+	+
Antraquinones	+	+	-	+	+
Cardiac glycosides	+	-	-	-	+
Coumarins	+	-	+	+	+
Flavonoids	+	+	+	+	+
Saponins	+	-	+	+	+
Phlobatannins	+	-	+	-	+
Tannins	+	+	-	+	+
Terpenoids	+	+	+	+	+

Table No. 2: Effect of different extract/fractions of *Bombax malabaricum*. in various in vitro antioxidant assay

Extract/ Fractions	% SC ₅₀ values		
	DPPH	ABTS	O-phenanthroline
Ethanol	6.98	4.69	198.4
Pet. Ether	200.06	48.31	101.9
DCM	50.12	31.64	91.63
Butanol	7.19	4.06	298.40
Aqueous	41.04	51.93	189.83




Fig. 1: Effects of different extract/fractions of *Bombax malabaricum*. in in vitro DPPH antioxidant assay.




Fig. 2: Effects of different extract/fractions of *Bombax malabaricum*. in in vitro ABTS radical scavenging assay.




Fig. 3: Effects of different extract/fractions of *Bombax malabaricum*. in in vitro O-phenanthroline assay

Table 3: Estimation of Total Antioxidant Capacity of different extract/fractions of *Bombax malabaricum*.

Extract/Fraction	Concentration equivalent to Ascorbic acid		
	(2mg/ml)	(5mg/ml)	(10 mg/ml)
Ethanol	535.36	1035.83	1816.07
Pet. ether	204.64	367.74	454.64
DCM	215.36	437.74	836.79
Butanol	635.83	1069.17	908.45
Aqueous	269.88	417.74	498.93

REFERENCES

- Halliwell B: Establishing the significance and optimal intake of dietary antioxidants; The biomarker concept. *Nutr. Rev* 1999; 57: 104-113.
- Hogg N: Free radicals in diseases. *Semin. In Reprod. Endocrin.*, 1998, 16, 241-88.
- Shinde: Effect of Free Radicals & Antioxidants on Oxidative Stress: A Review *Journal of Dental & Allied Sciences* 2012;1(2):63-6.
- Ratnam, D.V. Ankola, D.D. Bhardwaj, V.; Sahana, D.K.; Kumar: M.N.V.R. Role of antioxidants in prophylaxis and therapy: A pharmaceutical perspective. *J. Control Release.* 2006,113, 189-207.
- Hazra B, Biswas S, Mandal N: Antioxidant and free radical scavenging activity of *Spondiaspinnata*, *BMC Comp Alt Med*, 8, 2008, 63-64.
- Podsedek, A. Natural antioxidants and antioxidant activity of *Brassica* vegetables: A review. *LWT-Food Sci. Technol.* 2007, 40, 1-11.
- P. Vijayan, C. Raghu, G. Ashok, S. A. Dhanaraj, B. Suresh. Antiviral activity of medicinal plants of Nilgiris. *Indian J Med res.* 2004; 120, 24-29.
- K. Ananil, J. B. Hudson, C. De Souza, K. Akpaganal, G. H. N. Tower, J. T. Amason, M. Gbeassor. Investigation of medicinal plants of Togo for antiviral and antimicrobial activities. *Pharmaceutical Biology* 2004; 38, 1, 40-45.
- Baby Joseph and S. Justin Raj. A comparative study on various properties of five medicinally important plants. *International journal of Pharmacology*.2011; 1-6.
- Emdad Hossain, Subhash C Mandal, JK Gupta. Phytochemical screening and *In vivo* antipyretic activity of the methanol leaf extract of *Bombax malabaricum* DC(Bombaceae). *Tropical journal of Pharmaceutical research.* 2011; 10 (1): 55-60.
- Sanz MJ, Ferrandiz ML, Cejudo M et al. Influence of a series of natural flavonoids on free radical generating systems and oxidative stress. *Xenobiotica* 1994; 24: 589-599.
- The Wealth of India- a dictionary of Indian Raw Material and Industrial Products". Vol-II, Bombay: Popular Prakashan, 1976.
- Anonymous: Wealth of India, Volume II, Publication and Information Directorate, CSIR, New Delhi 1988; 177.
- Saleem R, Ahmad M, Hussain SA, Qazi AM, Ahmad SI, Qazi MH, Ali M, Faizi S, Akhtar S, Husnain SN. (1999). Hypotensive, hypoglycaemic and toxicological studies on the flavonol C-glycoside shamimin from *Bombax ceiba*. *Planta Med* 65:331-334.
- Shahat AA, Hassan RA, Nazif NM, Van Miert S, Pieters L, Hammuda FM, Vlietinck AJ. (2003). Isolation of mangiferin from *Bombax malabaricum* and structure revision of shamimin. *Planta Med* 69:1068-1070.
- Niranjana GS, Gupta PC. (1973). Anthocyanins from the flowers of *Bombax malabaricum*. *Planta Med* 24:196-199.
- Vijaya BRM, Kesava RM, Gunasekar D, Marthanda MM, Caux C, Bodo B. (2003). A new sesquiterpene lactone from *Bombax malabaricum*. *Chem Pharm Bull* 51:458-459.
- Agrawal GD, Rizvi SA, Gupta PC, Tewari JD. (1972). Study of a polysaccharide from the stamens of *Bombax malabaricum* flowers. *Planta Med* 21:293-303.
- Singh NP, McCoy M.T, Tice RR and Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells, *Exp. Cell Res*, 175, 1998; 184-191.
- Liotta LA. Tumor invasion and metastasis: Role of extra cellular matrix. Rhoads Memorial Award Lecture, *Cancer Research*, 46, 1986; 1-3.
- Steinmetz, K. A., & Potter, J. D. (1996). Vegetables, fruits, and cancer prevention: A review. *Journal of American Diet Association*, 96, 1027-1039.
- Rimm, E. B., Ascherio, A., Grovannucci, E., Spiegelman, D., Stampfer, M. J., & Willet, W. C. (1996). Vegetable, fruits, and cereal fiber intake and risk of coronary heart disease among men. *Journal of the American Medical Association*, 275, 447-451.
- Carini, R., Poli, G., Diazini, M. U., Maddix, S. P., Slater, T. F., & Cheesman, K. H. (1990). Comparative evaluation of the antioxidant activity of α -tocopherol, α -tocopherol polyethylene glycol 1000 succinate and α -tocopherol succinate in isolated hepatocytes and liver microsomal suspensions. *Biochemical Pharmacology*, 39, 1597-1601.
- Dorman, H. J. D., Peltoketo, A., Hiltunen, R., & Tikkanen, M. J. (2003). Characterization of the antioxidant properties of tde-odourised aqueous extracts from selected Lamiaceae herbs. *Food Chemistry*, 83,255-262.
- Kitts, D. D., Yuan, Y. V., Wijewickreme, A. N., & Hu, C. (2000). Antioxidant properties of a North American ginseng extract. *Molecular and Cellular Biochemistry*, 203, 1-10.
- Harborne JB: *Phytochemical methods*. London: London Chapman and Hall, Ltd; 1973:49-188.
- Trease GE, Evans WC: *Pharmacognosy*. London: Brailliar Tiridel Can Macmillian Publishers; 1989:11.
- Sofowora AE: *Medicinal plants and traditional medicine in Africa*, Volume 2. Ibadan, Nigeria: Spectrum books Ltd; 1993:289.
- Bansal P, Paul P, Nayak PG, Pannakal ST, Zou J-H, Laatsch H, et al. Phenolic compounds isolated from *Pilea microphylla* prevent radiation-induced cellular DNA damage. *Acta Pharm Sin B.* 2011;1: 226-35.
- Tachakittirungrod S, Okonogi S, Chowwanapoonpohn S. Study on antioxidant activity of certain plants in Thailand: Mechanism of antioxidant action of guava leaf extract. *Food Chem.* 2007; 103: 381-8.
- Prieto P, Pineda M, Aguilar MM. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Anal Biochem.* 1999; 269: 337-41.
- Besada A. A facile and sensitive spectrophotometric determination of ascorbic acid. *Talanta.* 1987; 34: 731-2.
- Berker KI, Guclu K, Tor I, Apak R. Comparative evaluation of Fe (III) reducing power-based antioxidant capacity assays in the presence of phenanthroline, batho-phenanthroline, tripyridyltriazine (FRAP) and ferricyanide reagents. *Talanta* 72 (2007) 1157-1165.
- Charanjeet Singh, Neeraj Kumar Saini, Hansraj Manda, Manmohan Singhal, Kamal Sachdeva. "Evaluation of antioxidant activity of *Terminalia arjuna* leaves extract. *Pharmacologyonline*, 2011, 1: 998-1006.
- Kaleem M and Asif MA, Antidiabetic and antioxidant activity of *Annona squamosa* extract in streptozotocin induced diabetic rats, *Journal of Singapore Medicine*, 2006; 47:670-676.
- Khlifi S, Hachimi YE, Khalil A, Es- Safi N and Abbouyi AE, *In vitro* antioxidant effect of *Globularia alypum* Linn. Hydromethanolic extract, *Indian Journal of Pharmacology*, 2005; 4; 27-34.
- Kharya MD, Bhardwaj S and Sharma A, Screening methods of antioxidant activity, *Pharmacog Reviews*, 2007; 232-237.
- Khandelwal KR, In *Practical Pharmacognosy*, Nirali Prakashan, Pune, 2nd Edition, 2000; 215-220.
- Kumar PV, Shashidhara S, Kumar MM and Sridhara BY, Effect of *Luffa echinata* on lipid peroxidation and free radical scavenging activity, *Journal of Pharmacy and Pharmacology*, 2000; 52; 891-898.
- Sathishkumar comparative analyses of non- enzymatic and enzymatic antioxidants of *Encostemmalittoraleblume*, *International Journal of Pharma and Bio Sciences* 2010; V1(2).