

Available online on 30.08.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Review Article

Ayurvedic Polyherbal Formulation and Diabetes: An Overview

Sughosh Vishweshwar Upasani*¹, Govinda Satish Bhandari¹, Manali Sughosh Upasani², Ashwinkumar Ashokrao Torne¹, Nitin Ghanshyamdas Haswani¹

1. SES's R. C. Patel Institute of Pharmacy, Shirpur, 425405, Dhule, Maharashtra, India.
2. KES's College of Pharmacy, Amalner, 425401, Jalgaon, Maharashtra, India.

ABSTRACT

At this era the world is suffering from many health dilemmas. Most of them are environmental, ecological, depending on several human races or ethnic background. Diabetes is one of them. Diabetes is chronic metabolic disorder which is group of diverse complications. The present study is overview of various 30 Ayurvedic polyherbal formulations and their significance as antidiabetic potential in diabetic rats. Models used are alloxan induced, streptozotocin induced and streptozotocin-nicotinamide induced diabetes. Plants have always been an exemplary source of drugs and many of the currently available drugs have been derived directly or indirectly from them. The Ayurveda and other Indian literature strongly believe; current scientific evidences, the use of medicinal plants in treatment of various human diseases is found significant. Diverse researches conducted in the last few decades on plants, mentioned in ancient literature or used traditionally for diabetes, have shown antidiabetic property. Further research should continue, especially in areas within nations that have received less attention.

Keywords: Ayurvedic polyherbal formulation, Diabetes, Antidiabetic, Survey, Preclinical, Madhumeh

Article Info: Received 06 July 2019; Review Completed 14 Aug 2019; Accepted 22 Aug 2019; Available online 30 Aug 2019



Cite this article as:

Upasani SV, Bhandari GS, Upasani MS, Torne AA, Haswani NG, Ayurvedic Polyherbal Formulation and Diabetes: An Overview, Journal of Drug Delivery and Therapeutics. 2019; 9(4-A):779-784 <http://dx.doi.org/10.22270/jddt.v9i4-A.3517>

*Address for Correspondence:

Mr. Sughosh Vishweshwar Upasani, SES's R. C. Patel Institute of Pharmacy, Shirpur, 425405, Dhule, Maharashtra, India.

DIABETES

At this era the world is suffering from many health dilemmas. Most of them are geographical, ecological, depending on a variety of human races or ethnic backgrounds. Diabetes is one of them. Diabetes is chronic metabolic disorder which is group of different complications. At present India is leading the world in number of diabetes, furthermore it having second ranking in number of Prediabetes.¹ As per International Diabetes Federation (IDF) Diabetes Atlas, the global Diabetes occurrence in the age group 20–79 years is estimated to be 6.6% for the year 2010 which convert into 285 million people suffering from Diabetes.^{1, 2, 3}

Signs and Symptoms

As per Indian Health Services (IHS), Australian Diabetes Society (ADS) and American Diabetes Association (ADA) following are some symptoms which observed - Constant hunger, Unexplained weight loss or weight gain, Flu-like symptoms, including weakness and fatigue, Blurred vision or visionary problems, Slow healing of cuts or bruises, Tingling or loss of feeling in hands or feet, Recurring gum or skin infections, and Recurring vaginal or bladder infections etc.¹⁻⁸

Risk Factors

Following are associated with insulin resistance and are risk factors for the development diabetes - Age \geq 30 years, Family history of diabetes, Sedentary lifestyle, Metabolic syndrome, Cardiovascular disease, Hypertension or family history of cardiovascular diseases, Overweight or obese, Women who have had gestational diabetes, had high birth weight babies (greater than 9 lbs.), and/or has Polycystic Ovarian Syndrome (PCOS), Middle age women who has Large bra cup size at age 20.^{1, 4-7}

TREATMENT AND AYURVEDIC APPROACH

The goals of early glucose-directed therapies are to normalize glucose levels, to prevent or delay progression to diabetes, and to prevent micro vascular complications. Plants have always been an exemplary source of drugs and many of the currently available drugs have been derived directly or indirectly from them. In India, diabetes has been known for a long time, but its incidence is not of the same magnitude across the subcontinent. The wide range of structures of the plant constituents, which appear to be the active hypoglycaemic principles, suggests different sites of action within the body. The plants provide a potential source

of hypoglycaemic drugs because many plants and plant derived compounds have been used in the treatment of diabetes. Several medicinal plants have found potential use as hypoglycaemic in the Indian system of medicines. A number of researches conducted in the last few decades on plants, mentioned in ancient literature or used traditionally for diabetes, have shown antidiabetic property. Among them, most off plants and their products (active natural principles and crude extracts) that have been mentioned used in the Indian traditional system of medicine have shown experimental or clinical antidiabetic activity.⁸ In Ayurved it is considered as 'Naasti Moolam Anaushadhim' - no plant present on Earth which isn't having medicinal properties. The Ayurved and other Indian literature strongly believe; current scientific evidences, the use of medicinal plants in

treatment of various human diseases is found significant. Diverse researches conducted in the last few decades on plants, mentioned in ancient literature or used traditionally for diabetes, have shown antidiabetic property.⁹ The Indian word for diabetes derived from Sanskrit literature Ayurved is "Madhumeha". The *Madhumeha* is a combination of two "Madhu" means "sweet or sweetness" as like honey and "Meha" means "excessive urination". All the renowned traditional texts of Ayurveda like Charaka Samhita (1000 B.C.), Sushruta Samhita (600 B.C.) and subsequent works refer to this disease under the term *Madhumeha* or *Ikshumeha* (*Ikshu* means Sugarcane).¹⁰ Indian plants which are most effective and the most commonly studied in relation to DM and their complications are ¹⁰⁻¹²

- | | | |
|-----------------------------------|----------------------------------|----------------------------------|
| 1. <i>Acacia Arabica</i> | 20. <i>Embelica officinalis</i> | 39. <i>Nelumbo nucifera</i> |
| 2. <i>Acacia nilotica</i> | 21. <i>Eucalyptus globulus</i> | 40. <i>Ocimum sanctum</i> |
| 3. <i>Aegle marmelos</i> | 22. <i>Eugenia jambolana</i> | 41. <i>Panax ginseng</i> |
| 4. <i>Allium cepa</i> | 23. <i>Enicostemma littorale</i> | 42. <i>Picrorrhiza kurroa</i> |
| 5. <i>Allium sativum</i> | 24. <i>Ficus bengalensis</i> | 43. <i>Phyllanthus niruri</i> |
| 6. <i>Aloe vera</i> | 25. <i>Ficus glomerulata</i> | 44. <i>Pterocarpus marsupium</i> |
| 7. <i>Andrographis paniculata</i> | 26. <i>Glycyrrhiza glabra</i> | 45. <i>Punica granatum</i> |
| 8. <i>Areca catechu</i> | 27. <i>Gymnema sylvestre</i> | 46. <i>Salacia reticulata</i> |
| 9. <i>Azadirachta indica</i> | 28. <i>Hibiscus rosa-sinesis</i> | 47. <i>Salacia oblonga</i> |
| 10. <i>Berginia ciliata</i> | 29. <i>Ipomoea batatas</i> | 48. <i>Swertia chirayita</i> |
| 11. <i>Beta vulgaris</i> | 30. <i>Lantana camara</i> | 49. <i>Syzgium cumini</i> |
| 12. <i>Biophytum sensitivum</i> | 31. <i>Magnifera indica</i> | 50. <i>Termenalia arjuna</i> |
| 13. <i>Bombax ceiba</i> | 32. <i>Memecylon umbellatum</i> | 51. <i>Termenalia chebulla</i> |
| 14. <i>Brassica juncea</i> | 33. <i>Momordica cymbalaria</i> | 52. <i>Tinospora cardifolia</i> |
| 15. <i>Caesalpinia bonducella</i> | 34. <i>Momordica charantia</i> | 53. <i>Trigonella foenum -</i> |
| 16. <i>Cajanus cajan</i> | 35. <i>Morus alba</i> | 54. <i>Vinca rosea</i> |
| 17. <i>Coccinia indica</i> | 36. <i>Musa sapientum</i> | 55. <i>Withania somnefera</i> |
| 18. <i>Caesalpinia bonducella</i> | 37. <i>Mucuna pruriens</i> | 56. <i>Zingiber officinalis</i> |
| 19. <i>Curcuma longa</i> | 38. <i>Murraya koenigii</i> | |

Methodology

The current study provides a collection of information on Ayurvedic Polyherbal formulation and their evaluation on diabetes. The appropriate literature, including books, journals, and reports, was reviewed. The relevant information was searched using various electronic catalogs (e.g., Google Scholar, Science Direct, etc.) and keywords such as "Ayurvedic Polyherbal formulation," "Polyherbal,"

"Ayurvedic formulation," "Indian," "diabetes," and "Madhumeh". It was difficult to include all the information. This study focused on information that would be easily accessible for researchers.¹³ Table number 01 is focusing some selected experimental preclinical antidiabetic activities. Figure number 01 and 02 will show frequency and number of plants used, and frequency of experimental model used respectively.

Table No. 01- Composition of Ayurvedic Polyherbal formulation and other details

No. of Plants	Ayurvedic Polyherbal formulation	Experimental Model	Extraction solvent	Standard drug used	Extraction Method	Ref
11	Aegle marmelos, Annona squamosa, Bougain villa, Cassia auriculata, Emblica officinale, Ficus carica, Hybiscus rosasinensis, Psidium guajava, Tenospora cardifolia, Terminalia chebula, Zingiber officinale	Alloxon Induced DM	Aqueous	Glibenclamide	Decoction	15
11	Aegle marmelos, Annona squamosa, Azadirachta indica, Cajanus cajan, Cinnamomum tamala Ferula assafoetida, Gymnema sylvestre, Moringa oleifera, Tamar indus indica, Trigonella foenum, Zingiber oficinale,	Streptozotocin Induced DM	Dried powder	Glibenclamide	Dried powder	16
10	Azadirachta indica, Curcuma longa, Gymnema sylvestre, Momordica charantia, Paneer pera, Piper nigrum, Swerita chirayita, Syzygium cumini, Terminalia bellirica, Trigonella foenum,	Alloxon Induced DM	Hydroalcoholic	Glibenclamide	Maceration	17
8	Acacia catechu, Cassia auriculata, Hemidesmus indicus, Holarrhena antidysenterica, Salacia reticulata, Strychno spotatrum, Trigonella graecum, Vetiveria zizanioides	Streptozotocin Induced DM	Dried powder	Glibenclamide	Dried powder	18
8	Azadirecta indica, Curcuma longa, Emblica officinale, Enicostemma littorale, Gymnema sylvestre, Momordica charatia, Syzygium cumini, Tinospora cordiofolia,	Streptozotocin Induced DM				19
6	Caesalpenia bonduc, Citrullus lanatus, Evolvulus alsinoide , Gymnema sylvestra, Tinospora cordiofolia, Withania coagulans,	Alloxon Induced DM	Ethanol	Glibenclamide	Soxlet	20
5	Andrographis paniculata, Gymnema sylvestere, Momordica charantia, Pterocarpus marsupium, Tinospora cordifolia	Alloxon Induced DM	Methanol		Maceration	21
5	Adiantum capillus, Astercantha longifolia, Callicarpa macrophylla, Ficus benghalensis, Melia azedarach	Alloxon Induced DM	Ethanol		Succesive solvent extraction by soxlet	22
5	Alstonia scholaris, Gymnema sylvestre, Holarrhena pubescens, Premna corymbosa, Solanum nigrum,	Alloxon Induced DM	Aqueous		Maceration	23
5	Emblica officinalis, Gymnema sylvestre, Terminalia arjuna, Tinospora cordifolia, Zingiber officinale	Nicotinamide Streptozotocin Induced DM	Supercritical CO2		Supercritical fluid extraction	24
5	Andrographis paniculata, Eugenia jambolana, Gymnema sylvestre, Momordica charantia, Myristica fragrans	Streptozotocin Induced DM	Hydroalcoholic	Glibenclamide	Tripple maceration	25
5	Camellia sinensis, Foeniculum vulgare, Macrotyloma uniflorum, Punica granatum, Trigonella foenum	Streptozotocin Induced DM	Ethanol	Glibenclamide	Soxlet	26
5	Gymnema sylvestre, Pterocarpus marsupium, Tinospora cordifolia Trigonella foenum	Streptozotocin Induced DM	Ethanol		Maceration	27
5	Cassia auriculata, Cinnamomum tamala, Ficus banghalensis, Mangifera indica, Trichosynthis diocia	Streptozotocin Induced DM	Aqueous		Maceration	28

5	Azadirachta indica, Gymnema sylvestre, Momordica charantia, Syzygium cumini, Trigonella foenum	Streptozotocin Induced DM	Aqueous	Glibenclamide	Morter paste	29
4	Allium sativum, Ocimum gratissimum, Vernonia amygdalina, and Zingiber officinale	Alloxon Induced DM	Aqueous		Cold maceration	30
3	Eugenia jambolana, Momordica charantia, Ocimum sanctum	Alloxon Induced DM	Hydroalcoholic		Soxlet	31
3	Piper nigrum, Ricinus Communis, Tribulus terrestris	Alloxon Induced DM	Ethanol	Glibenclamide	Cold maceration	32
3	Alternanthera sessilis, Amaranthus viridis, Boerhavia diffusa	Alloxon Induced DM	Aqueous	Glibenclamide	Decoction	33
3	Caesalpinia bonducella, Mucona puriens, Pongamia pinnata	Alloxon Induced DM	Aqueous	Glibenclamide	Continuous hot soxlet	34
3	Glycosmis pentaphylla, Mangifera indica, Tridax procumbens	Nicotinamide Streptozotocin Induced DM	Ethanol	Glibenclamide	Soxlet	35
3	Barleria Montana, Rotula aquatic, Schrebera swietenoides	Streptozotocin Induced DM	Methanol		Soxlet	36
3	Portulaca oleracea, Spilanthe safricana, Sida rhombifolia	Streptozotocin Induced DM	Aqueous	Glibenclamide	Maceration	37
3	Azadirachta indica, Boerhavia diffusa, Tribulus terrestris,	Streptozotocin Induced DM	Alcoholic	Glibenclamide	Maceration	38
3	Azadirachta indica, Andrographis paniculata, Moringa oleifera	Streptozotocin Induced DM	Hydroalcoholic		Maceration	39
3	Gymnema sylvestre, Syzygium cumini, Urtica dioica	Streptozotocin Induced DM	Methanol		Rotary shaking	40
3	Azadirachta indica, Bougainvillea spectabilis, Trigonella foenum	Streptozotocin Induced DM	Aqueous, chloroform, ethanol	Glibenclamide	Extraction	41
3	Azadirachta indica, Andrographis paniculata, Moringa oleifera	Streptozotocin Induced DM	Hydroalcoholic	Glibenclamide	Extraction	42
2	Camellia sinensis, Macrotyloma uniflorum	Nicotinamide Streptozotocin Induced DM	Methanol		Maceration	43
2	Annona squamosa, Phyllanthus emblica	Streptozotocin Induced DM	Aqueous	Glibenclamide	Decoction	44




Figure No. 01 - Number of Plants used




Figure No. 02 - Experimental Model used

DENOUEMENT

One valuable gift to human health is provided by nature in the form of medicinal plants in the locality and one of the significant ways in which humans directly reap the benefits provided by biodiversity.¹⁴ In this overview of Ayurvedic Polyherbal formulation found that it has a significant effect on the diabetes. According to editorial from "The Lancet" special issue on diabetes as health crisis, near about 366 million people worldwide currently have diabetes mellitus. Estimation is also that this number will increase by 552 million and about 470 million people with Prediabetes by 2030.⁴⁵ Thus, by triangulation, it is probably still possible to document most of the knowledge¹⁴, but further research should continue, especially in areas within nations that have received less attention.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

SOURCES OF FUNDS

Nil

ACKNOWLEDGMENT

The authors are thankful to Dr. S. J. Surana, Principal, R. C. Patel Institute of Pharmaceutical Education & Research, Shirpur-425405, Dhule, Maharashtra, India, Prof. R. G. Mali, Principal, KES's College of Pharmacy, Amalner-425401, Jalgaon, Maharashtra, India, for providing guidance in research work. I extend my thanks to my supportive friends, colleagues whose help improves my work and ability for it.

REFERENCES

- Upasani SV, Nandedkar RY., Patil SH, Shah VS., Clinical evaluation of mehant - an Ayurvedic polyherbal formulation in Prediabetes. *Int J Insti Pharm & L Sci* 2012; 2(3): 239-245.
- Misra. A Review of the Epidemiology of Diabetes in Rural India. *Diabetes Res. And Clin. Pract*, 2011; 303-311
- Mohan V. Incidence of Diabetes and Pre-Diabetes in a Selected Urban South Indian Population. (*Cups - 19*), *JAPI*, 56; 152-157
- ICMR Guidelines for Management of Type 2 Diabetes. 2005
- Aroda V. Approach to the Patient with Prediabetes. *J. Clin. Endocrinol Metab.*, 1993, Vol (9), 2008; 3259-3265
- USDHHS. Insulin Resistance and Pre-Diabetes. NIH Publication No. 09-4893, 2008.
- Ray. Breast Size and Risk of Type 2 Diabetes. *CMAJ*, Vol 178(3), 2008; 289-295
- Upasani SV, Nandedkar RY., Upasani MS, Prediabetes Metformin and Ayurvedic aspect - an overview", Lambert Academic Publications. ISBN 978-3-330-33376-5
- Bhoyar P. Herbal Antidiabetics:A Review.*Int.J.Res.Pharm.Sci.*,Vol 2(1),2011; 30-37
- Grover JK, Yadav S, Vats V. Medicinal plants of india with anti-diabetic potential. *J Ethnopharmacol* 2002; 81:81-100
- Manisha Modak , Priyanjali Dixit , Jayant Londhe , Saroj Ghaskadbi , and Thomas Paul A. Devasagayam, Indian Herbs and Herbal Drugs Used for the Treatment of Diabetes, *J. Clin. Biochem. Nutr.*, 40, 163-173, May 2007
- S. V. Upasani, P. V. Ingle, P. H. Patil, R. Y. Nandedkar, V. S. Shah1, S. J. Surana, Traditional Indian spices useful in Diabetes Mellitus - an updated review, *J. Pharm. BioSci.* 4(2013) 157-161
- MS Upasani, SV Upasani, VG Beldar, CG Beldar, PP Gujarathi, Infrequent use of medicinal plants from India in snakebite treatment, *Integr Med Res* 7 (2 0 1 8) 9-26, <https://doi.org/10.1016/j.imr.2017.10.003>
- SV Upasani, VG Beldar, AU Tatiya, MS Upasani, SJ Surana, DS Patil, Ethnomedicinal plants used for snakebite in India: a brief overview, *integr med res* 6 (2017), 114-130, <http://dx.doi.org/10.1016/j.imr.2017.03.001>
- Kasireddy PB, Arunkumar S, Namratha S, Swathi B, Manguluri P, Adapa V, Evaluation of anti-diabetic efficacy of polyherbal formulations in experimentally induced hyperglycemic rats., *International journal of research in pharmacy and chemistry.*, 6(4), 2016, 750-758.
- A Panda, S Jena, PK Sahu, S Nayak, P Padhi, Effect of Polyherbal Mixtures on the Treatment of Diabetes, 2013, <http://dx.doi.org/10.1155/2013/934797>
- Sharma N, Goyal A, Hypoglycemic Effect of Polyherbal Formulation in Alloxan Induced Diabetic Rats., *Chemistry Research Journal*, 2(4), 2017, 214-219
- S Tomy, S Johnson, UChander J, SCeline, TK Ujwala, CS Kandasamy, P Arulraj, R Venkatanarayanan, Anti-Diabetic Effect Of Polyherbal Formulation In Ogtt And Streptozotocin-Induced Diabetic Rat Model, *Int J Pharm Pharm Sci*, 2015, 7, 10, 216-219
- SS Patel, RS Shah, RK Goyal, Antihyperglycemic, antihyperlipidemic and antioxidant of Dihar a Ayurvedic polyherbal formulation in streptozocin induced diabetic rats, *Indian J of Experimental Biology*, 2009, 47, 564-570
- SM Mahajan, DT Baviskar, PM Chaudhari, Anti-Diabetic Activity Of Polyherbal Formulation On Alloxan Induced Diabetes, *IOSR J of P and Bio Sci*, 13, 1, 2018, 01-06
- Bhikshapathi DVRN, Rajesham VV, Pharmacological, Screening of Anti Diabetic Activity of Polyherbal Formulation In Wistar Albino Rats. *American Journal of PharmTech Research*, 8(3), 2018, 82-93
- Appalaraju G, Reddy S, Reddy V, Rajani G, Nithin, Raju AB, Anti diabetic activity of polyherbal extract on Streptozotocin

- induced diabetes in Wistar rats., Scholars Research Library Annals of Biological Research, 5(6), 2014, 39-46.
23. Thamizh SN, Hima KM, Sanjayakumar YR, Ranjini KR, Vasantha Kumar KG, Swamy GK, Hypoglycaemic and Antioxidant Activity of SPHAG - a Poly Herbal Formulation in Alloxan Induced Wistar Albino Rats., International Journal of Pharma Sciences and Research., 6(4), 2015, 767-772.
 24. Belhekar S, Chaudhari PD, Pandhare RB, Pawar AR, Effect of Polyherbal and Allopolyherbal Formulation on Streptozotocin-Nicotinamide Induced Diabetic Nephropathy in Rats, International Journal of Toxicological and Pharmacological Research, 8(3), 2016, 138-145.
 25. Tripathi N, Kumar V, Acharya S, Anti-Diabetic Activity of a Polyherbal Formulation in Streptozotocin Induced Type 2 Diabetic Rats., Journal of Natural Remedies., 16(4), 2016
 26. Gengiah K, Hari R, Anbu J, Antidiabetic antihyperlipidemic and hepato-protective effect of Gluconorm-5: A polyherbal formulation in streptozotocin induced hyperglycemic rats. Ancient Sci Life, 34, 2014, 23-32
 27. Yadav AP, Kori ML, Sagar R, Antidiabetic Activity of Polyherbal Formulation in Streptozotocin Induced Type II Diabetic Rats. Int. J. Pharm. Life Sci., 7(12), 2018, 5348-5353.
 28. Majumder P, Paridhavi M, A Novel PolyHerbal Formulation Hastens Diabetic Wound Healing with Potent Antioxidant Potential: A Comprehensive Pharmacological Investigation. Nat Prod Chem Res, 6, 2018, 324.
 29. Petchi, RR, Vijaya C, Parasuraman S, Antidiabetic activity of polyherbal formulation in streptozotocin - nicotinamide induced diabetic wistar rats., J Tradit Complement Med., 4(2), 2014 Apr, 108-17.
 30. Isitua CC, Akinyemi AJ, Akharaiyi FC, Olubiyi OO, Anadozie SO, Olayide II, Antihyperglycemic and Anti-hyperlipidemic Effect of Herbamed, A Herbal Formulation in Alloxan-Induced Diabetic Rats., Interventions Obes Diabetes., 1(4), 2005, 1-5.
 31. Kumar CH, Kumar S, Ishaq BM, Rani GU, Prakash KV, Antidiabetic Activity of A Polyherbal Preparation., Pharmacologyonline 2., 2010, 780-787.
 32. Baldi A, Goyal S, Hypoglycemic Effect of Polyherbal Formulation in Alloxan Induced Diabetic Rats, Pharmacologyonline, 3, 2011, 764-773.
 33. Shobana, G, Devi AG, Keerthana K, John NA, Jothi G, Hariharan G, Antidiabetic potential of poly herbal formulation on alloxan induced diabetic rats. Int. Res. J. Pharm., 9(8), 2018, 90-94.
 34. Selvam P, Evaluation of antidiabetic activity of a polyherbal formulation in alloxan induced diabetic rats, IJPSR, 1(9), 2010, 82-87.
 35. Petchi RR, Chockalingam V, Subramani P, Antidiabetic Activity of Polyherbal Formulation in Streptozotocin - Nicotinamide Induced Diabetic Wistar Rats., J Tradit Complement Med., 4(2), 2014, 108-117.
 36. Balakrishnaiah P, Satyanarayana T, Pharmaceutical and Biological Evaluations, 4(2), 2017, 111-117.
 37. Bruno MM, AmaMoor VJ, BiapaNya CP, Nanfack P, PiemeCA, Antioxidant and Synergistic Antidiabetic Activities of a Three- Plant Preparation Used in Cameroon Folk Medicine., International Scholarly Research Notices., Volume 2017.,7.
 38. Saxena N, Argal A, Evaluation of Antidiabetic Activity of a Herbal Formulation., Ec Pharmacology And Toxicology, 6(5), 2018, 349-355.
 39. Mahendra KS, Development and Evaluation of Antidiabetic Potential of Polyherbal Formulation in Streptozotocin Induced Animal Model, Int J cell Sci & mol boil., 5(2), 2018, 001-008.
 40. Gyawali R, Acharya G, Pokharel M, Shah AB, Silwal A, Evaluation of Antidiabetic Polyherbal Formulations., ARJMCS, Oct, 1(1), 2015, 33-37
 41. Gupta PP, Haider J, Yadav RP, Pal U. Preclinical evaluation of antidiabetic activity of poly herbal plant extract in streptozotocin induced diabetic rats. The Journal of Phytopharmacology 2016;5(2):45-49.
 42. Mahendra K S, Vijay K S, S Prakash R. Development and Evaluation of Antidiabetic Potential of Polyherbal Formulation in Streptozotocin Induced Animal Model. Int J cell Sci & mol biol. 2018; 5(2): 555656. DOI: 10.19080/IJCSMB.2018.05.555656
 43. Dhanalaxmi MR, Reddy B, Antidiabetic Activity of Polyherbal Formulation in Streptozotocin - Nicotinamide Induced Diabetic Wistar Rats., World Journal of Pharmaceutical Sciences., 3(4), 2015, 743-748
 44. Chaudhari A, Sharma S, Evaluation of Antidiabetic Activity of Polyherbal Formulation in Streptozotocin-Induced Diabetic Rats., UK Journal of Pharmaceutical and Biosciences., 4(5), 2016, 01-06.
 45. Editorial, The lancet, Vol. 379, June 16, 2012, 2213 DOI: [https://doi.org/10.1016/S0140-6736\(12\)60960-X](https://doi.org/10.1016/S0140-6736(12)60960-X) (Accessed 25.08.2019)

JDDT