brought to you by

Himmelfarb Health Sciences Library, The George Washington University Health Sciences Research Commons

Pediatrics Faculty Publications

Pediatrics

2012

Evaluation of sedimentation rate methodology reveals an unusual pediatric subpopulation with lupus or lupus-like syndrome and hemolytic anemia

Edward C. Wong George Washington University

Deborah Yuan

Claas Hinze

Lawrence Jung

Follow this and additional works at: http://hsrc.himmelfarb.gwu.edu/smhs_peds_facpubs Part of the <u>Pediatrics Commons</u>

Recommended Citation

Wong, E. C., Yuan, D., Hinze, C., & Jung, L. (2012). Evaluation of sedimentation rate methodology reveals an unusual pediatric subpopulation with lupus or lupus-like syndrome and hemolytic anemia. Pediatric Rheumatology Online Journal, 10(Suppl 1), A122.

This Journal Article is brought to you for free and open access by the Pediatrics at Health Sciences Research Commons. It has been accepted for inclusion in Pediatrics Faculty Publications by an authorized administrator of Health Sciences Research Commons. For more information, please contact hsrc@gwu.edu.

POSTER PRESENTATION



Open Access

Evaluation of sedimentation rate methodology reveals an unusual pediatric subpopulation with lupus or lupus-like syndrome and hemolytic anemia

Edward C Wong^{*}, Deborah Yuan, Claas Hinze, Lawrence Jung

From 2011 Pediatric Rheumatology Symposium sponsored by the American College of Rheumatology Miami, FL, USA. 2-5 June 2011

Purpose

Sedimentaion rate is often used to manage pediatric patients with rheumatologic disease. Most management decisions are dependent on studies which have used Wintrobe or Westergren sedimentation rate methodologies. However, these methods suffer from the need for relatively large amounts of blood and long turn-around times. Determination of sedimentation rate using laser kinetic rate determination has allowed calibration to Westergren methods, low volume of blood needed for testing and very rapid results. We sought to compare the Wintrobe method to the ESR Stat method (kinetic method; Hema-Technologies, Lebanon NJ) to determine suitability of the ESR Stat method for patient testing.

Methods

We performed a prospective comparison between the traditional Wintrobe and ESR Stat sedimentation rates in consecutive pediatric patients at a tertiary care pediatric hospital. Wintrobe and ESR Stat sedimentation data was fitted using a logarithmic model. Outliers were defined as those samples with ESR Stat sedimentation rates greater



Children's National Medical Center, Washington, DC, USA



© 2012 Wong et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

than 80 mm/hr and Wintrobe sedimentation rates less than 30 mm/hr (normal or mildly elevated sedimentation rate). Retrospective chart review was performed on all patients undergoing testing.

Results

A total of 131 pediatric patients (with one patient undergoing repeat testing because of sedimentation rate discrepancy) were tested. Age range was 18 months to 34 years with 29% being male. A logarithmic model appeared to best fit the data ($\mathbb{R}^2 = 0.7768$) and is seen below. Of interest was the identification of four patients who had apparently normal or mildly elevated sedimentation rates by the Wintrobe method versus an extremely high (greater than 120 mm/hr) by the ESR Stat method. These patients were noted to have lupus or lupus-like syndrome and a history of hemolytic anemia. Non-outlier samples were from patients who did not this combination of disease morbidities.

Conclusion

Discrepancies in Wintrobe and ESR Stat sedimentation rates may identify a subgroup of with lupus (or lupuslike syndrome) and a history of hemolytic anemia. Careful consideration of methodology is needed when sedimentation rate testing is performed on pediatric lupus patients.

Disclosure

Edward C. Wong: None; Deborah Yuan: None; Claas Hinze: None; Lawrence Jung: None.

Published: 13 July 2012

doi:10.1186/1546-0096-10-S1-A122 Cite this article as: Wong *et al*: Evaluation of sedimentation rate methodology reveals an unusual pediatric subpopulation with lupus or lupus-like syndrome and hemolytic anemia. *Pediatric Rheumatology* 2012 10(Suppl 1):A122.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit