

Himmelfarb Health Sciences Library, The George Washington University Health Sciences Research Commons

Global Health Faculty Publications

Global Health

4-2016

Validity of home-based sonographic diagnosis of obstetric risk factors by auxiliary nurse midwives in rural Nepal

Naoko Kozuki

Luke C. Mullany

Subarna K. Khatri

Ram Ghimire

Sharma Paudel

See next page for additional authors

Follow this and additional works at: http://hsrc.himmelfarb.gwu.edu/sphhs_global_facpubs

 Part of the [International Public Health Commons](#), [Maternal and Child Health Commons](#), and the [Nursing Midwifery Commons](#)

APA Citation

Kozuki, N., Mullany, L. C., Khatri, S. K., Ghimire, R., Paudel, S., Tielsch, J. M., & +5 additional authors (2016). Validity of home-based sonographic diagnosis of obstetric risk factors by auxiliary nurse midwives in rural Nepal. *Lancet Global Health*, 4, special issue (). [http://dx.doi.org/10.1016/S2214-109X\(16\)30028-6](http://dx.doi.org/10.1016/S2214-109X(16)30028-6)

This Journal Article is brought to you for free and open access by the Global Health at Health Sciences Research Commons. It has been accepted for inclusion in Global Health Faculty Publications by an authorized administrator of Health Sciences Research Commons. For more information, please contact hsrc@gwu.edu.

Authors

Naoko Kozuki, Luke C. Mullany, Subarna K. Khatri, Ram Ghimire, Sharma Paudel, James M. Tielsch, and +5 additional authors

Validity of home-based sonographic diagnosis of obstetric risk factors by auxiliary nurse midwives in rural Nepal



Naoko Kozuki, Luke C Mullany, Subarna K Khatri, Ram K Ghimire, Sharma Paudel, Karin Blakemore, Christine Bird, James M Tielsch, Steven C LeClerq, Joanne Katz

Abstract

Background Approximately 2·3 million fetal, neonatal, or maternal deaths occur annually during the intrapartum period or on the day of birth. Several risk factors for intrapartum-related complications require ultrasonography for accurate diagnosis, but sonographic services are scarce in low-resource settings. In this study, we aimed to assess the feasibility of community-based ultrasonography conducted by auxiliary nurse midwives to identify basic obstetric risk factors, in rural Sarlahi District, Nepal. We aimed to assess the validity at which these health workers can detect non-cephalic position, multiple gestation, and placenta previa and to explore whether sonographic diagnosis could improve outcomes for women with these high-risk conditions.

Methods Three auxiliary nurse midwives (ANM) received two 1-week ultrasound trainings at Tribhuvan University Teaching Hospital in Kathmandu, Nepal. Women at 32 weeks' or more gestation were enrolled in the study, and received ultrasonography from the ANMs during home visits. ANMs used ultrasonography to identify non-cephalic position, multiple gestation, and placenta previa. Images were saved and later reassessed by gold standard ultrasonographers to determine the validity of the ANM assessment. We also compared adverse outcomes in non-cephalic or multiple gestation births from the study group with those in a comparison group of women who did not receive ultrasonography from our study, but the study was not powered to detect a difference.

Findings We collected data from 815 women. The kappa statistics for diagnosis of non-cephalic position were 0·92, 0·98, and 0·94, respectively, for the three ANMs against the gold standard. Sensitivity, specificity, positive predictive value, and negative predictive value were between 90% and 100% for all ANMs. For multiple gestation pregnancies (n=6), the ANMs were in perfect agreement with both the gold standard reading and maternal postpartum self-report. Two cases of placenta previa were detected, and the gold standard was in agreement with both. There were 4 adverse outcomes out of 19 women (21%) with non-cephalic or multiple gestation pregnancies in the study group compared with 10 out of 36 (28%) in the comparison group. This difference was not significant (p=0·586).

Interpretation Our findings suggest that it is feasible for ANMs to conduct ultrasonography to identify basic obstetric risk factors in low-income settings. The difference in birth outcomes in non-cephalic and multiple births, comparing those antenatally diagnosed by ultrasound and those who were not, was not significant; the sample size was too small to detect a difference. Further investigation is warranted to determine whether sonographic diagnosis of selected obstetric factors could contribute to improved care-seeking and health outcomes in low-income settings.

Funding Cappsco Data for Life Award, SonoSite Soundcaring Program, National Institutes of Health/National Institute of Child Health and Human Development, Bill & Melinda Gates Foundation.

Copyright © Kozuki et al. Open Access article distributed under the terms of CC BY.

Declaration of interests

We declare no competing interests.

Published Online
April 8, 2016

Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA (N Kozuki PhD, L C Mullany PhD, SC LeClerq MPH, J Katz ScD); Nepal Nutrition Intervention Project – Sarlahi, Lalitpur, Nepal (S K Khatri MD); Tribhuvan University Teaching Hospital, Kathmandu, Nepal (R K Ghimire MD); The Johns Hopkins Hospital, Baltimore, MD, USA (C Bird RDMS, K Blakemore MD); George Washington University Milken Institute School of Public Health, Washington, DC, USA (J M Tielsch PhD)

Correspondence to:
Naoko Kozuki, 615 N Wolfe Street, W5019, Baltimore, MD 21205, USA
nkozuki@jhu.edu