

## CONTINUING MEDICAL EDUCATION

## ARTICLE

# Role of infection control in combating antibiotic resistance

A C Whitelaw, MB BCh, MSc, FCPATH (SA) (Micro)

*Division of Medical Microbiology, Department of Pathology, Faculty of Medicine and Health Sciences, Stellenbosch University and National Health Laboratory Service, Tygerberg Hospital, Cape Town, South Africa***Corresponding author:** A C Whitelaw ([awhitelaw@sun.ac.za](mailto:awhitelaw@sun.ac.za))

Infection control has been identified as one of the key interventions in controlling the threat of antibiotic resistance. Reducing the transmission of multidrug-resistant organisms (MDROs) reduces the need for broad-spectrum antibiotics in particular, while interventions that decrease the risk of infection have an impact on the use of any antibiotic. Hand hygiene remains the cornerstone of decreasing the transmission of MDROs. Alcohol-based hand rubs are a cheap, effective and convenient means of performing hand hygiene. Patients colonised or infected with MDROs should be placed on contact precautions, although implementation remains challenging in resource-limited environments. Screening for certain MDROs may play a role in curbing transmission of these organisms. If implemented, screening must be part of a comprehensive infection control strategy. In resource-limited settings, the costs and potential benefits of screening programmes need to be carefully weighed up. Care bundles have been shown to reduce the incidence of common healthcare-associated infections, including catheter-associated urinary tract infection, ventilator-associated pneumonia, central line-associated bloodstream infection and surgical site infection. These bundles are relatively inexpensive, and can play an important role in reducing antibiotic use and improving clinical outcomes.

*S Afr Med J* 2015;105(5):421. DOI:10.7196/SAMJ.9650

Antimicrobial resistance (AMR) is now well recognised as a global health threat<sup>[1]</sup> that impacts on human health and may potentially have a major effect on the global economy.<sup>[2]</sup> The United States Centers for Disease Control and Prevention (CDC) has identified four core actions to

combat this challenge, i.e. surveillance, antibiotic stewardship, improved drugs and diagnostics, and preventing spread.<sup>[3]</sup> While infection control is a very broad topic, this article focuses on measures to prevent the spread of multidrug-resistant organisms (MDROs) in healthcare settings and interventions to prevent infection.

## Transmission of multidrug-resistant organisms

Transmission of any infectious agent (whether multidrug resistant or not) implies the presence of a source of the organism, a susceptible host, and a transfer mechanism. Most infectious agents in the healthcare setting are transmitted from humans (most commonly patients, but also healthcare workers (HCWs) or visitors), although transmission from environmental sources has also been well described.<sup>[4,5]</sup> The greater the number of patients colonised with a particular MDRO (the 'colonisation pressure'), the greater the chance of transmission.<sup>[5]</sup> Colonised HCWs have also been implicated in the transmission of infectious agents,<sup>[6]</sup> but are thought to be a far less common source.<sup>[5]</sup>

Although organisms can be transmitted to any patient in hospital, certain patients are at higher risk of colonisation (and therefore of potential infection), including those with underlying medical conditions, those with compromised defences (such as indwelling devices, burns), and those who have undergone recent surgery.

There are three broad mechanisms by which organisms can be transmitted in healthcare settings:<sup>[4]</sup> contact, droplet transmission and airborne transmission. Droplet transmission refers to organisms spread by large respiratory droplets (e.g. pertussis, diphtheria, certain respiratory viruses), while airborne transmission applies to organisms spread on droplet nuclei that remain suspended for prolonged periods (e.g. *Mycobacterium tuberculosis*, measles, varicella). Neither

of these are a major route of transmission for MDROs, except for MDR tuberculosis, which is not the focus of this article.

Contact spread is the most important means of transmission for the vast majority of MDROs in healthcare settings, and can be either direct or indirect. Direct contact transmission occurs when an infected or colonised person comes into physical contact with a susceptible host. Indirect contact transmission implies the presence of some intermediate vehicle – either human or an inanimate object. Although transmission by items such as contaminated equipment, shared toys and poorly sterilised equipment is possible, the most important means of transmission is via the hands of HCWs.

## Prevention of transmission

As the hands of HCWs are the most important means of transmission of MDROs, the importance of hand hygiene cannot be overstated. A number of studies have shown that improvements in hand hygiene are associated with lower healthcare-associated infection rates, and/or reductions in MDRO transmission and acquisition.<sup>[7-10]</sup> Nonetheless, compliance with hand hygiene recommendations is often poor, with studies reporting rates as low as 20 - 30%.<sup>[11-13]</sup>

The adequate disinfection of hands is easily accomplished by using an alcoholic hand rub, with or without additional disinfectants, e.g. chlorhexidine. The alcohol is responsible for the immediate killing of transient flora on the hands, while the additional agents may exert a residual effect, limiting reacquisition of organisms for a limited period (although the evidence is mainly based on *in vitro* simulations, and the additional benefit of preventing transmission has not been studied well).

Alcohol hand rubs are superior to both ordinary soap and water and medicated soap and water with regard to the efficacy of removal of micro-organisms. This is based on numerous *in vitro* simulations and clinical evidence.<sup>[14]</sup> As an example, a 6-year observation study at a tertiary centre in the USA compared 3 years of medicated soap use with 3 years of alcoholic hand rub use, and showed reductions in acquisition of both methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant Enterococcus (VRE) (21% and

# Your 5 Moments for Hand Hygiene

<b>1</b>	<b>BEFORE TOUCHING A PATIENT</b>	<b>WHEN?</b> Clean your hands before touching a patient when approaching him/her. <b>WHY?</b> To protect the patient against harmful germs carried on your hands.
<b>2</b>	<b>BEFORE CLEAN/ ASEPTIC PROCEDURE</b>	<b>WHEN?</b> Clean your hands immediately before performing a clean/aseptic procedure. <b>WHY?</b> To protect the patient against harmful germs, including the patient's own, from entering his/her body.
<b>3</b>	<b>AFTER BODY FLUID EXPOSURE RISK</b>	<b>WHEN?</b> Clean your hands immediately after an exposure risk to body fluids (and after glove removal). <b>WHY?</b> To protect yourself and the health-care environment from harmful patient germs.
<b>4</b>	<b>AFTER TOUCHING A PATIENT</b>	<b>WHEN?</b> Clean your hands after touching a patient and her/his immediate surroundings, when leaving the patient's side. <b>WHY?</b> To protect yourself and the health-care environment from harmful patient germs.
<b>5</b>	<b>AFTER TOUCHING PATIENT SURROUNDINGS</b>	<b>WHEN?</b> Clean your hands after touching any object or furniture in the patient's immediate surroundings, when leaving – even if the patient has not been touched. <b>WHY?</b> To protect yourself and the health-care environment from harmful patient germs.

World Health Organization | Patient Safety | SAVE LIVES Clean Your Hands

Fig. 1. The World Health Organization's 'Your 5 Moments for Hand Hygiene' campaign.<sup>[14,16]</sup>

41% reductions, respectively).<sup>[15]</sup> This may be due to the greater efficacy of alcohol and improved compliance with hand hygiene, with hand rubs being more easily accessible. Alcohol hand rubs can be manufactured very easily and cheaply if commercial products are not available. One suggested recipe is as follows:<sup>[14]</sup>

- ethanol 96% v/v, 833.3 mL
- hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) 3%, 41.7 mL
- glycerol 98%, 14.5 mL.

Make up to 1 000 mL with distilled water or boiled and cooled water and shake gently to mix the contents.

H<sub>2</sub>O<sub>2</sub> is added to help eliminate contaminating spores in the bulk solutions, and glycerol acts as an emollient to improve

acceptability. Neither of these items plays a role in the actual hand disinfection process.

Other important factors to remember with regard to the use of alcohol-based hand rubs are:

- If hands are visibly soiled, they should first be cleaned with soap and water – before using hand rubs.
- Alcohol is less effective at removing spores; hence, when caring for patients with suspected or confirmed *Clostridium difficile* infections, medicated soap is preferred.

The World Health Organization is promoting the 'Your 5 Moments for Hand Hygiene' campaign (Fig. 1),<sup>[14,16]</sup> which recommends that hand hygiene be practised as follows:

- before patient contact
- before a clean or aseptic procedure

- after contact with body fluids
- after patient contact
- after contact with the patient's environment.

## Measures beyond hand hygiene

Other measures have been advocated to reduce the risk of transmission of MDROs. The CDC<sup>[4]</sup> recommends that all patients colonised or infected with epidemiologically important pathogens (including MDROs) that are spread by contact be placed on contact precautions. This consists of isolating the patient in a room (or placing the patient with others with the same organism), use of gloves for all patient contact, and use of gowns/aprons during patient contact. Where isolation rooms are limited, patients with MDROs may need to be nursed in open wards. However, even there, every effort should be made to implement the glove and gown components of contact precautions. These recommendations are based on expert opinion and knowledge of the transmission mechanism, but there is limited evidence that contact precautions make a significant difference to transmission of MDROs, over and above good hand hygiene and adherence to basic infection control principles.<sup>[5,17-19]</sup> Many studies include multiple interventions (improved staffing levels, education campaigns, hand hygiene campaigns) in addition to contact precautions, and it is difficult to identify which has the greatest impact.

There have been concerns about the impact of isolation on the mental wellbeing of patients.<sup>[20]</sup> The additional burden of having patients on contact precautions may lead to a reduction in HCW compliance with these precautions.<sup>[21]</sup> Even given these concerns, there is evidence that implementation of contact precautions does improve hand hygiene compliance,<sup>[19]</sup> and it is still recommended for patients with MDROs. However, the importance of adequate staffing to sustain contact precautions and standard infection control practices cannot be overstated.

## Screening and decolonisation

There is good evidence that active screening of preoperative patients for MRSA, with decolonisation of carriers, results in reductions in postoperative sepsis caused by MRSA and other pathogens.<sup>[22]</sup> The same effect has been described where all patients were decolonised with nasal mupirocin and chlorhexidine washes without the added cost of screening.<sup>[23]</sup> Chlorhexidine washes alone have also been shown to reduce the

acquisition of MDROs and development of healthcare-associated bacteraemia by 23% and 28%, respectively.<sup>[24]</sup> This raises the question of whether the application of intranasal mupirocin has benefit over chlorhexidine washes. The effect of chlorhexidine washes or wipes on infections with MDR Gram-negative bacteria specifically has not been as well described, and this, together with the descriptions of emerging chlorhexidine resistance, has led to some concerns about the implementation of routine chlorhexidine washes as a decolonisation strategy.<sup>[25]</sup>

Surveillance cultures for carbapenem-resistant Enterobacteriaceae (CRE) have been advocated in a number of reports and recommendations as part of an overall strategy to combat this specific MDRO.<sup>[26,27]</sup> Screening has been a part of successful CRE control measures described in a variety of settings.<sup>[28,29]</sup> However, as before, all these studies have included other interventions and it is very difficult to determine the effect of surveillance cultures alone on CRE control. There is evidence that CRE can be controlled without the use of surveillance cultures,<sup>[18]</sup> and in resource-limited settings in particular it may be more appropriate to increase overall infection control capacity than to focus on screening.

Decolonisation strategies have not been well described for any of the resistant Gram-negative organisms. Some studies have demonstrated emergence of resistance during CRE decolonisation regimens (which usually include colistin),<sup>[30-32]</sup> and this, allied to limited evidence of its efficacy, strongly suggests that decolonisation for multiresistant Gram-negative bacteria should not be attempted. If active screening for CRE carriage is carried out, the main intervention for carriers consists of isolation/cohorting and enhanced infection control measures. The duration of isolation for CRE (and for many other MDROs) is not well defined. CRE can be carried for  $\geq 12$  months,<sup>[33]</sup> and if patients are identified as CRE carriers, it may be valuable to be able to identify them on readmission to hospital. Screening to document elimination of carriage can be attempted; however, it has been suggested that there should be at least two negative cultures (and possibly a negative polymerase chain reaction test) before a known CRE carrier patient can be regarded as no longer carrying this organism.<sup>[34]</sup> This is an area where more research is required.

## Prevention of infection

While many infection control interventions focus on reducing the transmission of organisms, it is as important to identify measures to reduce the risk of infection. Fewer infections translate into reductions in antibiotic use and better patient outcomes. Many professional bodies and groups have adopted the use of 'care bundles', whereby different interventions focused on the same infection are combined and practised as a standard of care. In South Africa, this is being driven by the 'Best Care ... Always' (BCA) campaign (<http://www.bestcare.org.za/home>). The four infections addressed by the BCA bundles are catheter-associated urinary tract infection (CA-UTI), ventilator-associated pneumonia (VAP), central line-associated bloodstream infection (CLABSI), and surgical site infection (SSI).

Although there are some differences in the components of the bundles drawn up by different groups, and there is room for addition or modification, the components of the BCA bundles are as follows:

### CA-UTI

- Avoid unnecessary catheterisation
- Insert catheters using an aseptic technique
- Maintain catheters based on recommended guidelines
- Review the need for catheterisation daily and remove the catheter as soon as possible.

### VAP

- Elevation of the head of the bed to 45° (or 30° if 45° not possible)
- Subglottic drainage of secretions
- Assess readiness for extubation daily
- Oral care and cleaning with chlorhexidine
- Initiation of enteral feeding within 24 - 48 hours of admission.

### CLABSI

- Maximal barrier precautions when inserting line
- Chlorhexidine antiseptic when inserting line
- Selection of optimal catheter insertion site (weighing infection and complication risks)
- Hand hygiene when caring for line
- Review the need for line daily, and remove as soon as possible.

### SSI

- Appropriate antibiotic prophylaxis (timing, choice, duration)
- Appropriate hair removal (clippers or depilatory creams, not shaving)
- Postoperative glucose control
- Maintain postoperative normothermia.

Although some of the abovementioned components may be difficult to implement, the majority are relatively simple, require minimal resources, and when implemented can result in a dramatic reduction in infection. The incidence of SSI after colorectal surgery was 2% in patients who received all six components of a bundle, and 17% in those who received only one component.<sup>[35]</sup> A multicentre before-and-after study in Korea showed a reduction in VAP rates from 4.08/1 000 ventilator-days to 1.16/1 000 ventilator-days after implementation of a VAP bundle.<sup>[36]</sup> Reductions in VAP rates after implementing bundles have been associated with reduced ventilation-days and length of stay in the intensive care unit (ICU).<sup>[37]</sup>

Implementation of CLABSI bundles has resulted in reductions in CLABSI rates of up to 56%;<sup>[38]</sup> despite different studies using different bundles, there is a consistent reduction in the rates of these infections.<sup>[38-40]</sup> Introduction of bundles for CLABSI, VAP and CA-UTI in ICUs of a Saudi Arabian hospital resulted in reductions in the rates of all three infections (86% for VAP, 63% for CLABSI and 50% for CA-UTI).<sup>[41]</sup>

## Conclusion

Infection control has a major role to play in combating the threat of AMR, and many interventions that have proven benefit are relatively inexpensive. Chief among these is hand hygiene. The challenge remains the appropriate implementation of these interventions, and sustaining them once immediate threats are over. Infection control should no longer be seen as the sole responsibility of the infection control practitioner, but that of every HCW.

### References

1. World Health Organization. Antimicrobial Resistance: Global Report on Surveillance 2014. <http://www.who.int/drugresistance/documents/surveillance-report/en/> (accessed 8 April 2015).
2. O'Neil J. Review on Antimicrobial Resistance. Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations 2014. [http://www.jpiamr.eu/wp-content/uploads/2014/12/AMR-Review-Paper-Tackling-a-crisis-for-the-health-and-wealth-of-nations\\_1-2.pdf](http://www.jpiamr.eu/wp-content/uploads/2014/12/AMR-Review-Paper-Tackling-a-crisis-for-the-health-and-wealth-of-nations_1-2.pdf) (accessed 8 April 2015).
3. US Centres for Disease Control and Prevention. Antibiotic Resistance Threats in the United States, 2013. <http://www.cdc.gov/drugresistance/threat-report-2013/> (accessed 8 April 2015).
4. Siegel JD, Rhinehart E, Jackson M, Chiarello L; the Healthcare Infection Control Practices Advisory Committee. Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings, June 2007. <http://www.cdc.gov/hicpac/pdf/isolation/Isolation2007.pdf> (accessed 8 April 2015).
5. Siegel JD, Rhinehart E, Jackson M, Chiarello L; the Healthcare Infection Control Practices Advisory Committee. Management of multidrug-resistant organisms in healthcare settings, 2006. *Am J Infect Control* 2007;35:S165-S193.
6. Ben-David D, Mermel LA, Parenteau S. Methicillin-resistant *Staphylococcus aureus* transmission: The possible importance of unrecognized healthcare worker carriage. *Am J Infect Control* 2008;36(2):93-97. <http://dx.doi.org/10.1016/j.ajic.2007.05.013>

7. Kaier K, Hagist C, Frank U, Conrad A, Meyer E. Two time-series analyses of the impact of antibiotic consumption and alcohol-based hand disinfection on the incidences of nosocomial methicillin-resistant *Staphylococcus aureus* infection and *Clostridium difficile* infection. *Infect Control Hosp Epidemiol* 2009;30(4):346-353. [http://dx.doi.org/10.1086/596605]
8. Stone SP, Fuller C, Savage J, et al. Evaluation of the national Cleanyourhands campaign to reduce *Staphylococcus aureus* bacteraemia and *Clostridium difficile* infection in hospitals in England and Wales by improved hand hygiene: Four year, prospective, ecological, interrupted time series study. *BMJ* 2012;3(344):e3005. [http://dx.doi.org/10.1136/bmj.e3005]
9. Helder OK, Brug J, van Goudoever JB, Looman CW, Reiss IK, Kornelisse RE. Sequential hand hygiene promotion contributes to a reduced nosocomial bloodstream infection rate among very low-birth weight infants: An interrupted time series over a 10-year period. *Am J Infect Control* 2014;42(7):718-722. [http://dx.doi.org/10.1016/j.ajic.2014.04.005]
10. De Angelis G, Cataldo MA, De Waure C, et al. Infection control and prevention measures to reduce the spread of vancomycin-resistant enterococci in hospitalized patients: A systematic review and meta-analysis. *J Antimicrob Chemother* 2014;69(5):1185-1192. [http://dx.doi.org/10.1093/jac/dkt525]
11. Marra AR, Edmond MB. New technologies to monitor healthcare worker hand hygiene. *Clin Microbiol Infect* 2014;20(1):29-33. [http://dx.doi.org/10.1111/1469-0691.12458]
12. Tenna A, Stenehjem EA, Margoles L, Kacha E, Blumberg HM, Kempker RR. Infection control knowledge, attitudes, and practices among healthcare workers in Addis Ababa, Ethiopia. *Infect Control Hosp Epidemiol* 2013;34(12):1289-1296. [http://dx.doi.org/10.1086/673979]
13. Block L, Habicht R, Oluyadi FO, et al. Variability in hand hygiene practices among internal medicine interns. *Am J Infect Control* 2013;41(11):1107-1108. [http://dx.doi.org/10.1016/j.ajic.2013.03.303]
14. World Health Organization (WHO). WHO guidelines on hand hygiene in healthcare 2009. [http://whqlibdoc.who.int/publications/2009/9789241597906\\_eng.pdf](http://whqlibdoc.who.int/publications/2009/9789241597906_eng.pdf) (accessed 8 April 2015).
15. Gordin FM, Schultz ME, Huber RA, Gill JA. Reduction in nosocomial transmission of drug-resistant bacteria after introduction of an alcohol-based handrub. *Infect Control Hosp Epidemiol* 2005;26(7):650-653.
16. Sax H, Allegranzi B, Uçkay I, Larson E, Boyce J, Pittet D. 'My five moments for hand hygiene': A user-centred design approach to understand, train, monitor and report hand hygiene. *Journal of Hospital Infection* 2007;67:9-21.
17. Huskins WC, Huckabee CM, O'Grady NP, et al. Intervention to reduce transmission of resistant bacteria in intensive care. *N Engl J Med* 2011;364:1407-1418. [http://dx.doi.org/10.1056/NEJMoa1000373]
18. Kim NH, Han WD, Song KH, et al. Successful containment of carbapenem-resistant Enterobacteriaceae by strict contact precautions without active surveillance. *American Journal of Infection Control* 2014;42(12):1270-1273. [http://dx.doi.org/10.1016/j.ajic.2014.09.004]
19. Harris AD, Pinales L, Belton B, et al. Universal glove and gown use and acquisition of antibiotic-resistant bacteria in the ICU: A randomized trial. *J Am Med Assoc* 2013;310(15):1571-1580.
20. Bearman G, Stevens MP. Control of drug-resistant pathogens in endemic settings: Contact precautions, controversies, and a proposal for a less restrictive alternative. *Curr Infect Dis Rep* 2012;14(6):620-626. [http://dx.doi.org/10.1007/s11908-012-0299-8]
21. Dhar S, Marchaim D, Tansek R, et al. Contact precautions: More is not necessarily better. *Infect Control Hosp Epidemiol* 2014;35(3):213-221. [http://dx.doi.org/10.1086/675294]
22. Van Rijen M, Bonten M, Wenzel R, Kluytmans J. Mupirocin ointment for preventing *Staphylococcus aureus* infections in nasal carriers. *Cochrane Database Syst Rev* 2008;8(4):CD006216. [http://dx.doi.org/10.1002/14651858.CD006216.pub2]
23. Huang SS, Septimus E, Kleinman K, et al. Targeted versus universal decolonization to prevent ICU infection. *N Engl J Med* 2013;368(24):2255-2265. [http://dx.doi.org/10.1056/NEJMoa1207290]
24. Climo MW, Yokoe DS, Warren DK, et al. Effect of daily chlorhexidine bathing on hospital-acquired infection. *N Engl J Med* 2013;368(6):533-542. [http://dx.doi.org/10.1056/NEJMoa1113849]
25. Tacconelli E, Cataldo MA, Dancer SJ, et al. ESCMID guidelines for the management of the infection control measures to reduce transmission of multidrug-resistant Gram-negative bacteria in hospitalized patients. *Clin Microbiol Infect* 2014;20(Suppl 1):1-55. [http://dx.doi.org/10.1111/1469-0691.12427]
26. Carmeli Y, Akova M, Cornaglia G, et al. Controlling the spread of carbapenemase-producing Gram-negatives: Therapeutic approach and infection control. *Clin Microbiol Infect* 2010;16(2):102-111. [http://dx.doi.org/10.1111/j.1469-0691.2009.03115.x]
27. Centers for Disease Control and Prevention (CDC). Guidance for control of infections with carbapenem-resistant or carbapenemase-producing Enterobacteriaceae in acute care facilities. *MMWR Morb Mortal Wkly Rep* 2009;58(10):256-260.
28. Poulou A, Voulgari E, Vrioni G, et al. Imported *Klebsiella pneumoniae* carbapenemase-producing *K. pneumoniae* clones in a Greek hospital: Impact of infection control measures for restraining their dissemination. *J Clin Microbiol* 2012;50(8):2618-2623. [http://dx.doi.org/10.1128/JCM.00459-12]
29. Schwaber MJ, Carmeli Y. An ongoing national intervention to contain the spread of carbapenem-resistant Enterobacteriaceae. *Clin Infect Dis* 2014;58(5):697-703. [http://dx.doi.org/10.1093/cid/cit795]
30. Oren I, Sprecher H, Finkelstein R, et al. Eradication of carbapenem-resistant Enterobacteriaceae gastrointestinal colonization with non-absorbable oral antibiotic treatment: A prospective controlled trial. *Am J Infect Control* 2013;41(12):1167-1172. [http://dx.doi.org/10.1016/j.ajic.2013.04.018]
31. Lubbert C, Fauchoux S, Becker-Rux D, et al. Rapid emergence of secondary resistance to gentamicin and colistin following selective digestive decontamination in patients with KPC-2-producing *Klebsiella pneumoniae*: A single-centre experience. *International Journal of Antimicrobial Agents* 2013;42(6):565-570. [http://dx.doi.org/10.1016/j.ijantimicag.2013.08.008]
32. Brink AJ, Coetzee J, Corcoran C, et al. Emergence of OXA-48 and OXA-181 carbapenemases among Enterobacteriaceae in South Africa and evidence of *in vivo* selection of colistin resistance as a consequence of selective decontamination of the gastrointestinal tract. *J Clin Microbiol* 2013;51(1):369-372. [http://dx.doi.org/10.1128/JCM.02234-12]
33. Zimmerman FS, Assous MV, Bdolah-Abram T, Lachish T, Yinnon AM, Wiener-Well Y. Duration of carriage of carbapenem-resistant Enterobacteriaceae following hospital discharge. *Am J Infect Control* 2013;41(3):190-194. [http://dx.doi.org/10.1016/j.ajic.2012.09.020]
34. Feldman N, Adler A, Molshatzki N, et al. Gastrointestinal colonization by KPC-producing *Klebsiella pneumoniae* following hospital discharge: Duration of carriage and risk factors for persistent carriage. *Clinical Microbiology and Infection* 2013;19(4):E190-E196. [http://dx.doi.org/10.1111/1469-0691.12099]
35. Waits SA, Fritze D, Banerjee M, et al. Developing an argument for bundled interventions to reduce surgical site infection in colorectal surgery. *Surgery* 2014;155(4):602-606. [http://dx.doi.org/10.1016/j.surg.2013.12.004]
36. Eom JS, Lee MS, Chun HK, et al. The impact of a ventilator bundle on preventing ventilator-associated pneumonia: A multicenter study. *Am J Infect Control* 2014;42(1):34-37. [http://dx.doi.org/10.1016/j.ajic.2013.06.023]
37. Rello J, Afonso E, Lisboa T, et al. A care bundle approach for prevention of ventilator-associated pneumonia. *Clin Microbiol Infect* 2013;19(4):363-369. [http://dx.doi.org/10.1111/j.1469-0691.2012.03808.x]
38. Jeffries HE, Mason W, Brewer M, et al. Prevention of central venous catheter-associated bloodstream infections in pediatric intensive care units: A performance improvement collaborative. *Infect Control Hosp Epidemiol* 2009;30(7):645-651. [http://dx.doi.org/10.1086/598341]
39. Miller MR, Niedner ME, Huskins WC, et al. Reducing PICU central line-associated bloodstream infections: 3-year results. *Pediatrics* 2011;128(5):e1077-e1083. [http://dx.doi.org/10.1542/peds.2010-3675]
40. Schulman J, Stricof R, Stevens TP, et al. Statewide NICU central-line-associated bloodstream infection rates decline after bundles and checklists. *Pediatrics* 2011;127(3):436-444. [http://dx.doi.org/10.1542/peds.2010-2873]
41. Al-Tawfiq JA, Amalraj A, Memish ZA. Reduction and surveillance of device-associated infections in adult intensive care units at a Saudi Arabian hospital, 2004-2011. *Int J Infect Dis* 2013;17(12):e1207-e1211. [http://dx.doi.org/10.1016/j.ijid.2013.06.015]