



Sacred Heart University
DigitalCommons@SHU

Speech-Language Pathology Faculty Publications

Speech-Language Pathology

1-2016

Brief Report: A Mobile Application to Treat Prosodic Deficits in Autism Spectrum Disorder and Other Communication Impairments: A Pilot Study

Elizabeth Schoen Simmons
Yale Child Study Center, elizabeth.schoen@yale.edu

Rhea Paul
Sacred Heart University, paulr4@sacredheart.edu

Frederick Shic
Yale Child Study Center

Follow this and additional works at: http://digitalcommons.sacredheart.edu/speech_fac

 Part of the [Speech Pathology and Audiology Commons](#)

Recommended Citation

Simmons, E. S., Paul, R. & Shic, F. (2016). Brief report: a mobile application to treat prosodic deficits in autism spectrum disorder and other communication impairments: a pilot study. *Journal of Autism and Developmental Disorders*, 46(1), 320-327. doi: 10.1007/s10803-015-2573-8

This Peer-Reviewed Article is brought to you for free and open access by the Speech-Language Pathology at DigitalCommons@SHU. It has been accepted for inclusion in Speech-Language Pathology Faculty Publications by an authorized administrator of DigitalCommons@SHU. For more information, please contact ferribyp@sacredheart.edu, lysobeyb@sacredheart.edu.

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

ArticleTitle	Brief Report: A Mobile Application to Treat Prosodic Deficits in Autism Spectrum Disorder and Other Communication Impairments: A Pilot Study	
Article Sub-Title		
Article CopyRight	Springer Science+Business Media New York (This will be the copyright line in the final PDF)	
Journal Name	Journal of Autism and Developmental Disorders	
Corresponding Author	Family Name	Simmons
	Particle	
	Given Name	Elizabeth Schoen
	Suffix	
	Division	Child Study Center
	Organization	Yale University
	Address	40 Temple Street, Suite 7D, 06510, New Haven, CT, USA
	Email	elizabeth.schoen@yale.edu
Author	Family Name	Paul
	Particle	
	Given Name	Rhea
	Suffix	
	Division	Department of Speech-Language Pathology
	Organization	Sacred Heart University
	Address	Fairfield, CT, USA
	Email	
Author	Family Name	Shic
	Particle	
	Given Name	Frederick
	Suffix	
	Division	Child Study Center
	Organization	Yale University
	Address	40 Temple Street, Suite 7D, 06510, New Haven, CT, USA
	Email	
Schedule	Received	
	Revised	
	Accepted	
Abstract	This study examined the acceptability of a mobile application, <i>SpeechPrompts</i> , designed to treat prosodic disorders in children with ASD and other communication impairments. Ten speech-language pathologists (SLPs) in public schools and 40 of their students, 5–19 years with prosody deficits participated. Students received treatment with the software over eight weeks. Pre- and post-treatment speech samples and student engagement data were collected. Feedback on the utility of the software was also obtained. SLPs implemented the software with their students in an authentic education setting. Student engagement ratings indicated students' attention to the software was maintained during treatment. Although more testing is	

warranted, post-treatment prosody ratings suggest that *SpeechPrompts* has potential to be a useful tool in the treatment of prosodic disorders.

Keywords (separated by '-') Autism - Technology - Intervention - Prosody - Speech

Footnote Information

2 **Brief Report: A Mobile Application to Treat Prosodic Deficits**
3 **in Autism Spectrum Disorder and Other Communication**
4 **Impairments: A Pilot Study**

5 Elizabeth Schoen Simmons¹ · Rhea Paul² · Frederick Shic¹

6
7 © Springer Science+Business Media New York 2015

8 **Abstract** This study examined the acceptability of a
9 mobile application, *SpeechPrompts*, designed to treat pro-
10 sodic disorders in children with ASD and other commu-
11 nication impairments. Ten speech-language pathologists
12 (SLPs) in public schools and 40 of their students,
13 5–19 years with prosody deficits participated. Students
14 received treatment with the software over eight weeks. Pre-
15 and post-treatment speech samples and student engagement
16 data were collected. Feedback on the utility of the software
17 was also obtained. SLPs implemented the software with
18 their students in an authentic education setting. Student
19 engagement ratings indicated students' attention to the
20 software was maintained during treatment. Although more
21 testing is warranted, post-treatment prosody ratings suggest
22 that *SpeechPrompts* has potential to be a useful tool in the
23 treatment of prosodic disorders.

25 **Keywords** Autism · Technology · Intervention ·
26 Prosody · Speech

27 **Introduction**

28 For the majority of individuals with autism spectrum dis-
29 order (ASD) who acquire spoken language, expressive
30 prosody—the rhythm, stress, and intonation of speech—is
31 **AQ1** among the most noticeable and chronic impairments

(Baltaxe and Simmons 1985; DeMyer et al. 1973; Kanner 32
1971; Lyons et al. 2014; Rutter and Lockyer 1967; Shri- 33
berg et al. 2001). Prosodic deficits have been shown to 34
impact how listeners perceive the social and communica- 35
tive competence of high-functioning individuals with ASD 36
(Paul et al. 2005) and those with intellectual disability 37
(Shriberg and Widder 1990). Deficits in these supraseg- 38
mental features of speech also impede social interaction 39
and limit participation in vocational, recreational and 40
learning activities (Lewis et al. 2004; Wilson and Warton 41
2006). Prosodic deficits are also observed in children with 42
other communication disorders, as well as those with ASD 43
(Catterall et al. 2006; Marshall et al. 2009; Stojanovik et al. 44
2007; Wells and Peppé 2003). 45

A limited number of intervention strategies to treat these 46
deficits exist, with the majority of these lacking empirical 47
support. Diehl and Paul (2009) and Peppé (2009) reviewed 48
current prosodic intervention literature and reported that 49
methodological issues (e.g., small sample sizes) made it 50
difficult to interpret and generalize the findings. 51

The proliferation of mobile technology, including 52
tablets and smartphones, provides speech-language 53
pathologists (SLPs) with another medium to deliver inter- 54
vention. A recent survey of approximately 300 school- 55
based SLPs (Fernandes 2011) reported that a majority 56
owned either a tablet or smartphone and used their personal 57
device during intervention sessions with students. Emerg- 58
ing research suggests higher levels of student engagement 59
during sessions that use technology than those using tra- 60
ditional materials (American Speech-Language Hearing 61
Association 2011). 62

A small body of literature suggests that mobile tech- 63
nology is a valuable tool in the treatment of communica- 64
tion deficits and behavioral issues commonly observed in 65
students with ASD and other communication disorders. 66

A1 ✉ Elizabeth Schoen Simmons
A2 elizabeth.schoen@yale.edu

A3 ¹ Child Study Center, Yale University, 40 Temple Street, Suite
A4 7D, New Haven, CT 06510, USA

A5 ² Department of Speech-Language Pathology, Sacred Heart
A6 University, Fairfield, CT, USA

67 Increased frequency of peer initiations and response to peer
68 bids were observed after iPod training in a group of ado-
69 lescents with autism using an iPod Touch loaded with an
70 augmentative and alternative communication (AAC)
71 application (Carpenter 2012). In a single subject study, the
72 use of an iPad was shown to decrease levels of challenging
73 behavior while increasing academic engagement in two
74 students with autism spectrum disorder (Neely et al. 2013).
75 While the literature suggests using technology can improve
76 engagement, there is a dearth of research regarding the
77 utility of technology for improving specific communication
78 skills, such as prosody, in these populations.

79 The present study's primary aim was to assess the
80 acceptability of an application, *SpeechPrompts*, for mobile
81 devices in the treatment of prosodic disorders in school-age
82 children with ASD and those with other communication
83 impairments. A secondary aim was to provide preliminary
84 evaluation of the potential utility of this application for
85 improving prosody skills in students with prosodic deficits.

86 Methods

87 Participants

88 *Speech-Language Pathologists*

89 Inclusion criteria for SLPs included: (1) licensure by the
90 department of public health in the State of Connecticut (2)
91 certification by the American-Speech-Language-Hearing
92 Association and (3) caseloads including students who had
93 prosodic difficulties. Ten (10) SLPs were enrolled in this
94 pilot study. Each was asked to complete an online survey to
95 collect information about work setting, familiarity with
96 tablet devices and any training already received on assis-
97 tive technology (see Table 1).

98 *Student Participants*

99 Each SLP recruited four students from her caseload who
100 met the following inclusion criteria: (1) enrollment in
101 speech and language intervention as part of special edu-
102 cation services, (2) speech containing full sentences, and
103 (3) exhibiting prosodic difficulties secondary to ASD or
104 other communication disorder. A total of 40 students, aged
105 5 through 19 years, met study criteria and were enrolled for
106 participation. Approximately 67.5 % of the students had a
107 school-based classification of ASD on their individualized
108 education plan (IEP); the remainder were classified with
109 other impairments (e.g., speech and language impairment,
110 intellectual disability, multiple disabilities, traumatic brain
111 injury). Diagnostic information was not available at an
112 individual level for all students due to the study's IRB

Table 1 SLPs' clinical experiences

	N = 10 (%)
Current employment setting*	
Preschool	30
Elementary school	80
Middle school	40
High school	20
Years in current position	
0–5 years	20
6–10 years	40
11–15 years	20
16–20 years	0
≥21 years	20
Experience with tablets (e.g., iPads)	
Minimal experience	20
Some experience	20
Significant experience	60

* Percentage >100 as a subset of SLPs work in more than one setting

format; therefore, a subset analysis for 12 students with 113
ASD who had linkable diagnostic and study data is provided 114
in the appendices for greater specificity of information 115
for students with ASD. A wide distribution in the ages 116
of students was included to determine whether both 117
younger and older students would be engaged with the 118
software. A majority of the students (72.5 %) were assessed 119
as having impairments in two or more prosodic 120
domains as rated by their SLP. See Table 2. 121

122 Procedures

123 *Software*

124 *SpeechPrompts* was developed for iOS devices (e.g., iPad); 124
its main function was to provide a visual representation of 125
the prosodic features of speech. It contained two primary 126
features. The *VoiceMatch* feature allowed the SLP to 127
record a short target phrase, then view a waveform visu- 128
alization of the phrase. The student would then attempt to 129
produce a waveform matching the target by adjusting his/ 130
her speaking rate and/or stress (see Fig. 1). The second 131
feature, *VoiceChart*, provided real-time feedback on 132
speaking volume by displaying visual cues to monitor and 133
adjust the volume of speech. Slider controls were used by 134
the SLP to adjust the target speaking thresholds during 135
instruction. This feature had customizable visuals for 136
younger and older participants (i.e., teddy bears and written 137
words, respectively) (see Fig. 2). 138

139 The software was designed with usage-tracking
140 embedded within the application. This tool automatically

Table 2 Student participant characteristics

	N = 40
Gender	
Male	31
Female	9
Mean age in years (<i>SD</i>)	9.63 (3.70)
Grade level	
Elementary (PreK–4th)	22
Middle school (5th–8th)	13
High school (9th–12th)	5
Diagnosis based on IEP ^a	
ASD	27
Speech and language impairment	7
Intellectual disability	3
Traumatic brain injury	1
Multiple disabilities	1
Other health impairment	1
Number of students with prosodic impairments, by domain, as rated by SLP ^b	
Rate/rhythm	27
Stress	29
Volume	28

^a Individualized education plan

^b A subset of students were rated as having impairments in more than one prosodic domain




Fig. 1 Screenshot of waveforms generated by *VoiceMatch* feature. The *top* waveform is a sentence produced by the SLP while the bottom waveform is the student’s production of the same target sentence. The small microphone, scissors and speaker icons control recording, editing and volume functionality within the app




Fig. 2 Screenshot of *VoiceChart* with customizable visual supports and volume thresholds. The *top* half of the window provides the visual feedback. On the *left* is a teddy bear for younger students and on the *right* written words for older students. The *bottom* half of the window allows the SLP to move the sliders to set an appropriate speaking volume level

141 compiled usage statistics for each SLP including duration
 142 of treatment sessions, frequency of application use, and
 143 ranges of features accessed during each session. The
 144 application was designed in collaboration with the authors
 145 and a small software company. The authors received no
 146 financial compensation from the company.

Speech Samples

147

Five-minute speech samples were collected by each SLP, pre- 148
 and post-treatment, from student participants; these samples 149
 were audio recorded for later coding. A topic prompt, *tell me* 150
about your family and everyone who lives with you, was 151

152 provided. The SLP rated each sample on the following pro-
 153 sodic features (a) rate, (b) stress in words, (c) stress in sen-
 154 tences and (d) intensity. Each SLP also provided a global
 155 intonation summary rating for each sample. A scale of 0
 156 (typical prosody), 1 (mildly atypical prosody), or 2 (clearly
 157 atypical prosody) was used for these ratings.

158 *Speech Sample Reliability*

159 A randomly selected 20 % of speech samples were re-coded
 160 by a second coder blind to whether the sample was collected
 161 pre-treatment or post-treatment. Inter-rater reliability was
 162 established using Cohen's Kappa coefficient. Inter-rater
 163 agreement of 0.68 was obtained across the prosodic param-
 164 eters of global intonation, rate, and stress, indicating sub-
 165 stantial agreement (Viera and Garrett 2005). Inter-rater
 166 reliability could not be established for intensity as sample
 167 collection did not include calibration for baseline intensity.

168 *SLP Training*

169 Each SLP received an iPad 2 (iOS 6.0) preloaded with
 170 *SpeechPrompts*. A 20-min training tutorial was delivered
 171 by the research coordinator, which covered the use of the
 172 main features, enabled the SLP to navigate through the
 173 application and to answer any questions that arose during the
 174 tutorial session. The coordinator was available for the dura-
 175 tion of the study to provide technical assistance as needed.

176 *Intervention*

177 The *SpeechPrompts* software was presented to the students
 178 as part of their speech and language services that took place
 179 in their local school. The SLPs were instructed to use the
 180 application with four selected students at least once each
 181 week for 8 weeks.

182 *Student Engagement Questionnaire*

183 Each SLP completed a rating scale to assess the student's
 184 engagement while using the software following each
 185 treatment session. For each student, SLPs rated (1) enjoy-
 186 ment of the software, (2) attention while using the appli-
 187 cation, (3) consistent attempts to produce responses and (4)
 188 off-task behavior. Numerical ratings ranged from 1
 189 (Strongly Agree/Highly Engaged) through 5 (Strongly
 190 Disagree/Not engaged).

191 *Post-Study Questionnaire*

192 Each SLP completed a questionnaire containing Likert-
 193 scale ratings and open-ended questions regarding experi-
 194 ences with the software at study conclusion.

Results 195

Software Utilization 196

The mean number of sessions, or how many times the SLPs
 197 used the software, across student participants ranged from
 198 1 to 12 sessions with a mean of 4.7 sessions ($SD = 2.79$)
 199 although they had been asked to use the software at least
 200 one time a week for 8 weeks (see Discussion). Session
 201 length ranged from five to 90 min with a mean of
 202 21.25 min ($SD = 11.82$ min). Feature usage from the data-
 203 tracking component of the software revealed that *Voice-*
 204 *Match* and *VoiceChart* features were used 52.9 and 47.1 %
 205 of time spent with the software, respectively. 206

To ascertain whether clinical experience was related to
 207 software utilization (i.e., frequency and duration of inter-
 208 vention sessions), bivariate Pearson's correlations were
 209 computed between the SLPs' number of years in their
 210 current position and both the total number of intervention
 211 sessions conducted as well as total number of treatment
 212 minutes completed. Since the number of treatment minutes
 213 was highly correlated with number of treatment sessions
 214 ($r = .81, p = .005$), only treatment minutes was used in
 215 this analysis. There was no significant relationship between
 216 number of SLPs' years in current position and total number
 217 of treatment minutes received by student participants
 218 ($r = .259, p = .470$). 219


Student Engagement 220

A total of 188 student engagement questionnaires were
 221 collected. The number of students with mean scores ≤ 3
 222 across sessions in each engagement category, indicating
 223 high levels of engagement, were tallied to derive pro-
 224 portions. These proportions suggest that the students
 225 enjoyed the *SpeechPrompts* sessions (92.5 %; 37/40 stu-
 226 dents; $M = 1.66, SD = 0.67$), maintained attention dur-
 227 ing the sessions (87.5 %; 35/40 students; $M = 1.74,$
 228 $SD = 0.80$), provided consistent responses to stimuli
 229 (87.5 %; 35/40 students; $M = 1.78, SD = 0.80$) and did
 230 not produce maladaptive behaviors (85.0 %; 34/40 stu-
 231 dents, $M = 1.79, SD = 0.93$) during the sessions. Ratings
 232 were stable on the questionnaires from the first to final
 233 sessions (see Fig. 3). 234

SLP Feedback 235

Post-study surveys completed by all participating SLPs
 236 revealed that the majority (≥ 80 %) found the software (1)
 237 enjoyable, (2) easy to use (3) functional and (4) resulted in
 238 positive changes to students' prosody. All of the SLPs
 239 ($N = 10; 100$ %) reported feeling comfortable recom-
 240 mending the software to colleagues. 241

Fig. 3 Mean student engagement ratings from the first session to the last session are plotted over time. SLPs rated student's engagement from 1 (highly engaged) to 5 (not engaged). No student received a rating of 4 or 5. Low, stable ratings across sessions illustrate high engagement throughout the duration of treatment. Diminishing maladaptive behaviors during the course of treatment are also illustrated here




242 **Speech Sample Ratings**

243 Pre- and post-treatment prosody ratings were assigned to
 244 speech samples obtained from 32 of the 40 student partici-
 245 pants. Speech samples were not collected from the remaining
 246 8 students due to absenteeism, clinician error and equipment
 247 malfunction. A mean pre-treatment prosody rating was calcu-
 248 lated across the four main prosodic categories: rate, stress in
 249 words, stress in sentences and intensity. Students' mean
 250 prosody rating ranged from 0.25 to 2.00 with an average mean
 251 rating of 1.08 (*SD* = 0.44) across these constructs. Paired *t*-
 252 tests were used to compare pre- and post-treatment prosody

253 ratings for the four broad prosodic categories and the sum-
 254 mary category. A lower mean score, indicating improved
 255 prosodic performance, was observed in each domain (Stress
 256 in Words, $p = .012$, $d = 0.48$; Stress in Sentences, $p = .001$;
 257 $d = 0.77$; Intensity, $p = .001$, $d = 0.77$; Global Intonation,
 258 $p = .001$, $d = 0.71$) with the exception of *Rate* ($p = .100$).
 259 Figure 4 illustrates the prosody ratings for each prosodic
 260 category. No relationship was observed between change in
 261 the Global Intonation prosody rating from pre-treatment to
 262 post-treatment and number of treatment minutes received
 263 ($r = .16$; $p = .394$), potentially reflecting heterogeneity of
 264 learning in the sample.

Fig. 4 Pre- and post-prosody ratings derived from speech samples coded by SLPs. Error bars represent ± 1 SE. *ns *p* value



265 **ASD Specific Findings**

266 The same analyses were completed for a subset of 12
 267 participants, for whom diagnosis and treatment data could
 268 be linked, are reported in the appendices. The mean num-
 269 ber of intervention sessions across these participants ranged
 270 from 2 to 10 sessions with a mean of 5.83 sessions
 271 ($SD = 2.41$). Session length ranged from 10 to 30 min
 272 with a mean session lasting 25.99 min ($SD = 6.25$).

273 **Discussion**

274 The primary aim of this study was to evaluate the feasi-
 275 bility and acceptability of *SpeechPrompts*, a mobile
 276 application that provides a visual representation of the
 277 suprasegmentals of the speech signal to treat prosodic
 278 deficits. Although not designed to meet the standards of a
 279 randomized controlled trial, this study meets criteria for an
 280 adequate intervention research report based on the guide-
 281 lines defined by Reichow et al. (2008), with quality indi-
 282 cators (including description of both participant and
 283 interventionist, operational and replicable descriptions of
 284 dependent measures, a clear link between the research
 285 question and data analysis, and use of appropriate units of
 286 measurement) well documented within this report.

287 Results of this pilot study suggest that SLPs were able to
 288 use the application in an authentic educational setting with
 289 students who exhibit prosodic impairments. SLPs from our
 290 study reported a high level of familiarity with tablets, as
 291 other reports on the use of mobile technology among
 292 clinicians suggest (Fernandes 2011). Even those SLPs who
 293 reported little experience were able to utilize the applica-
 294 tion with their students.

295 Although prosodic impairments are observed in multiple
 296 clinical populations (Staum 1987; Wells and Peppé 2003;
 297 Catterall et al. 2006), the majority of students who partic-
 298 ipated in this study had a diagnosis of ASD. The experience
 299 of children with other clinical diagnoses in our sample,
 300 however, suggested that this application might be useful for
 301 a range of disorders, not solely ASD. Measures of student
 302 engagement reported by the SLPs suggest that the appli-
 303 cation captures student attention, is enjoyable and elicits
 304 consistent responses in a diverse group of students. Stable
 305 student engagement ratings suggest that students continued
 306 to attend to the software and provided responses throughout
 307 treatment, not only during the first session, suggesting the
 308 results were unlikely due to a “novelty” effect alone.
 309 Moreover, maladaptive behaviors were reported to dimin-
 310 ish over the course of treatment.

311 Lastly, data collected from SLPs about their responses
 312 to the software at the end of the study indicated that they
 313 liked the software, thought it was functional and enjoyable

for their students and that they felt comfortable recom-
 mending the application to colleagues.

A secondary aim of this research was to assess the effi-
 cacy of the software when implemented by licensed clini-
 cians in authentic settings. Although preliminary in nature,
 results suggest that *SpeechPrompts*, even in low doses, can
 be useful in the treatment of prosodic impairment in stu-
 dents with communication disorders, as evidenced by
 changes in prosodic functioning documented in this sample.

Although asked to use *SpeechPrompts* at least once a
 week for 8 weeks, most SLPs used it less than this, perhaps
 because of conflicting demands from other IEP goals. The
 relatively positive changes seen in prosodic ratings of
 speech, even at this low dose of intervention, suggest that
 use of *SpeechPrompts* has a potential for efficacy, although
 caution is warranted in interpreting the results, since SLPs
 were not blind to treatment status. Nonetheless, the question
 of adequate dosage remains an unanswered question for this
 intervention, as it does for many speech-language inter-
 ventions, and further research is needed to resolve it.

Additionally, it may be possible to use the application to
 address prosodic production while working on other language
 goals. For example, the *VoiceChart* feature could be used
 while practicing conversational skills. *VoiceMatch* feature
 could be used while teaching specific language targets. Again,
 more research is needed to determine whether working on
 multiple goals simultaneously is an effective strategy.

Our primary goal was to assess acceptability; therefore, no
 intervention control group was included, limiting our ability
 to measure the efficacy of the *SpeechPrompts* treatment. Still,
 improvements from pre- to post-treatment were observed,
 suggesting a more controlled trial is warranted. Subsequent
 iterations of our work will address this omission as well as the
 need for (1) secondary, blind clinical observation ratings
 obtained independently of the treating clinician to control for
 bias; (2) a measure of treatment fidelity to ensure SLPs are
 using the software appropriately; (3) more nuanced statistical
 analyses addressing how individual characteristics (e.g. IQ or
 treatment dosages) impact outcome measures; (4) in-depth
 examination of the relationship between changes in prosody
 and treatment dosages; and (5) new application capabilities
 for addressing other prosodic domains such as pitch and for
 providing more in-depth visualizations of speech.

Although further research is needed to rigorously eval-
 uate the efficacy of the application, preliminary results
 suggest that *SpeechPrompts* provides SLPs with an addi-
 tional tool in their repertoire to address mild to moderate
 prosodic difficulties commonly observed in children with
 ASD and with other communication impairments, for
 which there are currently few validated treatment approa-
 ches. This research adds to the sparse literature regarding
 the treatment of prosody deficits (Peppé 2009) in school
 age students with ASD and other communication disorders.

367 **Acknowledgments** This project was supported by the Small Business
 368 Innovation Research (SBIR) program of the Department of
 369 Education (ED-IES-12-C-0043; ED-IES-13-C-0046). We would like
 370 to thank Robert and Michael Tedesco of HandHold Adaptive LLC for
 371 the provision of the software, Carla Wall for her assistance with data
 372 collection and the speech-language pathologists and students who
 373 made the study possible.

374 **Appendixes**

375 See Tables 3, 4.
 376 See Fig. 5.


Table 3 ASD subset characteristics

	<i>n</i> = 12
Gender	
Male	11 (91.67 %)
Female	1 (8.33 %)
Mean age in years (SD)	8.25 (3.25)
Age range	6–12 years

Table 4 ASD subset prosody ratings

<i>n</i> = 12	Mean pre-treatment rating (SD)	Mean post-treatment rating (SD)	<i>p</i>	<i>d</i>
Rate	0.50 (0.67)	0.33 (0.49)	.116	–
Stress in words	0.50 (0.52)	0.42 (0.51)	.339	–
Stress in sentences	1.33 (0.49)	0.92 (0.51)	.017	0.80
Intensity	0.75 (0.87)	0.33 (0.65)	.017	0.90
Global intonation	1.17 (0.58)	0.75 (0.62)	.017	0.81

Fig. 5 Mean student engagement ratings from the first session to the last session are plotted over time for subset of 15 students with ASD. SLPs rated student’s engagement from 1 (highly engaged) to 5 (not engaged). No student received a rating of 4 or 5. Low, stable ratings across sessions illustrate high engagement throughout the duration of treatment



References

American Speech-Language Hearing Association. (2011). *Applications (apps) for speech-language pathology practice*. Retrieved from <http://www.asha.org/SLP/schools/Applications-for-Speech-Language-Pathology-Practice>

Baltaxe, C., & Simmons, J. (1985). Prosodic development in normal and autistic children. In E. Schopler & G. Mesibov (Eds.), *Communication problems in autism* (pp. 95–125). New York: Plenum Press.

Carpenter, L. A. (2012). *The effect of a peer-mediated intervention on the social communicative behavior of adolescents with autism using a dynamic display voice output communication aid*. Unpublished manuscript, Department of Special Education, California State University, Los Angeles, CA.

Catterall, C., Howard, S., Stojanovik, V., Szczerbinski, M., & Wells, B. (2006). Investigating prosodic ability in Williams syndrome. *Clinical Linguistics and Phonetics*, 20, 531–538.

DeMyer, M., Barton, S., DeMyer, W., Norton, J., Allen, J., & Stelle, R. (1973). Prognosis in autism: A follow-up study. *Journal of Autism and Childhood Schizophrenia*, 3, 199–246.

Diehl, J., & Paul, R. (2009). The assessment and treatment of prosodic disorders and neurological theories of prosody. *International Journal of Speech-Language Pathology*, 11, 287–292.

Fernandes, B. (2011). iTherapy: The revolution of mobile devices within the field of speech therapy. *SIG 16 Perspectives on School-Based*, 12, 35–40.

Kanner, L. (1971). Follow-up of eleven autistic children, originally reported in 1943. *Journal of Autism and Childhood Schizophrenia*, 2, 119–145.

Lewis, B., Freebairn, B., Hansen, A., & Iyengar, S. (2004). School-age follow-up of children with childhood apraxia of speech. *Language, Speech, and Hearing Services in Schools*, 35, 122–140.

Lyons, M., Simmons, E., & Paul, R. (2014). Prosodic development in middle childhood and adolescence in high functioning autism. *Autism Research*, 7, 181–196.

Marshall, C. R., Harcourt-Brown, S., Ramus, F., & van der Lely, H. J. K. (2009). The link between prosody and language skills in children with specific language impairment (SLI) and/or dyslexia. *International Journal of Language & Communication Disorders*, 44, 466–488.

Author Proof

- 419 Neely, L., Rispoli, M., Camargo, S., Davis, H., & Boles, M. (2013).
 420 The effect of instructional use of an iPad® on challenging
 421 behavior and academic engagement for two students with
 422 autism. *Research in Autism Spectrum Disorders*, 7, 509–516.
- 423 Paul, R., Shriberg, L. D., McSweeney, J., Cicchetti, D., Klin, A., &
 424 Volkmar, F. (2005). Brief report: Relations between prosodic
 425 performance and communication and socialization ratings in
 426 high functioning speakers with autism spectrum disorders.
 427 *Journal of Autism and Developmental Disorders*, 35, 861–869.
- 428 Peppé, S. (2009). Why is prosody in speech-language pathology so
 429 difficult? *International Journal of Speech-Language Pathology*,
 430 11, 259–271.
- 431 Reichow, B., Volkmar, F. R., & Cicchetti, D. V. (2008). Development
 432 of the evaluative method for evaluating and determining
 433 evidence-based practices in autism. *Journal of Autism and
 434 Developmental Disorders*, 38, 1311–1319.
- 435 Rutter, M., & Lockyer, L. (1967). A five to fifteen year follow-up
 436 study of infantile psychosis. I: Description of sample. *British
 437 Journal of Psychiatry*, 113, 1169–1182.
- 438 Shriberg, L., Paul, R., McSweeney, J., Klin, A., Cohen, D., &
 439 Volkmar, F. (2001). Speech and prosody characteristics of
 440 adolescents and adults with high-functioning autism and
 Asperger syndrome. *Journal of Speech, Language, and Hearing
 Research*, 44, 1097–1115.
- Shriberg, L. D., & Widder, C. J. (1990). Speech and prosody
 characteristics of adults with mental retardation. *Journal of
 Speech and Hearing Research*, 33, 627–653.
- Staum, M. (1987). Music notation to improve the speech prosody of
 hearing impaired children. *Journal of Music Therapy*, 24,
 146–159.
- Stojanovik, V., Setter, J., & van Weijk, L. V. (2007). Intonation
 abilities of children with Williams syndrome: A preliminary
 investigation. *Journal of Speech, Language & Hearing
 Research*, 50, 1606–1617.
- Viera, A., & Garrett, J. (2005). Understanding interobserver agree-
 ment: The Kappa Statistic. *Family Medicine*, 37, 360–363.
- Wells, B., & Peppé, S. (2003). Intonation abilities in children with
 speech and language impairments. *Journal of Speech, Language,
 and Hearing Research*, 46, 5–20.
- Wilson, D., & Warton, T. (2006). Relevance and prosody. *Journal of
 Pragmatics*, 38, 1559–1579.

Journal : **10803**

Article : **2573**

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details Required	Author's Response
AQ1	Please note that section heading " Introduction " is added as per journal style.	
AQ2	Please provide the Author contribution statement	