

Effect of body mass and physical activity volume and intensity on pedometry-measured activity energy expenditure in rural black South Africans in the Limpopo Province

Ian Cook (BA (Phys Ed) Hons, BSc (Med) Hons)¹

Marianne Alberts (PhD)²

Estelle V Lambert (PhD)³

¹ Physical Activity Epidemiology Laboratory, University of Limpopo (Turffloep Campus), Polokwane

² Department of Chemical Pathology, University of Limpopo (Turffloep Campus), Polokwane

³ MRC/UCT Research Unit for Exercise Science and Sports Medicine, Department of Human Biology, University of Cape Town

Abstract

Objectives. We developed a novel approach to investigate patterns of pedometry-measured total weekly activity energy expenditure (EE_{Act}) in rural black South Africans in the Limpopo Province.

Design. We analysed 7-day pedometry data in 775 subjects (female: $N=508$; male: $N=267$). Variance components models for EE_{Act} were used to estimate the variance explained by body mass (BM), total weekly steps (volume) and estimated intensity ($kcal.kg^{-1}.step^{-1}$). Univariate General Linear Models, adjusting for age, BM and physical activity (PA) volume, were used to determine if EE_{Act} was primarily affected by volume or intensity.

Results. BM (13.1%), PA intensity (24.4%) and PA volume (56.9%) explained 94.4% of the variance in EE_{Act} . Adjusted EE_{Act} did not differ between sexes ($78 kcal.week^{-1}$, $p=0.2552$). There were no significant differences across activity categories (sedentary to very active) for adjusted EE_{Act} ($62 - 287 kcal.week^{-1}$, $p>0.1$). Adjusted EE_{Act} for 6 - 7 days of compliance ($\geq 10\ 000 steps.day^{-1}$) differed significantly from 1 - 2 days of compliance ($266 - 419 kcal.week^{-1}$, $p<0.04$). Obese (body mass index $\geq 30 kg.m^{-2}$) and normal weight (body mass index $18.5 - 24.9 kg.m^{-2}$) women did not differ significantly across activity categories for EE_{Act} ($200 - 592 kcal.week^{-1}$, $p>0.30$).

Conclusions. We have highlighted an intensity effect for days of compliance and at very active ambulatory levels ($\geq 12\ 500 steps.day^{-1}$). A volume effect appeared to dominate between sexes, across activity categories and weight-by-activity categories. It is important that *post hoc* statistical adjustments be made for body mass and PA volume when comparing EE_{Act} across groups.

Introduction

Pedometry is considered a valid and reliable objective measure of free-living physical activity (PA).¹ However, a disadvantage of this methodology is that the primary measure which is usually reported, namely the number of steps, provides no information as to the intensity of the ambulation. Consequently, it is not possible to disentangle the effects of volume and intensity of PA on outcome variables if statistical analyses only consider the number of steps (volume).

We recently reported high ambulation levels (average steps day^{-1}) for a rural African population in transition but could not provide definitive data pertaining to the intensity at which steps were accumulated.² The study used a piezo-electric based pedometer (NL-2000) which stored both the number of steps and activity energy expenditure (EE_{Act}).² The NL-2000 is produced by the Suzuken-Kenz Company (<http://www.suzuken-kenz.com>) for a North American distributor (New Lifestyles, <http://www.new-lifestyles.com>) and is identical in function to the Suzuken-Kenz e-STEP products (personal communication: Hitoshi Ozawa, Suzuken-Kenz, 17-05-2007). Therefore calibration results for the NL-2000 would also be applicable to the Suzuken-Kenz e-STEP products.³⁻⁵ Moreover, because the Suzuken-Kenz range includes the Lifecorder EX, which has been validated,^{6,7} and the technology in this high-end product is essentially the same as the lower-end products (except for the download capacity to a personal computer), the algorithms for all Suzuken-Kenz products would be identical. For instance, the NL-2000 (e-STEP) (personal communication: Operations Manager, New Lifestyles, 02-09-2005) and the Lifecorder EX⁶ sense steps when there are three or more acceleration pulses for four consecutive seconds and calculate EE_{Act} using body mass (W) and an intensity dependent factor (Ka) such that $EE_{Act} (kcal) = Ka \times W (kg)$.

CORRESPONDENCE:

Ian Cook
Physical Activity Epidemiology Laboratory
University of Limpopo (Turffloep Campus)
PO Box 459
Fauna Park
0787 Polokwane
South Africa
Tel+fax: +27 15 268 2390
E-mail: ianc@ul.ac.za

TABLE I. Descriptive characteristics for rural and urban women

| | Residence | | p^{\dagger} |
|--|----------------|----------------|---------------|
| | Female (N=508) | Male (N=267) | |
| Continuous variables | | | |
| Age (years) | 40.1 ± 20.7 | 28.4 ± 17.6 | <0.0001 |
| Body-mass-index (kg.m ⁻²) | 26.6 ± 6.4 | 21.2 ± 3.9 | <0.0001 |
| Average steps.day ⁻¹ | 11 086 ± 4 538 | 14 028 ± 5 434 | <0.0001 |
| Average activity EE (kcal.day ⁻¹) | 393 ± 189 | 491 ± 213 | <0.0001 |
| Intensity factor (kcal.kg ⁻¹ .step ⁻¹ x1000) | 0.58 ± 0.28 | 0.65 ± 0.41 | 0.0032 |
| Categorical variables * | | | |
| Body mass index classification | | | |
| Normal weight (<25 kg.m ⁻²) | 47.8 (243) | 87.6 (234) | <0.0001 |
| Obese (≥30 kg.m ⁻²) | 27.2 (138) | 4.1 (11) | <0.0001 |
| Physical activity classification | | | |
| Inactive (<5000 steps.day ⁻¹) | 10.2 (52) | 3.0 (8) | 0.0006 |
| Active (≥10 000 steps.day ⁻¹) | 59.4 (302) | 77.9 (208) | <0.0001 |
| Completion of secondary school (≥Grade 12) | 16.5 (58) | 14.1 (29) | 0.5420 |
| Ownership of motor vehicle (Yes) | 21.2 (99) | 18.1 (45) | 0.3911 |
| Electricity available inside house (Yes) | 75.4 (353) | 65.3 (162) | 0.0055 |
| Water collected outside dwelling (Yes) | 8.5 (40) | 8.1 (20) | 0.9363 |

Data are reported as mean ± SD for all continuous variables and % (N) for * categorical variables, † p-values evaluate female v. male differences.

Consequently, the EE_{Act} displayed on the NL-2000 output is a function of the PA intensity (Ka), PA volume (number of steps) and the individual's body mass. It is thus not possible to ascertain if an EE_{Act} difference between two individuals is due to increased PA (volume and/or intensity) or because of body mass differences. One approach to circumvent this problem is to use statistical methods to adjust for body mass and PA volume to ascertain whether EE_{Act} differences between two individuals are possibly intensity dependent. Therefore the objective of this study was to explore the patterns of pedometry-measured total weekly EE_{Act} by statistically adjusting for body mass and PA volume to determine if PA intensity could be an important factor in explaining the high ambulatory levels in a rural African setting.

Methods

This analysis uses data for which the study protocol, subjects, field site, sample size and measurements have been described in detail elsewhere.² Briefly, 830 participants from the Dikgale Health and Demographic Surveillance System field site (DHDSS)⁸⁻¹¹ were conveniently recruited and contacted twice over a 9-day period between January 2005 and December 2007. On the first occasion, subjects were recruited and completed the informed consent, relevant sections of a health questionnaire and provided anthropometric data. Standard anthropometric measurements and interviews were performed by trained, local fieldworkers and included measures of stature (nearest 1 cm) and body mass (nearest 1 kg). We categorised subjects using body mass index (BMI = body mass ÷ stature², underweight: <18.5 kg.m⁻², normal weight: 18.5 - 24.9 kg.m⁻², overweight: 25 - 29.9 kg.m⁻², obese: ≥30 kg.m⁻²).¹² Finally, subjects were instructed on the required procedures for wearing the pedometer over 9 consecutive days. We used piezo-electric pedometers (NL-2000, New Lifestyles Inc., Kansas City, MO, USA) not affected by pedometer tilt or adiposity level¹³ to objectively measure PA. Data for day 1 and day 9 were omitted because these were incomplete days. The pedometer was worn on the right waist, securely attached

to a nylon belt and sealed with surgical tape. The pedometers could be removed for sleeping and bathing purposes by unclipping the nylon belt. Ambulation PA volume was defined as the average steps.day⁻¹ or steps.week⁻¹. Energy expenditure was defined as total activity energy expenditure.week⁻¹ (EE_{Act}, kcal.week⁻¹). We calculated a PA intensity factor (I_{FR}, kcal.kg⁻¹.step⁻¹) from the total weekly steps, total weekly EE_{Act} and body mass. Public health indices (thresholds) for steps.day⁻¹ were defined as follows:¹⁴ sedentary: <5 000 steps.day⁻¹, low active: 5 000 - 7 499 steps.day⁻¹, somewhat active: 7 500 - 9 999 steps.day⁻¹, active: 10 000 - 12 499 steps.day⁻¹, and very active: ≥12 500 steps.day⁻¹. A summary variable was created indicating the number of days a subject was compliant or not for 0 - 7 days (≥10 000 steps.day⁻¹). Subjects received a small honorarium on completion of the study. The study was approved by the Ethics Committee of the University of Limpopo (Turffloop Campus).

Statistical analysis

Descriptive statistics comprised means and 95% confidence intervals (95% CI) or one standard deviation (SD). Independent t-tests were used to compare variables across gender. Variance components were estimated for inter-individual variance (body mass = kg, PA volume = total weekly steps and I_{FR} = kcal.kg⁻¹.step⁻¹) and residual (intra-individual) variance. The variance components were also expressed as a percentage of the total variance. Inter-individual variance represents true variation between subjects while intra-individual variance represents unexplained variation within subjects. To identify additional variables that could affect the inter-individual variance we entered age and stature as covariates and sex, village and season as fixed factors. Multiple linear regression models, using backward selection, were used to examine the relative importance of PA volume, PA intensity and body mass to EE_{Act}. In addition to PA and body mass variables, age and sex were included in all initial models. Significance for variable entry into and exit out of the model were set at p=0.05 and p=0.1, respectively. Univariate General Linear Models (GLM) were used to compare ambulation (steps.day⁻¹) and




Fig. 1. Ambulatory and energy expenditure levels for males and females. Steps.day⁻¹ adjusted for age and body mass. Kcal.week⁻¹ adjusted for age, body mass and steps.week⁻¹. * Steps.day⁻¹: males > females, $p < 0.0001$.

EE_{Act} (kcal.week⁻¹) across gender, activity categories (sedentary to very active), days of non-compliance/compliance with public health guidelines ($\geq 10\,000$ steps.day⁻¹), and obese- or normal-weight inactive ($< 7\,500$ steps.day⁻¹), active (10 000 - 12 499 steps.day⁻¹) and very active ($\geq 12\,500$ steps.day⁻¹) participants. All initial models adjusted for age and body mass. Additional EE_{Act} models were also constructed which adjusted for age, body mass and steps.week⁻¹. *Post hoc* multiple comparison analyses (Sidak's *t*-test) assessed group differences. Data were analysed using appropriate statistical software (SPSS version 17.0.2). Significance for all inferential statistics was set at $p < 0.05$.

Results

We excluded 14 outliers identified during exploratory data analysis. Because of very few obese males in the sample (Table I), only adult (19 - 65 years) female subjects were used in the obese/normal weight comparison across activity categories.

The sizeable variance attributed to body mass (13.1%), PA volume (total weekly steps, 56.9%) and I_{FR} (kcal.kg⁻¹.step⁻¹, 24.4%) suggested further analysis was warranted to determine if EE_{Act} group differences persist, possibly due to an intensity effect, by adjusting for body mass and PA volume. The 5.5% error variance was likely due to the rather crude I_{FR} that was calculated. Entering age or stature as covariates, and sex, village or season as fixed factors, made no difference to variances ($< 1\%$). Because the data were collected over 2 years in a number of villages that differ in terms of infrastructure and access to public transport, it was important to test whether season and village could explain part of the variance. PA volume, I_{FR} and body mass were significant predictors of EE_{Act} (model and coefficients: $p < 0.0001$, model adjusted $R^2 = 0.814$) and part correlations were 0.887, 0.355 and 0.259, respectively. The multiple linear regression analysis revealed no collinearity between variables (variance inflation factors < 1.2).

The adjusted R^2 for the partially (age and body mass) and fully (age, body mass and steps.wk⁻¹) adjusted EE_{Act} GLM models ranged from 0.130 to 0.564 and 0.690 to 0.695, respectively ($p < 0.0001$). Males accumulated significantly more steps than females (687 steps.day⁻¹, $p < 0.0001$). (Fig. 1). The fully adjusted EE_{Act} was not significantly different (78 kcal.wk⁻¹, $p = 0.2552$) between sexes, suggesting no intensity effect (Fig. 1). However, adjusting for age and body mass only resulted in a significant difference (224 kcal.




Fig. 2. Ambulatory and energy expenditure levels across activity categories. Steps.day⁻¹ adjusted for age and body mass. Kcal.week⁻¹ adjusted for age, body mass and steps.week⁻¹. Steps.day⁻¹: all activity categories significantly different, $p < 0.0001$.

week⁻¹, $p = 0.0017$), suggesting that volume plays a more significant role in EE_{Act} differences between sexes.

Age and body mass-adjusted differences in steps across categories (sedentary - very active) were significant for all pairwise comparisons (1 783 steps.day⁻¹ to 9 710 steps.day⁻¹, $p < 0.0001$) (Fig. 2). Adjusted for age and body mass, the middle three categories (low-active to active) resulted in relatively similar EE_{Act}, ($p > 0.1$), but not so for two the extreme EE_{Act} categories (sedentary v. very active, $p < 0.04$). However, the fully adjusted model yielded no significant differences across activity categories, suggesting a significant volume effect ($p > 0.1$) (Fig. 2).

Age and body mass-adjusted steps.day⁻¹ for 6 - 7 days of compliance with public health guidelines ($\geq 10\,000$ steps.day⁻¹) were significantly different to all other levels of compliance ($p < 0.003$) but there were no significant differences between 0 days (non-compliance) and 1 - 5 days of compliance ($p > 0.4$) (Fig. 1). Adjusted for age and body mass, EE_{Act} differed significantly for days of compliance ≥ 6 compared with non-compliance (0 days) and lower levels of compliance (1 - 5 days) ($p < 0.03$). Significant differences persisted only for EE_{Act} for 6 - 7 days of compliance versus 1 - 2 days of compliance ($p < 0.04$), suggesting an intensity effect only for 6 - 7 days of compliance (Fig. 3).

The EE_{Act}, adjusted for age and body mass of highly active normal weight and obese women was significantly higher than inactive normal weight and obese women (difference: 2 224 - 2 403 kcal.wk⁻¹, $p < 0.0001$). Adjusted for age, body mass and steps.week⁻¹, obese and normal weight adult females did not differ in EE_{Act} for the three PA categories, suggesting that there are no statistical differences in PA intensity (Fig. 4). However, there was a tendency for normal weight females to have higher EE_{Act} (200 kcal.week⁻¹ to 592 kcal.week⁻¹, $p > 0.30$).

Discussion

This is a novel study reporting for the first time volume and intensity effects from data obtained using pedometers. The analysis has highlighted an intensity effect for days of compliance and especially at very active ambulatory levels ($\geq 12\,500$ steps.day⁻¹). Interestingly, there did not seem to be a significant intensity effect between sexes, activity categories or obese versus normal weight across activity




Fig. 3. Ambulatory and energy expenditure levels across number of days compliant ($\geq 10\,000$ steps.day⁻¹). Steps.day⁻¹ adjusted for age and body mass. Kcal.week⁻¹ adjusted for age, body mass and steps.week⁻¹. * Steps.day⁻¹: 6 days and 7 days significantly different to all other days, $p < 0.003$. † Kcal.week⁻¹: 1 - 2 days significantly different to 6 - 7 days, $p < 0.04$.




Fig. 4. Ambulatory and energy expenditure levels for normal weight and obese women across activity categories. Kcal.week⁻¹ adjusted for age, body mass and steps.week⁻¹.

categories once age, body mass and accumulated steps had been adjusted for.

Differences in accumulated steps between males and females have been reported.^{15,16} Our results suggest that it is the difference in PA volume, and not PA intensity, that explains the difference in EE_{Act} between males and females from this rural, African setting. Although average step totals increased significantly across activity categories, adjusted EE_{Act} did not increase accordingly. We also did not find markedly increased adjusted EE_{Act} between obese and normal weight females across activity categories, which is in agreement with findings of similar gross EE for walking and jogging at the same speed between normal weight and overweight/obese women, once adjusted for body mass and free fat mass.¹⁷ Furthermore, non-compliance or compliance on 1 - 5 days of the week with public health guidelines ($\geq 10\,000$ steps.day⁻¹)¹⁸ did not seem to reveal differences in EE_{Act} . However, complying on 6 - 7 days, required significant increases in volume and intensity. These results suggest that public health PA

guidelines of at least 5 times per week, 30 minutes per session,¹⁹ which equates to approximately $10\,000$ steps.day⁻¹,²⁰⁻²² were likely met through increases in accumulated steps throughout the day instead of increasing PA intensity. It was interesting that these findings would seem to provide non-intervention, free-living support for the feasibility of promoting moderate intensity PA such that the lack of a vigorous intensity requirement would not be a barrier to increasing PA. In other words, our results suggest that within this rural African population, walking behaviours are naturally modelled according to public health PA guidelines. However, highly active groups such as those achieving $\geq 12\,500$ steps on 6 or more days a week, required increases in intensity. This finding is in accord with the significantly higher accelerometer-measured moderate-to-vigorous PA recorded for subjects achieving $\geq 11\,762$ steps.day⁻¹ compared with those achieving $\leq 8\,123$ steps.day⁻¹; 68.6 min v. 23.6 min, respectively ($p = 0.000$).²² Le Masurier *et al.* also reported higher moderate-to-vigorous activity for subjects achieving $\geq 10\,000$ steps.day⁻¹ compared with $< 10\,000$ steps.day⁻¹ whether accumulated in bouts ≥ 1 -, ≥ 5 - or ≥ 10 min (difference: $p < 0.05$).²⁰ Recently, Dugas *et al.* suggested that PA intensity and not PA volume was a greater determinant of adiposity in young, black South African urban dwellers.²³ Our results would suggest that the PA volume is the dominant contributor to EE_{Act} in rural dwellers. Moreover, we have shown that average steps.day⁻¹ is significantly associated with adiposity levels in rural African women ($r = -0.20$, $p = 0.032$).²⁴

Several limitations must be acknowledged. Firstly, we could not compare our statistical adjustment against actual volume and intensity measures. Future analyses using uni-axial accelerometer data to ascertain at which intensity levels steps are being accumulated would provide more definitive answers as to the relative importance of PA volume and intensity, specifically within the context of a rural African setting. Secondly, the absolute EE_{Act} values reported in this study should be carefully interpreted because treadmill calibration studies for the NL-2000 suggest an overestimation of approximately 25% for EE_{Act} .³

In conclusion, we have highlighted an intensity effect for 6 - 7 days of compliance and at very active ambulatory levels. A volume effect appeared to dominate between sexes, across activity categories and weight-by-activity categories and would suggest that public health messages in this specific rural setting should focus on maintaining PA volume through daily living rather than advocating increases in PA intensity. It is important that *post hoc* statistical adjustments be made for body mass and PA volume when comparing EE_{Act} data across groups.

Acknowledgements

The Research Development and Administration Division of the University of Limpopo (Turloop Campus) and the Thuthuka Programme of the National Research Foundation supported this study.

REFERENCES

1. Bassett DR, Strath SJ. Use of pedometers to assess physical activity. In: Welk GJ ed. Physical activity assessments for health-related research. Champaign, IL: Human Kinetics 2002:163-177.
2. Cook I, Alberts M, Brits JS, Choma S, Mkhonto SS. Descriptive epidemiology of ambulatory activity in rural, black South Africans. *Med Sci Sports Exerc* 2010; (In press).
3. Crouter SE, Schneider PL, Karabulut M, Bassett DR. Validity of 10 electronic pedometers for measuring steps, distance, and energy cost. *Med Sci Sports Exerc* 2003;35:1455-1460.

-
4. Schneider PL, Crouter SE, Lukajic O, Bassett DR. Accuracy and reliability of 10 pedometers for measuring steps over a 400-m walk. *Med Sci Sports Exerc* 2003;35:1779-1784.
 5. Schneider PL, Crouter SE, Bassett DR. Pedometer measures of free-living physical activity: comparison of 13 models. *Med Sci Sports Exerc* 2004;36:331-335.
 6. Kumahara H, Schutz Y, Ayabe M, *et al.* The use of uniaxial accelerometry for the assessment of physical-activity-related energy expenditure: a validation study against whole-body indirect calorimetry. *Br J Nutr* 2004;91:235-243.
 7. McClain JJ, Craig CL, Sisson SB, Tudor-Locke C. Comparison of Life-corder EX and ActiGraph accelerometers under free-living conditions. *Appl Physiol Nutr Metab* 2007;32:753-761.
 8. Alberts M, Burger S. INDEPTH DSS Site Profiles: Dikgale DSS, South Africa. In: Sankoh OA, Kahn K, Mwageni E, Ngom P, Nyarko P, eds. *Population and Health in Developing Countries. Volume 1. Population, Health, and Survival at INDEPTH Sites.* Ottawa: IDRC, 2002: 207-211.
 9. Alberts M, Urdal P, Steyn K, *et al.* Prevalence of cardiovascular diseases and associated risk factors in a rural black population of South Africa. *Eur J Cardiovasc Prev Rehabil* 2005;12:347-354.
 10. Cook I, Alberts M, Burger S, Byass P. All-cause mortality trends in Dikgale, rural South Africa, 1996-2003. *Scand J Public Health* 2008;36:753-760.
 11. Alberts M, Burger S, Tollman SM. The Dikgale field site. *S Afr Med J* 1999;89:851-852.
 12. Puoane T, Steyn K, Bradshaw D, *et al.* Obesity in South Africa: The South African Demographic and Health Survey. *Obes Res* 2002;10:1038-1048.
 13. Crouter SE, Schneider PL, Bassett DR. Spring-levered versus piezo-electric pedometer accuracy in overweight and obese adults. *Med Sci Sports Exerc* 2005;37:1673-1679.
 14. Tudor-Locke C, Bassett DR. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med* 2004;34:1-8.
 15. McCormack G, Giles-Corti B, Milligan R. Demographic and individual correlates of achieving 10,000 steps/day: use of pedometers in a population-based study. *Health Promot J Austr* 2006;17:43-47.
 16. Tudor-Locke C, Johnson WD, Katzmarzyk PT. Accelerometer-determined steps per day in US Adults. *Med Sci Sports Exerc* 2009;41:1384-1391.
 17. Le Cheminant JD, Heden T, Smith J, Covington NK. Comparison of energy expenditure, economy, and pedometer counts between normal weight and overweight or obese women during a walking and jogging activity. *Eur J Appl Physiol* 2009;106:675-682.
 18. Tudor-Locke C, Hatano Y, Pangrazi RP, Kang M. Revisiting "How Many Steps Are Enough?". *Med Sci Sports Exerc* 2008;40:S537-S543.
 19. Haskell WL, Lee IM, Pate RR, *et al.* Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39:1423-1434.
 20. Le Masurier GC, Sidman CL, Corbin CB. Accumulating 10,000 steps: does this meet current physical activity guidelines? *Res Q Exerc Sport* 2003;74:389-394.
 21. Hultquist CN, Albright C, Thompson DL. Comparison of walking recommendations in previously inactive women. *Med Sci Sports Exerc* 2005;37:676-683.
 22. Macfarlane DJ, Chan D, Chan KL, Ho EY, Lee CC. Using three objective criteria to examine pedometer guidelines for free-living individuals. *Eur J Appl Physiol* 2008;104:435-444.
 23. Dugas LR, Carstens MA, Ebersole K, *et al.* Energy expenditure in young adult urban informal settlement dwellers in South Africa. *Eur J Clin Nutr* 2009;63:805-807.
-