

Dartmouth College

Dartmouth Digital Commons

Dartmouth Scholarship

Faculty Work

5-24-2011

Boundary Quasi-Orthogonality and Sharp Inclusion Bounds for Large Dirichlet Eigenvalues

A. H. Barnett

Dartmouth College

A. Hassell

Australian National University

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Mathematics Commons](#)

Dartmouth Digital Commons Citation

Barnett, A. H. and Hassell, A., "Boundary Quasi-Orthogonality and Sharp Inclusion Bounds for Large Dirichlet Eigenvalues" (2011). *Dartmouth Scholarship*. 2046.

<https://digitalcommons.dartmouth.edu/facoa/2046>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

BOUNDARY QUASI-ORTHOGONALITY AND SHARP INCLUSION BOUNDS FOR LARGE DIRICHLET EIGENVALUES*

A. H. BARNETT[†] AND A. HASSELL[‡]

Abstract. We study eigenfunctions ϕ_j and eigenvalues E_j of the Dirichlet Laplacian on a bounded domain $\Omega \subset \mathbb{R}^n$ with piecewise smooth boundary. We bound the distance between an arbitrary parameter $E > 0$ and the spectrum $\{E_j\}$ in terms of the boundary L^2 -norm of a normalized trial solution u of the Helmholtz equation $(\Delta + E)u = 0$. We also bound the L^2 -norm of the error of this trial solution from an eigenfunction. Both of these results are sharp up to constants, hold for all E greater than a small constant, and improve upon the best-known bounds of Moler–Payne by a factor of the wavenumber \sqrt{E} . One application is to the solution of eigenvalue problems at high frequency, via, for example, the method of particular solutions. In the case of planar, strictly star-shaped domains we give an inclusion bound where the constant is also sharp. We give explicit constants in the theorems, and show a numerical example where an eigenvalue around the 2500th is computed to 14 digits of relative accuracy. The proof makes use of a new quasi-orthogonality property of the boundary normal derivatives of the eigenmodes (Theorem 1.3), of interest in its own right. Namely, the operator norm of the sum of rank 1 operators $\partial_n \phi_j \langle \partial_n \phi_j, \cdot \rangle$ over all E_j in a spectral window of width \sqrt{E} —a sum with about $E^{(n-1)/2}$ terms—is at most a constant factor (independent of E) larger than the operator norm of any one individual term.

Key words. Laplacian, Helmholtz, Dirichlet, eigenvalue, eigenfunction, particular solutions

AMS subject classifications. 35P15, 47A55, 65N35, 65N15

DOI. 10.1137/100796637

1. Introduction and main results. The computation of eigenvalues and eigenmodes of Euclidean domains is a classical problem (in two dimensions this is the “drum problem,” reviewed in [21, 34]) with a wealth of applications to engineering and physics, including acoustic, electromagnetic and optical cavity and resonator design, microlasers [35], and data analysis [30]. It also has continued interest in mathematical community in the areas of quantum chaos [37, 3] and spectral geometry [18]. Let ϕ_j be a sequence of orthonormal eigenfunctions and E_j the respective eigenvalues ($0 < E_1 < E_2 \leq E_3 \leq \dots$ counting multiplicities) of $-\Delta$, where $\Delta := \sum_{m=1}^n \partial^2 / \partial x_m^2$ is the Laplacian in a bounded domain $\Omega \in \mathbb{R}^n$, $n \geq 2$, with Dirichlet boundary condition. That is, ϕ_j satisfies

$$(1.1) \quad (\Delta + E_j)\phi_j = 0 \quad \text{in } \Omega,$$

$$(1.2) \quad \phi_j = 0 \quad \text{on } \partial\Omega,$$

$$(1.3) \quad \|\phi_j\|_{L^2(\Omega)} = 1.$$

We will call the spectrum $\sigma := \{E_j\}_{j=1}^\infty$. Many of the applications mentioned demand high frequencies, that is, mode numbers j from 10^2 to as high as 10^6 . An efficient

*Received by the editors May 27, 2010; accepted for publication (in revised form) March 14, 2011; published electronically May 24, 2011.

<http://www.siam.org/journals/sinum/49-3/79663.html>

[†]Department of Mathematics, Dartmouth College, Hanover, NH 03755 (ahb@math.dartmouth.edu). This author’s work was supported by NSF grant DMS-0811005 and a Visiting Fellowship to ANU in February 2009 as part of the ANU 2009 *Special Year on Spectral Theory and Operator Theory*.

[‡]Mathematical Sciences Institute, Australian National University, Canberra, ACT 0200, Australia (hassell@maths.anu.edu.au). This author’s work was supported by Australian Research Council Discovery grant DP0771826.




FIG. 1.1. Tension $t[\tilde{u}_{\min}]$ versus energy E for the domain shown on the right. \tilde{u}_{\min} is the optimal trial Helmholtz solution lying in the span of a numerical basis set (see section 7). (a) Low frequency, showing the minima corresponding to the lowest 20 Dirichlet eigenvalues. (b) Medium-high frequency, showing a similar interval starting at eigenvalue number $j \approx 2552$; note the new vertical scale. (c) Density plot of eigenfunction $\phi_j \approx \tilde{u}_{\min}$ corresponding to the eigenvalue $E_j = 10005.02135797 \dots$ shown by the dot in (b) (black indicates large values of $|\phi_j|^2$, white zero).

solution of the problem thus requires specialized numerical approaches that scale with wavenumber better than conventional discretization methods.

The goal of this paper is to bound the errors of approximate eigenvalues and eigenfunctions computed using trial functions that satisfy exactly the homogeneous Helmholtz equation in Ω . As we will review below, such computational methods have proven to be very powerful. Recently one of the authors [4] improved upon the classical eigenvalue bound of Moler and Payne [24] by a factor of the wavenumber; however, this result has limited utility since it applies only to Helmholtz parameters lying in neighborhoods of σ of unknown size. In the present paper we go well beyond this result by giving new theorems, which (i) hold for *all* Helmholtz parameters (greater than an $O(1)$ constant), (ii) retain the improved high-frequency asymptotic behavior of [4] and show that this behavior is sharp, and (iii) improve upon the best known eigenfunction estimates, again by a factor of the wavenumber. To achieve this we make use of a new form of quasi-orthogonality of the eigenfunctions on the boundary, Theorem 1.3, of independent interest.

Before presenting our results, we need to review some known inclusion bounds and their importance for applications. Given an energy parameter¹ $E > 0$, let u be a nontrivial solution to the homogeneous Helmholtz equation $(\Delta + E)u = 0$ in Ω with no imposed boundary condition, and define its boundary error norm (or “tension”)

$$(1.4) \quad t[u] := \frac{\|u\|_{L^2(\partial\Omega)}}{\|u\|_{L^2(\Omega)}}.$$

Clearly, $t[u] = 0$ implies that E is an eigenvalue. It is reasonable to expect that if $t[u]$ is small for some Helmholtz solution u , then E is close to an eigenvalue. Moler and Payne [24] (building upon [16]) quantified this: there is a constant C_{MP} depending only on the domain, such that

$$(1.5) \quad d(E, \sigma) \leq C_{\text{MP}} E t[u],$$

¹The Helmholtz parameter E may be interpreted as energy, or as the square of frequency, depending on the application.

where $d(E, \sigma) := \min_j |E_j - E|$ denotes the distance of E from the spectrum.

An important application is solving (1.1)–(1.3) via global approximation methods, including the method of particular solutions (MPS) [9, 4]. One writes a trial eigenmode $u = \sum_{n=1}^N c_n \xi_n$ via basis functions ξ_n which are closed-form Helmholtz solutions in Ω but which need not satisfy any particular boundary condition. By adjusting the coefficients c_n (via a generalized eigenvalue [4] or singular value problem [8]) one may minimize $t[u]$ at fixed E ; by repeating this in a search for E values where the minimum $t[u]$ is very small, as illustrated by Figure 1.1(a) and (b), one may then locate approximate eigenvalues whose error is bounded above by (1.5). (This is sometimes called the method of a priori-a posteriori inequalities [21, sec. 16].)

Due to the work of Betcke and Trefethen [9] and others, such methods have enjoyed a recent revival, at least in $n = 2$, due to their high (often spectral) accuracy and their efficiency at high frequency when compared to direct discretization methods such as finite elements. For example, in various domains, 14 digits may be achieved in double precision arithmetic [9], and with an MPS variant known as the scaling method, tens of thousands of eigenmodes as high as $j \sim 10^6$ have been computed [3, 5]. (There are also successful variants [14, 15] by Descloux and Tolley, Driscoll, and others, in which subdomains are used, which we will not pursue here.)

If we instead interpret u as the solution error for an interior Helmholtz boundary-value problem (solved, for instance, via MPS or boundary integral methods), then (1.5) states that the interior error is controlled by the boundary error; this aids the numerical analysis of such problems [22, 6]. Similar estimates (which, however, rely on impedance boundary conditions) enable the analysis of least-squares nonpolynomial finite element methods [25, Thm 3.1]. Improving such estimates could thus be of general benefit for the numerical solution of Helmholtz problems.

Recently one of the authors [4] observed numerical evidence that (1.5) is not sharp for large E , and showed that there is a constant C_B depending only on Ω such that, for each $\varepsilon > 0$,

$$(1.6) \quad d(E, \sigma) \leq C_B(1 + \varepsilon)\sqrt{E}t[u]$$

holds whenever E lies in some open, possibly disconnected, subset of the real axis containing σ . This is an improvement over (1.5) by a factor of the wavenumber \sqrt{E} , which in problems of interest can be as high as 10^3 . However, since the proof relied on analytic perturbation in the parameter E , there was no knowledge about the *size* of this (ε -dependent) subset, hence no way to know in a given practical situation whether the error bound holds. The point of the present work is then to remedy this problem by removing any restriction to an unknown subset, and also to extend the \sqrt{E} improvement to bounds on approximate eigenfunctions.

We assume the domain $\Omega \subset \mathbb{R}^n$ has unit area (or volume for $n > 2$), and obeys the following rather weak geometric condition.

CONDITION 1. *The domain $\Omega \subset \mathbb{R}^n$ is bounded, with piecewise smooth boundary in the sense of Zelditch and Zworski [38]. This means that Ω is given by an intersection*

$$\Omega = \bigcap_{i=1}^N \{\mathbf{x} \mid f_i(\mathbf{x}) > 0\},$$

where the f_i are smooth functions defined on a neighborhood of $\overline{\Omega}$ such that

- $\nabla f_i \neq 0$ on the set $\{f_i = 0\}$,
- $\{f_i = f_j = 0\}$ is an embedded submanifold of \mathbb{R}^n , $1 \leq i < j \leq N$, and

- Ω is locally Lipschitz, i.e., for any boundary point $\mathbf{x}_0 \in \partial\Omega$, there is a Euclidean coordinate system z_1, \dots, z_n and a Lipschitz function k of $n-1$ variables such that in some neighborhood of \mathbf{x}_0 , we have

$$(1.7) \quad \partial\Omega = \{z_n = k(z_1, \dots, z_{n-1})\}.$$

Our main result on eigenvalue inclusion is the following.

THEOREM 1.1. *Let $\Omega \subset \mathbb{R}^n$ be a domain satisfying Condition 1. Then there are constants C, c depending only on Ω , such that the following holds. Let $E > 1$ and suppose u is a nontrivial solution of $(\Delta + E)u = 0$ in $C^\infty(\Omega)$, with $t[u] := \|u\|_{L^2(\partial\Omega)} / \|u\|_{L^2(\Omega)}$. Then,*

$$(1.8) \quad d(E, \sigma) \leq C\sqrt{E}t[u],$$

and for the normalized Helmholtz solution u_{\min} minimizing $t[u]$ at the given E ,

$$(1.9) \quad c\sqrt{E}t[u_{\min}] \leq d(E, \sigma) \leq C\sqrt{E}t[u_{\min}].$$

Remark 1.1. The estimate (1.9) states that (1.8) is sharp, i.e., using $t[u]$ alone one cannot localize the spectrum any more tightly than this, apart from optimizing the constants c and C .

Remark 1.2. The existence of a minimizer for $t[u]$ follows from Lemma 3.1, in the case that E is not a Dirichlet eigenvalue (and is trivial when E is a Dirichlet eigenvalue). The lower bound on the distance to the spectrum in (1.9) is of use when the numerical scheme is known to produce a good approximation to u_{\min} .

We will also prove the following corresponding bound on the error of the trial eigenfunction u , which improves by a factor \sqrt{E} the previously best known result (Moler and Payne [24, Thm. 2]).

THEOREM 1.2. *Let Ω be as in Theorem 1.1. Then there is a constant C depending only on Ω , such that the following holds. Let $E > 1$, let E_j be the eigenvalue nearest to E , and let E_k be the next nearest distinct eigenvalue. Suppose u is a solution of $(\Delta + E)u = 0$ in $C^\infty(\Omega)$ with $\|u\|_{L^2(\Omega)} = 1$, and let \hat{u}_j be the projection of u onto the E_j eigenspace. Then,*

$$(1.10) \quad \|u - \hat{u}_j\|_{L^2(\Omega)} \leq C \frac{\sqrt{E}t[u]}{|E - E_k|}.$$

Remark 1.3. The left-hand side above is equal to $\sin \theta$, where θ is the subspace angle between u and the E_j eigenspace (this viewpoint is elaborated in [9, sec. 6]). Similarly, when E_j is a simple eigenvalue, we may write $\|u - \phi_j\|_{L^2(\Omega)} = 2 \sin(\theta/2)$.

Remark 1.4. This result is also sharp, in a certain sense; see Remark 4.1.

To conclude this introduction, we present some key ingredients of the proofs. Define the boundary functions of the eigenmodes by

$$(1.11) \quad \psi_j(s) := \partial_n \phi_j(s), \quad s \in \partial\Omega,$$

where $\partial_n = \mathbf{n} \cdot \nabla$ is the usual normal derivative. Our main tools will be two theorems stating that boundary functions ψ_j lying close in eigenvalue are almost orthogonal. The first is the following new result which we prove in section 2.

THEOREM 1.3 (spectral window quasi-orthogonality). *Let $\Omega \subset \mathbb{R}^n$ be a domain satisfying Condition 1. There exists a constant C_Ω depending only on Ω such that the operator norm bound*

$$(1.12) \quad \left\| \sum_{|E_j - E| \leq E^{1/2}} \psi_j \langle \psi_j, \cdot \rangle \right\|_{L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)} \leq C_\Omega E$$

holds for all $E \geq 1$. (Here, $\langle \cdot, \cdot \rangle$ denotes the inner product in $L^2(\partial\Omega)$.)

Remark 1.5. By Weyl's Law [17, Chap. 11] there are $O(E^{(n-1)/2})$ terms in the above sum. Since each term already has norm $\geq cE$ [28, 19], the theorem expresses essentially complete mutual orthogonality, up to a constant. Only the scaling of the window width with E is important: the theorem also holds for a window $|E_j - E| \leq cE^{1/2}$ for any fixed c (C_Ω will then depend on c as well as Ω). On the other hand, one could not expect it to hold over a spectral window of width $O(E^\beta)$ for $\beta > 1/2$, since the boundary functions are approximately band-limited to spatial wavenumber $E^{1/2}$ and thus no more than $O(E^{(n-1)/2})$ of them could be orthogonal on the boundary.

The second result is a pairwise estimate on the inner product of boundary functions lying close in eigenvalue, with respect to a special inner product. (Here, $\mathbf{x}(s)$ refers to the location of boundary point s relative to a fixed origin, which may or may not be inside Ω .)

THEOREM 1.4 (pairwise quasi-orthogonality). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, and let $S := \frac{1}{2} \sup_{\mathbf{x} \in \Omega} \|\mathbf{x}\|$. Then, for all $i, j \geq 1$,*

$$(1.13) \quad \left| \int_{\partial\Omega} (\mathbf{x}(s) \cdot \mathbf{n}(s)) \psi_i(s) \psi_j(s) ds - 2E_i \delta_{ij} \right| \leq S^2 (E_i - E_j)^2.$$

Remark 1.6. Theorem 1.4 was proved by the first-named author in [3, Appendix B]. It may be viewed as an off-diagonal generalization of a theorem of Rellich [28] which gives the $i = j$ case. The boundary weight $\mathbf{x} \cdot \mathbf{n}$ (also known as the Morawetz multiplier) is the only one known that gives quadratic growth away the diagonal yet also gives nonzero diagonal elements.

Note that neither of the above quasi-orthogonality theorems implies the other. We also note that Bäckér et al. derived a completeness property of the boundary functions in a (smoothed) spectral window [2, eq. (53)], that is closely related to Theorem 1.3.

After proving Theorem 1.3 in section 2, we combine it with a boundary operator defined in section 3 to prove the main theorems in section 4. In section 5 we state and prove a variant of Theorem 1.1 for strictly star-shaped planar domains, which has an optimal constant C . This builds on Theorem 1.4 combined with the Cotlar–Stein lemma (see Lemma 5.2). In the main theorems (Theorems 1.1, 1.2, and 5.1), the domain-dependent constants are not explicit; we discuss their explicit values in section 6. We present a high-accuracy numerical example using the MPS, and sketch some of the implementation aspects in section 7. Finally, we conclude in section 8.

2. Quasi-orthogonality in an eigenvalue window. Here we prove Theorem 1.3 using a “ TT^* ” argument.” We need the fact that the upper bound $\|\psi_j\|_{L^2(\partial\Omega)}^2 \leq CE_j$ on eigenmode normal derivatives, proved, for example, in [19], generalizes to quasimodes living in an $O(E^{1/2})$ spectral window. The proof is almost the same as in [19].

LEMMA 2.1. Let $\Omega \subset \mathbb{R}^n$ satisfy Condition 1. Let $E > 1$, and let

$$(2.1) \quad \phi := \sum_{|E_j - E| \leq E^{1/2}} c_j \phi_j$$

with real coefficients c_j , and $\sum_j c_j^2 = \|\phi\|_{L^2(\Omega)}^2 = 1$. Then,

$$(2.2) \quad \|\partial_n \phi\|_{L^2(\partial\Omega)}^2 \leq C_\Omega E,$$

where the constant C_Ω depends only on Ω .

Proof. To prove this we need the following lemma, proved in Appendix A, stating that for any piecewise smooth domain (in the sense of Condition 1) there is a smooth vector field that is outgoing at each boundary point.

LEMMA 2.2. Let Ω satisfy Condition 1. Then there exists a smooth vector field \mathbf{a} , defined on a neighborhood of $\overline{\Omega}$, such that

$$(2.3) \quad \mathbf{a} \cdot \mathbf{n} \geq 1$$

almost everywhere on $\partial\Omega$.

The main tool for proving Lemma 2.1 is the identity

$$(2.4) \quad \int_{\partial\Omega} (D\phi) \partial_n \phi = - \int_{\Omega} \phi [\Delta, D] \phi + \int_{\Omega} (D\phi) (\Delta + E) \phi - \int_{\Omega} \phi D(\Delta + E) \phi$$

for any first order differential operator D , which follows² from Green's 2nd identity, the definition of the commutator, and $\phi|_{\partial\Omega} = 0$. Choosing $D := \mathbf{a} \cdot \nabla$, where \mathbf{a} is as in Lemma 2.2, we notice that the left-hand side of (2.4) bounds the left-hand side of (2.2), since

$$(2.5) \quad \int_{\partial\Omega} \psi_j^2 \leq \int_{\partial\Omega} (\mathbf{a} \cdot \mathbf{n}) \psi_j^2$$

by Lemma 2.2. We may now bound each of the terms on the right-hand side of (2.4). Defining $C_a = \sup_{\mathbf{x} \in \Omega} |\mathbf{a}(\mathbf{x})|$, we first need

$$(2.6) \quad \|D\phi\|_{L^2(\Omega)}^2 \leq C_a^2 \int_{\Omega} \|\nabla \phi\|^2 = -C_a^2 \int_{\Omega} \phi \Delta \phi = C_a^2 \sum_j |c_j|^2 E_j \leq C_a^2 F,$$

where $F := E + E^{1/2}$ is the upper end of the energy window. Similarly,

$$(2.7) \quad \begin{aligned} \|D(\Delta + E)\phi\|_{L^2(\Omega)}^2 &\leq C_a^2 \int_{\Omega} \|\nabla(\Delta + E)\phi\|^2 \\ &= C_a^2 \sum_{ij} c_i c_j \int_{\Omega} (\Delta + E)\phi_i (-\Delta)(\Delta + E)\phi_j \\ &= C_a^2 \sum_j c_j^2 E_j (E - E_j)^2 \leq C_a^2 E F. \end{aligned}$$

Using Cauchy-Schwarz, the sum of the last two terms in (2.4) is then bounded by $2C_a \sqrt{EF}$. For the first term on the right-hand side of (2.4), we use, in Einstein

²The computation, involving a total of three derivatives, is justified for our class of domains, since Dirichlet eigenfunctions are in $H^{3/2}(\Omega)$ for any Lipschitz Ω ; see [13, Theorem B, p. 164]. Rellich-type computations are also justified on Lipschitz domains in [1].

notation, $[\Delta, D] = \partial_{ii}(a_j \partial_j \cdot) - a_j \partial_{ii}$. After several steps, using integration by parts and $\phi|_{\partial\Omega} = 0$, we get

$$(2.8) \quad - \int_{\Omega} \phi [\Delta, D] \phi = 2 \int_{\Omega} (\partial_i a_j) (\partial_i \phi) \partial_j \phi + \int_{\Omega} (\partial_{ii} a_j) \phi \partial_j \phi.$$

The constants $C'_a := \sup_{\mathbf{x} \in \Omega} \|\mathbb{A}(\mathbf{x})\|_2$, where the matrix $\mathbb{A} \in \mathbb{R}^{n \times n}$ has entries $\partial_i a_j$, and $C''_a := \sup_{\mathbf{x} \in \Omega, j=1, \dots, n} |\Delta a_j(\mathbf{x})|$, exist and are finite. Then (2.8) is bounded by $2C'_a F + C''_a F^{1/2}$. Adding all bounds on terms in (2.4) and using (2.5), we get

$$(2.9) \quad \|\partial_n \phi\|_{L^2(\partial\Omega)}^2 \leq 2(C_a + C'_a)F + C''_a \sqrt{F},$$

which is bounded by a constant times E for $E > 1$. \square

Proof of Theorem 1.3. Consider the coefficient vector $\mathbf{c} := \{c_j\} \in \mathbb{R}^N$ appearing in (2.1), where N is the number of eigenvalues (counting multiplicity) in the spectral window. Define the linear operator $T : \mathbb{R}^N \rightarrow L^2(\partial\Omega)$ by

$$(2.10) \quad T\mathbf{c} = \sum_j c_j \psi_j.$$

Lemma 2.1 states that $\|T\|_{l^2 \rightarrow L^2(\partial\Omega)} \leq (C_{\Omega} E)^{1/2}$. Thus $\|TT^*\|_{L^2(\partial\Omega)} \leq C_{\Omega} E$. But TT^* is the operator in the statement of Theorem 1.3, which completes its proof. \square

3. Relating tension to a boundary operator. In this section, we show, following Barnett [4], that the tension $t[u]$ is related to the operator norm of a natural boundary operator.

For E a noneigenvalue of Ω , let $\mathcal{K}(E) : L^2(\partial\Omega) \rightarrow L^2(\Omega)$ be the solution operator (Poisson kernel) for the interior Dirichlet boundary-value problem,

$$(3.1) \quad (\Delta + E)u = 0 \quad \text{in } \Omega,$$

$$(3.2) \quad u = f \quad \text{on } \partial\Omega,$$

that is, $u = \mathcal{K}f$. (For existence and uniqueness for L^2 data on a Lipschitz boundary, see, for example, [23, Thm. 4.25].) Since the eigenbasis is complete in $L^2(\Omega)$, we may write $u = \sum_{j=1}^{\infty} c_j \phi_j$. We evaluate each c_j by applying Green's 2nd identity,

$$(3.3) \quad (E - E_j)(\phi_j, u)_{L^2(\Omega)} = \int_{\Omega} (u \Delta \phi_j - \phi_j \Delta u) = \int_{\partial\Omega} (f \psi_j - \phi_j \partial_n u) ds,$$

thus $c_j = \langle \psi_j, f \rangle / (E - E_j)$. The solution operator may therefore be written as a sum of rank-1 operators,

$$(3.4) \quad \mathcal{K}(E) = \sum_{j=1}^{\infty} \frac{\phi_j \langle \psi_j, \cdot \rangle}{E - E_j}.$$

By the definition (1.4) we have, now for any u satisfying $(\Delta + E)u = 0$ in Ω , that $t[u]^{-1} \leq \|\mathcal{K}(E)\|$. Since $\|\mathcal{K}^* \mathcal{K}\| = \|\mathcal{K}\|^2$, then by defining the boundary operator in $L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)$,

$$(3.5) \quad A(E) := \mathcal{K}(E)^* \mathcal{K}(E),$$

we have an estimate on the tension that will be the main tool in our analysis,

$$(3.6) \quad t[u]^{-2} \leq \|A(E)\|.$$

Inserting (3.4) into (3.5) and using orthogonality (or see [4, sec. 3.1]), we have that A also may be written as the sum of rank-1 operators,

$$(3.7) \quad A(E) = \sum_{j=1}^{\infty} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2}.$$

This sum is conditionally convergent: the sum of the operator norm of each term diverges. For instance, for $n = 2$, Weyl's law [17, Chap. 11] states that the density of eigenvalues E_j is asymptotically constant, but since $\|\psi_j\|^2 = \Omega(E_j)$ the sum of norms is logarithmically divergent; for $n > 2$ the divergence is worse. Despite this, we have the following, which improves upon the results of [4].

LEMMA 3.1. *Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, satisfy Condition 1, and let $E > 0$. Then*

$$(3.8) \quad \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2}$$

converges in the norm operator topology. Furthermore, the limit operator $A(E)$ is compact in $L^2(\partial\Omega)$.

Proof. This follows immediately from (4.8) in the proof of Lemma 4.2, which shows that the tail of the sum in (3.7) has vanishing operator norm. A is therefore also the norm limit of a sequence of finite-rank operators. \square

4. Proof of Theorems 1.1 and 1.2. In the previous section we related tension to the norm of a boundary operator which itself can be written as a sum involving mode boundary functions. Here we place upper bounds on $\|A(E)\|$ in order to prove Theorems 1.1 and 1.2. First we note that when E is an eigenvalue, Theorem 1.1 is trivially satisfied, since $t[u_{\min}] = 0$. When E is a noneigenvalue, formula (3.7) enables us to split up contributions from different parts of the Dirichlet spectrum,

$$(4.1) \quad A(E) = A_{\text{near}}(E) + A_{\text{far}}(E) + A_{\text{tail}}(E),$$

where

$$(4.2) \quad A_{\text{near}}(E) = \sum_{|E_j - E| \leq E^{1/2}} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2},$$

$$(4.3) \quad A_{\text{far}}(E) = \sum_{E/2 \leq E_j \leq 2E, |E_j - E| > E^{1/2}} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2},$$

$$(4.4) \quad A_{\text{tail}}(E) = \sum_{E_j < E/2} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2} + \sum_{E_j > 2E} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2}.$$

It is sufficient (due to the operator triangle inequality) to bound the norms of these three terms independently. We first tackle the “far” and “tail” terms.

LEMMA 4.1. *There is a constant C dependent only on Ω such that*

$$(4.5) \quad \|A_{\text{far}}(E)\| \leq C \quad \text{for all } E > 1.$$

Proof. For any $E > 1$, consider the spectral interval $I_m := [E + mE^{1/2}, E + (m + 1)E^{1/2}]$. For any such interval lying in $[E/2, 2E]$ we may apply Theorem 1.3, with E

replaced by at most $2E$, to bound $\|\sum_{E_j \in I_m} \psi_j \langle \psi_j, \cdot \rangle\|$ by $2C_\Omega E$. For terms in (4.3) associated with this interval, the denominators are no less than $m^2 E$. Thus

$$(4.6) \quad \left\| \sum_{E_j \in I_m} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2} \right\| \leq \frac{2C_\Omega}{m^2}.$$

Covering $[E + E^{1/2}, 2E]$ by summing over $m = 1, 2, \dots$ gives a constant, since $\sum m^{-2} = \pi^2/6$. The same argument applies for intervals covering $[E/2, E - E^{1/2}]$. \square

LEMMA 4.2. *There is a constant C dependent only on Ω such that*

$$(4.7) \quad \|A_{\text{tail}}(E)\| \leq CE^{-1/2} \quad \text{for all } E > 1.$$

Proof. Consider a spectral interval $I_m := [2^m E, 2^{m+1} E]$. We may cover this with at most $2^{m/2-1} E^{1/2} + 1$ windows of half-width at most $2^{m/2} E^{1/2}$; for each of these windows Theorem 1.3 applies to bound $\|\sum_{E_j \in I_m} \psi_j \langle \psi_j, \cdot \rangle\|$ by $C_\Omega 2^{m+1} E$. For each $E_j \in I_m$, the denominator is no smaller than $(2^{m-1} E)^2$. Thus

$$(4.8) \quad \left\| \sum_{E_j \in I_m} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2} \right\| \leq (2^{m/2-1} E^{1/2} + 1) \frac{C_\Omega 2^{m+1} E}{(2^{m-1} E)^2} \\ = C_\Omega (2^{-m/2-2} E^{-1/2} + 2^{-m+1} E^{-1}).$$

The infinite sum over $m = 1, 2, \dots$ gives

$$(4.9) \quad \left\| \sum_{E_j > 2E} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2} \right\| \leq C_\Omega \left(\frac{E^{-1/2}}{4(\sqrt{2}-1)} + 2E^{-1} \right) \leq CE^{-1/2} \quad \text{for all } E > 1.$$

We treat the interval $(0, E/2)$ similarly, using a sequence of intervals $J_m := [2^{-m-1} E, 2^{-m} E]$. Each such interval may be covered by at most $2^{-(m+3)/2} E^{1/2} + 1$ windows of half-width $2^{-(m+1)/2} E^{1/2}$. For each $E_j \in J_m$, the denominator is no smaller than $E^2/4$. In a similar manner as before, the operator norm of the partial sum associated with J_m is then $O(2^{-m} E^{-1/2})$, thus the infinite sum over m is $O(E^{-1/2})$. Note that Theorem 1.3 does not apply for $E < 1$, but that there are $O(1)$ such E_j values and each contributes $O(E^{-1})$. This proves Lemma 4.2. \square

Proof of Theorem 1.1. Examining the “near” term (4.2), we use Theorem 1.3 on the sum of numerators, and get a bound by taking the minimum denominator,

$$(4.10) \quad \|A_{\text{near}}(E)\| \leq \frac{C_\Omega E}{d(E, \sigma)^2} \quad \text{for all } E > 1.$$

Using this and Lemmas 4.1 and 4.2 to sum the terms in (4.1) gives

$$(4.11) \quad \|A(E)\| \leq \frac{C_\Omega E}{d(E, \sigma)^2} + C \quad \text{for all } E > 1.$$

From Lemma B.1, an upper bound on the distance to the spectrum, we see that the second term is bounded by at most a constant times the first, so it may be absorbed into it to give

$$(4.12) \quad \|A(E)\| \leq \frac{CE}{d(E, \sigma)^2} \quad \text{for all } E > 1.$$

Combining this with (3.6) proves (1.8), hence also the second inequality in (1.9). The first inequality in (1.9) simply follows from the fact that, since A is a sum of positive operators,

$$(4.13) \quad t[u_{\min}]^{-2} = \|A(E)\| \geq \left\| \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2} \right\| = \frac{\|\psi_j\|^2}{d(E, \sigma)^2},$$

where E_j is the eigenvalue closest to E . Using the lower bound $\|\psi_j\|^2 \geq cE_j$ from [19] (4.13) becomes

$$(4.14) \quad d(E, \sigma) \geq c\sqrt{E_j}t[u_{\min}].$$

With a change of constant, E_j may be replaced here by E to give the first inequality in (1.9), since Lemma B.1 ensures that E_j is relatively close to E . (The lemma is not useful for E less than some constant and $E_j < E$, but then the ratio E/E_j is still bounded by a constant because $E_j \geq E_1$.)

Proof of Theorem 1.2. We next prove the eigenfunction error bound (1.10), first considering E a noneigenvalue. We denote the boundary data by $U := u|_{\partial\Omega}$. From orthogonality, then using the formula for the c_i coefficients below (3.3), we get

$$(4.15) \quad \begin{aligned} \|u - \hat{u}_j\|_{L^2(\Omega)}^2 &= \sum_{E_i \neq E_j} |(\phi_i, u)_{L^2(\Omega)}|^2 = \sum_{E_i \neq E_j} \frac{|\langle \psi_i, U \rangle|^2}{(E - E_i)^2} \\ &\leq \left\| \sum_{E_i \neq E_j} \frac{\psi_i \langle \psi_i, \cdot \rangle}{(E - E_i)^2} \right\| \|U\|_2^2. \end{aligned}$$

The operator in the last expression is identical to (3.7) except with the E_j -eigenspace terms omitted. Therefore, its norm may be bounded in the same way as that of $A(E)$, the only difference being that the $d(E, \sigma)$ introduced in (4.10) is replaced by $\min_{E_i \neq E_j} |E - E_i| = |E - E_k|$. Thus the bound analogous to (4.12) is

$$\left\| \sum_{E_i \neq E_j} \frac{\psi_i \langle \psi_i, \cdot \rangle}{(E - E_i)^2} \right\| \leq \frac{CE}{(E - E_k)^2} \quad \text{for all } E > 1,$$

and inserting this and $\|U\|_{L^2(\partial\Omega)} = t[u]\|u\|_{L^2(\Omega)} = t[u]$ into (4.15) gives (1.10). Finally, if E is an eigenvalue, i.e., $E = E_j$, the solution operator (3.4) is undefined, since a solution to (3.1)–(3.2) exists if and only if f is orthogonal to the normal derivative functions in the E -eigenspace. This can be seen by applying Green's 2nd identity to ϕ , any function in the E -eigenspace, and u , giving $\langle \partial_n \phi, U \rangle = 0$. However, the solution coefficients c_i for which $E_i \neq E$ are uniquely defined by the same formula as before. Thus (4.15) and the rest of the proof carries through.

Remark 4.1. Theorem 1.2 is sharp, as can be seen in the following way: if u is such that $t[u]$ is close to $t[u_{\min}]$ (say, less than $2t[u_{\min}]$), then we have, by combining (1.9) and (1.10),

$$(4.16) \quad \|u - \hat{u}_j\|_{L^2(\Omega)} \leq C \frac{|E - E_j|}{|E - E_k|}.$$

Apart from the value of the constant, one cannot expect to do better than this. For example, if E is midway between E_j and E_k , then the error $\|u - \hat{u}_j\|_{L^2(\Omega)}$ cannot be expected to be better than $1/\sqrt{2}$.

5. Star-shaped planar domains. The purpose of this section is to strengthen Theorem 1.1 in the special case of star-shaped domains in $n = 2$. We take weighted boundary functions

$$(5.1) \quad \psi_j^{(s)}(s) := (\mathbf{x}(s) \cdot \mathbf{n}(s)) \partial_n \phi_j(s), \quad s \in \partial\Omega,$$

and our boundary inner product as

$$(5.2) \quad \langle f, g \rangle_s := \int_{\partial\Omega} (\mathbf{x}(s) \cdot \mathbf{n}(s))^{-1} f(s) g(s) ds,$$

hence norm $\|f\|_s := \sqrt{\langle f, f \rangle_s}$, and $t_s[u] := \|U\|_s / \|u\|_{L^2(\Omega)}$. The significance of the weight $(\mathbf{x} \cdot \mathbf{n})$ is twofold: it is strictly positive for strictly star-shaped domains, and also turns the inner product in (1.13) into $\langle \psi_i^{(s)}, \psi_j^{(s)} \rangle_s$, enabling us to benefit from pairwise quasi-orthogonality. The Rellich theorem $\|\psi_j^{(s)}\|_s^2 = 2E_j$ states that, with this special weight, there is no fluctuation in the L^2 -norms of the boundary functions. As shown in [4], the function $t_s[u_{\min}]$ versus E has slope $1/\|\psi_j^{(s)}\|_s^2$ in the neighborhood of E_j (this arises from dominance of a single term in (5.5) below). Hence these slopes are predictable *independently* of the particular form of each mode ϕ_j . This enables us to get the following eigenvalue inclusion result analogous to Theorem 1.1.

THEOREM 5.1. *Let $\Omega \subset \mathbb{R}^2$ be a strictly star-shaped bounded domain with piecewise smooth boundary. Then there are constants c_1, c_2, c_3 depending only on Ω , such that the following holds. Let $E > 1$, and suppose u is a nontrivial solution of $(\Delta + E)u = 0$ in $C^\infty(\Omega)$, with $c_2 t_s[u]^2 < 1$. Let $F := E + \sqrt{E}$. Then,*

$$(5.3) \quad d(E, \sigma) \leq \sqrt{2F} t_s[u] \frac{1 + c_1 \sqrt{F} t_s[u]}{1 - c_2 t_s[u]^2}.$$

For the Helmholtz solution u_{\min} minimizing $t_s[u]$ at the given E ,

$$(5.4) \quad \sqrt{2(E - c_3 E^{1/2})} t_s[u_{\min}] \leq d(E, \sigma).$$

Remark 5.1. In the limit of high frequency $E \gg 1$ and small tension $t_s[u] \ll E^{-1/2}$, the right-hand side of (5.3) and the left-hand side of (5.4) are both $\sqrt{2E}(1 + o(1))t_s$. Combining them proves that both the power of E and the constant $\sqrt{2}$ are sharp.

Remark 5.2. Notice that this theorem is not applicable for all E since there may be large spectral gaps where $c_2 t_s[u]^2 < 1$ cannot be satisfied. Due to the numerator, it becomes far from optimal when $t_s[u]$ is $O(E^{-1/2})$ or larger. In these respects it is less general than Theorem 1.1, even though it gives better bounds in the small tension limit.

The main tool used in the proof of Theorem 5.1 is the pairwise quasi-orthogonality result, Theorem 1.4, together with the Cotlar–Stein lemma, which we state here for the special case of self-adjoint operators:

LEMMA 5.2 (Cotlar–Stein lemma [12, 32, 11]). *Let $\{T_j\}_{j \in J}$ be a countable set of bounded self-adjoint operators, $J \subset \mathbb{N}$. Then*

$$\left\| \sum_{j \in J} T_j \right\| \leq \max_{j \in J} \sum_{i \in J} \sqrt{\|T_i T_j\|}.$$

Proof of Theorem 5.1. The weighted equivalent of (3.7) is the operator

$$(5.5) \quad A^{(s)}(E) = \sum_{j=1}^{\infty} \frac{\psi_j^{(s)} \langle \psi_j^{(s)}, \cdot \rangle_s}{(E - E_j)^2},$$

which, by analogy with (3.6), satisfies

$$(5.6) \quad t_s[u]^{-2} \leq \|A^{(s)}(E)\|_s.$$

The lower bound (5.4) follows by analogy with (4.13)–(4.14), using $\|\psi_j^{(s)}\|_s^2 = 2E_j$, and $E_j \geq E - c_3 E^{1/2}$ from Lemma B.1.

Using the same splitting into “near,” “far,” and “tail” parts as in section 4, we can bound the norm of the “near” part in a new way, as follows.

LEMMA 5.3. *There is a constant $c_1 > 0$ depending only on Ω such that*

$$\|A_{\text{near}}^{(s)}(E)\|_s \leq \frac{2F}{d(E, \sigma)^2} + \frac{\sqrt{2}c_1 F}{d(E, \sigma)} \quad \text{for all } E > 1.$$

The first term in this bound will arise simply from the single term in the sum (5.5) with E_j nearest to E . The second term requires more work, as we now show.

Proof. Let $J = \{j : |E_j - E| \leq E^{1/2}\}$. Using $T_j = \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2}$ in Lemma 5.2 gives

$$(5.7) \quad \left\| \sum_{j \in J} T_j \right\|_s \leq \max_{j \in J} \frac{\|\psi_j^{(s)}\|_s^{1/2}}{|E - E_j|} \sum_{i \in J} \frac{(\langle \psi_i^{(s)}, \psi_j^{(s)} \rangle_s \|\psi_i^{(s)}\|_s)^{1/2}}{|E - E_i|}.$$

Applying quasi-orthogonality (Theorem 1.4) for the inner product, and $\|\psi_j^{(s)}\|_s^2 = 2E_j$, and separating diagonal ($i = j$) from off-diagonal terms, we get

$$(5.8) \quad \left\| \sum_{j \in J} \frac{\psi_j \langle \psi_j, \cdot \rangle}{(E - E_j)^2} \right\|_s \leq \max_{j \in J} \frac{2E_j}{(E - E_j)^2} + \sqrt{2}S \max_{j \in J} \frac{E_j^{1/4}}{|E - E_j|} \sum_{i \in J} \frac{E_i^{1/4} |E_i - E_j|}{|E - E_i|}.$$

Here S is as in Theorem 1.4. The first term is bounded by $2F/d(E, \sigma)^2$. Using $|E_i - E_j| \leq |E_i - E| + |E - E_j|$ bounds the second term by

$$(5.9) \quad \begin{aligned} & \sqrt{2}S \max_{j \in J} \frac{E_j^{1/4}}{|E - E_j|} \sum_{i \in J} E_i^{1/4} \left(1 + \frac{|E - E_j|}{|E - E_i|} \right) \\ & \leq \sqrt{2FS} |J| \max_{j \in J} \left(\frac{1}{|E - E_j|} + \frac{1}{d(E, \sigma)} \right) \leq \frac{2\sqrt{2FS}|J|}{d(E, \sigma)}. \end{aligned}$$

Recall Weyl's law for the asymptotic density of eigenvalues, which states that, for $n = 2$ and $\text{vol } \Omega = 1$,

$$(5.10) \quad N(E) := \#\{j : E_j < E\} = \frac{1}{4\pi} E + R(E),$$

where the remainder is $R(E) = O(\sqrt{E})$ (see [29]; for the case of piecewise-smooth boundary, see [31, eq. (0.3)]). Since the remainder is bounded for small E , there is a

constant C_w such that $|R(E)| \leq C_w \sqrt{E}$ for all $E > 1$. Thus $|J|$, the number of terms in the “near” window, is bounded by

$$|J| \leq \left(\frac{1}{4\pi} + 2C_w \right) \sqrt{F}.$$

Inserting this into (5.9) proves Lemma 5.3, and we may take $c_1 = 2S(1/4\pi + 2C_w)$. \square

Completion of the proof of Theorem 5.1. The proofs of analogously weighted versions of Lemmas 4.1 and 4.2 are unchanged. So we may combine them with Lemma 5.3 and (5.6) to get, for some constant c_2 ,

$$t_s[u]^{-2} \leq \|A^{(s)}(E)\|_s \leq \frac{2F}{d(E, \sigma)^2} + \frac{\sqrt{2}c_1 F}{d(E, \sigma)} + c_2 \quad \text{for all } E > 1.$$

Multiplying through by $d(E, \sigma)^2$ we solve the quadratic inequality for $d(E, \sigma)$,

$$d(E, \sigma) \leq \frac{c_1 F / \sqrt{2} + \sqrt{c_1^2 F^2 / 2 + 2(t_s[u]^{-2} - c_2)F}}{t_s[u]^{-2} - c_2}.$$

Using the subadditivity of the square root completes the proof of (5.3). \square

6. Discussion of explicit constants. For the practical application of Theorems 1.1 and 1.2, it is important to have an explicit value for the constant C (from the discussion after (4.15) we notice that C in the two theorems is the same).

We now compute an explicit value of this C that holds for all $E > 1$. Examining (2.9) we see that a choice of constant in Lemma 2.1, and hence Theorem 1.3, that holds for all $E > 1$ is $C_\Omega = 4(C_a + C'_a) + \sqrt{2}C''_a$. To compute this we need sup norms of the value, and first and second derivative, of a vector field \mathbf{a} as in Lemma 2.2.

The proof of Lemma 2.2 shows such a construction; the values will depend on the size of the vectors \mathbf{a}_{x_i} and the choice of partition of unity used to cover $\overline{\Omega}$. The vectors \mathbf{a}_{x_i} will be large (order $1/\epsilon$) if Ω has corners with angles less than ϵ or greater than $2\pi - \epsilon$. We note that a numerical procedure for this construction could be useful.

In some special cases, a simpler prescription for the vector field can be given:

- For strictly star-shaped domains in \mathbb{R}^n , we may choose $\mathbf{a} = \mathbf{x} / \inf_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})$, which gives $C_a = \sup_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n}) / \inf_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})$, $C'_a = 1 / \inf_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})$, and $C''_a = 0$.
- For a domain with C^2 boundary, let $\delta > 0$ be the largest number such that for each $\mathbf{x}_0 \in \partial\Omega$, a ball of radius δ can be placed within Ω so as to be tangent to $\partial\Omega$ at \mathbf{x}_0 . We may then choose $\mathbf{a} = (1 - r/\delta)^2 \mathbf{n}_r$, for $r < \delta$, $\mathbf{a} = \mathbf{0}$ otherwise, where the coordinate r is the distance from $\partial\Omega$, and \mathbf{n}_r is the unit vector in the local decreasing r direction. This gives constants $C_a = 1$ and $C'_a = 2/\delta$. C''_a depends on δ and an upper bound on the rate of change of surface curvature. (Also note that a slight modification of the proof of Theorem 1.3 would allow estimation purely in terms of C_a and C'_a , but with a doubling of the numerical constants.)

Summing the terms of the form (4.6) above and below E we have that the constant in Lemma 4.1 is $2\pi^2 C_\Omega / 3$. Similarly, using (4.9) and its equivalent for $(0, E/2)$ gives the constant in Lemma 4.2 as $C_\Omega \left(\frac{1}{4(\sqrt{2}-1)} + \frac{1}{4-\sqrt{2}} + 6 \right) < 7C_\Omega$. Summing these two constants gives a constant C in (4.11) as $14C_\Omega$. A choice of constant in (4.12) is then $C_\Omega + 14C_\Omega \max[E_1^2, C_d^2]$, where from Appendix B we have $C_d = 2\sqrt{E_1}$, and the max accounts for the case $1 < E \leq E_1$. Finally, the constant in (1.8) is the square root of this, $C = \sqrt{C_\Omega(1 + 14 \max[E_1, 4]E_1)}$.

Requiring that the above estimates hold for all $E > 1$ caused nonoptimality in the choice of constant. It is more sensible in high-frequency applications to use a better constant which is approached for $E \gg 1$, and small tension $t \ll 1$. We now give this explicitly. In this limit, in (2.9), F tends to E , and we drop lower-order terms to get $C_\Omega = 2(C_a + C'_a)$, which in the star-shaped case is

$$(6.1) \quad C_\Omega = 2 \frac{1 + \sup_{\partial\Omega}(\mathbf{x} \cdot \mathbf{n})}{\inf_{\partial\Omega}(\mathbf{x} \cdot \mathbf{n})} \quad \text{for } E \gg 1, \Omega \text{ star-shaped.}$$

If tension is small (i.e., E is not in a large spectral gap), the second term in (4.12) is negligible, so we may approximate the constant in (1.8) as

$$(6.2) \quad C = \sqrt{C_\Omega} \quad \text{for } E \gg 1, t \ll 1.$$

Remark 6.1. The limiting constant (6.2) does not reflect the limiting slopes of the graph $t[u_{\min}]$ versus E near eigenvalues. These slopes are known [4] to be $1/\|\psi_j\|^2$, which is bounded by $(2C'_a E_j)^{-1}$ [19].

We end by discussing the constants c_1 and c_2 in Theorem 5.1. Constant c_2 may be estimated easily, as above, using the weighted versions of Lemmas 4.1 and 4.2. In the proof of Lemma 5.3, c_1 involves the Weyl constant C_w ; we know of no explicit estimates for C_w in the literature (the closest we know are estimates of the form $|R(E)| < C\sqrt{E} \ln E$ with explicit constants [26, 10]). However, these constants are effectively irrelevant for practical purposes, when $E \gg 1$ and $t \ll E^{-1/2}$, since in these limits, one may replace (5.3) by $d(E, \sigma) \leq \sqrt{2Et_s}[u]$ and still have an error bound very close to that given by the full expression.

7. Numerical example. In Figure 1.1(c) we show a planar nonconvex domain given by the radial function $r(\theta) = 1 + 0.3 \cos[3(\theta + 0.2 \sin \theta)]$. The domain is star-shaped and smooth (we will not address numerical issues raised by corners here; see [16, 14, 9, 15, 3, 7]). For high-frequency eigenvalue problems, a convenient computational basis of Helmholtz solutions are “method of fundamental solutions” basis functions $\xi_n(\mathbf{x}) = Y_0(\sqrt{E}|\mathbf{x} - \mathbf{y}_n|)$, where Y_0 is the irregular Bessel function of order zero, and $\{\mathbf{y}_n\}_{n=1}^N$ are a set of “charge points” in $\mathbb{R}^2 \setminus \Omega$. The latter were chosen by a displacement of the boundary parametrization $\mathbf{x}(\theta)$, $0 < \theta \leq 2\pi$, in the imaginary direction (see [6]); specifically $\mathbf{y}_n = \mathbf{x}(2\pi n/N - 0.025i)$.

We compute the data plotted in Figure 1.1(a) and (b) as follows. At each E , $t[\tilde{u}_{\min}]$ is given by the square root of the smallest generalized eigenvalue of a generalized eigenvalue problem (GEVP) involving $N \times N$ symmetric real dense matrices F and G (the basis representations of the boundary and interior norms, respectively). Both matrices are evaluated using M -point periodic trapezoidal quadrature in θ , that is, quadrature points $\mathbf{x}_m = \mathbf{x}(2\pi m/M)$, $m = 1, \dots, M$, and weights $w_m = 2\pi|\mathbf{x}'(2\pi m/M)|/M$. For instance, $F = P^*P$, where $P \in \mathbb{R}^{M \times N}$ has elements

$$(7.1) \quad P_{mn} = \sqrt{w_m} \xi_n(\mathbf{x}_m),$$

and G is similarly found [4, sec. 4.1] using P and the matrices $P^{(1)}$ and $P^{(2)}$ whose entries are the x_1 - and x_2 -derivatives of those in P . Since this GEVP is numerically singular, regularization was first performed, similarly to [8, sec. 6], by restricting to an orthonormal basis for the numerical column space of $[P; P^{(1)}; P^{(2)}]$ comprising the left singular vectors with singular values at least 10^{-14} times the largest singular value.

For low frequencies (Figure 1.1(a)), 8-digit accuracy requires $N = 100$ basis functions and $M = 200$ quadrature points. For higher frequencies corresponding to 40

wavelengths across the domain (Figure 1.1(b) and (c)), it requires $N = 400$ and $M = 500$, and the above GEVP procedure takes three seconds per E value.³ Very small ($< 10^{-8}$) tensions cannot be found this way, and instead are best approximated via the GSVD [8]: the optimal tension at a given E is the lowest generalized singular value of the matrix pair (P, Q) , where matrix Q has entries

$$(7.2) \quad Q_{mn} = \sqrt{\frac{\mathbf{x}_m \cdot \mathbf{n}_m}{2E}} \sqrt{w_m} \frac{\partial \xi_n}{\partial n}(\mathbf{x}_m),$$

where \mathbf{n}_m is the normal at \mathbf{x}_m , and regularization as before. Note that $G = Q^*Q$ well approximates the interior norm in the subspace with zero Dirichlet data, due to the Rellich formula (case $i = j$ of Theorem 1.4).

Any single-variable function minimization algorithm may then be used to search for a local minimum of $t[\tilde{u}_{\min}]$ versus E ; we prefer iterated fitting of a parabola to $t[\tilde{u}_{\min}]^2$ at three nearby E values, which converges typically in five iterations. Using this with the GSVD (with $N = 500$, $M = 700$, i.e., six points per wavelength on $\partial\Omega$, and taking eight seconds per iteration), we find the tension

$$(7.3) \quad t[\tilde{u}_{\min}] = 2.2 \times 10^{-12} \quad \text{at } E = 10005.0213579739.$$

This is shown by the dot in Figure 1.1(b). The GSVD right singular vector gives the basis coefficients of the corresponding trial function \tilde{u}_{\min} , which is plotted in Figure 1.1(c) (this took 34 seconds to evaluate on a square grid of size 0.005, i.e., 1.3×10^5 points).

Armed with datapoint (7.3), what can we deduce about a Dirichlet eigenpair of Ω using our new theorems, and how much better are they than previous results? The constant in the Moler–Payne bound (1.5) is $C_{\text{MP}} = q_1^{-1/2}$ where q_1 is the lowest eigenvalue of a Stekloff eigenproblem on Ω [20, eq. (2.11)]. Since Ω is star-shaped, the bound $q_1 \geq E_1^{1/2} \inf_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n}) / 2 \sup_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})$ from [20, Table I] applies, giving $C_{\text{MP}} = 1.31$ as a valid choice. Thus (1.5) states that there is an eigenvalue E_j a distance no more than 2.9×10^{-8} from the above E . On the other hand, (6.2) and (6.1) give the constant in Theorem 1.1 as $C = 2.9$. Applying the theorem gives a distance from the spectrum of no more than 6.3×10^{-10} . Taking the small-tension limit of the star-shaped planar result (5.3), and recomputing the weighted tension $t_s[\tilde{u}_{\min}]$ from section 5, we get an even smaller distance of 3.5×10^{-10} , that is, an error of ± 3 in the last digit of (7.3). The latter is a 80-fold improvement over Moler–Payne. (Also, see [4] for an example at higher frequency with three digits of improvement.)

How good an approximation is \tilde{u}_{\min} to the eigenfunction ϕ_j ? Using the observation that the next nearest eigenvalue is $E_k = 10007.339 \dots$, the eigenfunction bound of Moler and Payne [24] gives an L^2 -error of 1.2×10^{-8} . With the same data, using the constant C above, Theorem 1.2 gives an L^2 -error of 2.7×10^{-10} , a 50-fold improvement over that achievable with previously known theorems.

8. Conclusions. We have improved, by a factor of the wavenumber, the Moler–Payne bounds on Dirichlet eigenvalues and eigenfunctions which have been the standard for the last 40 years. This makes rigorous the conjecture based on numerical observations in [4]. We expect this to be useful since high-frequency wave and eigenvalue calculations are finding more applications in recent years. Of independent interest is a new quasi-orthogonality result in a spectral window (Theorem 1.3).

³All computation times are reported for a laptop with 2GHz Intel Core Duo processor and 2GB RAM, running MATLAB 2008a on a Linux kernel.

For numerical utility, throughout we have been explicit with constants, and have specified a lower bound on E for which the estimates hold (this being stronger than merely a “big- O ” asymptotic estimate). For star-shaped domains we strengthened the inclusion bounds (Theorem 5.1), achieving a sharp power of E and sharp constant, in the limit of small tension, when tension is weighted by a special geometric function. This weight allowed pairwise quasi-orthogonality to be used, but since an upper bound for the number of eigenvalues in a \sqrt{E} window is needed, this is only useful for $n = 2$ (planar domains).

We applied our theorems to a numerical example, enabling close to 14 digit accuracy in a high-lying eigenvalue, and 10 digits in the eigenfunction. Both are two digits beyond what could be claimed with previously known theorems.

Our estimate $C\sqrt{E}t[u]$ on the distance to the spectrum is sharp (up to constants) if the tension $t[u]$ (or $t_s[u]$) is comparable to $t[u_{\min}]$ (or $t_s[u_{\min}]$). However, numerically, one generally has access to other properties of u (e.g., its normal derivative) which could give more detailed information about the spectrum. For example, the powerful “scaling method” [36, 3] is able to locate many eigenvalues using an operator computed at a single energy E . In another direction, Still [33, Thm. 4] obtains improved inclusion bounds when the approximate eigenvalue is given by a Rayleigh quotient; in this case, the bound is proportional to $t[u]^2$, but the scaling for large energy is worse, being E^2 .

An open problem whose solution would have practical benefits is the generalization of our results to Neumann and Robin boundary conditions, and to multiple subdomains with different trial functions on each subdomain and least-square errors on artificial boundaries [14, 15, 7] (these are known as Trefftz or nonpolynomial finite element methods).

Appendix A. Proof of Lemma 2.2.

Proof. For every boundary point x , we can find a constant vector field \mathbf{a}_x having the property (2.3) in a neighborhood U_x of x . (Take a multiple of the vector field ∂_{z_n} in the Euclidean coordinate system used in (1.7).) By compactness we can find a finite number of such neighborhoods $U_i = U_{x_i}$, $i = 1, \dots, N$, covering $\partial\Omega$. We can add to this collection of open sets one additional set U_0 , whose closure does not meet $\partial\Omega$, yielding an open cover of $\bar{\Omega}$. Let ϕ_i , $i = 0, \dots, N$, be a smooth partition of unity subordinate to this open cover. Then

$$\mathbf{a} = \sum_{i=1}^N \phi_i \mathbf{a}_{x_i}$$

is a vector field with the required property. \square

Appendix B. Upper bound on distance to spectrum.

LEMMA B.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. Let E_1 be the lowest Dirichlet eigenvalue of Ω . Then for any $E > E_1$,*

$$d(E, \sigma) \leq C_d E^{1/2}, \quad C_d = 2\sqrt{E_1}.$$

Remark B.1. The result becomes interesting only for $E > 2(1 + \sqrt{2})E_1$. Bounds on E_1 exist as follows. If Ω contains a Euclidean ball of radius r , then E_1 is less than or equal to $E_1(B(0, r))$, which is equal to $j_{n/2-1,1}^2/r^2$, where $j_{n/2-1,1}$ is the first positive zero of the Bessel function $j_{n/2-1}$. For $n = 2$ we have $j_{0,1} = 2.4048\dots$ and for $n = 3$ we have $j_{1/2,1} = \pi$. Also, E_1 is greater than or equal to the first eigenvalue of the ball having the same n -volume as Ω , by the Faber–Krahn inequality [27].

Proof. Choose a wavevector $\mathbf{k} \in \mathbb{R}^n$ with $|\mathbf{k}|^2 = E - E_1$, and consider the trial function $u : \Omega \rightarrow \mathbb{C}$ defined by $u(\mathbf{x}) := \phi_1(\mathbf{x})e^{i\mathbf{k} \cdot \mathbf{x}}$, where ϕ_1 is the normalized first Dirichlet eigenmode of Ω with eigenvalue E_1 . We calculate

$$(\Delta + E)u = 2i\mathbf{k} \cdot \nabla \phi_1 e^{i\mathbf{k} \cdot \mathbf{x}}.$$

Since u is in the domain of Ω , has norm $\|u\|_{L^2(\Omega)} = 1$, and

$$(B.1) \quad \|(\Delta + E)u\|^2 = 4 \int_{\Omega} |\mathbf{k} \cdot \nabla \phi_1|^2 d\mathbf{x} \leq 4(E - E_1)E_1 < 4EE_1,$$

we see that u is an $O(E^{1/2})$ quasi-mode. On the other hand, writing $u = \sum_{j=1}^{\infty} a_j \phi_j$, we have

$$(B.2) \quad \begin{aligned} \|(\Delta + E)u\|^2 &= \left\| \sum_j a_j (E - E_j) \phi_j \right\|^2 = \sum_j |a_j|^2 (E - E_j)^2 \\ &\geq d(E, \sigma)^2 \sum_j |a_j|^2 = d(E, \sigma)^2. \end{aligned}$$

Combining (B.1) and (B.2) completes the proof. \square

Acknowledgments. The authors are grateful for discussions with Dana Williams, Timo Betcke, and Chen Hua.

REFERENCES

- [1] A. ANCONA, *A note on the Rellich formula in Lipschitz domains*, Publ. Math., 42 (1998), pp. 223–237.
- [2] A. BÄCKER, S. FÜRSTBERGER, R. SCHUBERT, AND F. STEINER, *Behaviour of boundary functions for quantum billiards*, J. Phys. A, 35 (2002), pp. 10293–10310.
- [3] A. H. BARNETT, *Asymptotic rate of quantum ergodicity in chaotic Euclidean billiards*, Comm. Pure Appl. Math., 59 (2006), pp. 1457–1488.
- [4] A. H. BARNETT, *Perturbative analysis of the method of particular solutions for improved inclusion of high-lying Dirichlet eigenvalues*, SIAM J. Numer. Anal., 47 (2009), pp. 1952–1970.
- [5] A. H. BARNETT AND T. BETCKE, *Quantum mushroom billiards*, Chaos, 17 (2007), 043125.
- [6] A. H. BARNETT AND T. BETCKE, *Stability and convergence of the method of fundamental solutions for Helmholtz problems on analytic domains*, J. Comput. Phys., 227 (2008), pp. 7003–7026.
- [7] T. BETCKE, *A GSVD formulation of a domain decomposition method for planar eigenvalue problems*, IMA J. Numer. Anal., 27 (2007), pp. 451–478.
- [8] T. BETCKE, *The generalized singular value decomposition and the method of particular solutions*, SIAM J. Sci. Comput., 30 (2008), pp. 1278–1295.
- [9] T. BETCKE AND L. N. TREFETHEN, *Reviving the method of particular solutions*, SIAM Rev., 47 (2005), pp. 469–491.
- [10] H. CHEN, *Irregular but non-fractal drums, and n-dimensional Weyl conjecture*, Acta Math. Sinica (N.S.), 11 (1995), pp. 168–178.
- [11] A. COMECH, *Cotlar-Stein Almost Orthogonality Lemma*, preprint, available online at <http://www.math.tamu.edu/~comech/papers/CotlarStein/CotlarStein.pdf> (2007).
- [12] M. COTLAR, *A combinatorial inequality and its applications to L^2 -spaces*, Rev. Mat. Cuyana, 1 (1955), pp. 41–55.
- [13] D. JERISON AND C. E. KENIG, *The inhomogeneous Dirichlet problem in Lipschitz domains*, J. Funct. Anal., 130 (1995), pp. 161–219.
- [14] J. DESCLOUX AND M. TOLLEY, *An accurate algorithm for computing the eigenvalues of a polygonal membrane*, Comput. Methods Appl. Mech. Engrg., 39 (1983), pp. 37–53.
- [15] T. A. DRISCOLL, *Eigenmodes of isospectral drums*, SIAM Rev., 39 (1997), pp. 1–17.
- [16] L. FOX, P. HENRICI, AND C. MOLIER, *Approximations and bounds for eigenvalues of elliptic operators*, SIAM J. Numer. Anal., 4 (1967), pp. 89–102.

- [17] P. R. GARABEDIAN, *Partial Differential Equations*, John Wiley & Sons Inc., New York, 1964.
- [18] C. GORDON, D. WEBB, AND S. WOLPERT, *Isospectral plane domains and surfaces via Riemannian orbifolds*, *Invent. Math.*, 110 (1992), pp. 1–22.
- [19] A. HASSELL AND T. TAO, *Upper and lower bounds for normal derivatives of Dirichlet eigenfunctions*, *Math. Res. Lett.*, 9 (2002), pp. 289–305.
- [20] J. R. KUTTLER AND V. G. SIGILLITO, *Inequalities for membrane and Stekloff eigenvalues*, *J. Math. Anal. Appl.*, 23 (1968), pp. 148–160.
- [21] J. R. KUTTLER AND V. G. SIGILLITO, *Eigenvalues of the Laplacian in two dimensions*, *SIAM Rev.*, 26 (1984), pp. 163–193.
- [22] Z. C. LI, *The Trefftz method for the Helmholtz equation with degeneracy*, *Appl. Numer. Math.*, 58 (2008), pp. 131–159.
- [23] W. C. H. MCLEAN, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, Cambridge, UK, 2000.
- [24] C. B. MOLER AND L. E. PAYNE, *Bounds for eigenvalues and eigenvectors of symmetric operators*, *SIAM J. Numer. Anal.*, 5 (1968), pp. 64–70.
- [25] P. MONK AND D.-Q. WANG, *A least-squares method for the Helmholtz equation*, *Comput. Methods Appl. Mech. Engrg.*, 175 (1999), pp. 121–136.
- [26] Y. NETRUSOV AND Y. SAFAROV, *Weyl asymptotic formula for the Laplacian on domains with rough boundaries*, *Comm. Math. Phys.*, 253 (2005), pp. 481–509.
- [27] G. PÓLYA AND G. SZEGÖ, *Isoperimetric Inequalities in Mathematical Physics*, *Ann. Math. Stud.* 27, Princeton University Press, Princeton, NJ, 1951.
- [28] F. RELICH, *Darstellung der Eigenwerte von $\Delta u + \lambda u = 0$ durch ein Randintegral*, *Math. Z.*, 46 (1940), pp. 635–636.
- [29] Y. SAFAROV AND D. VASSILIEV, *The asymptotic distribution of eigenvalues of partial differential operators*, *Transl. Math. Monogr.* 155, American Mathematical Society, Providence, RI, 1997.
- [30] N. SAITO, *Data analysis and representation on a general domain using eigenfunctions of Laplacian*, *Appl. Comput. Harmon. Anal.*, 25 (2008), pp. 68–97.
- [31] R. SEELEY, *An estimate near the boundary for the spectral function of the Laplace operator*, *Amer. J. Math.*, 102 (1980), pp. 869–902.
- [32] E. M. STEIN, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, in *Monographs in Harmonic Analysis, III*, Princeton University Press, Princeton, NJ, 1993.
- [33] G. STILL, *Computable bounds for eigenvalues and eigenfunctions of elliptic differential operators*, *Numer. Math.*, 54 (1988), pp. 201–223.
- [34] L. N. TREFETHEN AND T. BETCKE, *Computed eigenmodes of planar regions*, *Contemp. Math.* 412, American Mathematical Society, Providence, RI, 2006, pp. 297–314.
- [35] H. E. TURECI, H. G. L. SCHWEFEL, P. JACQUOD, AND A. D. STONE, *Modes of wave-chaotic dielectric resonators*, *Progress in Optics*, 47 (2005), pp. 75–137.
- [36] E. VERGINI AND M. SARACENO, *Calculation by scaling of highly excited states of billiards*, *Phys. Rev. E*, 52 (1995), pp. 2204–2207.
- [37] S. ZELDITCH, *Quantum ergodicity and mixing of eigenfunctions*, in *Encycl. Math. Phys.*, Vol. 1, Elsevier, Oxford, 2006, pp. 183–196.
- [38] S. ZELDITCH AND M. ZWORSKI, *Ergodicity of eigenfunctions for ergodic billiards*, *Comm. Math. Phys.*, 175 (1996), pp. 673–682.