

9-1-1996

The Mass Distribution in the Nearby Universe

Luiz N. da Costa
European Southern Observatory

Wolfram Freudling
European Southern Observatory

Gary Wegner
Dartmouth College

Riccardo Giovanelli
Cornell University

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>

 Part of the [Cosmology, Relativity, and Gravity Commons](#)

Recommended Citation

da Costa, Luiz N.; Freudling, Wolfram; Wegner, Gary; and Giovanelli, Riccardo, "The Mass Distribution in the Nearby Universe" (1996). *Open Dartmouth: Faculty Open Access Articles*. 2294.
<https://digitalcommons.dartmouth.edu/facoa/2294>

This Article is brought to you for free and open access by Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Faculty Open Access Articles by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

THE MASS DISTRIBUTION IN THE NEARBY UNIVERSE

LUIZ N. DA COSTA

European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching bei München, Germany
Observatório Nacional, Rua Gen. José Cristino 77, São Cristovão, Rio de Janeiro, Brazil

WOLFRAM FREUDLING

Space Telescope-European Coordinating Facility and European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching bei München, Germany

GARY WEGNER

Department of Physics and Astronomy, Dartmouth College, Hanover, NH 03755

RICCARDO GIOVANELLI AND MARTHA P. HAYNES

Center for Radiophysics and Space Research and National Astronomy and Ionosphere Center,¹ Cornell University, Ithaca, NY 14853

AND

JOHN J. SALZER

Department of Astronomy, Wesleyan University, Middletown, CT 06457

Received 1996 March 21; accepted 1996 June 18

ABSTRACT

We present a new reconstruction of the mass density and the peculiar velocity fields in the nearby universe using recent measurements of Tully-Fisher distances for a sample of late spirals. We find significant differences between our reconstructed fields and those obtained in earlier work: overdensities tend to be more compact while underdense regions, consisting of individual voids, are more abundant. Our results suggest that voids observed in redshift surveys of galaxies represent real voids in the underlying matter distribution. While we detect a bulk velocity of $\sim 300 \text{ km s}^{-1}$, within a top-hat window 6000 km s^{-1} in radius, the flow is less coherent than previously claimed, exhibiting a bifurcation toward the Perseus-Pisces and the Great Attractor complexes. This is the first time that this feature is seen from peculiar velocity measurements. The observed velocity field resembles, more closely than any previous reconstruction, the velocity field predicted from self-consistent reconstructions based on all-sky redshift surveys. This better match is likely to affect estimates of the parameter $\beta = \Omega^{0.6}/b$ and its uncertainty based on velocity-velocity comparisons.

Subject headings: cosmic microwave background — cosmology: observations — galaxies: distances and redshifts — photometry

1. INTRODUCTION

Fluctuations in the mass distribution on intermediate scales can be probed by peculiar velocity measurements, by measurements of anisotropies in the cosmic background radiation at the 1° scale and by distortions of galaxy images induced by weak field lensing. Despite several pitfalls, the peculiar motion approach still remains the most direct and the easiest to interpret, provided that large-scale structures grow gravitationally and fluctuations in the mass distribution induce the observed peculiar velocities of galaxies. Mapping the peculiar velocity field can thus provide information useful to constrain the mass power-spectrum, the relation between the galaxy and matter distributions, and the value of the cosmological density parameter Ω on large scales.

In recent years, considerable effort has been made to understand the nature of the peculiar velocity field and the determination of the parameter $\beta = \Omega^{0.6}/b$, where b is the linear biasing parameter. Measurements of β have been attempted either from the comparison of the observed fluctuations in the galaxy distribution with those obtained from dynamical studies or from the direct comparison of the measured peculiar velocity field to that predicted from dynam-

ically self-consistent reconstruction of the density field from all-sky redshift surveys (for a review see Dekel 1994). These estimates based on peculiar velocity measurements have generally led to higher values than those obtained on small scales by other methods (e.g., Strauss & Willick 1995). The reason for this inconsistency remains an open question.

Until recently, most of the work was based on compilations of distance measurements by different authors which were sparse, inhomogeneous and probed a shallow volume in a nonuniform way (Bertschinger et al. 1990; Dekel 1994). Although several new peculiar velocity surveys have been carried out (Willick 1990; Freudling, Martel, & Haynes 1991; Courteau et al. 1993; Mathewson, Ford, & Buchhorn 1992b; Wegner et al. 1996), none covers the whole sky uniformly. Efforts to homogenize existing datasets are currently underway (Willick et al. 1996), but error estimates for the distances of individual galaxies are likely to continue to be uncertain. This directly affects the estimation of the bias which might, in the case of the inhomogeneous Malmquist bias correction, lead to spurious detection of density fluctuations.

The need for a sample with uniform sky coverage and homogeneous measurements has prompted us to carry out an independent redshift-distance survey of Sbc-Sc galaxies, based on I -band Tully-Fisher (TF) distances, covering essentially the whole sky ($\text{decl.} \geq -45^\circ$ and $|b| \geq 10^\circ$). The sample has well-defined selection criteria, and new measurements were

¹ The National Astronomy and Ionosphere Center is operated by Cornell University under a cooperative agreement with the National Science Foundation.

carried out in order to assure the homogeneity of the data. This data set was complemented with galaxies drawn from Mathewson et al. (1992b), appropriately converted to our magnitude and width scale, yielding a combined sample of about 1300 field galaxies (hereafter the SFI sample) and 500 galaxies in clusters. Details on sample selection and the procedure adopted to combine the two datasets can be found in Giovanelli et al. (1994, 1995a, 1995b). Here, we present a summary of the results of a reconstruction of the three-dimensional velocity and density fields obtained from the SFI sample. We use the three-dimensional reconstruction method primarily to illustrate the nature of the sample as compared to those currently available. A more quantitative analysis will be the subject of future papers. In § 2, we briefly describe the reconstruction method and discuss the approach used to correct for the various biases that affect the distance estimates. Our main results are presented in § 3, while in § 4 we summarize our conclusions.

2. ANALYSIS OF THE SFI SAMPLE

Distances of galaxies were estimated using the direct *I*-band TF relation derived from the combination of data for 24 clusters (Giovanelli et al. 1996a, 1996b). However, errors in the distance measurements lead to biases which directly affect the peculiar velocity field and thus the derived density field. The biases result from the coupling of uncertainties in the distance estimate with inhomogeneities in the galaxy distribution, and the selection of the sample. The latter is particularly important for the Sc sample because the adopted redshift-dependent selection criteria (e.g., Giovanelli et al. 1994) implies that near each redshift limit outflowing galaxies are preferentially excluded from the sample. This effect becomes dominant at large distances leading to a spurious systematic infall of galaxies at the outer edge of the surveyed volume. In order to deal with this problem we have resorted to a numerical approach to estimate the bias field at each point in estimated distance space (Freudling et al. 1995). The method, which is based on mock catalogs with characteristics similar to the data, allows for the simultaneous correction of the homogeneous and inhomogeneous Malmquist bias, and biases introduced by selection effects. The mock catalogs are generated from the distribution of optically selected galaxies in real space. This distribution was obtained using an iterative method of reconstructing a dynamically self-consistent distribution of galaxies from all-sky redshift surveys (*Optical-IRAS* sample) using a quasi-linear theory (Freudling, da Costa, & Pellegrini 1994).

The bias correction is sensitive to the value adopted for the scatter in distance estimates. Attempts to use the scatter as determined for the TF relation of galaxies in clusters led to an overcorrection of the velocity field. In order to overcome this problem we have also used the mock catalogs to determine the scatter in the distance estimates of field galaxies, taking advantage of the homogeneity of the sample. This was done by comparing the observed radial peculiar velocities at large distances with those of mock catalogs, since these velocities are primarily produced by biases which depend on the assumed scatter. The scatter of the mock samples was adjusted so that the radial variation of the peculiar velocities at large distances in the mock catalog match the observations. The derived scatter of ≈ 0.4 mag depends on the H I line width and is in any case smaller than that obtained for galaxies in clusters (Giovanelli et al. 1996a). A more extensive discussion of the

motivation and the details for the adopted procedure will be presented in da Costa et al. (1996).

Distances were calculated from the TF relation and the radial component of the peculiar velocity $u = cz_{\text{CMB}} - d_c$, where d_c is the “fully” corrected distance estimate. Galaxies in clusters were assigned to a single object. A small fraction of galaxies ($\sim 6\%$) with small line widths ($\log w < 2.25$) were discarded because of the large fractional errors in the measurement of their widths. The final input sample used in the reconstruction of the density field consists of 1234 independent points.

The computed distances and peculiar velocities are the input data to reconstruct the density and three-dimensional peculiar velocity fields. For the reconstruction, we use the standard assumption of irrotational flow which allows calculation of the scalar velocity potential field from the integral along radial paths of the line-of-sight component of the peculiar velocity. Our algorithm is similar to that discussed by Dekel, Bertschinger, & Faber (1990). We adopt a grid in spherical coordinates of 20 equal shells out to a maximum radius of 7500 km s^{-1} and 64 latitude and longitude circles. The measured radial peculiar velocities were smoothed using a Gaussian tensor window with a fixed smoothing scale of $R_w = 900 \text{ km s}^{-1}$. Our methodology was extensively tested using mock catalogs with the same number of objects and selection criteria as the data, which were also used to determine the uncertainties in the reconstruction (da Costa et al. 1996).

3. RESULTS

The reconstructed density fluctuation field δ along the supergalactic plane on a cubic grid of spacing 500 km s^{-1} for a fully corrected model is displayed in Figure 1 (Plate L1), where we show in the top plane density contours in $\delta = 0.2$ intervals, in the middle plane surface density maps with the contour level proportional to the density contrast δ and in the bottom plane density contrast maps. We use a Cartesian supergalactic coordinate system with SGX-axis pointing to $L = 0^\circ$ and the SGY-axis pointing nearly to the North Galactic Pole.

From visual inspection of the map one can recognize some well-known structures in the nearby volume such as the Great Attractor (GA, $\text{SGX} \sim -2000 \text{ km s}^{-1}$, $\text{SGY} \sim -500 \text{ km s}^{-1}$), the Perseus-Pisces (PP) complex ($\text{SGX} \sim 6000 \text{ km s}^{-1}$, $\text{SGY} \sim -1000 \text{ km s}^{-1}$) and the large void ($\text{SGX} \sim 2500 \text{ km s}^{-1}$, $\text{SGY} \sim 0 \text{ km s}^{-1}$) between the GA and the PP. We can also see traces of the Coma/A1367 supercluster along $\text{SGY} \sim 7000 \text{ km s}^{-1}$, and the Cetus region ($\text{SGX} \sim 500 \text{ km s}^{-1}$, $\text{SGY} \sim -6000 \text{ km s}^{-1}$). These structures are located roughly at the same position as in the *IRAS* reconstructed density field (e.g., Strauss & Willick 1995).

In comparison to earlier work (e.g., Dekel 1994), better sampling of the northern hemisphere apparently leads to significant differences in the density field especially in the region not covered by MFB. Some important differences can be seen: (1) in the foreground of the Coma/A1367 supercluster, where now a large void is detected; (2) in the PP region, which appears as a compact high density peak; and (3) in the Great Attractor region, which looks less prominent. Examination of the reconstructed density field in three dimensions also suggests that the GA consists of different substructures, much in the way seen in redshift surveys (e.g., Willmer et al. 1995).

A striking feature of our reconstructed density field is the existence of several voids surrounded by tenuous structures. Among them is the nearby void separated from the voids in the foreground of the Coma/A1367 supercluster by a small amplitude coherent structure, which we identify as the Local Supercluster. It is, as expected, much less prominent than other structures within the volume. Other prominent voids are the void behind the GA ($SGX \sim -6500 \text{ km s}^{-1}$, $SGY \sim 1500 \text{ km s}^{-1}$) and the Sculptor void ($SGX \sim -2000 \text{ km s}^{-1}$, $SGY \sim -6000 \text{ km s}^{-1}$). Both voids have been detected in galaxy redshift maps (e.g., da Costa et al. 1988; Geller & Huchra 1989) but are now seen in the mass distribution with density contrasts reaching $\delta \approx -0.6$, as indicated by the contours in the top plane.

Several other positive and negative fluctuations are seen near the outer edge of our survey volume. Comparison with the *Optical-IRAS* and *IRAS* density fields indicate that most are likely to be real structures. However, it is important to emphasize that the error in the density field derived from mock catalogs, which is smaller than 0.2 within 5000 km s^{-1} , grows rapidly with radius. At 6000 km s^{-1} the error is estimated to be about 0.4, corresponding to two contour levels. Therefore, the amplitude and location of structures beyond 5000 km s^{-1} are very poorly determined. This is also true for the Coma/A1367 and PP superclusters. However, although PP itself is poorly mapped, its influence in the velocity field extends over a large volume and into the region where the errors are small and the results insensitive to the details of the inhomogeneous bias correction.

The density field reconstructed with the SFI sample is characterized by overdensities which are compact and underdensities which are large, have regular shape and have a high density contrast. This is in marked contrast to earlier reconstructions. Our results suggest the existence of real voids in the matter distribution, qualitatively similar to those observed in galaxy redshift surveys (e.g., da Costa et al. 1994). This supports the contention that galaxies delineate real voids in the underlying matter distribution which may impose severe constraints in the amplitude of the mass power-spectrum (e.g., Piran et al. 1993).

In Figure 2, we show the velocity field along the supergalactic plane superimposed to the density field shown in Figure 1. This figure should be compared to a similar map presented by Dekel (1994). In contrast to Dekel's result, our velocity field is qualitatively similar to the predicted *IRAS* velocity field (e.g., Yahil 1988) showing a region where the flow bifurcates towards the GA and toward the PP complex. If true, this resolves a longstanding discrepancy between predicted and measured velocity fields. Moreover, an infall into PP is clearly observed while none is seen in Dekel's (1994) reconstruction. This contrasts with the interpretation of Willick (1990) who claims that "the principal feature of the velocity field in the PP region is a coherent streaming towards the Local Group." Willick's smaller sky coverage of the PP region could well be the cause of this difference.

On the opposite side of the figure, we find that the GA is less important than originally believed. Instead of being responsible for most of the motion in the nearby volume, there is strong evidence that it is smaller and less massive than originally estimated by Lynden-Bell et al. (1988). This is probably due to the larger surveyed volume and better sampling of the PP region. Note, in particular, that the amplitude of the backflow into the GA is small. Instead, the flow field in




FIG. 2.—Projected components (from the SFI sample) of the three-dimensional peculiar velocity field along the supergalactic plane superimposed to the density field. The contour spacing is 0.2 in δ , with the heavy solid line indicating $\delta = 0$. The coordinate system is as described in the text. The major structures seen in the map are the Great Attractor, the Perseus-Pisces complex, the Cetus region, and the foreground of the Coma/A1367 supercluster.

the upper left-hand side of Figure 2 could be explained by a mass concentration beyond the survey volume pulling the GA. This would explain the lack of an unambiguous signature for backside infall in the survey of Burstein, Faber, & Dressler (1990). Finally, the volume-weighted bulk velocity as measured within a top-hat window 6000 km s^{-1} in radius is about 300 km s^{-1} in the direction $L \sim 160^\circ$ and $B \sim -20^\circ$, where L and B are the supergalactic coordinates. These values are very close to those reported by Dekel (1994). However, the flow shows an overall shear unlike the coherence suggested by Mathewson, Ford, & Buchhorn (1992a) and Courteau et al. (1993).

There are several possible explanations for the observed differences between the results of our reconstruction and those of earlier work (e.g., Dekel 1994). The uniform sampling over the sky and depth is certainly a major factor, as pointed out above. Differences in the assumed scatter properties and methods for inhomogeneous Malmquist bias correction could also contribute to the position-dependent differences observed. Differences in the relative zero points between different data sets in the sample used by Dekel (1994) could be another reason for the observed discrepancies. A detailed comparison of the two datasets will be required to better understand the reason for the discrepancies.

4. CONCLUSIONS

The picture that emerges from our reconstruction brings together the somewhat fragmentary view previously held. The main results may be summarized as follows:

1. The positive fluctuations of the matter density field tend to be more compact while well-defined voids are more abundant than those obtained in previous reconstructions.

2. Comparison with redshift maps suggests that galaxies delineate real voids in the mass distribution. These voids tend to be separated by relative low-density structures which correspond to the filamentary and wall-like structures observed in the galaxy distribution.

3. The GA is significantly less prominent than previously claimed. There is also some indication for sub-structures.

4. An outside mass concentration may be pulling the GA. This induced motion overshadows the backflow and explains the lack of convincing evidence of such a flow.

5. The PP supercluster makes a significant contribution to the flow in its neighborhood leading to an infall into the supercluster seen for the first time from peculiar velocity measurements.

6. The measured mean bulk velocity within a top-hat window of 6000 km s^{-1} is about 300 km s^{-1} . The flow shows an overall shear which may be consistent with the existence of a mass concentration outside the observed volume.

Our results should affect conclusions drawn earlier regarding the value of β , or at least the accuracy with which it can be determined, because of the better agreement between the velocity fields reconstructed from peculiar velocity measurements and from the galaxy distribution. The results may also affect the conclusions regarding the biasing of galaxies relative

to the mass and the mass power spectrum. These topics will be pursued in more detail in future papers.

We would like to thank A. Dekel for many useful discussions and to the anonymous referee for his constructive comments. L. N. dC. would like to thank the hospitality of the Institut d'Astrophysique and the Hebrew University where part of this work was carried out. G. W. thanks for the hospitality of the European Southern Observatory, NSF grant AST93-47714, and the Alexander von Humboldt foundation for partial support. We acknowledge the financial support from US National Science Foundation grants AST94-20505 to R. G., AST92-18038 and AST90-23450 to MPH and from the Research Corporation to J. J. S. The results presented in this paper are based on observations carried out at the European Southern Observatory (ESO), the National Astronomy and Ionosphere Center (NAIC), the National Radio Astronomy Observatory (NRAO), the Kitt Peak National Observatory (KPNO), the Cerro Tololo Interamerican Observatory (CTIO), the Observatory of Paris at Nançay and the Michigan-Dartmouth-MIT Observatory (MDM). NAIC, NRAO, KPNO, and CTIO are, respectively, operated by Cornell University, Associated Universities, Inc., and Associated Universities for Research in Astronomy, all under cooperative agreements with the National Science Foundation.

REFERENCES

- Bertschinger, E., Dekel, A., Faber, S. M., Dressler, A., & Burstein, D. 1990, *ApJ*, 364, 370
 Burstein, D., Faber, S. M., & Dressler, A. 1990, *ApJ*, 354, 18
 Courteau, S., Faber, S. M., Dressler, A., & Willick, J. A. 1993, *ApJ*, 412, L51
 da Costa, L. N., et al. 1988, *ApJ*, 327, 544
 da Costa, L. N., et al. 1994, *ApJ*, 424, L1
 da Costa, L. N., et al. 1996, in preparation
 Dekel, A. 1994, *ARA&A*, 32, 371
 Dekel, A., Bertschinger, E., & Faber, S. M. 1990, *ApJ*, 364, 349
 Freudling, W., da Costa, L. N., & Pellegrini, P. S. 1994, *MNRAS*, 268, 943
 Freudling, W., da Costa, L. N., Wegner, G., Giovanelli, R., Haynes, M. P., & Salzer, J. J. 1995, *AJ*, 110, 920
 Freudling, W., Martel, H., & Haynes, M. P. 1991, *ApJ*, 377, 349
 Geller, M. G., & Huchra, J. P. 1989, *Science*, 246, 897
 Giovanelli, R., Haynes, M. P., Chamaroux, P., da Costa, L. N., Freudling, W., Salzer, J. J., & Wegner, G. 1995a, in *IAU Symp. 168, Examining the Big Bang and Diffuse Background Radiations*, ed. M. Kafatos (Dordrecht: Reidel), 183
 Giovanelli, R., Haynes, M. P., Salzer, J. J., Wegner, G., da Costa, L. N., & Freudling, W. 1994, *AJ*, 107, 2036
 ———. 1995b, *AJ*, 110, 1059
 Giovanelli, R., et al. 1996a, in preparation
 ———. 1996b, in preparation
 Lynden-Bell, D., Faber, S. M., Burstein, D., Davies, R. L., Dressler, A., Terlevick, R. J., & Wegner, G. 1988, *ApJ*, 326, 19
 Mathewson, D. S., Ford, V. L., & Buchhorn, M. 1992a, *ApJ*, 389, L5
 ———. 1992b, *ApJS*, 81, 413 (MFB)
 Piran, T., Lecar, M., Goldwirth, D., da Costa, L., & Blumenthal, G. 1993, *MNRAS*, 265, 681
 Strauss, M., & Willick, J. A. 1995, *Phys. Rep.*, 261, 271
 Wegner, G., Colless, M., Baggley, G., Davies, R. L., Bertschinger, E., Burstein, D., McMahan, Jr., R. K., & Saglia, R. P. 1996, preprint
 Willick, J. A. 1990, *ApJ*, 351, L5
 Willick, J., Courteau, S., Faber, S. M., Burstein, D., Dekel, A., & Kollat, T. 1996, *ApJ*, 457, 460
 Willmer, C. N. A., da Costa, L. N., Pellegrini, P. S., Fairall, A., & Latham, D. 1995, *AJ*, 109, 61
 Yahil, A. 1988, in *Large-Scale Structure of the Universe*, ed. V. C. Rubin & G. V. Coyne (Princeton: Princeton Univ. Press), 219




FIG. 1.—Density field along the supergalactic plane reconstructed from the SFI sample. We show density contours in intervals of $\delta = 0.2$, surface density maps with the height proportional to δ and density contrast maps. Bright yellow represents overdensities and dark blue underdense regions, respectively. See text or Fig. 2 for the identification of the structures.

DA COSTA et al. (see 468, L6)