

2006

# A Solidification Phenomenon in Random Packings

L. Bowen

*Indiana University - Bloomington*

R. Lyons

*Indiana University - Bloomington*

C. Radin

*University of Texas at Austin*

P. Winkler

*Dartmouth College*

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Mathematics Commons](#)

---

## Recommended Citation

Bowen, L.; Lyons, R.; Radin, C.; and Winkler, P., "A Solidification Phenomenon in Random Packings" (2006). *Open Dartmouth: Faculty Open Access Articles*. 2470.

<https://digitalcommons.dartmouth.edu/facoa/2470>

This Article is brought to you for free and open access by Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Faculty Open Access Articles by an authorized administrator of Dartmouth Digital Commons. For more information, please contact [dartmouthdigitalcommons@groups.dartmouth.edu](mailto:dartmouthdigitalcommons@groups.dartmouth.edu).

## A SOLIDIFICATION PHENOMENON IN RANDOM PACKINGS\*

L. BOWEN<sup>†</sup>, R. LYONS<sup>†</sup>, C. RADIN<sup>‡</sup>, AND P. WINKLER<sup>§</sup>

**Abstract.** We prove that uniformly random packings of copies of a certain simply connected figure in the plane exhibit global connectedness at all sufficiently high densities, but not at low densities.

**Key words.** sphere packing, solidification

**AMS subject classifications.** 52C25, 52C17

**DOI.** 10.1137/050647785

**1. Introduction.** The densest way to cover a large area with nonoverlapping unit disks is as in Figure 1, in which the disk centers form the vertices of a triangular lattice.




FIG. 1. *The densest packing of unit disks in the plane.*

A *packing* is a collection of congruent copies of a subset with pairwise disjoint interiors. See [5] for a proof that the above packing is indeed the densest possible for unit disks.

It is an old unsolved problem to understand whether densest packings of spheres, simplices, or other shapes, in a Euclidean or hyperbolic space of any dimension, exhibit crystallographic symmetry, such as that of Figure 1. This is the spirit, for instance, of Hilbert’s eighteenth problem; see [5, 10] for background.

Using physics models of two- and three-dimensional matter as a guide, we are tempted to try to gain insight into densest packings by considering packings at densities below the maximum. (For an example concerning spheres in  $\mathbb{R}^3$ , see [9].) In

\*Received by the editors December 16, 2005; accepted for publication (in revised form) July 19, 2006; published electronically November 21, 2006. This research was partially supported by the National Science Foundation under grants DMS-0406017 and DMS-0352999. The authors also thank the Banff International Research Station for support at a workshop where this research was begun.  
<http://www.siam.org/journals/sima/38-4/64778.html>

<sup>†</sup>Department of Mathematics, Indiana University, Bloomington, IN 47405 (lpbowen@indiana.edu, rdlyons@indiana.edu).

<sup>‡</sup>Department of Mathematics, University of Texas, Austin, TX 78712 (radin@math.utexas.edu).

<sup>§</sup>Department of Mathematics, Dartmouth College, Hanover, NH 03755 (peter.winkler@dartmouth.edu).

effect, we are emphasizing not so much that densest packings and low density packings differ by their *symmetry*, but rather that they differ in some fundamental geometric fashion. Indeed, it is commonly suggested in the physics literature (see, for instance, [1]) that two-dimensional models of matter do not exhibit crystallographic symmetry, and it is sometimes said by mathematicians that in high-dimensional Euclidean space, densest packings of spheres may not have crystallographic symmetry. So perhaps it is appropriate to re-examine the precise manner in which densest packings differ fundamentally from low density packings, and to use packings at less than optimum density as a guide.

In this work we replace round disks with deformed disks, which are copies of a “zipper” tile; see Figure 2. This tile can cover the plane completely, in which case the packing has density 1, and is completely connected in any sense. What we show is that even at somewhat lower densities, the uniform random packing still has rigid structure; in particular it has a form of connectedness associated with site percolation [7]. What this means for packing large but finite boxes (with torus boundary conditions) is that the necessary gross irregularities of most packings at such high densities disconnect the packings, if at all, along fault lines whose density tends to 0 as the size of the box tends to infinity. Although we define “uniform random packing” of the plane by limits of measures on packings of finite boxes, the key to our proof is to examine isometry-invariant probability measures on packings of the whole plane and to show that the ones that maximize “degrees of freedom per tile” are unique for high densities.



We show that at high density in our model there is a nonzero probability of an infinite linked component, and that this probability is zero at low density. Thus, there are different “phases” of the packings [2]. (This is closely related to continuum percolation, where one looks at overlapping disks with random independent centers, but our methods are quite different.)

Although we believe that such a result also holds for packings of disks or of spheres—pairs of which would be called “linked” if sufficiently close—we are able to prove the result only for our tiles, which are shaped to allow three well-defined levels of pairwise separation. (We discuss the appropriate notion of linking for collections of disks in the last section of the paper.) It is generally understood that true crystalline symmetry is not seen below optimal density in two dimensions—see [11]—so the form of connectedness we use may be useful in understanding the role of geometry in Hilbert’s problem.

**2. Description of the tile.** We consider packings by a deformed disk denoted by  $t$ , referred to as “the tile” and depicted in Figure 2. In this section, we define it precisely.


Let  $H$  be a regular hexagon of area 1. Let  $r$  be the radius of the in-circle of  $H$ . Let  $D$  be a disk concentric with the in-circle and of radius  $r + \rho$ , where  $0 < \rho \ll 1$  is a number we shall choose more precisely later. We shall construct the shape  $t$  by modifying  $H$  as follows;  $D$  will be called the *shadow disk* of  $t$ .

As shown in Figure 2, the tile  $t$  equals  $H$  with each side modified by a “fringe” and each corner modified by a hook and inlet, where a hook is about half an element of the fringe. As shown in Figure 3, the fringe height is  $2\rho$ . The elements of the fringe have two different size “necks,” one of size  $\rho^2$  and one of size  $2\rho^2$ , allowing neighboring tiles to be linked in either of two well-defined modes, “tightly linked” and “loosely linked”; the former is illustrated in Figure 4 and the latter in Figure 5. We say that two tiles  $t$  are *linked* (tightly or loosely) if, when one is held fixed, the other can be moved continuously only by a bounded amount (without overlapping the first). A

FIG. 2. *The zipper tile.*FIG. 3. *Close-up view of the fringe.*

*tight* link is one that permits no movement of one tile while fixing the other, while a link that is not tight is called *loose*. A key feature of our model is that when two tiles are tightly linked, any motion of one would necessitate a corresponding motion of the other. As we shall explain, the uniform probability distribution on packings of the plane at given density is a limit of such a distribution on packings of larger and larger tori. In our model, these distributions on packings of finite tori are concentrated on packings with the largest number of degrees of freedom, and therefore, roughly speaking, the fewest tiles bound by tight links. This gives us useful control on the packings in the support of our distributions.

A tile is called *fully linked on one side* if it is linked with another tile on that side in such a way that either they are tightly linked, and the line joining their centers goes through the midpoint of the sides of the corresponding hexagons, or they are not tightly linked but can be moved continuously so that their shadow disks touch each other. A tile is *fully linked* if it is fully linked on all sides. We note that the fully tightly linked packing (Figure 4) corresponds to a tiling by the original hexagon and has density 1, and that the tile has area 1 by construction.

FIG. 4. *Tightly linked tiles.*

**3. Statement of results.** To state our results we need some notation. Let  $X$  be the space of all packings of the plane by the tile  $t$ . Given a compact subset  $K$  of the plane and two packings of the plane, we consider the distance between the two packings with respect to  $K$  to be the Hausdorff distance between the unions of the tiles in the respective packings intersected with  $K$ . Then  $X$  is endowed with the topology of Hausdorff convergence on compact subsets;  $X$  is compact. Intuitively, two packings are close in  $X$  if they are close in the Hausdorff sense in a large ball centered at the origin. We shall define a probability measure on  $X$  that is “uniform” on the set of all packings of a fixed density. For this, we shall need the space  $X_n$  of all packings by the tile of the  $n \times n$  torus  $\mathbb{R}^2/(n\mathbb{Z})^2$ .

For any integer  $m$ , let  $X_{n,m} \subset X_n$  consist of those packings which contain exactly  $m$  tiles ( $X_{n,m}$  is empty if  $m$  is large enough). To each tile, we assign the set of six unit vectors based on its center and pointing toward the center of each of its edges. Through this assignment, we can view  $X_{n,m}$  as a subset of  $T_n^m/\Sigma_m$ , where  $T_n$  is the unit tangent bundle of the  $n \times n$  torus modulo a  $2\pi/6$  rotation and the symmetric group  $\Sigma_m$  acts by permuting the factors.

When  $m/n^2$  is small,  $X_{n,m}$  is  $(3m)$ -dimensional. However, when  $m/n^2$  is sufficiently large, the dimension of  $X_{n,m}$  inside  $T_n^m/\Sigma_m$  is less than  $3m$ . This is because




FIG. 5. Loosely linked tiles.

at least two tiles in any packing of  $X_{n,m}$  will have to be tightly linked, so that it is impossible to move one continuously without moving the other. Thus, it is useful to decompose  $X_{n,m}$  into a (finite) disjoint union of sets  $X_{n,m,k}$  of packings containing exactly  $k$  tight links. Generically, the dimension of  $X_{n,m,k}$  is  $3(m-k)$ . The dimension can be strictly less than this if the packings are jammed in the sense of [4], although this fact will not be important for us. The *top dimension* of  $X_{n,m}$  means the maximum dimension of all  $X_{n,m,k}$ . Let  $\mu_{n,m}$  be the probability measure on  $X_{n,m}$  obtained by normalizing the Hausdorff measure on  $X_{n,m}$  in the top dimension of  $X_{n,m}$  with respect to the natural metric inherited from  $T_n^m/\Sigma_m$ . We interpret  $\mu_{n,m}$  as being a uniform measure. The fact that  $\mu_{n,m}$  is supported on those packings with the fewest tight links will be crucial in the analysis to follow.

Let  $\tilde{X}_n$  be the space of all  $(n \times n)$ -periodic packings of the plane. In other words,  $\tilde{X}_n$  consists of those packings that are preserved under translations by  $n\mathbb{Z} \times n\mathbb{Z}$ . Under the quotient map, this space is naturally identified with  $X_n$ . Therefore, we can view the measures  $\mu_{n,m}$  as living on  $\tilde{X}_n \subset X$ .

For a fixed density  $d \in [0, 1]$ , let  $\mu^{(d)}$  be any measure obtained as the weak\* limit of measures of the form  $\mu_{n,m}$  such that  $n \rightarrow \infty$  and  $m/n^2 \rightarrow d$ . (Note that  $m/n^2$  is the density of every packing in the support of  $\mu_{n,m}$  and  $d$  is the average density

of a packing chosen with respect to  $\mu^{(d)}$ ; see Lemma 5.1.) A priori,  $\mu^{(d)}$  may not be unique, although we shall prove that it is for large enough  $d$ .

A *linked component* of a packing is a maximal subpacking in which for every two tiles  $t, t'$ , there is a sequence  $t = t_1, t_2, \dots, t_n = t'$  such that  $t_i$  is linked to  $t_{i+1}$  ( $i = 1, \dots, n - 1$ ). A *tightly linked component* is defined similarly, except that we require  $t_i$  to be tightly linked to  $t_{i+1}$ .

We say that a measure on the space  $X$  of packings is invariant if it is preserved under the full isometry group of the plane. All the measures we consider are probability measures unless stated otherwise.

Let  $\lambda_0$  be the unique invariant measure on tilings (packings that cover  $\mathbb{R}^2$ ) by our tile. Let  $\lambda_1$  be the unique invariant measure on packings by  $t$  such that all tiles are fully loosely linked, are as close as possible to each other, and the packing has hexagonal symmetry. Write  $\lambda_s := s\lambda_1 + (1 - s)\lambda_0$ .

Our main results are the following.

**THEOREM 3.1.** *There exists  $0 < d_1 < 1$  such that if  $d \geq d_1$ ,  $\mu^{(d)}$  is unique and equals  $\lambda_s$ , where  $s := (1 - d)/(1 - d_1)$ .*

**COROLLARY 3.2.** *The  $\mu^{(d)}$ -probability that the origin is inside a tile belonging to an infinite linked component is nonzero for  $d \geq d_1$ .*

**PROPOSITION 3.3.** *For some  $d_2 > 0$ , the probability (with respect to any  $\mu^{(d)}$  for any  $d < d_2$ ) that the origin is inside an infinite linked component is zero.*

#### 4. Tile properties.

**LEMMA 4.1.** *For small  $\rho$ , if tiles  $t_1$  and  $t_2$  are not tightly linked and do not overlap, then the distance between their centers is at least  $2r + 2\rho$ .*

*Proof.* Consider the line segment from the center of  $t_1$  to the center of  $t_2$ . If this segment traverses near a corner of  $t_1$  or  $t_2$ , then it must be longer than  $2r + 2\rho$  for small enough  $\rho$ . Suppose it crosses a fringe of  $t_1$  and of  $t_2$ . If the tiles are not linked, then the claim is obvious. If they are linked, then to minimize the distance, it must be that their fringes match up (so they are fully linked on one side). Thus, they can come closest to each other when pushed flat up against each other so that their shadow disks touch. In this case, the distance between the centers is exactly  $2r + \rho$ .  $\square$

We shall say that two tiles are *densely loosely linked* if they are loosely linked and their shadow disks touch. There is a unique invariant measure on maximally dense packings by congruent disks [3]. Hence the probability measure  $\lambda_1$  that we defined earlier is the unique invariant measure on packings by  $t$  such that all tiles are fully and densely loosely linked. Let  $d_1$  be the density of such a packing.

Given a tile  $t$  in a packing  $P$ , we denote by  $V(t)$  the Voronoi cell of the center of  $t$  with respect to the centers of the other tiles; that is,  $V(t)$  is the open set of points closer to the center of  $t$  than to the center of any other tile. We denote the area of a region  $A$  of the plane by  $|A|$ .

**LEMMA 4.2.** *The following holds for small enough  $\rho > 0$ . For any packing  $P$ , if  $t \in P$  is a tile that has no tight links, then the area of  $V(t)$  is at least  $1/d_1$ . Moreover, equality holds iff the configuration of tiles determining  $V(t)$  is congruent to a corresponding configuration of a packing in the support of  $\lambda_1$ .*

*Proof.* For a tile  $t$ , let  $H(t)$  denote the hexagon from which  $t$  is created. For  $x > 0$ , let  $H_x(t)$  denote the homothetic copy  $\frac{r+x}{r}H(t)$  about the center of  $H(t)$ .

Suppose  $t$  is a tile of  $P$  without any tight links. Consider the rays  $R_1, \dots, R_6$  from the center of the hexagon  $H(t)$  through each of its six vertices. These rays divide the plane into six sectors,  $S_1, \dots, S_6$ .

By construction, if  $t$  and  $t_1$  are loosely linked, then  $|H_\rho(t) \cap H_\rho(t_1)| = O(\rho^2)$ : The hexagon interiors do not intersect if they are parallel, while if they are not parallel, they can intersect only very slightly at a corner. The openings at which a corner can enter have area  $O(\rho^2)$  as  $\rho \rightarrow 0$ .

Thus, we have proved that whenever  $t$  is loosely linked in the sector  $S_i$ , then  $|V(t) \cap S_i| \geq |H_\rho(t)|/6 - \delta_1$ , with  $\delta_1 = O(\rho^2)$  as  $\rho \rightarrow 0$ .

Similarly, if  $t$  and  $t_2$  are not linked at all, then  $|H_{2\rho}(t) \cap H_{2\rho}(t_2)| = O(\rho^2)$ : Again, their interiors do not intersect if they are parallel, while if they are not parallel, they can intersect only very slightly at a corner. So there exists  $\delta_2 > 0$  such that whenever  $t$  is not linked in the sector  $S_i$ , we have  $|V(t) \cap S_i| \geq |H_{2\rho}(t)|/6 - \delta_2$ , with  $\delta_2 = O(\rho^2)$  as  $\rho \rightarrow 0$ .

Therefore, if  $t$  has no tight links but is not fully linked, then

$$|V(t)| \geq j \left( \frac{|H_\rho(t)|}{6} - \delta_1 \right) + (6 - j) \left( \frac{|H_{2\rho}(t)|}{6} - \delta_2 \right)$$

for some  $j$  with  $0 \leq j \leq 5$ . Given that  $\delta_1, \delta_2$  are of order  $\rho^2$  while  $|H_{2\rho}(t)| - |H_\rho(t)|$  is of order  $\rho$ , for  $\rho$  small enough we may conclude that  $|V(t)| > |H_\rho(t)|$  in this case.

On the other hand, the geometry of a tile is such that for small  $\rho$ , if  $t_1$  and  $t_2$  are two tiles loosely linked to  $t$ , then  $t_1$  cannot be tightly linked to  $t_2$ . Now suppose that  $t$  is fully loosely linked. Then the Voronoi cell of the center of  $t$  is determined by six tiles  $t_1, \dots, t_6$  all loosely linked to  $t$  and all with the property that their shadow disks  $D, D_1, \dots, D_6$  do not overlap (by the previous lemma). It follows [6] that  $|V(t)| \geq |H_\rho(t)|$ , with equality iff each of the disks  $D_1, \dots, D_6$  touches  $D$ . But there is only one way in which this can occur (up to isometry). So  $V(t) = H_\rho(t)$  in this case. This implies that the configuration  $t, t_1, \dots, t_6$  is congruent to a corresponding configuration of a packing in the support of  $\lambda_1$ .  $\square$

It is easy to see that given  $\rho > 0$ , there exists  $\varepsilon > 0$  such that for any finite component  $c$  of tightly linked tiles in any packing, the union  $V_c$  of the Voronoi cells of the centers of the tiles of  $c$  has area at least  $j_c + \varepsilon \text{Per}_c$ . Here  $j_c$  is the number of tiles in  $c$  and  $\text{Per}_c$  is the perimeter of the union of hexagons corresponding to  $c$ . Let  $\varepsilon$  be the largest such constant. Let  $\delta > 0$  be such that the area of the Voronoi cell in the fully densely loosely linked packing equals  $1 + \varepsilon \text{Per}_1 + \delta$ , where  $\text{Per}_1$  is the perimeter of the hexagon of a single tile. Since  $\varepsilon = \rho + O(\rho^2)$  and  $\delta = O(\rho^2)$ , we have the following.

LEMMA 4.3. *For sufficiently small  $\rho$ , there are  $\varepsilon, \delta > 0$  such that for any finite tightly linked component  $c$ ,*

$$d_1 = \frac{1}{1 + \text{Per}_1 \varepsilon + \delta},$$

$$|V_c| \geq j_c + \varepsilon \text{Per}_c,$$

and

$$\delta \leq \varepsilon/100.$$

**5. High density.** Recall that  $X$  is the compact space of all packings of the plane by the tile (with the topology of Hausdorff-metric convergence on compact subsets). Let  $\widetilde{M}$  be the space of isometry-invariant Borel probability measures on  $X$ . For any  $\mu \in \widetilde{M}$ , we denote by  $|\mu| := \mu(A_0)$  the *density* of  $\mu$ , where  $A_0$  is the set of all packings  $P \in X$ , one of whose tiles contains the origin. Since a tile is the closure of its interior,  $A_0$  is a closed set.



LEMMA 5.1. *If  $\mu_i \in \widetilde{M}$  converges to  $\mu$  in the weak\* topology, then  $|\mu_i|$  converges to  $|\mu|$ .*

*Proof.* Let  $\widehat{P}$  denote the union of tiles in a packing,  $P$ . For any invariant probability measure  $\nu$  and any  $z \in \mathbb{R}^2$ , we have

$$|\nu| = \int \mathbf{1}_{\{0 \in \widehat{P}\}} d\nu(P) = \int \mathbf{1}_{\{z \in \widehat{P}\}} d\nu(P).$$

Integrating over  $z$  in a unit-area disk,  $D$ , with respect to the Lebesgue measure and using Fubini's theorem gives the identity  $|\nu| = \int |\widehat{P} \cap D| d\nu(P)$ . Since the function  $P \mapsto |\widehat{P} \cap D|$  is continuous on  $X$ , the lemma follows.  $\square$

Recall that  $\lambda_0$  is the unique invariant measure on tilings by our tile, so that  $|\lambda_0| = 1$ . Recalling that  $d_1$  is the density of a fully densely loosely linked tiling, fix a density  $d$  with  $d_1 \leq d \leq 1$ . Let  $\mu_N$  be the uniform measure on configuration of tiles at density  $d_N$  in an  $N \times N$  torus, where  $d_N \rightarrow d$  as  $N \rightarrow \infty$ . To prove Theorem 3.1, we shall show that the weak\* limit of  $\mu_N$  exists and equals  $\lambda_s$ , where  $s := (1-d)/(1-d_1)$ .

We shall use several lemmas that depend on the following notation. Given a packing  $P \in X$ , let

- $t_P$  be the tile of  $P$  such that the origin belongs to  $V(t_P)$  (this exists as long as the origin is not on the boundary of a Voronoi cell),
- $K_P$  be the tightly linked component containing  $t_P$ ,
- $j_P$  be the number of tiles in  $K_P$ , and
- $f(P) := 3/j_P$  if  $j_P$  is finite and  $t_P$  contains the origin, and 0 otherwise.

Thus  $f(P)$ , in a sense, measures the number of degrees of freedom per tile near the origin.

LEMMA 5.2. *If  $\nu$  is any measure in  $\widetilde{M}$ , then  $\int f d\nu(f) \leq 3|\nu|$ , with equality iff  $t_P$  has no tight links for  $\nu$ -almost every packing  $P$ .*

The proof is immediate.

LEMMA 5.3. *If a sequence  $\langle \nu_n \rangle \subset \widetilde{M}$  converges to  $\nu$  in the weak\* topology, then  $\int f d\nu_n$  converges to  $\int f d\nu$ .*

Lemma 5.3 is proved in a manner similar to Lemma 5.1.

Given a finite tightly linked component  $c$ , let the congruence class of  $c$  be  $C$  and let  $X_C \subset X$  be the space of all packings  $P$  for which  $t_P$  exists and  $K_P$  is in  $C$ . Let  $X''$  be the space of all packings  $P$  with density 1, where “density” refers to the usual concept of the limit of the proportion of the area of  $P$  inside a large disk centered at the origin as the radius tends to infinity. Let  $X' \subset X$  be the space of all packings  $P$  such that  $K_P$  is infinite and either the density of  $P$  is less than 1, the density is not defined, or  $t_P$  does not exist. Thus,  $X$  is the disjoint union of  $X'$ ,  $X''$  and the collection of  $X_C$  for all  $C$ .

Let  $\nu$  be any invariant probability measure with density  $d$ . Let  $\nu_C$  be  $\nu$  conditioned on  $X_C$ ,  $\nu'$  be  $\nu$  conditioned on  $X'$ , and  $\nu''$  be  $\nu$  conditioned on  $X''$ . Since  $\lambda_0$  is the only invariant probability measure with support in  $X''$ , we have  $\nu'' = \lambda_0$ . Thus,

$$\nu = \nu(X')\nu' + \nu(X'')\lambda_0 + \sum_C \nu(X_C)\nu_C.$$

Define the density  $|\omega| := \omega(A_0)$  as before, but for any (invariant or noninvariant) probability measure  $\omega$  on  $X$ . We have

$$d = |\nu| = \nu(X')|\nu'| + \nu(X'') + \sum_C \nu(X_C)|\nu_C|$$

and

$$\int f \, d\nu = \sum_C \nu(X_C) \int f \, d\nu_C = \sum_C \nu(X_C) \frac{3}{j_C} |\nu_C|,$$

where  $j_C$  is the number of tiles in  $C$ .

LEMMA 5.4. *Let  $\nu \in \widetilde{M}$  and  $C$  be a finite-component class. Suppose that  $0 \leq s \leq 1$  is such that  $|\lambda_s| = |\nu_C|$ . Then  $\int f \, d\lambda_s \geq \int f \, d\nu_C$ . Moreover, equality holds only if  $j_C = 1$ .*

*Proof.* As in the proof of Lemma 5.1, we have that  $|\nu_C| = \int j_C / |V(t_P)| \, d\nu_C(P)$ .

First suppose that  $j_C = 1$ . Then  $|\nu_C| = \int 1 / |V(t_P)| \, d\nu_C(P) \leq d_1$  by Lemma 4.2. This means that  $s = 1$  and  $\int f \, d\nu_C = 3 |\nu_C| = 3 |\lambda_s| = \int f \, d\lambda_s$ .

Now assume that  $j = j_C > 1$  and put  $p := \text{Per}_C$ . By definition,

$$\int f \, d\lambda_s = s \int f \, d\lambda_1 + (1 - s) \int f \, d\lambda_0 = s \int f \, d\lambda_1 = 3s d_1.$$

Since  $\nu_C(f) = 3|\nu_C|/j_C = 3|\lambda_s|/j_C = 3(sd_1 + 1 - s)/j$ , it suffices to show that

$$s d_1 > \frac{s d_1 + (1 - s)}{j},$$

which is equivalent to

$$s(j d_1 - d_1 + 1) > 1.$$

Now  $s d_1 + (1 - s) = |\nu_C| \leq \frac{j}{j + \varepsilon p}$ , where  $\varepsilon$  is from Lemma 4.3. Solving for  $s$  gives

$$s \geq \frac{1 - \frac{j}{j + \varepsilon p}}{1 - d_1},$$

whence it is enough to show that

$$(j d_1 - d_1 + 1) \frac{1 - \frac{j}{j + \varepsilon p}}{1 - d_1} > 1.$$

This boils down to

$$d_1(p\varepsilon + 1) > 1.$$

Now,  $j > 1$  implies that  $p \geq (7/6)\text{Per}_1$ , where  $\text{Per}_1$  is the perimeter of a single tile. Since  $\varepsilon/100 > \delta$  (by Lemma 4.3), this implies that  $p\varepsilon + 1 > 1 + \varepsilon\text{Per}_1 + \delta = 1/d_1$ , proving the last inequality.  $\square$

LEMMA 5.5. *We have  $\int f \, d\nu \leq \int f \, d\lambda_s$  for all  $\nu \in \widetilde{M}$  with  $|\nu| = |\lambda_s|$ . Equality holds only if*

- $\nu(X_C) = 0$  for every component class  $C$  with  $j_C > 1$ , and
- whenever  $\nu(X_C) > 0$  and  $j_C = 1$ , we have  $|\nu_C| = d_1$ .

*Proof.* Recall that

$$\int f \, d\nu = \sum_C \nu(X_C) \int f \, d\nu_C.$$

For each component class  $C$ , let  $s_C$  be defined as follows:

- If there exists  $s \in [0, 1]$  such that  $|\nu_C| = sd_1 + (1 - s)$ , then set  $s_C := s$ ;
- otherwise, set  $s_C := 1$ .

Let  $\omega_C := s_C\lambda_1 + (1 - s_C)\lambda_0$  and

$$\sigma := (\nu(X') + \nu(X''))\lambda_0 + \sum_C \nu(X_C)\omega_C.$$

From the previous lemma, if  $|\nu_C| \geq d_1$ , then  $\int f d\nu_C \leq \int f d\omega_C$ , with equality only if  $j_C = 1$ . If  $|\nu_C| < d_1$ , then  $s_C = 1$  and

$$\int f d\nu_C = \frac{3|\nu_C|}{j_C} < 3d_1 = \int f d\omega_C.$$

Summing up, we obtain

$$\begin{aligned} \int f d\sigma &= \sum_C \nu(X_C) \int f d\omega_C \\ &\geq \sum_C \nu(X_C) \int f d\nu_C = \int f d\nu. \end{aligned}$$

Moreover, equality holds only if  $\nu(X_C) = 0$  for every component  $C$  with  $j_C > 1$  and  $|\nu_C| = d_1$  whenever  $j_C = 1$ . Since  $|\omega_C| \geq |\nu_C|$ , we have

$$\begin{aligned} |\sigma| &= \nu(X') + \nu(X'') + \sum_C \nu(X_C) |\omega_C| \\ &\geq \nu(X') |\nu'| + \nu(X'') + \sum_C \nu(X_C) |\nu_C| \\ &= |\nu| \\ &= |\lambda_s|. \end{aligned}$$

Since  $\sigma$  and  $\lambda_s$  are both convex combinations of  $\lambda_0$  and  $\lambda_1$ , this implies that  $\int f d\sigma \leq \int f d\lambda_s$  with equality iff  $\sigma = \lambda_s$ . Thus,  $\int f d\nu \leq \int f d\lambda_s$ . In the equality case we must have  $\int f d\nu = \int f d\sigma = \int f d\lambda_s$  and  $\sigma = \lambda_s$ . This implies that  $\nu(X_C) = 0$  if  $j_C > 1$  and  $|\nu_C| = d_1$  if  $j_C = 1$ .  $\square$

LEMMA 5.6. *Let  $\nu \in \widetilde{M}$ . If  $|\nu| = |\lambda_s|$ , then  $\int f d\nu \leq \int f d\lambda_s$ . Equality holds iff  $\nu = \lambda_s$ .*

(Informally,  $\lambda_s$  uniquely maximizes the number of degrees of freedom per tile for invariant measures of a fixed density.)

*Proof.* The previous lemma implies  $\int f d\nu \leq \int f d\lambda_s$ . Assume  $\int f d\nu = \int f d\lambda_s$ ; then

$$\nu = \nu(X')\nu' + \nu(X'')\lambda_0 + \nu(X_C)\nu_C,$$

where  $C$  is the component of size 1 and  $|\nu_C| = d_1$ . This gives  $\int f d\nu = \nu(X_C)3d_1 = \int f d\lambda_s = 3sd_1$ . Hence  $\nu(X_C) = s$ . Since  $\nu'$  has density strictly less than  $1 = |\lambda_0|$  but  $|\nu| = |\lambda_s|$ , we must have  $\nu(X') = 0$ . That is,

$$\nu = \nu(X'')\lambda_0 + \nu(X_C)\nu_C.$$

Since  $\nu$  and  $\lambda_0$  are isometry invariant,  $\nu_C$  must also be isometry invariant. By Lemma 4.2,  $\lambda_1$  is the unique isometry-invariant measure with support in  $X_C$  and with density  $d_1$ . Hence  $\nu_C = \lambda_1$ . This implies  $\nu = \lambda_s$ , and the proof is finished.  $\square$

*Proof of Theorem 3.1.* It is easy to see that one can pack the  $N \times N$  torus in such a way that there is a large region of tightly linked tiles and a large region of densely loosely linked tiles, and in such a way that the interface between the two regions has a density which approaches zero as  $N$  tends to infinity, and the density  $d_N$  of the packing  $P_N$  tends to  $d$ . Let  $\omega_N$  be the invariant measure supported on isometric copies of  $P_N$  (a pull-back of  $P_N$  to the plane). Then  $\omega_N$  tends to  $\lambda_s$  in the weak\* topology. By Lemma 5.3, this implies that  $\int f d\omega_N \rightarrow \int f d\lambda_s$ .

Now  $\mu_N$ , the uniform measure of density  $d_N$  on the  $N \times N$  torus, satisfies  $\int f d\mu_N \geq \int f d\omega_N$ . This is because  $\mu_N$  is by definition supported on packings with the maximal number of degrees of freedom for the given density  $d_N$ . Hence  $\liminf_N \int f d\mu_N \geq \liminf_N \int f d\omega_N = \int f d\lambda_s$ .

Therefore, if  $\mu_\infty$  is any weak\* subsequential limit of  $\langle \mu_N \rangle_N$ , then  $\int f d\mu_\infty \geq \int f d\lambda_s$ . But  $d_N \rightarrow d$ , so  $|\mu_\infty| = |\lambda_s|$  by Lemma 5.1. The previous lemma now implies that  $\mu_\infty = \lambda_s$ .  $\square$

Returning to the discussion of the introduction, we note that from simulations of hard disks, one would expect the corollary to hold even for a range of densities below  $d_1$ , but we do not know how to prove this.

**Remark on higher dimensions.** The basic features of our argument can be generalized to dimension 3 or higher, except for our use in Lemma 4.2 of [6] on the minimal Voronoi region in disk packings in the plane. It would be of interest if this part of our proof could be replaced with an argument insensitive to dimension, since the Voronoi regions of, say, the spheres in a face-centered cubic lattice do not minimize volume per site [8].

**6. Low density.** In this final section, we confirm the intuition that at low densities there will be no infinite loosely linked component. It is obvious that there is no infinite *tightly* linked component at densities smaller than  $d_1$ .

We begin with a lemma that holds for any tile shape (in fact, for any *collection* of shapes and sizes, as long as each can be fit into a disk of some fixed radius  $s$ , and “density” is interpreted as the number of tiles per unit area).

LEMMA 6.1. *For small enough density  $d$ , if a packing  $P$  is drawn from  $\mu^{(d)}$ , then the probability that the disk  $B_R$  of radius  $R$  about the origin contains more than  $9R^2d$  tile centers goes to zero as  $R \rightarrow \infty$ .*

*Proof.* Let  $s$  be the radius of the smallest disk containing the tile (in our case,  $s$  is about  $2^{1/2} \cdot 3^{-3/4} \cdot (1+2\rho)$ ) and choose

$$0 < d < \frac{.05}{13\pi s^2} ;$$

for our zipper tiles with small enough  $\rho$ ,  $d \leq .003$  suffices. Let  $T$  be the set of tiles whose centers fall in  $B_R$ ,  $k := \lceil \pi R^2 d \rceil$ , and  $\ell > 9R^2d$ . Letting  $\mu^{(d)}(\cdot)$  denote the probability of an event with respect to the measure  $\mu^{(d)}$ , we shall show that

$$\frac{\mu^{(d)}(|T| = \ell)}{\mu^{(d)}(|T| = k)} \leq \gamma^{\ell-k}$$

for some constant  $\gamma < 1$ . It then follows that

$$\mu^{(d)}(|T| > 9R^2d) \leq \mu^{(d)}(|T| = k) \sum_{\ell = \lceil 9R^2d \rceil}^{\infty} \gamma^{\ell-k} \leq \mu^{(d)}(|T| = k) \frac{\gamma^{\lceil (9-\pi)R^2d \rceil}}{1-\gamma} \rightarrow 0$$

as  $R \rightarrow \infty$ , as desired.

The measure  $\mu^{(d)}$  is the limit of uniform distributions of configurations on the  $N \times N$  torus  $\mathbf{T}_N$ , in turn obtainable by choosing a sequence of  $n = \lfloor N^2 d \rfloor$  points from the Lebesgue distribution  $\lambda$  on  $\mathbf{T}_N^n$  as the centers of the tiles, orienting each tile independently and uniformly at random, and finally conditioning on no overlap. We denote by  $\lambda(|T|=j)$  the a priori probability that exactly  $j$  points fall inside  $B_R$  (which we take to be some fixed disk in the torus).

Let  $\Phi$  be the event that there is no overlap among the tiles whose centers lie in  $B_R$ , and  $\Psi$  be the event that there is no overlap involving any tile whose center falls outside  $B_R$ . Then

$$\frac{\mu^{(d)}(|T|=\ell)}{\mu^{(d)}(|T|=k)} = \frac{\lambda(|T|=\ell)}{\lambda(|T|=k)} \cdot \frac{\lambda(\Phi \mid |T|=\ell)}{\lambda(\Phi \mid |T|=k)} \cdot \frac{\lambda(\Psi \mid |T|=\ell \wedge \Phi)}{\lambda(\Psi \mid |T|=k \wedge \Phi)},$$

and our job is to bound the three fractions on the right.

For the first, we note that  $|T|$  is binomially distributed in the measure  $\lambda$ ; hence

$$\begin{aligned} \frac{\lambda(|T|=\ell)}{\lambda(|T|=k)} &= \frac{\binom{n}{\ell} \left(\frac{\pi R^2}{N^2}\right)^\ell \left(1 - \frac{\pi R^2}{N^2}\right)^{n-\ell}}{\binom{n}{k} \left(\frac{\pi R^2}{N^2}\right)^k \left(1 - \frac{\pi R^2}{N^2}\right)^{n-k}} \leq \frac{(n-k)!/(n-\ell)!}{\ell!/k!} \left(\frac{k}{1 - \frac{k}{n}}\right)^{\ell-k} \\ &< \frac{(n-k)^{\ell-k}}{(\ell/e)^{\ell-k}} \cdot \frac{k^{\ell-k}}{(n-k)^{\ell-k}} = \left(\frac{ek}{\ell}\right)^{\ell-k} \leq 0.95^{\ell-k} \end{aligned}$$

for large  $R$ .

The next fraction is easy: Since we may throw the first  $k$  centers into  $B_R$ , then for the remaining  $\ell - k$ , we have that

$$\frac{\lambda(\Phi \mid |T|=\ell)}{\lambda(\Phi \mid |T|=k)}$$

is the probability that the additional  $\ell - k$  centers do not cause a collision, which is at most 1.

For the (inverse of) the third fraction, we throw  $n - \ell$  centers into the region outside  $B_R$ , then throw the remaining  $\ell - k$ . A new point, if it lands at a distance greater than  $2s$  from any previous point or from the disk  $B_R$ , causes no new overlap, and at each stage there are fewer than  $n - k$  points already placed. Hence

$$\begin{aligned} \frac{\lambda(\Psi \mid |T|=k \wedge \Phi)}{\lambda(\Psi \mid |T|=\ell \wedge \Phi)} &> \left(\frac{N^2 - \pi(R + 2s)^2 - (n-k)4\pi s^2}{N^2 - \pi R^2}\right)^{\ell-k} \\ &\geq \left(1 - 4\pi s^2 d - \frac{4\pi s R + 4\pi s^2}{N^2 - \pi R^2}\right)^{\ell-k} > (1 - 13s^2 d)^{\ell-k} \end{aligned}$$

for  $N \gg R$ .

Putting the inequalities together, we have

$$\frac{\mu^{(d)}(|T|=\ell)}{\mu^{(d)}(|T|=k)} \leq \left(\frac{.95}{1 - 13s^2 d}\right)^{\ell-k} = \gamma^{\ell-k},$$

where  $\gamma := .95/(1 - 13s^2 d) < 1$  by choice of  $d$ .  $\square$

PROPOSITION 6.2. For some  $d_2 > 0$ , the  $\mu^{(d)}$ -probability that the origin is inside an infinite connected component of loosely linked tiles is zero for  $d < d_2$ .

*Proof.* Let  $d \in (0, .003)$  be a density to be chosen later. Let  $P$  be a packing drawn from  $\mu^{(d)}$ ; we aim to show that the probability that the origin is connected by a loosely linked chain of tiles of  $P$  to some point at distance  $R$  approaches zero as  $R \rightarrow \infty$ .

We again choose some large radius  $R$  and let  $T$  be the set of tiles of  $P$  whose centers fall inside the disk  $B_R$ .




FIG. 6. An unlikely configuration of tiles in and around  $B_R$ .

Fix the positions of the tiles of  $P \setminus T$  (the black tiles of Figure 6) and consider the space of packings having these tiles plus  $n$  tiles whose centers fall in  $B_R$ . We think of this space as being a subset of  $T_1(B_R)^n / \Sigma_n$ , where  $T_1(B_R)$  is the unit tangent bundle of  $B_R$  (modulo a  $2\pi/6$  rotation to take into account the symmetries of the tile) and the symmetric group acts by permuting the factors.

If  $\alpha_n$  is the volume (in  $T_1(B_R)^n / \Sigma_n$ -space) of this space and  $m < n$ , then by packing  $n - m$  tiles into  $B_R$  and then the remaining  $m$  in the leftover space, we have

$$\alpha_n \geq \frac{1}{\binom{n}{n-m}} \alpha_{n-m} \frac{1}{m!} [\pi(R - 2s)^2 - n\pi(2s)^2]^m,$$

where  $s$  is, as before, the radius of the circle circumscribing a tile. This takes into account possible intrusion of tiles in  $P - T$  into  $B_R$ , and the fact that a tile center at point  $x$  can exclude nearby centers, but only within distance  $2s$  of  $x$ .

Let  $\beta$  denote the “wiggle room” of a tile  $t$  loosely linked to a stationary tile  $t'$ , that is, the three-dimensional volume of the space of positions of  $t$ ; then  $\beta = O(\rho^3)$  (but we use only that  $\beta$  is bounded by a constant). If a packing “percolates,” that is, contains a chain of loosely linked tiles connecting the center to the boundary of

$B_R$ , let  $t_1, \dots, t_m$  be a shortest such chain (the dark grey tiles of Figure 6). Note that  $m \geq R/(2s)$ . For each  $i > 2$ , the tile  $t_i$  is linked to one of the three sides of  $t_{i-1}$  farthest from the side of  $t_{i-1}$  linked to  $t_{i-2}$ , and has wiggle room at most  $\beta$  with respect to  $t_{i-1}$ . Accounting for the orientation of  $t_1$  and allowing the remaining  $n-m$  tile centers to fall anywhere in  $B_R$ , we have that the  $3n$ -dimensional volume of the set of percolating packings is bounded by  $(\pi/3) \cdot 6 \cdot 3^{m-2} \cdot \beta^{m-1} \cdot \alpha_{n-m} < 3^m \beta^{m-1} \alpha_{n-m}$ .

Comparing with the lower bound for  $\alpha_n$ , we find that given  $|T| = n \leq 9dR^2$ , the probability of percolation is less than

$$\frac{3^m \beta^{m-1} \alpha_{n-m}}{\alpha_n} \leq \frac{3^m \beta^{m-1} n!/(n-m)!}{[\pi(R-2s)^2 - n\pi(2s)^2]^m} < \left( \frac{27\beta dR^2}{\pi[(R-2s)^2 - 36dR^2s^2]} \right)^m / \beta,$$

which goes to zero as  $R$  (thus also  $m$ ) increases, for suitably chosen  $d$ . Since we know from Lemma 6.1 that  $\mu^{(d)}(|T| \leq 9dR^2)$  approaches 1 as  $R \rightarrow \infty$ , the proposition follows.  $\square$

A more careful argument would prove Proposition 6.2 for any density below  $1/(4\pi(2/3\sqrt{3})) = .2067^+$  for sufficiently small  $\rho$ , but clearly the probability of percolation will remain 0 for much higher densities than that.

**7. A conjecture.** We have shown that high-density random packings of zipper tiles in the plane contain an infinite loosely linked component with positive probability, while low-density random packings do not. What happens in the case of ordinary disks, where there is no apparent linking mechanism? We believe, but cannot prove, the following.

**CONJECTURE.** *Suppose  $\mu^{(d)}$  is defined as above, but for geometric disks of radius 1. Join two centers by an edge if their distance is at most  $2 + \varepsilon$  for some fixed  $\varepsilon \ll 1$ . Then for sufficiently high-density  $d$  below the maximum, the graph resulting from a configuration drawn from  $\mu^{(d)}$  will contain an infinite connected component a.s.*

This connectedness property can in fact be proven by a standard Peierls-type argument for large  $\varepsilon$ . This may be known already, though we do not know a reference; it is a straightforward extension of the traditional percolation proof to a situation with a new length scale given by the size of the disks. In general, there is some parameter set of  $(\varepsilon, d) \subset (0, \infty) \times (0, 1)$  for which there is an infinite component. For small  $d$  or for large  $\varepsilon$ , the problem is quite similar to continuum percolation, where one connects by an edge two points of a Poisson point process if their distance is at most  $r$ . Because of homotheties, one may fix the intensity of the point process to be 1. Then there is a phase transition in  $r$ . Our situation is quite different in that we really have two parameters, due to the size of the disks, but our conjecture is that there is a phase transition in  $d$  for every  $\varepsilon$  nevertheless.

There is a fundamental difference between the connectedness property for small  $\varepsilon$  and for large  $\varepsilon$ . An infinite set of disks connected or linked in the sense of small  $\varepsilon$  would resist shearing in a sense not true for a set linked only in the sense of large  $\varepsilon$ . We prove the connectedness property for the zipper model for small  $\varepsilon$ , using in an essential way special features of the nonconvex zipper tiles.

**Acknowledgment.** We gratefully acknowledge learning of [11] from an anonymous referee.

## REFERENCES

- [1] B. J. ALDER AND W. G. HOOVER, *Numerical Statistical Mechanics*, in *Physics of Simple Liquids*, H. N. V. Temperley, J. S. Rowlinson, and G. S. Rushbrooke, eds., John Wiley, New York, 1968, pp. 79–113.
- [2] L. BOWEN, R. LYONS, C. RADIN, AND P. WINKLER, *Fluid/solid transition in a hard-core system*, *Phys. Rev. Lett.*, 96 (2006), paper 025701.
- [3] L. BOWEN, C. HOLTON, C. RADIN, AND L. SADUN, *Uniqueness and symmetry in problems of optimally dense packings*, *Math. Phys. Electron. J.*, 11 (2005), paper 1.
- [4] A. DONEV, S. TORQUATO, F. H. STILLINGER, AND R. CONNELLY, *Jamming in hard sphere and disk packings*, *J. Appl. Phys.*, 95 (2004), pp. 989–999.
- [5] L. FEJES TÓTH, *Regular Figures*, Macmillan, New York, 1964.
- [6] L. FEJES TÓTH, *Lagerungen in der Ebene auf der Kugel und im Raum*, 2nd ed., Springer-Verlag, Berlin, 1972, pp. 62–63.
- [7] G. GRIMMETT, *Percolation*, 2nd ed., Springer-Verlag, Berlin, 1999.
- [8] T. C. HALES AND S. MCLAUGHLIN, *A Proof of the Dodecahedral Conjecture*, <http://www.arxiv.org/abs/math?papernum=9811079> (2002).
- [9] H. KOCH, L. SADUN, AND C. RADIN, *Most stable structure for hard spheres*, *Phys. Rev. E*, 72 (2005), paper 016708.
- [10] C. RADIN, *Orbits of orbs: Sphere packing meets Penrose tilings*, *Amer. Math. Monthly*, 111 (2004), pp. 137–149.
- [11] T. RICHTHAMMER, *Translation-Invariance of Two-Dimensional Gibbsian Systems of Particles with Internal Degrees of Freedom*, preprint, 2006. Available online from <http://www.arxiv.org/abs/math.PR/0603140>.