

Intensive home hemodialysis: the best treatment in the best system

Citation for published version (APA):

Cornelis, T., Kooman, J. P., & van der Sande, F. M. (2011). Intensive home hemodialysis: the best treatment in the best system. *Nephrology Dialysis Transplantation*, 26(9), 3067-3068.
<https://doi.org/10.1093/ndt/gfr348>

Document status and date:

Published: 01/09/2011

DOI:

[10.1093/ndt/gfr348](https://doi.org/10.1093/ndt/gfr348)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

only in specific circumstances such as pregnancy, after non-renal solid organ transplantation or in elderly may intensive home HD be strongly indicated [2, 3], but there should be an ‘intensive HD first’ strategy at the time of education of the predialysis patient [4]. Home HD promotes autonomy and independence and may therefore further increase quality of life (QoL) in addition to the multiple benefits of optimized uraemic toxin clearance and volume control secondary to intensive HD as compared to other dialysis strategies [2].

Piccoli mentions two potential clinical concerns for intensive HD. The first refers to the increased risk of vascular access problems in the context of more frequent cannulation in intensive HD. The recent FHN trial did indeed reveal an increased need for vascular access procedures in the short daily hemodialysis (SDHD) group as compared to the conventional HD group [5]. However, the reason for this observation remains unclear as it may well have been that it was purely due to more frequent visits to the HD unit with more frequent vascular access control. The second potential clinical problem mentioned by Piccoli is the depletion of small molecules and trace elements and an increased risk of inflammation. However, excellent phosphate control, improvement of malnutrition and suppression of inflammation are well-studied advantages of intensive HD and we therefore do not consider the abovementioned as a potential clinical problem for intensive HD [2].

We support the idea, suggested by Piccoli, of interchangeability between different dialysis regimes as long as the patient follows one of the intensive HD regimes, whether it is SDHD, alternate night nocturnal HD or nightly nocturnal HD. Indeed, previous studies have shown that QoL is higher if patients are provided with the choice of the dialysis modality and it is therefore possible that providing home HD patients with the autonomy of alternating between different intensive HD modalities may further increase QoL.

Finally, it is indeed time now for the governments and for health care insurance companies to obtain the insight that end-stage renal disease (ESRD) patients should be treated with the best treatment in the best system by changing the reimbursement system. This may enable these patients to have the highest QoL and physical activity and thereby also the best social functioning and economic productivity. This may further reduce the financial burden of the treatment of ESRD patients.

Conflict of interest statement. None declared.

Intensive home hemodialysis: the best treatment in the best system

Sir,

With great interest, we read the recently published editorial comment by Dr Piccoli entitled ‘The never-ending search for the perfect dialysis. Should we move from the best treatment to the best system?’ [1].

We are convinced that intensive haemodialysis (HD) may be the best dialysis treatment in the best system. Not

Department of Internal
Medicine, Division of
Nephrology, Maastricht
University Medical Center,
Maastricht, Netherlands
E-mail: tom.cornelis@mumc.nl

Tom Cornelis
Jeroen P. Kooman
Frank M. van der Sande

1. Piccoli GB. The never-ending search for the perfect dialysis. Should we move from the best treatment to the best system? *Nephrol Dial Transplant* 2011; 26: 1128–1131

2. Perl J, Chan CT. Home hemodialysis, daily hemodialysis, and nocturnal hemodialysis: core curriculum 2009. *Am J Kidney Dis* 2009; 54: 1171–1184
3. Cornelis T, Rioux JP, Bargman JM *et al*. Home dialysis is a successful strategy in nonrenal solid organ transplant recipients with end-stage renal disease. *Nephrol Dial Transplant* 2010; 25: 3425–3429
4. Cornelis T, Kooistra MP, Kooman J *et al*. Education of end-stage renal disease patients on dialysis modality selection: “intensive hemodialysis first”. *Nephrol Dial Transplant* 2010; 25: 3129–3130
5. The FHN Trial Group. In-center hemodialysis six times per week versus three times per week. *N Engl J Med* 2010; 363: 2287–2300

doi: 10.1093/ndt/gfr348