



ISSN: 2348-1900

Plant Science Today

<http://www.plantsciencetoday.online>

Review Article

Positive and negative effects of nanoparticles on plants and their applications in agriculture

Pooja Goswami, Sonali Yadav & Jyoti Mathur*

Department of Bioscience and Biotechnology, Banasthali University, P.O. Banasthali Vidyapith, Rajasthan 304 022, India

Article history

Received: 07 February 2019

Accepted: 06 April 2019

Published: 30 May 2019

Abstract

Nanotechnology is the promising field with its wide applications in biotechnology, pharmaceutical science, drug targeting, nano-medicine and other research areas. This review highlights the positive and negative impact of nanoparticles on plants and its wide applications in agricultural sciences. Effect of NPs in terms of seed germination, growth promotion and enhancement of metabolic rate has been evaluated by several scientific researches. However, NPs also exert their negative effects such as suppression of plant growth, inhibition of chlorophyll synthesis, photosynthetic efficiency etc. Effects of NPs can be either positive or negative it depending upon the plant species and type of nanoparticles used & its concentration. Modern nano-biotechnological tools have a great potential to increase food quality, global food production, plant protection, detection of plant and animal diseases, monitoring of plant growth nano-fertilizers, nano-pesticide, nano-herbicides and nano-fungicides.

Publisher

Horizon e-Publishing Group

Keywords: Nanoparticles; Agro-waste; Nano-application; plant protection; biogenic nanoparticles.

Citation: Goswami P, Yadav S, Mathur J. Positive and negative effects of nanoparticles on plants and their applications in agriculture. *Plant Science Today* 2019;6(2):232-242. <https://doi.org/10.14719/pst.2019.6.2.502>

Copyright: © Goswami et al (2019). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>).

*Correspondence

Jyoti Mathur

✉ contact.srivastava@gmail.com

Indexing: Plant Science Today is covered by Scopus, Web of Science, BIOSIS Previews, ESCI, CAS, AGRIS, CABI, Google Scholar, etc. Full list at <http://www.plantsciencetoday.online>

Introduction

Nanoscience is a developing concept in the agricultural area. In recent times, rapid progress in nanotechnology has made it possible to synthesize engineered nanoparticles (ENPs) with multiple types, sizes, and morphologies (1). Nanoparticles (NPs) are characterized by unique physical and chemical features like, surface area, pore size, particle morphology, and reactivity. Another name of NPs is “magic bullets” due to their intensive applications in agricultural field. Nanoparticles can

be used as nanofertilizers, nano-pesticides and herbicides which are useful to increase crops productivity, to control excessive uses of chemicals fertilizers and also increase survivability against biotic stress. They regulate plant development and increase metabolic activity (2-4). NPs may affect on the plant growth species either positively or negatively depending on its type and concentration used. It is important to consider the interaction and impact of NPs in agricultural, for example Zn which expresses toxic effects whereas silicon is detected as valuable for flora (5). Several investigators



Fig. 1. Synthesis of silicon nanoparticles from agro-waste (Sugarcane bagasse)

depicted that NPs have potential to improve the germination rate, growth and productivity. Higher amount of NPs occur in the soil rather than water and air (6). The influence of NPs on agronomic, atmosphere or society can be both progressive and destructive.

Synthesis of Nanoparticles

Nanoparticles can be synthesized in three ways: physical, chemical, and biological. Commonly physical and chemical methods used for the synthesis of nanoparticles. Due to aforesaid reasons, biological synthesis pathway prefer over the physical and chemical methods for the synthesis of NPs.

Major disadvantage of Physical and chemical methods is that physical methods needs a more time to achieve thermal stability, consuming a lot of energy, occupying large space in case of furnace. In case of chemical synthesis of NPs employs harsh reducing agent like sodium borohydride, sodium citrate and organic solvents etc. These chemical reagents pose toxicity issues along with environment issues. Due to toxic impacts of the physical and chemical methods, biological method prefer for the synthesis of NPs. Production of NPs involves use of bacteria, fungi, algae, plant and agro-waste. Agronomy is a backbone of our economy but now agricultural sector facing many problems like environmental issue, urbanization, sustainable uses of resources, climate changes. NPs can be synthesized through green technology by using algae, plant extract and agricultural waste. It would be a cost-effective, ecofriendly and easy to handle as compare to physical and chemical methods (7,8). So, these materials can be used for improvement of crop productivity and minimize agronomy problems. Plants and agro-waste tender a superior option for the biogenic NPs as the protocols involving plant source are free from toxicants. The various NPs can be synthesized through plant extract like *Aloe vera*, *Allium sativum*, *Chenopodium album*, *Azadirachta indica*, *Citrus sinensis*, *Eclipta prostrata*, *Vitex negundo*, *Mangifera indica*, *Glycine max* and from algae such as *Sargassum muticum*, *Chlamydomonas spp.*, *Euglena gracillis*, *Chlorella vulgaris* and from agriculture waste, sugarcane bagasse (Fig. 1), Egg shell, rice husk, mango peel, corn cob bamboo leaves etc. (Table 1 and 2).

Interaction and Translocation of Nanoparticles in plants

Interaction and uptake of NPs by the plant system based upon the nature of NPs (Size, shape and concentration). The NPs lies in size (40-50 nm), can penetrate the plant tissues. NPs interact with plant roots system with different methods, which are specialized in absorption of nutrient and water. NPs of 2 nm size can be applied exogenously through leaves and will cross cuticular pore and stomata (29,30). The presence of symbiotic bacteria, fungi, algae also influences the uptake of NPs into plant system (31,32). Here are certain processes and carriers by which NPs can enter into the plant system and cross the plasma membrane such as endocytosis, pore-formation, carrier proteins, Plasmodesmata and ion channels. Endocytosis is most suitable way for NPs to enter into plant cell. The interaction and translocation of NPs depend upon the size, shape, concentration and plant variety. The response of NPs depends upon the plant species, age and internal and external conditions. NPs can damage the cell wall and plasma membranes through penetration into the cell (33). Cell wall is first barrier as a molecular sieve which permits only minor NPs to enter into the cell. Small sized NPs move into the cell by root system and large sized NPs enter through flower stigma and stomata with the help of endocytosis and symplast (34,35). After penetrating the cell, NPs can move only by two ways such as the apoplast and the symplast (Fig. 2).

Phytotoxicity and survival mechanism induced by NPs

NPs enter into the cell and interrupt Electron transport system (ETS) cycle of chloroplast and mitochondria and triggered oxidative burst due to increase of reactive oxygen species (ROS) concentration (36, 37). ROS have both hydroxyl radical and superoxide free radical and non-radical singlet like oxygen and hydrogen peroxidase (37). It work as an transporter molecule, excess level of ROS cause adversative impacts in cell known as oxidative stress (38-42). When cells are exposed to excess levels of ROS induce destruction of DNA, oxidation of protein, lipid peroxidation and membrane damage finally caused programmed cell death (43). So, metal and metal based NPs can induce oxidative stress in many plant species. ROS lead to oxidative

destruction in chloroplast and mitochondrial DNA damage and DNA molecule causes change in the encoded protein that result in malfunction and entire inactivation of the determined proteins (44). Various study showed that *Ricinus communis* seed under the treatment of AgNPs increase ROS

production which involved in antioxidant defense mechanism (Superoxide dismutase enzyme (SOD), peroxidase (POD) activity and phenolic acids). Production of phenol increase defense against pathogen. Due to ROS generation, increase root length up to a certain level (45-48).

Table 1. Nanoparticles synthesized from different plants species

Plant	Type of Nanoparticles	References
<i>Aloe vera</i>	Au, Ag, Cu, ZnO	(10,20)
<i>Allium sativum</i>	Ag	(11)
<i>Anacardium occidentale</i>	Au/Ag bimetallic	(12)
<i>Azadirachta indica</i>	Au/Ag bimetallic	(13)
<i>Chenopodium album</i>	Ag, Au	(14)
<i>Citrus sinensis</i>	Ag	(15)
<i>Eclipta prostrata</i>	Ag	(16)
<i>Terminalia catappa</i>	Au	(17)
<i>Vitex negundo</i>		(18)
<i>Mangifera indica</i>		(19)
<i>Lycopersicum esculentum</i>	Zn	(21)
<i>Glycine max</i>	Pd	(22)

Table 2. NPs from Agro waste of wild growing plants

Agro waste	NPs	References
Tea leaf extract (<i>Camellia sinensis</i>)	Ag	(23)
Egg shell	Au	(24)
Banana peel extract	Au	(25)
Papaya fruit extract	Ag	(26)
Sugarcane bagasse, Cassava periderm, Maize stalk	Si	(9)
Grape waste	Au	(27)
Tea waste	Fe	(28)




Fig. 2. Interaction, translocation and mechanism induced by nanoparticles in plants, from Kurpea (33)

Table 3. Positive and negative effect of different nanoparticles on various plant species

Nanoparticles	Plant species	Effects	References
AgNPs		Positive effect	
	<i>Lolium multifolium, Eruca sativa</i>	Enhance plant growth	(50)
	<i>Zea mays</i>	Negative effect	
		At low concentration root and shoot length increase but at high concentration decrease the length	(51)
	<i>Sorghum bicolor</i>	Reduced root growth, root length and biomass	(50)
	<i>Lolium multiform</i>	Broken epidermis and root cap	(52)
	<i>Baccopa monnieri</i>	Different concentration affected the seed germination	(53)
<i>Vigna radiata</i>	Decrease shoot weight and increase biomass. Reduction in growth		
<i>Triticum aestivum</i>			
ZnNPs		Positive effect	
	<i>Cucumis sativa</i>	Increase micronutrients	
	<i>Cicer arietum</i>	Increase dry shoot weight	(70)
	<i>Cyamopsis tetragonoloba</i>	Increase biomass, root and shoot length	(84)
	<i>Vigna radiata</i>	Increase germination, root and shoot growth	(101)
	<i>Arachis hypogaea</i>	Increase seed germination at low concentration but decrease at high concentration	(67)
ZnO		Negative effect	
	<i>Cucumis sativa</i>	Inhibited root growth	(70)
	<i>Triticum aestivum</i>	Inhibition of chlorophyll synthesis leading to reduce photosynthetic efficiency	(68)
	<i>Brassica napus</i>	Reduced germination and simultaneously inhibit root growth	(102)
SiO ₂		Positive effect	
	<i>Lycopersicum esculentum</i>	At low concentration enhance seed germination of tomato	(63)
	<i>Glycine max</i>	Promoter effect on germination	(64)
Si	<i>Ocimum basilicum</i>	Increase dry and fresh weight of leaf	(66)
TiO ₂		Positive effect	
	<i>Vigna radiata, Arabidopsis thaliana, Foenicum vulgare, Lemna minor, Triticum aestivum</i>	enhance germination, plant growth chlorophyll content	(82)
AlO		Positive effect	
	<i>Triticum aestivum</i>	Significantly improved elongation	(83)
		Negative effect	
	<i>Nicotiana tabacum</i>	Decrease plant growth and development	(83)
Cu		Positive effect	
	<i>Triticum aestivum</i>	Increase in germination percentage	(57)
Fe		Positive effect	
	<i>Triticum aestivum</i>	Reduction in root length	(57)
FeO		Negative effect	
	<i>Arabidopsis thaliana</i>	Inhibitory effects on development	(52)
CNTs		Positive effect	
	<i>Hordium vulgare, Glycine max, Zea mays</i>	Increase seed germination	(62)
MWCNTs			
	<i>Lycopersicum esculentum, Nicotiana tabacum</i>	Increase plant growth, height, flower number	(61)

Positive and Negative Effects of various NPs on Plant Growth and Seed Germination

The main objective of nanomaterials in agronomy is to promote plant protection, productivity and

reduce nutrient losses (7). They can also be used to reduce the toxicity of metals and increase crop productivity without high cost of energy (8). The extensive uses of NPs promote seed germination,

plant development and can change its morphology with abiotic stress shown in (Table 3).

Silver (Ag) NPs

AgNPs shows both progressive and destructive influence on plants growth and development. The extensive use of AgNPs leads to an enhancement in peroxidase, catalase activities and synthesis of antioxidants complexes like carotenoids and proline. It also enhance plant growth and seed germination in *Lolium multifolium* and *Eruca sativa* (50) but in *Vigna radiata*, *Sorghum bicolor*, inhibit the root length at high concentration and increase at low concentration (51,52). In *Triticum aestivum*, AgNPs decrease shoots weight and increase biomass. Green synthesized AgNPs increase the rate of germination of *Boswellia ovalifoliolata*, improved growth rate (root and shoot length, leaf area) and enhanced biochemical traits (enzymes, protein content antioxidant, carbohydrates, chlorophyll) in *Brassica juncea*, *Phaseolus vulgaris* and *Zea mays*. The seeds of *Oryza sativa* treated with 20 nm AgNO₃ which shows minor negative effects on seed germination (Fig. 3). The silver NPs synthesized by green route also shows significant impact on *Trigonella foenum-graecum* seedlings (53).



Fig. 3. Effect of Silver nanoparticles on *Oryza sativa*, Thuesombat (49)

Carbon Nanotubes (CNTs)

Carbon-based nanomaterials have gained high attention due to its unique properties. CNTs are classified into two categories *i.e.* single-walled carbon nanotubes (SWNT) and multi-walled carbon nanotubes (MWNTs). CNTs have the tendency to penetrate the wall of cell and deliver to cells. Single-walled-CNTs (SWCNTs) act as transporters for transmit DNA fragments. Numerous studied shows that multi-walled-CNTs (MWCNTs) have tendency to improved development of plant and uptake efficiency of essential nutrients (54-56). MWCNTs regulated genes expression in *Hordeum vulgare*. CNTs show the progressive role in growth and germination rate, but have negative influence in various floras (57). MWCNTs improved peroxidase, dehydrogenase activity, gene expression (58,59)

and increase the rate of germination in *Oryza sativa*, *Glycine max*, *Brassica juncea*, *Phaseolus mungo* and *Lycopersicum esculentum*. Water soluble CNTs influence the *Triticum aestivum* development in sunny and shady environments (60-62).

Silicon Dioxide (SiO₂) NPs

NPs influence the plant growth and seed germination and it depends upon the concentration. Nano-SiO₂ increase sprouting in *Lycopersicum esculentum* at low concentration (63). Nano-SiO₂ and nano-titanium dioxide (nano-TiO₂) enhanced seed sprouting of *Glycine max* through increasing nitrate reductase (64) which improved the ability of seed to absorb water and nutrients. Due to Nano-SiO₂, plant can grow despite in environment stress and the use of nano-SiO₂ enhance leaves weight, accumulation of proline, amino acids, nutrients, chlorophyll, and action of enzymes (65). SiO₂NPs also enhance the growth of plant, rate of photosynthesis, PSII potential activity, photochemical efficiency, stomata conductance, transpiration rate, ETS. Leaf dry and fresh weight, chlorophyll and proline content of *Ocimum basilicum* increased by silica NPs (66).

Zinc Oxide (ZnO) NPs

ZnO NPs show positive impact on *Arachis hypogaea*, *Glycine max*, *T. aestivum* and *Allium cepa* at low concentration (67-69). Only *Cucumis sativus* germination was increase when different concentrations of ZnO NPs were applied on different plant species like *Cucumis sativus*, *Medicago sativa* and *Lycopersicum esculentum* (70). In *Cymopsistetra gonoloba*, ZnO NPs promoted an important enhancement like improve plant biomass, plant growth, chlorophyll content, synthesis of protein (71). Nano ZnO added in MS media promoted embryogenesis, plantlets regeneration, plant development and some enzymes (proline, superoxide dismutase, catalase, and peroxidase) which are promote to survive in biotic stress (72). ZnO NPs show positive effect on *Fagopyrum esculentum* (Fig. 4). The presence of ZnO NPs improved the antioxidant properties, photosynthetic efficiency and enhanced proline accumulation that provides stability to plants (74).

Gold (Au) NPs

AuNPs increase the amount of leaves, plant height, and chlorophyll content and promoted the photochemical reaction, increase NADPH + H⁺ and ATP to convey CO₂ fixation, increases sugar amount in treated seedling (75,76). Cellular and physiological process (lipid metabolic rate, glycolysis, transport system, DNA transcription, synthesis of protein, photosynthesis rate) increased due to treatment of AuNPs. Green synthesized AuNPs from *Tiliacora sp.* show minimum toxicity in rice seedlings (Fig. 5) which can be potentially used as nanocarriers in agriculture.




Fig. 4. Scanning electron microscope images of Buckwheat (*Fagopyrum esculentum*) root surface under control (left) and treatment (right) with ZnO nanoparticles (1,000 mgL⁻¹) at a magnification of ×1,000 (a), ×5,000 (b), and ×150,000 (c); Lee (73)




Fig. 5. Effect of different concentration of gold nanoparticles on *Oryza sativa*; Ndeh (77)

Titanium Dioxide (TiO₂) NPs

TiO₂ NPs initiate progressive growth, development and productivity of crop under abiotic pressure in *Triticum aestivum* plant. Various studies of TiO₂ NPs were done on bacteria, algae, planktons, fish, mice, but remain incomplete on plant. Enzymes activity (glutamine synthase, glutamate dehydrogenase, glutamate-pyruvate transaminase and nitrate reductase), which are involve in nitrogen metabolism was direct regulated by TiO₂ NPs. Conversion of Inorganic nitrogen to organic nitrogen in the form of protein and chlorophyll and enhancement weight of plant was

also done with the help of titanium oxide (78,79). TiO₂ NPs protect the chloroplast from extreme light through activity of peroxidase, superoxide dismutase catalase, antioxidant enzymes (80,81). TiO₂ NPs act as a photo substance which is used in pigment production, photosynthesis rate enhancer etc. and also increase germination rate and plumule growth in *Brassica napus* (Fig. 6).

Aluminium (Al₂O₃) NPs

In *Nicotiana tabacum*, Al₂O₃ NPs show negative impacts like decrease root length and seedling growth at higher concentration while in *Triticum*

aestivum did not show any toxicity (83). In *Oryza sativa* and *Sorghum bicolor*, excess Al induce iron (Fe) deficiency symptoms (84) and it does not affect the seed germination but helps in new root development and seedling establishment (85). After 2-3 days the initiation of seed germination the root growth inhibition was detected (86). Nano-scale alumina did not have any significant effect on inhibition or increment of shoot length and dry biomass.




Fig. 6. Titanium oxide effect on *Brassica napus* seedlings, Farooqui (3)

Iron oxide (Fe₃O₄) NPs

Uptake and translocation of Fe₃O₄NPs in *Hordeum vulgare* plant which promoted the plant growth probably due to enhancement in photosynthetic efficiency. No phytotoxic effects were recorded at high dose (87). Iron oxide NPs show significant effect on root elongation of *Lactuca sativa* seedlings. FeNPs could be used as a fertilizer (88, 89). The Iron oxide NPs shows positive and negative effects on *Arabidopsis thaliana*. These NPs did not show any effects at the lower concentration but at higher concentration reduce the shoot and root length in plants.

Nanoparticles influence on molecular level in plants

Seedlings of *Oryza sativa* were exposed to various concentrations of AgNPs which increase hydrogen peroxidase formation, lipid peroxidation, proline accumulation and decrease sugar content in shoot and root. High dose of AgNPs increase reactive oxygen species generation which cause cytotoxicity and decrease mitochondrial membrane potential (90). In *Triticum aestivum*, high concentration of AgNPs induced the metallothionein gene (MT) while did not show any expression of the MT gene at low concentration. ZnO NPs up regulated 660 genes and down regulated 826 genes in *Arabidopsis thaliana* which are related to different stress, cell organization. TiO₂NPs showed mild changes in genes as compared to ZnONPs (91,92). In *Arabidopsis thaliana*, genes which are related to oxidative stress, sulfur assimilation, glutathione and proline biosynthesis (ATPS, APR, CS, GCL, P5CS1, and P5GS2) were up regulated by high concentrations

of CuONPs (93). Reduced the hydrogen peroxidase content and expression of photosynthetic genes (petA and psaA) were increased at high concentration of Fe₃O₄NPs. Receptor molecules sense the occurrence of exterior or interior particles and modify its own confirmation then message transferred to the receptor molecules which bind to DNA and leads to change in the transcription and control the gene expression. Thus, nanoparticles have good potential to change the cell signaling pathways. Different proteins translate via different RNA and these proteins are directly associated with metabolic processes by accessing of the RNA expression. When plants are exposed to various nanoparticles several reports have shown an up regulation or down regulation of genes which are related to plant stress (94). Other reports depicted that plant DNA can be damaged under the presence of nanoparticles which may result in a low level of RNA expression and ultimately the death of the organism. NPs used as a carrier for transport of genetic material into the plant cell.

Application of nanoparticles on sustainable agriculture

Nanoparticles has the ability to transform the agricultural and food industry through various innovative tools for the disease management, disease detection and enhancing the capacity of plants to absorb nutrients (7). The main objective of NPs is to increase crop efficiency with low input of fertilizers, pesticides, and herbicides among others chemical compounds with same salts. NPs also called as “magic bullets” which aims to target particular cellular organelles in plant to pass their content (8) and also contain nano-pesticide, nano-fertilizers and herbicides properties to control plant diseases and promote plant growth.

NPs as fertilizers

Uses of Nano-fertilizers can improve crop yield. Contented and reactivity of phytohormones are main index of plant physiology. CeO₂ NPs improve the biochemical aspects like amino acids, non-reducing sugar, fatty acids, and phenolic complex in numerous plants. Nano-fertilizers increase nutrient productivity and provide the nutrient according to the need of crops. Many studies depicted that chemical fertilizers are less efficient compare to nano fertilizers (7,8,87).

Nano-herbicides

Nano-herbicides used to eradicate weeds and their propagation. Very minor amount of nanoparticles is capable to amalgam with loam and eliminate weeds from nature without any toxicity and avoid the development of unwanted plant (87).

Nano-Pesticides

Porous hollow silica NPs (PHSNs) with validamycin (pesticide) as an effective transport structure of water-soluble pesticide to control many diseases.

Nano-emulsions were valuable to constructions of pesticides then it might be active in contradiction of the numerous pests in agronomy (95). Likewise, oil-loaded NPs were more beneficial on behalf of preparations of nano-pesticides (96). Nano silica can be effectively used as a nano pesticide.

Nano insecticide

Encapsulation of nano is presently the greatest skill used for defense host plant in contradiction of pest. Nanosilicon can be used as insect repellent. After applied the silicon on the surface of plant cause insects death by physical mean. Insect repellent activity of Poly- ethylene glycol-coated NPs used as insecticide against adult *Tribolium castaneum* insect. SiNPs loaded with validamycin (pesticide) can be used as efficient pesticide (96).

NPs as a Plant diseases control

Ag and SiNPs have potential to control the many plant diseases. Silver nanoparticles (AgNPs) can be used as antimicrobial negotiator which prevents the action to microbes (97,98). These nanoparticles has countless prospective in the controlling of plant syndromes comparatively to artificial fungicides (99,100). ZnO and MgO NPs are active antibacterial agents. Therefore, usage of NPs has been measured an alternative and operative method which is environmentally friendly and cost effective for the control of pathogenic microorganism.

Conclusion

Nanotechnology have huge prospective and profits in the field of agronomy and biotechnology. As discussed in the present review the possibility of negative impact of nanoparticles on plants along with positive implications, it is necessary to carefully examine the concentration of nanoparticles and its interaction with plants. Such knowledge is important before the use of nanoparticles in future research work in plant sciences. The size of NPs also plays an important role in the interaction and accumulation inside the plants. Although there are new approaches for plants improvement using NPs has shown the successful effect even though NPs has shown many lethal effects on plant growth and survivability rate. Therefore, the research on NPs especially related to interaction with plants and their impacts on physiological, biochemical and molecular impacts need to be carried out.

Acknowledgements

The authors are grateful to Prof. Aditya Shastri, Vice Chancellor, Banasthali Vidyapith and Prof. Dipjyoti Chakraborty, Head, Bioscience and Biotechnology Department, Banasthali Vidyapith for providing their necessary support and encouragement.

Authors' contribution

JM conceived the idea of writing the review and designed the content. PG and SY collected the literature and prepared the article. All authors contributed to the content of the manuscript and approved the final version.

Competing Interest

The authors declare that they have no competing interest.

References

- Hao Y, Zhang Z, Rui Y, Ren J. Effect of different nanoparticles on seed germination and seedling growth in rice. Annual International Conference on Advanced Material Engineering 2016;1-8. <https://doi.org/10.2991/ame-16.2016.28>
- Siddiqui MH, Al-Wahaibi MH. Role of nano-SiO₂ in germination of tomato (*Lycopersicon esculentum* seeds Mill.). Saudi J. Biol. Sci. 2014;21(1):13-7. <https://doi.org/10.1016/j.sjbs.2013.04.005>
- Farooqui AR, Tabassum HE, Ahmad AS, Mabood AB, Ahmad AD, Ahmad IZ. Role of nanoparticles in growth and development of plant. Int. J. Pharma. Bio. Sci. 2016; 7(4):22-37.
- Husein A, Siddiqui KS. Carbon and fullerene nanomaterials in plant system. J. Nanobiotechnology. 2014;12(1):16. <https://doi.org/10.1186/1477-3155-12-16>
- Mirshekali H, Hadi HA, Amirnia R, Khodaverdiloo H. Effect of zinc toxicity on plant productivity, chlorophyll and Zn contents of sorghum (*Sorghum bicolor*) and common lambsquarter (*Chenopodium album*) Int. J. Agr. Sci. 2012; 2(3):247-54.
- Gottschalk F, Sonderer T, Scholz RW, Nowack B. Modeled environmental concentrations of engineered nanomaterials (TiO₂, ZnO, Ag, CNT, and Fullerenes) for different regions. Environ. Sci. Technol. 2009; 43(24):9216-22. <https://doi.org/10.1021/es9015553>
- Ingale AG, Chaudhari AN. Biogenic synthesis of nanoparticles and potential applications: An eco-friendly approach. J. Nanomed. Nanotechnol. 2013;4:165. <https://doi.org/10.4172/2157-7439.1000165>
- Rai M, Ingle A. Role of nanotechnology in agriculture with special reference to management of insect pests. Appl. Microbiol. Biotechnol. 2012;94(2):287-93. <https://doi.org/10.1007/s00253-012-3969-4>
- Mohd NK, Wee NN, Azmi AA. Green synthesis of silica nanoparticles using sugarcane bagasse. AIP Conference Proceedings. 2017;1885(1). <https://doi.org/10.1063/1.5002317>
- Karimi J, Mohsenzadeh S. Rapid green and eco-friendly biosynthesis of copper nanoparticles using lower extract of *Aloe vera*. Synth. React. Inorg. M. 2014;45(6):895-8. <https://doi.org/10.1080/15533174.2013.862644>
- Rastogi L, Arunachalam J. Sunlight based irradiation strategy for rapid green synthesis of highly stable silver nanoparticles using aqueous garlic (*Allium sativum*) extract and their antibacterial potential. Mater. Chem. Phys. 2011;129(1-2):558-63. <https://doi.org/10.1016/j.matchemphys.2011.04.068>
- Sheny DS, Philip D, Mathew J. Rapid green synthesis of palladium nanoparticles using the dried leaf of *Anacardium occidentale*. Philip. Spectrochim. Acta A. 2012; 91:35-8. <https://doi.org/10.1016/j.saa.2012.01.063>

13. Shankar SS, Rai A, Ahmad A, Sastry M. Rapid synthesis of Au, Ag, and bimetallic Au core-Ag shell nanoparticles using Neem (*Azadirachta indica*) leaf broth. *J. Colloid. Interface. Sci.* 2004;275(2):496-502. <https://doi.org/10.1016/j.jcis.2004.03.003>
14. Dwivedi AD, Gopal K. Biosynthesis of silver and gold nanoparticles using *Chenopodium album* leaf extract. *Colloids Surf. A.* 2010;369(1-3):27-33. <https://doi.org/10.1016/j.colsurfa.2010.07.020>
15. Kaviya S, Santhanalakshmi J, Viswanathan B, Muthumary J, Srinivasan K. Biosynthesis of silver nanoparticles using *Citrus sinensis* peel extract and its antibacterial activity. *Spectrochim. Acta A.* 2011; 79(3):594-8. <https://doi.org/10.1016/j.saa.2011.03.040>
16. Rajakumar G, Rahuman AA. Larvicidal activity of synthesized silver nanoparticles using *Eclipta prostrata* leaf extract against filariasis and malaria vectors. *Acta Trop.* 2011;118(3):196-203. <https://doi.org/10.1016/j.actatropica.2011.03.003>
17. Kumar V, Yadav SK. Plant-mediated synthesis of silver and gold nanoparticles and their applications. *J. Chem. Technol. Biotechnol.* 2009;84(2):151-7. <https://doi.org/10.1002/jctb.2023>
18. Prabhu D, Arulvasu C, Babu G, Manikandan R, Srinivasan P. Biologically synthesized silver nanoparticles from leaf extract of *Vitex negundo* L. induce growth-inhibitory effect on human colon cancer cell line HCT15. *Process Biochem.* 2013;48(2):317-24. <https://doi.org/10.1016/j.procbio.2012.12.013>
19. Yang N, Wei Hong, Hao. Biosynthesis of Au nanoparticles using agricultural waste mango peel extract and its in vitro cytotoxic effect on two normal cells. *Mater. Lett.* 2014; 134:67-70. <https://doi.org/10.1016/j.matlet.2014.07.025>
20. Manickam S, Venkatachalam M, Parthasarathy. Green Synthesis of Nano Particles from *Aloe vera* Extract – Review Paper. *Imp. J. Interdiscip. Res.* 2016; 2(10):1570-5.
21. Sutradhar P, Saha M. Green synthesis of zinc oxide nanoparticles using tomato (*Lycopersicon esculentum*) extract and its photovoltaic application. *J. Exp. Nanosci.* 2016;11(5):314-27. <https://doi.org/10.1080/17458080.2015.1059504>
22. Petla RK, Vivekanandhan S, Misra M, Mohanty AK, Satyanarayana N. Soybean (*Glycine max*) leaf extract based green synthesis of palladium nanoparticles. *J. Biomater. Nanobiotechnol.* 2012;3(1):14-19. <https://doi.org/10.4236/jbnb.2012.31003>
23. Loo YY, Chieng BW, Nishibuchi M, Radu S. Synthesis of silver nanoparticles by using tea leaf extract from *Camellia sinensis*. *Int. J. Nanomedicine.* 2012;7:4263-7. <https://doi.org/10.2147/IJN.S33344>
24. Devi PS, Banerjee S, Chowdhury SR, Kumar GS. Eggshell membrane: a natural biotemplate to synthesize fluorescent gold nanoparticles. *RSC Adv.* 2012;2(30):11578-85. <https://doi.org/10.1039/C2RA21053C.s>
25. Bankar A, Joshi B, Kumar AR, Zinjarde S. Banana peel extract mediated synthesis of gold nanoparticles. *Colloids Surf. B.* 2010;80(1):45-50. <https://doi.org/10.1016/j.colsurfb.2010.05.029>
26. Jain D, Daima HK, Kachhwaha S, Kothari SL. Synthesis of plant-mediated silver nanoparticles using papaya fruit extract and evaluation of their antimicrobial activities *Dig. J. Nanomater. Bios.* 2009; 4(3):557-63.
27. Krishnaswamy K, Vali H, Orsat V. Value-adding to grape waste: green synthesis of gold nanoparticles. *Int. J. Food Eng.* 2014;142:210-20. <https://doi.org/10.1016/j.jfoodeng.2014.06.014>
28. Gottimukkala KSV, Harika P, Zamare D. Green synthesis of iron nanoparticles using green tea leaves extract. *J. Nanomedicine Biotherapeutic Discov.* 2017;7:151. <https://doi.org/10.4172/2155-983X.1000151>
29. Eichert T, Goldbach HE. Equivalent pore radii of hydrophilic foliar uptake routes in stomatous and leaf surfaces—further evidence for a stomatal pathway. *Physiol. Plant.* 2008; 132(4):491-502. <https://doi.org/10.1111/j.1399-3054.2007.01023.x>
30. Schwab F, Zhai G, Kern M, Turner A, Schnoor JL, Wiesner MR. Barriers, pathways and processes for uptake, translocation and accumulation of nanomaterials in plants. *Nanotoxicology.* 2016;10(3):257-78. <https://doi.org/10.3109/17435390.2015.1048326>
31. Feng Y, Cui X, He S, Dong G, Chen M, Wang J, Lin X. The role of metal nanoparticles in influencing arbuscular mycorrhizal fungi effects on plant growth. *Environ. Sci. Technol.* 2013; 47(16):9496-504. <https://doi.org/10.1021/es402109n>
32. Wang Z, Xie X, Zhao J, Liu X, Feng W, White JC, Xing B. Xylem- and phloem-based transport of CuO nanoparticles in maize (*Zea mays* L.). *Environ. Sci. Technol.* 2012; 46(8):4434-41. <https://doi.org/10.1021/es204212z>
33. Kurepa J, Paunesku T, Vogt S, Arora H, Rabatic BM, Lu J, Wanzer MB, Woloschak GE, Smalle JA. Uptake and distribution of ultrasmall anatase TiO₂ alizarin red. S nanoconjugates in *Arabidopsis thaliana*. *Nano. Lett.* 2010; 10(7):2296-302. <https://doi.org/10.1021/nl903518f>
34. Ma X, Geisler-Lee J, Deng Y and Kolmakov A. Interactions between engineered nanoparticles (ENPs) and plants: phytotoxicity, uptake and accumulation. *Sci. Total Environ.* 2010;408(16):3053-61. <https://doi.org/10.1016/j.scitotenv.2010.03.031>
35. Nair R, Varghese SH, Nair BG, Maekawa T, Yoshida Y, Kumar DS. Nanoparticulate material delivery to plants. *Plant. Physiol.* 2010;179(3):154-63. <https://doi.org/10.1016/j.plantsci.2010.04.012>
36. Hossain Z, Mustafa G, and Komatsu S. Plant responses to nanoparticle stress. *Int. J. Mol. Sci.* 2015; 16(11):26644-53. <https://doi.org/10.3390/ijms161125980>
37. Dimkpa CO, McLean JE, Martineau N, Britt DW, Haverkamp R, Anderson AJ. Silver nanoparticles disrupt wheat (*Triticum aestivum* L.) growth in a sand matrix. *Environ. Sci. Technol.* 2013;47(2):1082-90. <https://doi.org/10.1021/es302973y>
38. Faisal M, Saquib Q, Alatar AA, Al-Khedhairi AA, Hegazy AK, Musarrat J. Phytotoxic hazards of NiO-nanoparticles in tomato: a study on mechanism of cell death. *J. Hazard. Mater.* 2013;15(250):318-32. <https://doi.org/10.1016/j.jhazmat.2013.01.063>
39. Jiang HS, Qiu XN, Li GB, Li W, Yin LY. Silver nanoparticles induced accumulation of reactive oxygen species and alteration of antioxidant systems in the aquatic plant *Spirodela polyrhiza*. *Environ. Toxicol. Chem.* 2014; 33(6):1398-405. <https://doi.org/10.1002/etc.2577>
40. Pakrashi S, Jain N, Dalai S, Jayakumar J, Chandrasekaran PT, Raichur AM, Chandrasekaran N, Mukherjee A. *In vivo* genotoxicity assessment of titanium dioxide nanoparticles by *Allium cepa* root tip assay at high exposure concentrations. *PLoS One* 2014;49(2):877-89. <https://doi.org/10.1371/journal.pone.0087789>
41. Cvjetko P, Milošić A, Domijan AM, Vrčec IV, Tolić S, Štefanić PP, Letofsky-Papst I, Tkalec M, Balen B. Toxicity of silver ions and differently coated silver nanoparticles in *Allium cepa* roots. *Ecotox. Environ. safe.* 2017;137:18-28. <https://doi.org/10.1016/j.ecoenv.2016.11.009>
42. Gill SS, Tuteja N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant. Physiol. Biochem.* 2010;48(12):909-30. <https://doi.org/10.1016/j.plaphy.2010.08.016>

43. Van Breusegem F, Dat JF. Reactive oxygen species in plant cell death. *Plant Physiol.* 2006;141(2):384-90. <https://doi.org/10.1104/pp.106.078295>
44. Atha DH, Wang H, Petersen EJ, Cleveland D, Holbrook RD, Jaruga P, Dizdaroglu M, Xing B, Nelson BC. Copper oxide nanoparticle mediated DNA damage in terrestrial plant models. *Environ. Sci. Technol.* 2012;46(3):1819–27. <https://doi.org/10.1021/es202660k>
45. Belava VN, Panyuta OO, Yakovleva GM, Pysmenna YM, Volkogon MV. The effect of silver and copper nanoparticles on the wheat- *Pseudocercospora herpotrichoides* patho system. *Nanoscale Res. Lett.* 2017; 12(1):250. <https://doi.org/10.1186/s11671-017-2028-6>
46. Saha N, Gupta S. Low-dose toxicity of biogenic silver nanoparticles fabricated by *Swertia chirata* on root tips and flower buds of *Allium cepa*. *J. Hazard. Mater.* 2017; 15:330:18-28. <https://doi.org/10.1016/j.jhazmat.2017.01.021>
47. Oukarroum A, Barhoumi L, Pirastru L, Dewez D. Silver nanoparticle toxicity effect on growth and cellular viability of the aquatic plant *Lemna gibba*. *Environ. Toxicol. Chem.* 2013; 32(4):902-7. <https://doi.org/10.1002/etc.2131>
48. Sharma P, Jha B, Dubey RS and Pessaraki M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Am. J. Bot.* 2012; Article ID 217037, 26 pages. <http://dx.doi.org/10.1155/2012/217037>
49. Thuesombat P, Hannongbua S, Akasit S, Chadchawan S. Effect of silver nanoparticles on rice (*Oryza sativa* L. cv. KDML 105) seed germination and seedling growth. *Ecotoxicol. Environ. Saf.* 2014;104:302–9. <http://dx.doi.org/10.1016/j.ecoenv.2014.03.022>
50. Vannini C, Domingo G, Onelli E, Prinsi B, Marsoni M, Espen L, Bracale M. Morphological and proteomic responses of *Eruca sativa* exposed to silver nanoparticles or silver nitrate. *PLoS ONE.* 2013;8(7):68752. <http://dx.doi.org/10.1371/journal.pone.0068752>
51. Krishnaraj C, Ramachandran R, Mohan K, Kalaichelvan PT. Optimization for rapid synthesis of silver nanoparticles and its effect on phytopathogenic fungi. *Spectrochim. Acta A. Mol. Biomol. Spectrosc.* 2012; 93:95–99. <https://doi.org/10.1016/j.saa.2012.03.002>
52. Fageria NK, Baligar VC, and Wright RJ. Iron nutrition of plants: An overview on the chemistry and physiology of its deficiency and toxicity. *Pesqui. Agropecu. bras.* 1990; 25(4):553-70.
53. Jasim B, Thomas R, Mathew J, Radhakrishnan EK. Plant growth and diosgenin enhancement effect of silver nanoparticles in Fenugreek (*Trigonella foenum-graecum* L.). *Saudi. Pharm. J.* 2017;25(3):443–7. <https://doi.org/10.1016/j.jsps.2016.09.012>
54. Srinivasan C, Saraswathi R. Nano-agriculture-carbon nanotubes enhance tomato seed germination and plant growth. *Curr. Sci.* 2010; 99(3):273–5.
55. Tiwari DK, Dasgupta-Schubert N, Cendejas LV, Villegas J, Montoya LC, García SB. Interfacing carbon nanotubes (CNT) with plants: enhancement of growth, water and ionic nutrient uptake in maize (*Zea mays*) and implications for nanoagriculture. *Appl. Nanosci.* 2014;4(5):577-91. <https://doi.org/10.1007/s13204-013-0236-7>
56. Villagarcia H, Dervishi E, de Silva K, Biris AS, Khodakovskaya MV. Surface chemistry of carbon nanotubes impacts the growth and expression of water channel protein in tomato plants. *Small.* 2014; 6-8(15):2328-34. <https://doi.org/10.1002/sml.201102661>
57. Husein A, Siddiqi KS. Carbon and fullerene nanomaterials in plant system. *J. Nanobiotechnology.* 2014;12(1):16. <https://doi.org/10.1186/1477-3155-12-16>
58. Khodakovskaya MV, De Silva K, Biris AS, Dervishi E, Villagarcia H. Carbon nanotubes induce growth enhancement of tobacco cells. *ACS Nano.* 2012; 6(3):2128–35. <https://doi.org/10.1021/nn204643g>
59. Lahiani MH, Dervishi E, Chen J, Nima Z, Gaume A, Biris AS, Khodakovskaya MV. Impact of carbon nanotube exposure to seeds of valuable crops. *Appl. Mater. Interfaces.* 2013; 5(16):7965–73. <https://doi.org/10.1021/am402052x>
60. Tripathi S, Sarkar S. Influence of water soluble carbon dots on the growth of wheat plant. *Appl. Nanosci.* 2015; 5(5):609-16. <https://doi.org/10.1007/s13204-014-0355-9>
61. Morla S, Rao CR, Chakrapani R. Factors affecting seed germination and seedling growth of tomato plants cultured in vitro conditions. *J. Chem. Biol. Phys.* 2011; 1(2):328–34.
62. Lu C, Zhang C, Wen J, Wu G, Tao M. Research on the effect of nanometer materials on germination and growth enhancement of *Glycine max* and its mechanism. *Soybean Science.* 2002; 21(3):168-71.
63. Haghghi M, Afifipour Z, Mozafarian M. The effect of N-Si on tomato seed germination under salinity levels. *JBES.* 2012; 6(16):87-90.
64. Zheng L, Hong F, Lu S, Liu C. Effect of nano-TiO₂ on strength of naturally aged seeds and growth of spinach. *Biol. Trace. Elem. Res.* 2005;104(1):83-91. <https://doi.org/10.1385/BTER:104:1:083>
65. Siddiqui MH, Al-Wahaibi MH, Faisal M, Al Sahli AA. Nano-silicon dioxide mitigates the adverse effects of salt stress on *Cucurbita pepo* L. *Environ. Toxicol. Chem.* 2014; 33(11):2429-37. <https://doi.org/10.1002/etc.2697>
66. Kalteh M, Alipour ZT, Ashraf S, Marashi Aliabadi M, Falah Nosratabadi A. Effect of silica nanoparticles on Basil (*Ocimum basilicum*) under salinity stress. *Journal of chemical health and risks.* 2018;4(3). <https://doi.org/10.22034/JCHR.2018.544075>
67. Prasad TN, Sudhakar P, Sreenivasulu Y, Latha P, Munaswamy V, Reddy KR, Sreeprasad TS, Sajanlal PR, Pradeep T. Effect of nanoscale zinc oxide particles on the germination, growth and yield of peanut. *J. Plant. Nut.* 2012; 35(6):905-27. <https://doi.org/10.1080/01904167.2012.663443>
68. Ramesh M, Palanisamy K, Babu NK. Effects of bulk and nano-titanium dioxide and zinc oxide on physio-morphological changes in *Triticum aestivum* Linn. *Int. J. Glob. Sci.* 2014; 3(2):415-22.
69. Raskar SV and Laware SL. Effect of zinc oxide nanoparticles on cytology and seed germination in onion. *Int. J. Curr. Microbiol. App. Sci.* 2014; 3(2):467-73.
70. de la Rosa G, López-Moreno ML, de Haro D, Botez CE, Peralta-Videa JR, Gardea-Torresdey JL. Effects of ZnO nanoparticles in alfalfa, tomato, and cucumber at the germination stage: root development and X-ray absorption spectroscopy studies. *Pure. Appl. Chem.* 2013; 85(12):2161–74. <https://doi.org/10.1351/pac-con-12-09-05>
71. Raliya R and Tarafdar JC. ZnO nanoparticle biosynthesis and its effect on phosphorous-mobilizing enzyme secretion and gum contents in cluster bean (*Cyamopsis tetragonoloba* L.). *J. Agric.* 2013;2(1):48–57. <https://doi.org/10.1007/s40003-012-0049-z>
72. Helaly MN, El-Metwally MA, El-Hoseiny H. Effect of nanoparticles on biological contamination of *in-vitro* cultures and organogenic regeneration of banana. *Aust. J. Crop Sci.* 2014; 8(4):612-24.
73. Lee S, Kim S, Kim S, Lee I. Assessment of phytotoxicity of ZnONPs on a medicinal plant *Fagopyrum esculentum*. *Environ. Sci. Pollut. R.* 2013;20(2):848-54. <https://doi.org/10.1007/s11356-012-1069-8>

74. Faizan M, Faraz A, Yusuf M, Zinc oxide nanoparticles-mediated changes in photosynthetic efficiency and antioxidant system of tomato plants. *Photosynthetica*. 2018; 56:678. <https://doi.org/10.1007/s11099-017-0717-0>
75. Arora S, Sharma P, Kumar S, Nayan R, Khanna PK, Zaidi MG. Gold-nanoparticles induced enhancement in growth and seed yield of *Brassica juncea*. *Plant Growth Regul.* 2012;66(3):303-10. <https://doi.org/10.1007/s10725-011-9649-z>
76. Gopinath K, Gowri S, Karthika V, Arumugam A. Green synthesis of gold nanoparticles from fruit extract of *Terminalia arjuna*, for the enhanced seed germination activity of *Gloriosa superba*. *J. Nanostructure Chem.* 2014; 4(3):115. <https://doi.org/10.1007/s40097-014-0115-0>
77. Ndeh N, Maensiri N, Maensiri D. The effect of green synthesis gold nanoparticles on rice germination and root. *Adv Nat Sci-Nanosci.* 2017;8:1-10. <https://doi.org/10.1088/2043-6254/aa724a>
78. Yang FL, Li XG, Zhu F, Lei CL. Structural characterization of nanoparticles loaded with garlic essential oil and their insecticidal activity against *Tribolium castaneum* (Herbst) (Coleoptera:Tenebrionidae). *J. Agric. Food Chem.* 2009; 57(21):10156-62. <https://doi.org/10.1021/jf9023118>
79. Pérez-de-Luque A. Interaction of Nanomaterials with Plants: What do we need for real applications in agriculture? *Front. Environ. Sci. Eng.* 2017;5:12. <https://doi.org/10.3389/fenvs.2017.00012>
80. Hong F, Zhou J, Liu C, Yang F, Wu C, Zheng L, Yang P. Effect of nano-TiO₂ on photochemical reaction of chloroplasts of spinach. *Biol. Trace Elem. Res.* 2005; 05:(1-3):269-79. <https://doi.org/10.1385/BTER:105:1-3:269>
81. Mahmoodzadeh H, Nabavi M, Kashefi H. Effect of nanoscale titanium dioxide particles on the germination and growth of canola (*Brassica napus*). *J. Ornamental Hort.* 2013; 3(1):25-32.
82. Scott NR, Chen H, Cui H. Nanotechnology applications and implications of agrochemicals towards sustainable agriculture and food systems. *J. Agric. Food Chem.* 2018; 66(26):6451-56. <https://doi.org/10.1021/acs.jafc.8b00964>
83. Foy CD, Fleming AL. Aluminium of two wheat genotype related to nitrate reductase activity. *J. Plant Nutr.* 1982; 5(11):1313-33. <https://doi.org/10.1080/01904168209363064>
84. Furlani PR, Clark RB. Screening sorghum for aluminium tolerance in nutrient solution. *Agron. J.* 1981; 73(4):587-94.
85. Nosko P, Brassard P, Kramer JR, Kershaw KA. The effect of aluminium on seed germination and early seedling establishment growth and respiration of white spruce (*Picea glauca*). *Can. J. Bot.* 1988;66(11):2305-10. <https://doi.org/10.1139/b88-313>
86. Nicholas DJD. Mineral nutrition of plants. *Annual. J. Plant Physiol.* 2003;12(1):63-90. <https://doi.org/10.1146/annurev.pp12.060161.000431>
87. Tombuloglu H, Slimani Y, Tombuloglu G, Almessiere M, Baykal A. Uptake and translocation of magnetite (Fe₃O₄) nanoparticles and its impact on photosynthetic genes in barley (*Hordeum vulgare* L.). *Chemosphere.* 2019. <https://doi.org/10.1016/j.chemosphere.2019.03.075>
88. Zhu H, Han J, Xiao JQ, Jin Y. Uptake, translocation, and accumulation of manufactured iron oxide nanoparticles by pumpkin plants. *J Environ. Monit.* 2008;10(6):713-7. <https://doi.org/10.1039/b805998e>
89. A farooqui, Tabassum H, Ahmad AS, Mabood AB, Ahmad AD, Ahmad IZ. Role of nanoparticles in growth and development of plants: a review. *Int. J. Pharma Bio. Sci.* 2016; 7(4):22-37. <https://doi.org/10.22376/ijpbs.2016.7.4.p22-37>
90. Prakash M, Gopalakrishnan N. Physiological and molecular level effects of silver nanoparticles exposure in rice (*Oryza sativa* L) seedlings. *Chemosphere* 2014;112:105-13. <https://doi.org/10.1016/j.chemosphere.2014.03.056>
91. Nair PM, Chung IM. Physiological and molecular level effects of silver nanoparticles exposure in rice (*Oryza sativa* L.) seedlings. *Chemosphere* 2014;112:105-13. <https://doi.org/10.1016/j.chemosphere.2014.03.056>
92. Wang Z, Xu L, Zhao J, Wang X, White JC, Xing B. CuO nanoparticle interaction with *Arabidopsis thaliana*: toxicity, parent-progeny transfer, and gene expression. *Environ. Sci. Technol.* 2016;50(11):6008-16. <https://doi.org/10.1021/acs.est.6b01017>
93. Atha H, Wang H, Petersen J. Copper oxide nanoparticle mediated DNA damage in terrestrial plant models. *Environ. Sci. Technol.* 2012;46(3):1819-27. <https://doi.org/10.1021/es202660k>
94. Wang L, Li X, Zhang G, Dong J, Eastoe J. Oil-in-water nano emulsions for pesticide formulations. *J. Colloid Interface Sci.* 2007; 314(1):230-5. <https://doi.org/10.1021/jf9023118>
95. Yang FL, Li XG, Zhu F, Lei CL. Structural characterization of nanoparticles loaded with garlic essential oil and their insecticidal activity against *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *J. Agric Food Chem.* 2009; 57(21):10156-62. <https://doi.org/10.1021/jf9023118>
96. Jo YK, Kim BH, Jung G. Antifungal activity of silver ions and nanoparticles on phytopathogenic fungi. *Plant Dis.* 2009;93(10):1037-43. <https://doi.org/10.1094/PDIS-93-10-1037>
97. Kumar V, Yadav SK. Plant-mediated synthesis of silver and gold nanoparticles and their applications. *J. Chem. Technol. Biotechnol.* 2008;84(2):151-7. <https://doi.org/10.1002/jctb.2023>
98. Swamy VS, Prasad R. Green synthesis of silver nanoparticles from the leaf extract of *Santalum album* and its antimicrobial activity. *J. Optoelectron. Adv. M.* 2012; 4(3):53-59.
99. Park HJ, Kim SH, Kim HJ, Choi SH. A new composition of nanosized silica-silver for control of various plant diseases. *Plant Pathol. J.* 2006; 22(3). <https://doi.org/10.5423/PPJ.2006.22.3.295>
100. Dhoke SK, Mahajan P, Kamble R, Khanna A. Effect of nanoparticles suspension on the growth of mung (*Vigna radiata*) seedling by foliar spray method. *Nanotechnology Development.* 2013; 3(1). <https://doi.org/10.4081/nd.s>
101. Zafar H, Ali A, Ali JS, Haq IU, Zia M. Effect of ZnO nanoparticles on *Brassica nigra* seedlings and stem explants growth dynamics and antioxidative response. *Front. Plant Sci.* 2016;7:535. <https://doi.org/10.3389/fpls.2016.00535>

