

Note di Matematica
Note Mat. **38** (2018) no. 2, 105–115.

ISSN 1123-2536, e-ISSN 1590-0932
doi:10.1285/i15900932v38n2p105

Derivation subalgebras of Lie algebras

F. Saeedi

Department of Mathematics, Mashhad Branch, Islamic Azad University, Mashhad, Iran.
saeedi@mshdiau.ac.ir

S. Sheikh-Mohseni

Department of Mathematics, Mashhad Branch, Islamic Azad University, Mashhad, Iran.
sh.mohseni.s@gmail.com

Received: 25.5.2018; accepted: 30.10.2018.

Abstract. Let L be a Lie algebra and I, J be two ideals of L . If $\text{Der}_J^I(L)$ denotes the set of all derivations of L whose images are in I and send J to zero, then we give necessary and sufficient conditions under which $\text{Der}_J^I(L)$ is equal to some special subalgebras of the derivation algebra of L . We also consider finite dimensional Lie algebra for which the center of the set of inner derivations, $Z(\text{IDer}(L))$, is equal to the set of central derivations of L , $\text{Der}_z(L)$, and give a characterisation of such Lie algebras.

Keywords: Derivation, central derivation, inner derivation

MSC 2000 classification: primary 16W25, secondary 17B40

Introduction

Let L be a Lie algebra over an arbitrary field F . Let L^2 and $Z(L)$ denote the derived algebra and the center of L , respectively. A *derivation* of L is an F -linear transformation $\alpha : L \rightarrow L$ such that $\alpha([x, y]) = [\alpha(x), y] + [x, \alpha(y)]$ for all $x, y \in L$. We denote by $\text{Der}(L)$ the vector space of all derivations of L , which itself forms a Lie algebra with respect to the commutator of linear transformations, called the *derivation algebra* of L . For all $x \in L$, the map $\text{ad}_x : L \rightarrow L$ given by $y \rightarrow [x, y]$ is a derivation called the *inner derivation* corresponding to x . Clearly, the space $\text{IDer}(L) = \{\text{ad}_x \mid x \in L\}$ of inner derivations is an ideal of $\text{Der}(L)$.

Let I and J be two ideals of L . By $\text{Der}^I(L)$ we mean the subalgebra of $\text{Der}(L)$ consisting of all derivations whose images are in I and by $\text{Der}_J(L)$, we mean the subalgebra of $\text{Der}(L)$ consisting of all derivations mapping J onto 0. The subalgebra $\text{Der}^I(L) \cap \text{Der}_J(L)$ is denoted by $\text{Der}_J^I(L)$.

The relationships between the structure of L and $\text{Der}(L)$ is studied by various authors in [1, 4, 6, 7, 8, 16, 19, 20]. Finding necessary and sufficient conditions under which subalgebras of $\text{Der}(L)$ coincide seems to be an interesting problem. There are some results in this regard for a Lie algebra L . Leger [7]

studied the equality $\text{Der}(L) = \text{IDer}(L)$. Tôgô [21], proved that if L is a Lie algebra over a field of characteristic zero such that $Z(L) \neq 0$, then

- (i) $\text{IDer}(L) = \text{Der}_z(L)$ if and only if L is a Heisenberg algebra ($L^2 = Z(L)$ and $\dim Z(L) = 1$), and
- (ii) $\text{Der}(L) = \text{Der}_z(L)$ if and only if L is abelian,

where $\text{Der}_z(L)$ is the subalgebra of $\text{Der}(L)$ consisting of all central derivations of L , that is, the set of all derivations of L mapping L into the center of L . It is easy to see that every element of $\text{Der}_z(L)$ sends the derived algebra of L to 0. In other word, $\text{Der}_z(L) = \text{Der}_{L^2}^{Z(L)}(L)$.

Also, there are many papers about the study of complete Lie algebras. A Lie algebra is said to be *complete*, if its center is zero and all its derivations are inner. The definition of complete Lie algebras was introduced by Jacobson in [5] and later studied in [7, 12, 9, 13, 10, 11, 14, 15, 17, 18, 22].

The aim of this paper is to investigate the equalities $\text{Der}_J^I(L) = Z(\text{IDer}(L))$ and $\text{Der}_J^I(L) = \text{IDer}(L)$ in some special cases. As consequences of our main result, we obtain the result of Tôgô in [21] for finite dimensional Lie algebras and we give a characterisation of Lie algebras satisfying $Z(\text{IDer}(L)) = \text{Der}_z(L)$. Notice that for any Lie algebra L , the center of the set of inner derivations, $Z(\text{IDer}(L))$, is always contained in $\text{Der}_z(L)$.

In Section 3, we study the relationships between $\text{Der}_J^I(L)$ and I, J up to isomorphism for some ideals of L and give a necessary and sufficient condition under which $\text{Der}_{J_1}^{I_1}(L) = \text{Der}_{J_2}^{I_2}(L)$. In Section 4, we discuss the equalities $Z(\text{IDer}(L)) = \text{Der}_J^I(L)$ and $\text{IDer}(L) = \text{Der}_J^I(L)$ in some special cases (Main Theorem and Corollaries). In Section 5, we will characterize all nilpotent Lie algebras of class 2 and naturally graded quasi-filiform Lie algebras satisfying $Z(\text{IDer}(L)) = \text{Der}_z(L)$. We also show that for filiform Lie algebras the above mentioned equality does not hold.

1 Preliminaries

Let A, B be two Lie algebras over a field F and $T(A, B)$ be the set of all linear transformations from A to B . Clearly, if B is an abelian Lie algebra, then $T(A, B)$ equipped with Lie bracket $[f, g](x) = [f(x), g(x)]$ for all $x \in A$ and $f, g \in T(A, B)$ is an abelian Lie algebra. Notice that in this equation the first Lie bracket is taken in $T(A, B)$ and the second Lie bracket is taken in B .

Given a Lie algebra L , the *lower central series* of L is defined as follows:

$$L = \gamma_1(L) \supseteq \gamma_2(L) \supseteq \cdots \supseteq \gamma_n(L) \supseteq \cdots,$$

where $\gamma_2(L) = L^2$ is the derived algebra of L and $\gamma_n(L) = [\gamma_{n-1}(L), L]$.

Also, the *upper central series* of L is defined as

$$\{0\} = Z_0(L) \subseteq Z_1(L) \subseteq \cdots \subseteq Z_n(L) \subseteq \cdots,$$

where $Z_1(L) = Z(L)$ is the center of L and $Z_{n+1}(L)/Z_n(L) = Z(L/Z_n(L))$.

A Lie algebra L is *nilpotent* if there exists a non-negative integer k such that $\gamma_k(L) = 0$. The smallest integer k for which $\gamma_{k+1}(L) = 0$ is called the *nilpotency class* of L . A Lie algebra L of dimension n is *filiform* if $\dim \gamma_i(L) = n - i$ for all $2 \leq i \leq n$. These algebras have maximal nilpotency class $n - 1$. The nilpotent Lie algebras of class $n - 2$ are called *quasi-filiform* and those whose nilpotency class is 1 are abelian.

Let L be a nilpotent Lie algebra of class k over the field of complex numbers \mathbb{C} . Put $S_i = L$, if $i \leq 1$; $S_i = \gamma_i(L)$, if $2 \leq i \leq k$; and $S_i = \{0\}$, if $i > k$. Then, with L we can associate naturally a graded Lie algebra with the same nilpotency class, noted by $\text{gr}L$ and defined by $\text{gr}L = \bigoplus_{i \in \mathbb{Z}} L_i$, where $L_i = \frac{S_i}{S_{i+1}}$. Because of the nilpotency of the algebra, the above gradation is finite, that is $\text{gr}L = L_1 \oplus L_2 \oplus \dots \oplus L_k$ with $[L_i, L_j] \subset L_{i+j}$, for $i + j \leq k$, $\dim L_1 \geq 2$ and $\dim L_i \geq 1$, for all $2 \leq i \leq k$. A Lie algebra L is said to be *naturally graded* if $\text{gr}L$ is isomorphic to L , denoted by $\text{gr}L = L$.

The following theorem of Gómez and Jiménez-Merchán [2] gives a classification of naturally graded quasi-filiform Lie algebras of dimension n .

Theorem 1. *Every naturally graded quasi-filiform Lie algebra of dimension n over the field of complex numbers \mathbb{C} is isomorphic to one of the following algebras:*

- (i) *If n is even to $L_{n-1} \oplus \mathbb{C}$, $\mathcal{T}_{(n,n-3)}$ or $\mathcal{L}_{(n,r)}$, with r odd and $3 \leq r \leq n - 3$.*
- (ii) *If n is odd to $L_{n-1} \oplus \mathbb{C}$, $Q_{n-1} \oplus \mathbb{C}$, $\mathcal{L}_{(n,n-2)}$, $\mathcal{T}_{(n,n-4)}$, $\mathcal{L}_{(n,r)}$, or $\mathcal{Q}_{(n,r)}$, with r odd and $3 \leq r \leq n - 4$. In the cases of $n = 7$ and $n = 9$, we add algebras $\varepsilon_{(7,3)}$, $\varepsilon_{(9,5)}^1$ and $\varepsilon_{(9,5)}^2$.*

Naturally graded quasi-filiform Lie algebras in Theorem 1 are defined in a basis $(X_0, X_1, \dots, X_{n-1})$ as follows.

Split:

$$\begin{aligned} L_{n-1} \oplus \mathbb{C} \ (n \geq 4) : \\ [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq n-3 \\ Q_{n-1} \oplus \mathbb{C} \ (n \geq 7, n \text{ odd}) : \\ [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq n-3 \\ [X_i, X_{n-2-i}] = (-1)^{i-1} X_{n-2}, \ 1 \leq i \leq \frac{n-3}{2} \end{aligned}$$

Terminal:

$$\begin{aligned} \mathcal{T}_{(n,n-3)} \ (n \text{ even}, n \geq 6) : \\ [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq n-3 \\ [X_{n-1}, X_1] = \frac{n-4}{2} X_{n-2} \\ [X_i, X_{n-3-i}] = (-1)^{i-1} (X_{n-3} + X_{n-1}), \\ 1 \leq i \leq \frac{n-4}{2} \\ [X_i, X_{n-2-i}] = (-1)^{i-1} \frac{n-2-2i}{2} X_{n-2}, \\ 1 \leq i \leq \frac{n-4}{2} \end{aligned}$$

$\varepsilon_{(9,5)}^1 :$

$$\begin{aligned} [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq 6 \\ [X_8, X_i] = 2X_{5+i}, \ 1 \leq i \leq 2 \\ [X_1, X_4] = X_5 + X_8 \\ [X_1, X_5] = 2X_6 \\ [X_1, X_6] = 3X_7 \\ [X_2, X_3] = -X_5 - X_8 \\ [X_2, X_4] = -X_6 \\ [X_2, X_5] = -X_7 \end{aligned}$$

$\varepsilon_{(7,3)} :$

$$\begin{aligned} [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq 4 \\ [X_6, X_i] = X_{3+i}, \ 1 \leq i \leq 2 \\ [X_1, X_2] = X_3 + X_6 \\ [X_1, X_i] = X_{i+1}, \ 3 \leq i \leq 4 \end{aligned}$$

Principal:

$$\begin{aligned} \mathcal{L}_{(n,r)} \ (n \geq 5, r \text{ odd}, 3 \leq r \leq 2\lfloor \frac{n-1}{2} \rfloor - 1) : \\ [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq n-3 \\ [X_i, X_{r-i}] = (-1)^{i-1} X_{n-1}, \ 1 \leq i \leq \frac{r-1}{2} \\ \mathcal{Q}_{(n,r)} \ (n \geq 7, n \text{ odd}, r \text{ odd}, 3 \leq r \leq n-4) : \\ [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq n-3 \\ [X_i, X_{r-i}] = (-1)^{i-1} X_{n-1}, \ 1 \leq i \leq \frac{r-1}{2} \\ [X_i, X_{n-2-i}] = (-1)^{i-1} X_{n-2}, \ 1 \leq i \leq \frac{n-3}{2} \end{aligned}$$

$\mathcal{T}_{(n,n-4)} \ (n \text{ odd}, n \geq 7) :$

$$\begin{aligned} [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq n-3 \\ [X_{n-1}, X_i] = \frac{n-5}{2} X_{n-4+i}, \ 1 \leq i \leq 2 \\ [X_i, X_{n-4-i}] = (-1)^{i-1} (X_{n-4} + X_{n-1}), \\ 1 \leq i \leq \frac{n-5}{2} \\ [X_i, X_{n-3-i}] = (-1)^{i-1} \frac{n-3-2i}{2} X_{n-3}, \\ 1 \leq i \leq \frac{n-5}{2} \\ [X_i, X_{n-2-i}] = (-1)^i (i-1) \frac{n-3-i}{2} X_{n-2}, \\ 2 \leq i \leq \frac{n-3}{2} \end{aligned}$$

$\varepsilon_{(9,5)}^2 :$

$$\begin{aligned} [X_0, X_i] = X_{i+1}, \ 1 \leq i \leq 6 \\ [X_8, X_i] = 2X_{5+i}, \ 1 \leq i \leq 2 \\ [X_1, X_4] = X_5 + X_8 \\ [X_1, X_5] = 2X_6 \\ [X_1, X_6] = X_7 \\ [X_2, X_3] = -X_5 - X_8 \\ [X_2, X_4] = -X_6 \\ [X_2, X_5] = X_7 \\ [X_3, X_4] = -2X_7 \end{aligned}$$

2 Special subalgebras of L and $\text{Der}(L)$

Let L be a Lie algebra over a field F . Clearly, $Z(L) = \bigcap_{\alpha \in \text{IDer}(L)} \text{Ker} \alpha$ and $L^2 = \sum_{\alpha \in \text{IDer}(L)} \text{Im} \alpha$. The following lemma is useful for the proof of our main results.

Lemma 1. *Let I, J be two ideals of Lie algebra L such that $I \subseteq Z(L)$.*

Then

- (i) $\text{Der}_J^I(L) \cong T(L/(L^2 + J), I)$ as vector spaces, and
- (ii) if $I \subseteq J$, then the above isomorphism turns into an isomorphism of Lie algebras, in particular $\text{Der}_J^I(L)$ is abelian.

Proof. For any $\alpha \in \text{Der}_J^I(L)$, the map $\psi_\alpha : L/L^2 + J \rightarrow I$ defined by $\psi_\alpha(x + L^2 + J) = \alpha(x)$ for all $x \in L$ is a linear transformation. It is easy to see that the map $\psi : \text{Der}_J^I(L) \rightarrow T(L/(L^2 + J), I)$ defined by $\psi(\alpha) = \psi_\alpha$ is a one-to-one and onto linear transformation. Moreover, if $I \subseteq J$, then ψ is a Lie isomorphism, from which the result follows. \square

For each Lie algebra L and ideal J of L , let $C_L(J) = \{x \in L \mid [x, y] = 0 \forall y \in J\}$ denote the centralizer of J in L .

Let \mathcal{D} be a subalgebra of $\text{Der}(L)$ containing $\text{IDer}(L)$. Then $E(L) = \bigcap_{\alpha \in \mathcal{D}} \text{Ker} \alpha$ and $U(L) = \sum_{\alpha \in C} \text{Im} \alpha$, where $C = C_{\mathcal{D}}(\text{Der}^{E(L)}(L))$, are ideals of L such that $E(L) \subseteq Z(L)$ and $L^2 \subseteq U(L)$. It is easy to see that

$$C = C_{\mathcal{D}}(\text{Der}^{E(L)}(L)) = \left\{ \alpha \in \mathcal{D} : \beta \circ \alpha = 0, \forall \beta \in \text{Der}^{E(L)}(L) \right\}.$$

In particular, we can let $\mathcal{D} = \text{Der}(L)$ or $\mathcal{D} = \text{Der}^{L^2}(L)$.

Now, if we let $\text{Der}_z(L) = \text{Der}^{Z(L)}(L)$, $\text{Der}_e(L) = \text{Der}^{E(L)}(L)$, then by invoking the previous lemma we obtain the following result.

Corollary 1. *Let L be a Lie algebra. Then*

- (i) $\text{Der}_z(L) \cong T(L/L^2, Z(L))$ as vector spaces and if $Z(L) \subseteq L^2$, then $\text{Der}_z(L)$ is isomorphic to the abelian Lie algebra $T(L/L^2, Z(L))$.
- (ii) $\text{Der}_e(L) \cong T(L/U(L), E(L))$ as vector spaces and if $E(L) \subseteq U(L)$, then $\text{Der}_e(L)$ is isomorphic to the abelian Lie algebra $T(L/U(L), E(L))$.
- (iii) If $\mathcal{D} = \text{Der}(L)$, then $\text{Der}_e(L)$ is isomorphic to the abelian Lie algebras $T(L/(L^2 + E(L)), E(L))$ and $T(L/(U(L) + E(L)), E(L))$.
- (iv) $\text{Der}_{Z(L)}^{Z(L)}(L)$ is isomorphic to the abelian Lie algebra $T(L/(L^2 + Z(L)), Z(L))$.

Proof. In view of the previous lemma it is enough to show that every element of L^2 and $U(L)$ is sent to zero by every element of $\text{Der}_z(L)$ and $\text{Der}_e(L)$, respectively. Also, if $\mathcal{D} = \text{Der}(L)$, then every element of $E(L)$ is sent to zero by every element of $\text{Der}_e(L)$. \square

Corollary 2. *Let L be a finite dimensional Lie algebra. Let I_1, I_2, J_1 and J_2 be ideals of L such that $I_1 \subseteq I_2 \subseteq Z(L)$, $J_2 \subseteq J_1$. Then $\text{Der}_{J_1}^{I_1}(L) \subseteq \text{Der}_{J_2}^{I_2}(L)$. Also $\text{Der}_{J_1}^{I_1}(L) = \text{Der}_{J_2}^{I_2}(L)$ if and only if $I_1 = I_2$, $J_1 = J_2$.*

Proof. It is obvious that if $I_1 \subseteq I_2, J_2 \subseteq J_1$, then $\text{Der}_{J_1}^{I_1}(L) \subseteq \text{Der}_{J_2}^{I_2}(L)$. Now suppose that $\text{Der}_{J_1}^{I_1}(L) = \text{Der}_{J_2}^{I_2}(L)$. Then by Lemma 1,

$$\dim T(L/(L^2 + J_1), I_1) = \dim T(L/(L^2 + J_2), I_2).$$

If $I_1 \subset I_2$ or $J_2 \subset J_1$, then $\dim T(L/(L^2 + J_1), I_1) < \dim T(L/(L^2 + J_2), I_2)$. By this contradiction we have $I_1 = I_2, J_1 = J_2$. The converse is clear. \square

Corollary 3. *Let L be a finite dimensional Lie algebra and I, J be two ideals of L such that $I \subseteq Z(L)$. Then $\text{Der}_J^I(L) = \text{Der}_z(L)$ if and only if $I = Z(L)$ and $J \subseteq L^2$.*

Proof. Since $I \subseteq Z(L)$, $\text{Der}_J^I(L) = \text{Der}_{J+L^2}^I(L)$. Then by Corollary 2, $\text{Der}_{J+L^2}^I(L) = \text{Der}_{L^2}^{Z(L)}(L)$ if and only if $I = Z(L), J + L^2 = L^2$ or, equivalently, $I = Z(L)$ and $J \subseteq L^2$. \square

3 Main theorem

The following lemma is useful for our study of $\text{Der}_J^I(L)$.

Lemma 2. *Let I, J be two ideals of Lie algebra L and $K/I = Z(L/I)$. Then*

$$\text{Der}_J^I(L) \cap \text{IDer}(L) \cong \frac{K \cap C_L(J)}{Z(L)}.$$

In particular, if $\text{Der}_J^I(L) \subseteq \text{IDer}(L)$, then

$$\text{Der}_J^I(L) \cong \frac{K \cap C_L(J)}{Z(L)}.$$

Proof. For each $x \in K \cap C_L(J)$, we have $\text{ad}_x \in \text{Der}_J^I(L) \cap \text{IDer}(L)$. Now, define the map

$$\begin{aligned} \psi : K \cap C_L(J) &\longrightarrow \text{Der}_J^I(L) \cap \text{IDer}(L) \\ x &\longmapsto \text{ad}_x. \end{aligned}$$

Clearly, ψ is a Lie epimorphism such that $\text{Ker}\psi = Z(L)$, as required. \square

Now, we are in the position to state and prove the main result of this section.

Main Theorem. *Let I, J be two ideals of a non-abelian Lie algebra L such that $I \subseteq Z(L)$. If $Z(L/I) = K/I$, then*

(i) $Z(\text{IDer}(L)) \subseteq \text{Der}_J^I(L)$ if and only if $K = Z_2(L) \subseteq C_L(J)$,

(ii) if $I \subseteq J$ and $\dim Z_2(L)/Z(L) < \infty$, then $Z(\text{IDer}(L)) = \text{Der}_J^I(L)$ if and only if $K = Z_2(L) \subseteq C_L(J)$ and $T(L/(L^2 + J), I) \cong Z_2(L)/Z(L)$. In particular, $\text{IDer}(L) = \text{Der}_J^I(L)$ if and only if $J \subseteq Z(L)$, $L^2 \subseteq I$ and $T(L/J, I) \cong L/Z(L)$.

Proof. (i) First suppose that $K = Z_2(L) \subseteq C_L(J)$. It is obvious that for each $x \in L$, we have that $x \in Z_2(L)$ if and only if $\text{ad}_x \in Z(\text{IDer}(L))$. Now, assume that $\text{ad}_x \in Z(\text{IDer}(L))$. Then

$$\text{ad}_x(y) = [x, y] \in [Z_2(L), L] = [K, L] \subseteq I$$

for all $y \in L$. Also, $\text{ad}_x(y) = [x, y] \in [Z_2(L), J] = 0$ for all $y \in J$, which implies that $\text{ad}_x \in \text{Der}_J^I(L)$.

Conversely, suppose that $Z(\text{IDer}(L)) \subseteq \text{Der}_J^I(L)$. Since $[K, L] \subseteq I \subseteq Z(L)$, it follows that $K \subseteq Z_2(L)$. On the other hand, $\text{ad}_x \in \text{Der}_J^I(L)$ for all $x \in Z_2(L)$, which implies that $[Z_2(L), L] \subseteq I$ and hence $Z_2(L) \subseteq K$. Thus $K = Z_2(L)$. Moreover, $[x, y] = 0$ for all $x \in Z_2(L)$ and $y \in J$, from which it follows that $[Z_2(L), J] = 0$, that is, $K = Z_2(L) \subseteq C_L(J)$.

(ii) Suppose $Z(\text{IDer}(L)) = \text{Der}_J^I(L)$. By Lemma 1,

$$T(L/(L^2 + J), I) \cong Z_2(L)/Z(L).$$

Conversely, we have

$$\text{Der}_J^I(L) \cong T(L/(L^2 + J), I) \cong \frac{Z_2(L)}{Z(L)} \cong Z(\text{IDer}(L)).$$

On the other hand, $K = Z_2(L) \subseteq C_L(J)$, hence $Z(\text{IDer}(L)) \subseteq \text{Der}_J^I(L)$. But $\dim Z_2(L)/Z(L) < \infty$ so that the equality holds.

Now, if $\text{IDer}(L) = \text{Der}_J^I(L)$, then $\text{IDer}(L) \subseteq \text{Der}^{Z(L)}(L)$, which implies that L is nilpotent of class 2. Thus $K = Z_2(L) = C_L(J) = L$. Hence $J \subseteq Z(L)$, $L^2 \subseteq I$ and $T(L/J, I) \cong L/Z(L)$. Conversely, if $J \subseteq Z(L)$ and $L^2 \subseteq I$, then $\text{IDer}(L) \subseteq \text{Der}_J^I(L)$ and $T(L/J, I) \cong L/Z(L)$, from which the equality holds. The proof is complete. \square

Corollary 4. *Let I, J be two ideals of a finitely generated non-abelian Lie algebra L with $Z(L) \neq 0$ such that $I \subseteq Z(L) \subseteq J$. Then $\text{Der}_J^I(L) = \text{IDer}(L)$ if and only if L is a finite dimensional nilpotent Lie algebra of class 2, $J = Z(L)$, $L^2 \subseteq I$ and $\dim I = 1$*

Proof. First we prove that if $L^2 \subseteq I$, then $T(L/Z(L), I) \cong L/Z(L)$ if and only if $\dim I = 1$. Since L is finitely generated and $L^2 \subseteq I \subseteq Z(L)$, by Lemma 1(i) in [3], L is a finite dimensional Lie algebra. Then it is obvious that $T(L/Z(L), I) \cong L/Z(L)$ if and only if $\dim I = 1$. Now by main theorem(ii) $\text{Der}_J^I(L) = \text{IDer}(L)$ if and only if $J \subseteq Z(L), L^2 \subseteq I$ and $T(L/J, I) \cong L/Z(L)$ or equivalently $J = Z(L), L^2 \subseteq I$ and $\dim I = 1$ \square

If we put $I = J = Z(L)$ in the above corollary, then we obtain the following result immediately.

Corollary 5. *Let L be a finitely generated Lie algebra with $Z(L) \neq 0$. Then $\text{Der}_{Z(L)}^{Z(L)}(L) = \text{IDer}(L)$ if and only if L is finite dimensional abelian, or L is finite dimensional nilpotent Lie algebra of class 2 and $\dim Z(L) = 1$.*

The following corollary is the result of Tôgô [21, Theorem 3] for Lie algebras of finite dimension.

Corollary 6. *Let L be a finite dimensional Lie algebra with $Z(L) \neq 0$. Then $\text{Der}_z(L) = \text{IDer}(L)$ if and only if $L^2 = Z(L)$ and $\dim Z(L) = 1$, in which case L is a Heisenberg algebra.*

Proof. Let $L^2 = Z(L)$, $\dim Z(L) = 1$. By putting $I = Z(L)$ and $J = L^2$ in the Corollary 4, we have $\text{Der}_z(L) = \text{IDer}(L)$.

Conversely suppose that $\text{Der}_z(L) = \text{IDer}(L)$. First we show that $L^2 = Z(L)$. If $\text{Der}_z(L) = \text{IDer}(L)$, then $L^2 \subseteq Z(L)$ and $\psi : \text{Der}_z(L) \rightarrow T(L/Z(L), L^2)$ defined by $\psi(\alpha) = \psi_\alpha$, where $\psi_\alpha(x + Z(L)) = \alpha(x)$ is a Lie isomorphism. Suppose on the contrary that $L^2 \subsetneq Z(L)$. Then $\dim \frac{L}{Z(L)} < \dim \frac{L}{L^2}$ and $\dim L^2 < \dim Z(L)$. Thus $\dim T(L/Z(L), L^2) < \dim T(L/L^2, Z(L))$, which is a contradiction. Now by Corollary 4, $\dim Z(L) = 1$. \square

4 Characterization

It is easy to see that a central derivation of a Lie algebra L commutes with every inner derivation. Also, for any Lie algebra L , the center of the Lie algebra of inner derivations is always contained in the set of central derivations of L , because $\text{Der}_z(L) = C_{\text{Der}(L)}(\text{IDer}(L))$. In this section we consider finite dimensional Lie algebras for which this lower bound is attained, that is $\text{Der}_z(L) = Z(\text{IDer}(L))$, and give a characterisation of such Lie algebras.

Theorem 2. *Let L be a finite dimensional Lie algebra with $Z(L) \neq 0$. If $\text{Der}_z(L) = Z(\text{IDer}(L))$, then $Z(L) \subseteq L^2$. Also $\text{Der}_z(L) = Z(\text{IDer}(L))$ if and only if $T(L/L^2, Z(L)) \cong Z_2(L)/Z(L)$.*

Proof. Let $\text{Der}_z(L) = Z(\text{IDer}(L))$. Then $\text{Der}_z(L) = \text{Der}_{Z(L)}^{Z(L)}(L)$ and $\text{Der}_{Z(L)}^{Z(L)}(L) \cong T(L/(L^2 + Z(L)), Z(L))$, by Corollary 1(iv). Therefore we have

$$\text{Der}_{Z(L)}^{Z(L)}(L) \cong T(L/(L^2 + Z(L)), L^2 \cap Z(L)),$$

because $\psi_\alpha(x + L^2 + Z(L)) \in L^2 \cap Z(L)$ for all $x \in L$. Thus $\dim(L^2 \cap Z(L)) = \dim Z(L)$. This implies that $Z(L) \subseteq L^2$.

Now, if $\text{Der}_z(L) = Z(\text{IDer}(L))$, by putting $I = Z(L)$ and $J = L^2$ in the main theorem, $T(L/L^2, Z(L)) \cong \frac{Z_2(L)}{Z(L)}$.

Conversely, if $T(L/L^2, Z(L)) \cong \frac{Z_2(L)}{Z(L)}$, then since for any Lie algebra L , $Z(\text{IDer}(L)) \subseteq \text{Der}_z(L)$ and $\dim \text{Der}_z(L) = \dim T(L/L^2, Z(L))$, we conclude that $\text{Der}_z(L) = Z(\text{IDer}(L))$. \square

Clearly, for any nontrivial abelian Lie algebra L , $Z(\text{IDer}(L)) \subsetneq \text{Der}_z(L)$.

Corollary 7. *Let L be a finite dimensional nilpotent Lie algebra of class 2. Then $\text{Der}_z(L) = Z(\text{IDer}(L))$ if and only if $L^2 = Z(L)$ and $\dim Z(L) = 1$, in which case L is a Heisenberg algebra.*

Proof. Let L be a finite dimensional nilpotent Lie algebra of class 2. Then $\text{IDer}(L)$ is an abelian Lie algebra. Now, the result follows from Corollary 6. \square

Corollary 8. *Let L be a nilpotent Lie algebra of finite dimension such that $\dim(Z_2(L)/Z(L)) = 1$. Then $Z(\text{IDer}(L)) \subsetneq \text{Der}_z(L)$. In particular, this holds if L is a filiform Lie algebra.*

Proof. It is well known that $\dim(L/L^2) \geq 2$ and $\dim Z(L) \geq 1$ for a nilpotent Lie algebra. Thus $\dim T(L/L^2, Z(L)) \geq 2$, so can not be isomorphic to $Z_2(L)/Z(L)$. Hence the result follows from Theorem 2. \square

Corollary 9. *Let L be a naturally graded quasi-filiform Lie algebra of dimension n over the field of complex numbers \mathbb{C} . Then $\text{Der}_z(L) = Z(\text{IDer}(L))$ if and only if $L = \mathcal{T}_{(n, n-3)}$ (n even, $n \geq 6$)*

Proof. Let L be an n -dimensional naturally graded quasi-filiform Lie algebra and $\text{Der}_z(L) = Z(\text{IDer}(L))$. It is well known that $1 \leq \dim(Z_2(L)/Z(L)) \leq 2$ for any quasi-filiform Lie algebra. Therefore by Theorem 2, $\dim Z(\text{IDer}(L)) = 2$ and hence $\dim(L/L^2) = 2$ and $\dim Z(L) = 1$.

If n is even, then by Theorem 1 L is isomorphic to one of the algebras $L_{n-1} \oplus \mathbb{C}$, $\mathcal{T}_{(n, n-3)}$, or $\mathcal{L}_{(n, r)}$, with r odd and $3 \leq r \leq n-3$. If $L = L_{n-1} \oplus \mathbb{C}$, then $\dim(L/L^2) = 3$, $\dim Z(L) = 2$. If $L = \mathcal{L}_{(n, r)}$, then $\dim(L/L^2) = \dim Z(L) = 2$.

Therefore equality does not hold. If $L = \mathcal{T}_{(n,n-3)}$, then $\dim(L/L^2) = 2$ and $\dim Z(L) = 1$. Since $\gamma_{n-3}(L) \subseteq Z_2(L)$ and $\dim \gamma_{n-3}(L) = 3$, $\dim Z_2(L) = 3$. Therefore $\dim Z(\text{IDer}(L)) = 2$. This implies that $\text{Der}_z(L) = Z(\text{IDer}(L))$.

If n is odd, then by Theorem 1 L is isomorphic to one of the algebras $L_{n-1} \oplus \mathbb{C}$, $Q_{n-1} \oplus \mathbb{C}$, $\mathcal{L}_{(n,n-2)}$, $\mathcal{T}_{(n,n-4)}$, $\mathcal{L}_{(n,r)}$, or $\mathcal{Q}_{(n,r)}$, with r odd and $3 \leq r \leq n-4$. If $L = L_{n-1} \oplus \mathbb{C}$ or $L = Q_{n-1} \oplus \mathbb{C}$, then $\dim(L/L^2) = 3$ and $\dim Z(L) = 2$. If $L = \mathcal{L}_{(n,n-2)}$, $L = \mathcal{L}_{(n,r)}$ or $L = \mathcal{Q}_{(n,r)}$, then $\dim(L/L^2) = \dim Z(L) = 2$. If $L = \mathcal{T}_{(n,n-4)}$, then it is easy to see that $Z(\text{IDer}(L)) = \langle ad_{X_{n-3}} \rangle$ and hence $\dim Z(\text{IDer}(L)) = 1$. Therefore for these Lie algebras equality does not hold.

In the case of $n = 7$ and $n = 9$, we have algebras $\varepsilon_{(7,3)}$, $\varepsilon_{(9,5)}^1$ and $\varepsilon_{(9,5)}^2$. For these algebras, $\dim Z(\text{IDer}(L)) = 1$ and so $Z(\text{IDer}(L)) \subsetneq \text{Der}_z(L)$. \square

References

- [1] J. DIXMIER, W. G. LISTER: *Derivations of nilpotent Lie algebras*, Proc. Amer. Math. Soc., **8**, (1957), 155–158.
- [2] J.R. GÓMEZ, A. JIMÉNEZ-MERCHÁN: *Naturally graded quasi-filiform Lie algebras*, J. Algebra **256**, (2002), 211–228.
- [3] B. Hartley, *Locally nilpotent ideals of a Lie algebra*. Proc. Cambridge Philos. Soc. **63** (1967), 257–272.
- [4] G. HOCHSCHILD: *Semisimple algebras and generalized derivations*, Amer. J. Math., **64**, (1942), 677–694.
- [5] N. JACOBSON: *Lie algebras*, Wiley Interscience, New York, 1962.
- [6] G. LEGER: *A note on the derivations of Lie algebras*, Proc. Amer. Math. Soc., **4**, (1953), 511–514.
- [7] G. LEGER: *Derivations of Lie algebras III*, Duke Math. J., **30**, (1963), 637–645.
- [8] G. LEGER, S. TÔGÔ: *Characteristically nilpotent Lie algebras*, Duke Math. J., **26**, (1959), 523–528.
- [9] D. J. MENG: *On complete Lie algebras*, Act. Sci. Mat. Univ. Mankai, **2**, (1985), 9–19. (Chinese)
- [10] D. J. MENG: *The uniqueness of the decomposition of the complete Lie algebras*, Ibid, **3**, (1990), 23–26. (Chinese)
- [11] D. J. MENG: *The complete Lie algebras with abelian nilpotent radical*, Act. Math., **34**, (1991), 191–202.
- [12] D. J. MENG: *Complete Lie algebras and Heisenberg algebras*, Comm. Algebra, **22**, (1994), 5509–5524.
- [13] D. J. MENG: *Some results on complete Lie algebras*, Comm. Algebra, **22**, (1994), 5457–5507.
- [14] D. J. MENG, S. P. WANG: *On the construction of complete Lie algebras*: J. Algebra, **176**, (1995), 621–637.
- [15] D. J. MENG, L. S. ZHU: *Solvable complete Lie algebras. I*, Comm. Algebra, **24**, n. 13, (1996), 4181–4197.

- [16] E. SCHENKMAN: *On the derivation algebra and the holomorph of a nilpotent Lie algebra*, Mem. Amer. Math. Soc., **14**, (1955), 15–22.
- [17] E. L. STITZINGER: *On Lie algebras with only inner derivations*, J. Algebra, **105**, (1987), 341–343.
- [18] Y. SU, L. ZHU: *Derivation algebras of centerless perfect Lie algebras are complete*, J. Algebra, **285**, (2005), 508–515.
- [19] S. TÔGÔ: *On the derivations of Lie algebras*, J. Sci. Hiroshima Univ. Ser. A, **19**, (1955), 71–77.
- [20] S. TÔGÔ: *On the derivation algebras of Lie algebras*, Canad. J. Math., **13**, (1961), 201–216.
- [21] S. TÔGÔ: *Derivations of Lie algebras*, J. Sci. Hiroshima Univ. Ser. A-I, **28**, (1964), 133–158.
- [22] S. TÔGÔ, *Outer derivations of Lie algebras*, Trans. Amer. Math. Soc., **128**, (1967), 264–276.

