

Determinants of quality of life in survivors of cardiac arrest

Citation for published version (APA):

Moulaert, V. R. M. P., Wachelder, E. M., Verbunt, J. A., Wade, D. T., & van Heugten, C. M. (2010). Determinants of quality of life in survivors of cardiac arrest. *Journal of Rehabilitation Medicine*, 42(6), 553-558. <https://doi.org/10.2340/16501977-0547>

Document status and date:

Published: 01/06/2010

DOI:

[10.2340/16501977-0547](https://doi.org/10.2340/16501977-0547)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

ORIGINAL REPORT

DETERMINANTS OF QUALITY OF LIFE IN SURVIVORS OF CARDIAC ARREST

Véronique R. M. P. Moolaert, MSc^{*1}, Esther M. Wachelder, MSc¹, Jeanine A. Verbunt, PhD^{1,2}, Derick T. Wade MD^{2,3} and Caroline M. van Heugten, PhD^{4,5}

From the ¹Adelante, Centre of Expertise in Rehabilitation and Audiology, Hoensbroek, ²Care and Public Health Research Institute, Maastricht University, Maastricht, The Netherlands, ³Oxford Centre for Enablement, Oxford, UK, ⁴School for Mental Health and Neuroscience and ⁵Department of Neuropsychology and Psychopharmacology, Maastricht University, Maastricht, The Netherlands

Objective: To study factors related to quality of life after a hypoxic period due to cardiac arrest.

Design: Retrospective cohort study.

Subjects: Eighty-eight survivors of out-of-hospital cardiac arrest, admitted to a Dutch academic hospital between 2001 and 2006.

Methods: Patients received a set of questionnaires at home. The main outcome measures were physical and mental quality of life (Medical Outcomes Study 36-item Short Form Health Survey; SF-36). Potential determinants were cognitive complaints, emotional problems (depression/anxiety), post-traumatic stress, fatigue, daily functioning and participation in society. Multiple linear regression analyses were performed with physical and mental quality of life as dependent variables.

Results: Sixty-three (72%) patients responded. Mean time since cardiac arrest was 36 months (standard deviation (SD) 19). Backward regression analyses showed that physical quality of life was significantly ($p < 0.001$, adjusted $R^2 = 0.531$) related to cognitive complaints ($\beta = -0.378$), instrumental daily life activities ($\beta = 0.262$), post-traumatic stress ($\beta = -0.246$) and fatigue ($\beta = -0.226$). Mental quality of life was significantly ($p < 0.001$, adjusted $R^2 = 0.664$) explained by anxiety/depression ($\beta = -0.609$), fatigue ($\beta = -0.177$) and cognitive complaints ($\beta = -0.175$).

Conclusion: Quality of life is related to cognitive complaints, fatigue, anxiety/depression, post-traumatic stress and difficulties in daily activities in survivors of out-of-hospital cardiac arrest. Rehabilitation programmes for this group should specifically address these topics.

Key words: heart arrest; quality of life; cognition disorders; hypoxia-ischaemia brain.

J Rehabil Med 2010; 42: 553–558

Correspondence address: Caroline van Heugten, Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, PO Box 616, NL-6200 MD, Maastricht, The Netherlands. E-mail: c.vanheugten@np.unimaas.nl

Submitted September 16, 2009; accepted January 28, 2010

INTRODUCTION

A person's quality of life (QoL) should be a major outcome measure in studies of survivors of cardiac arrest (1), since

reporting survival rates alone does not provide complete information about the success of resuscitation. Several prospective studies have reported that the long-term QoL of cardiac arrest survivors is generally acceptable to good (2–4). However, a substantial group of cardiac arrest survivors experience ongoing problems and a decreased QoL (5). Medical variables cannot reliably predict QoL after cardiac arrest. Although the performance of bystander cardiopulmonary resuscitation (CPR) seems to be associated with a better QoL (4), longer time delays during resuscitation and the type of emergency cardiac care were not related to post-cardiac arrest QoL (6–8). In order to help those survivors with long-term problems, it is important to investigate which factors influence their QoL, as this may identify potential targets for rehabilitation treatment.

Survivors of cardiac arrest are often categorized as “cardiac patients”, although they may also be regarded as “neurological patients” because a cardiac arrest can result in hypoxic brain injury. Indeed, approximately half of cardiac arrest survivors have enduring cognitive impairments, as was shown in a recent systematic review (9). The long-term consequences of hypoxic brain injury secondary to cardiac arrest may therefore be comparable to the long-term consequences of other types of brain damage, such as traumatic brain injury.

Studies on patients with varying other causes of brain injury have revealed that reduced QoL was associated with 3 important variables, namely cognitive impairments, emotional problems and fatigue. Cognitive impairments were related to the QoL of patients after stroke (10). The importance of emotional problems was shown in a study on patients with mild traumatic brain injury in which life satisfaction was related to post-traumatic stress and depression (11). Fatigue was associated with QoL in both stroke patients (12) and traumatic brain-injured patients (13). Similarly, multivariate regression analyses in a cohort of survivors of subarachnoid haemorrhage showed that mood, fatigue, and cognitive complaints were strongly associated with reduced scores on a stroke-specific QoL scale (14).

There are few studies that have investigated possible cognitive or psychosocial factors affecting QoL after cardiac arrest. One study showed that memory complaints were related to QoL in a study on survivors of out-of-hospital cardiac arrest (15). Another study identified that post-traumatic stress was common after cardiac arrest and was negatively related to QoL (16). In

addition, anxiety and depression were also reported frequently after cardiac arrest and affected perceived health improvement and social functioning (8). However, the amount of information on factors affecting QoL after cardiac arrest is limited and an integrative multivariate model is currently lacking.

The aim of this study was therefore to investigate potential determinants of QoL after a hypoxic period due to cardiac arrest. It was hypothesized that QoL would be affected by the same factors that influence QoL of other brain-injured patients, namely cognitive complaints, emotional problems and fatigue.

METHODS

Survivors of out-of-hospital cardiac arrest who had been admitted to the Department of Cardiology of Maastricht University Medical Centre (MUMC+), the Netherlands, between January 2001 and December 2006 were invited to participate in this retrospective cohort study. MUMC+ is an academic hospital in the southern part of the Netherlands and is the primary referral centre for a radius of 25 km, with a catchment area of approximately 200,000 inhabitants, which is served by a single ambulance service. The incidence of cardiac arrest in this area is 9.2 per 10,000 for people between 20 and 75 years old (17). Survival rates until discharge are, in general, low (18), being 7.4% in this specific region (19).

For inclusion in this study, participants had to be survivors aged 18 years or older, still alive at the time of the study, and with sufficient knowledge of the Dutch language to complete the questionnaires.

Patient data for potential participants for this study were retrieved from 2 hospital databases containing all out-of-hospital cardiac arrests in Maastricht and the surrounding area (17, 20). Furthermore, the admission registers from 2001 to 2006 of the Coronary Care Unit were hand-searched for admissions stating "cardiac arrest" or "resuscitation". After retrieving potential participants, data were checked for accuracy of the diagnosis and for survival until discharge. Next, the general practitioners of all potential participants were contacted to confirm whether the subjects were currently still alive. People thought to be alive received information about the study by post, together with the informed consent form and the set of questionnaires. The study protocol was approved by the medical ethics committee of MUMC+.

Other results about the respondents and their caregivers concerning their long-term functioning have been published elsewhere (5).

Assessment instruments

Socio-demographic and medical variables. The following socio-demographic variables were extracted from the notes: age, gender, marital status, living situation, educational and work situation. The medical variables extracted were: age at time of cardiac arrest, time since cardiac arrest, location of cardiac arrest, presence of witnesses, performance of bystander cardiopulmonary resuscitation, initial cardiac rhythm, number of defibrillations, interval from collapse to return of spontaneous circulation, (sub)acute interventions, left ventricular ejection fraction and cardiovascular history.

Quality of life. The Medical Outcomes Study 36-item Short Form Health Survey (SF-36), a 36-item generic QoL measure distinguishing 2 domains, namely physical health and mental health. The scores on the 2 domains range from 0 to 100. Higher scores represent a better experienced QoL. Norm values for the Dutch population are a mean (standard deviation (SD)) of 76.3 (SD 26) for the physical component and 78.0 (SD 23) for mental component (21).

Participation in society. The Community Integration Questionnaire (CIQ), a 15-item questionnaire that assesses the level of social participation in society in 3 domains: Home Integration, Social Integration and Productive Activity (22).

Basic activities of daily living. The Barthel Index (BI), a widely used instrument that measures basic (personal) activities of daily living (ADL). The instrument consists of 10 items and measures to what extent a person can perform basic ADL activities independently. High scores indicate a high level of independence (23).

Instrumental activities of daily living. The Frenchay Activities Index (FAI), an accepted measure for instrumental (extended) activities of daily living, that reports on the performance of 15 daily life activities (24).

Cognitive complaints. The Cognitive Failures Questionnaire (CFQ), a 25-item questionnaire on self-reported cognitive failures. A higher score represents the experience of more cognitive complaints (25).

Emotional problems. The Hospital Anxiety and Depression Scale (HADS), a 14-item questionnaire, designed to detect the presence of mild degrees of mood disorders in non-psychiatric hospital outpatients (26). The HADS consists of 2 sub-scales: depression and anxiety. Higher scores represent more emotional problems.

Post-traumatic stress. The Impact of Event Scale (IES), which measures psychological reactions that take place after a traumatic event. The IES consists of 15 items and explores the level of post-traumatic stress by asking for intrusive and avoidance symptoms (27, 28).

Fatigue. The Fatigue Severity Scale (FSS), a 9-item questionnaire that measures the impact of fatigue symptoms in daily activities, with a higher score indicating a more negative impact (29).

Statistical analysis

Descriptive statistics were used to present the demographic, medical and resuscitation characteristics of the study population. To investigate possible differences between the responders and the non-responders, the *t*-test and χ^2 test were used for age and gender, respectively.

Assessment instruments in which more than 15% of the items were missing, were not used in our analyses. When there were less than 15% missing items, the missing data were imputed by dividing the sum of the scale by the number of items answered and then multiplied by the total number of items of that measurement instrument.

Because of the non-parametric distribution of most of the variables, Spearman's rank correlations coefficients (2-tailed) were calculated to investigate relationships between the physical and mental QoL and the potential determinants. A *p*-value < 0.01 was considered statistically significant.

The influence of the potential determinants on QoL was determined with multiple linear regression analyses (backward method) for the physical and mental domains separately. The independent variables that were introduced were the variables with a significant correlation with QoL. A limit value for elimination in the regression analyses was set at *p* < 0.10. For the regression models a *p*-value of < 0.05 was considered statistically significant. The regression models were checked for independence of errors and absence of co-linearity and outliers.

Statistical analyses were performed using the software package SPSS version 15.0 and 16.0 (SPSS Inc., Chicago, IL, USA).

RESULTS

From the databases searched, 220 persons with a possible out-of-hospital cardiac arrest were identified. After excluding those who did not fulfil the inclusion criteria, or who had died, or who were lost to follow-up, 88 persons remained who were eligible for this study. This group received the set of questionnaires at home. Sixty-three persons responded by returning the questionnaires: a response rate of 72% (Fig. 1 for flow chart of patient inclusion).




Fig. 1. Patient inclusion flowchart.

There were no significant differences in gender ($t=-0.62$, $p=0.54$) or age ($\chi^2=0.04$, $p=0.84$) between the responders and the non-responders.

The group of responders consisted of 54 men (86%) and 9 women. Their mean age was 60.0 (SD 12.7) years. The mean

Table I. Demographic characteristics of the participants (n = 63)

	At time of cardiac arrest	Current situation
Age, years		
Mean (SD)	57.3 (12.5)	60.2 (12.7)
Range	18–81	20–85
Gender, n %		
Male		54 (86)
Female		9 (14)
Marital status, n %		
Married/living with partner	47 (75)	46 (73)
Unmarried/divorced/widowed	15 (24)	15 (24)
Unknown	1 (2)	2 (3)
Living situation, n %		
Home (independent)	54 (86)	53 (84)
Home (with help from others)	8 (13)	8 (13)
Sheltered housing	1 (2)	1 (2)
Nursing home	–	1 (2)
Education, n %		
Primary school		7 (11)
Secondary school		12 (19)
High school		30 (48)
University		6 (10)
Other/unknown		8 (13)
Working situation, n %		
Full-time/self-owned company	30 (48)	17 (27)
Part-time	5 (8)	10 (16)
Permanently unfit to work	5 (8)	6 (10)
Volunteer work	–	1 (2)
No job	–	1 (2)
Retired	17 (27)	22 (35)
Housewife	6 (10)	6 (10)

SD: standard deviation

time since cardiac arrest was 36.0 (SD 18.8) months. The demographic characteristics of the patients are presented in Table I. For the marital status, living situation and work situation both the results at the time of cardiac arrest and their current situation are shown. Table II shows information on the cardiac arrest, resuscitation, medical interventions and patient history.

Table III shows the scores on all the questionnaires. All outcome measures were not normally distributed, except the FSS and CIQ. When more than 15% of the items were missing, this person was not taken into account in the analyses; this was the case for 2 patients on the SF-36 per subdomain, CIQ, FAI, CFQ and FSS. In case of less than 15% missing values, data were imputed; this varied from one patient (1.6%) on the HADS to 8 patients (12.7%) on the CFQ. For 6 patients this was only one item, for one patient 2 items and for one patient 3 items on the CFQ.

The mean/median score (range) on the SF-36 in our population was 71.8/77.4 (28–100) on the SF-36 physical domain and 73.0/79.5 (26–100) on the SF-36 mental domain. The scores in our sample are somewhat lower than in the Dutch norm population (21).

Table II. Medical data of the participants

	n	
Age at time of cardiac arrest, mean (SD)	63	57.3 (12.5)
Time since cardiac arrest (months), mean (SD)	63	60.0 (18.8)
Location of cardiac arrest, n (%)	63	
Street/public area		30 (48)
Home		20 (32)
In the ambulance		4 (6)
Other		9 (14)
Witnessed cardiac arrest, n (%)	63	56 (89)
Bystander CPR, n (%)	63	
Yes, by lay person		26 (41)
Yes, by trained person		23 (37)
No		14 (22)
Initial cardiac rhythm, n (%)	62	
Ventricular Fibrillation		55 (89)
Ventricular tachycardia/Ventricular Fibrillation		6 (10)
Other rhythm		1 (2)
Number of defibrillations, median (range)	63	2 (0–12)
Interval collapse – ROSC (min), median (range)	47	10 (2–35)
(Sub)acute interventions, n (%)	63	
none/only catheterization		24 (38)
Catheterization with PCI		28 (44)
Coronary artery bypass graft		13 (21)
Implantable cardioverter defibrillator		25 (46)
Therapeutic hypothermia		17 (27)
LVEF <1 week after cardiac arrest, mean (SD)	58	47.5 (12.7)
Cardiovascular history, n (%)	63	
Myocardial infarction		19 (30)
Cardiac arrest		2 (3)
Cardiac interventions (PCI, CABG)		8 (13)
Claudicatio intermittens		4 (6)
Cerebrovascular accident		2 (3)
Diabetes mellitus		3 (5)

CPR: cardiopulmonary resuscitation; LVEF: left ventricular ejection fraction; ROSC: return of spontaneous circulation; PCI: percutaneous coronary intervention; SD: standard deviation.

Table III. Quality of life, participation, basic and instrumental activities of daily living, cognitive complaints, emotional problems, post-traumatic stress and fatigue of the participants

Domain	Measurement instrument	n	Cut-off	Mean (SD)	Median (range)
<i>Quality of life</i>					
	SF-36 physical	58	–	71.8 (19.6)	77.4 (28–100)
	SF-36 mental	55	–	73.0 (20.5)	79.5 (26–100)
<i>Participation in society</i>					
	Community Integration Questionnaire	61	–	14.7 (5.5)	15.0 (4–26)
<i>Basic activities of daily living</i>					
	Barthel Index	63	<15	19.4 (2.1)	20.0 (7–20)
<i>Instrumental activities of daily living</i>					
	Frenchay Activities Index	61	–	25.7 (8.7)	27.0 (6–41)
<i>Cognitive complaints</i>					
	Cognitive Failures Questionnaire	61	>43	29.8 (18.4)	30.0 (0–73)
<i>Emotional problems</i>					
	Hospital Anxiety and Depression Scale, total	63	>10	8.5 (6.4)	7.0 (0–25)
	– subscale anxiety	63	>8	4.4 (3.4)	4.0 (0–14)
	– subscale depression	63	>8	4.2 (3.6)	3.0 (0–12)
<i>Post-traumatic stress</i>					
	Impact of Event Scale	63	>25	13.0 (13.7)	8 (0–53)
<i>Fatigue</i>					
	Fatigue Severity Scale	61	>3	4.3 (1.6)	4.3 (1–7)

SD: standard deviation.

Table IV shows the Spearman’s rank correlations between the 2 SF-36 domains and the other variables. No significant correlations were found between the SF-36 domains and the CIQ and the BI.

Fig. 2 presents the results of the backward regression analyses. The variables with significant correlations in Table IV were introduced as independent variables.

For the SF-36 physical domain the model could significantly explain 53% of the variance ($p < 0.001$, adjusted $R^2 = 0.531$). Four variables contributed to the model: in order of impact, the CFQ ($\beta = -0.378$, $p = 0.001$), FAI ($\beta = 0.262$, $p = 0.013$), IES ($\beta = -0.246$, $p = 0.017$) and the FSS ($\beta = -0.226$, $p = 0.057$). The final model for the SF-36 mental domain accounted for 66%

Table IV. Spearman’s correlations between quality of life and the potential determinants

	SF-36 Physical	SF-36 Mental
CIQ	0.29	0.26
BI	0.22	0.10
FAI	0.44†	0.41*
FSS	-0.61†	-0.57†
CFQ	-0.55†	-0.40*
HADS	-0.48†	-0.73†
IES	-0.37*	-0.43†

* $p \leq 0.01$; † $p \leq 0.001$.

CIQ: Community Integration Questionnaire; BI: Barthel Index; FAI: Frenchay Activities Index; FSS: Fatigue Severity Scale; CFQ: Cognitive Failures Questionnaire; HADS: Hospital Anxiety and Depression Scale; IES: Impact of Event Scale; SF-36: Short Form 36.

of the variance ($p < 0.001$, adjusted $R^2 = 0.664$). In this model, mental QoL was related to the HADS ($\beta = -0.609$, $p < 0.001$), the FSS ($\beta = -0.177$, $p = 0.069$) and the CFQ ($\beta = -0.175$, $p = 0.081$).

DISCUSSION

This retrospective cohort study has shown that QoL can be at risk after out-of-hospital cardiac arrest. The QoL scores were slightly below that of the general Dutch population (21). This is in line with previous studies on QoL after cardiac arrest (6, 30). Mental QoL could be affected more than physical QoL, probably because significant motor impairments after cardiac arrest are rare. In addition, both cognitive and emotional functioning contribute to mental QoL. Both are known to be at risk after hypoxic brain injury (5, 31, 32).

The correlation matrix showed that QoL significantly correlated with instrumental daily activities, severity of fatigue, amount of cognitive complaints, level of mood disturbance (anxiety/depression) and degree of post-traumatic stress. In contrast, QoL was not related to participation in society and basic (personal) ADL functioning. That social participation was not related to QoL is surprising. However, this might be explained by the fact that the majority of the participants were




Fig. 2. Regression analyses: determinants of physical and mental quality of life.

no longer working and were over 60 years old, implying that their participation may already have been limited prior to the cardiac arrest. Basic ADL activities were hardly impaired and showed a clear ceiling effect, which probably explains the lack of any detectable influence.

The regression analyses revealed that physical QoL was significantly related to cognitive complaints, performance of instrumental daily activities, post-traumatic stress and fatigue. Mental QoL was primarily affected by anxiety/depression and, to a lesser extent, to fatigue and cognitive complaints. Both models could explain a substantial amount of the variance, 53% and 66%, respectively. Instrumental daily activities contributed only to physical QoL, probably because a minimal level of physical fitness is necessary to perform these daily activities. Post-traumatic stress also contributed to physical QoL only. In contrast, in a group of subarachnoid haemorrhage patients, comparable in the sense that they also have brain injury without clear physical impairments, post-traumatic stress disorders strongly predicted both mental and physical QoL after 13 months (33). Anxiety and depressive symptoms were strongly related to mental QoL alone, as would be expected since both are measures of mood disturbance. Cognitive complaints and fatigue were related to both physical and mental health. Fatigue is clearly a concept with both a mental and a physical aspect. That cognitive problems also affect physical health is an interesting finding, and possibly arises because cognitive impairments, for example in memory and planning, reduce the performance of complex physical activities.

There are some limitations to this study. First, the design was retrospective, which may have led to a selection bias and makes it difficult to draw causal relationships between QoL and the potential determinants. Secondly, not all potential participants actually participated. Although the response was relatively high, this may have led to selection bias. It is possible that some people were in such a poor condition that completing the questionnaires was not possible for them. On the other hand, it might also be that those without any complaints did not want to participate as they felt that there was "nothing wrong", or due to the fact that they were too busy with their daily life. Looking at the incidence rates in the region and the number of patients retrieved from the hospital files, the sample can however be regarded as highly representative. Thirdly, time since cardiac arrest varied between 1 and 6 years and therefore participants may have been in different phases of recovery. Fourthly, we used self-report. The self-reporting of cognitive complaints has to be interpreted with caution, especially in cases of severe hypoxic brain injury. However, self-report of mood and activities are probably valid. To overcome the weaknesses mentioned above, a larger, prospective study would be necessary, in which cognitive impairments should be objectified with neuropsychological tests. Such a study is currently being conducted (34).

In conclusion, this study has shown that QoL after cardiac arrest seems to be affected by multiple factors including cognitive complaints, fatigue, emotional problems, post-traumatic stress and difficulties in daily activities. A similar pattern has been shown in patients after stroke (10, 12, 14) and traumatic brain injury (11, 13). This confirms our hypothesis that a survivor

of a cardiac arrest should not only be regarded as a "cardiac patient", but also as a "brain-injured patient". Furthermore, our finding that QoL is related to a diversity of factors, suggests that multidisciplinary rehabilitation programmes focussing on these factors, should be developed and evaluated.

ACKNOWLEDGEMENTS

We are grateful to B. Eikemans, J. de Munter, M. de Munter and J. Partouns for their help in identifying potential participants and collecting medical data. S. Bekkers and A. Gorgels are thanked for facilitating this study.

The study is funded by the Netherlands Organization for Health Research and Development (ZON/MW) and Fonds Nuts/Ohra.

REFERENCES

1. Neumar RW, Nolan JP, Adrie C, Aibiki M, Berg RA, Bottiger BW, et al. Post-cardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A consensus statement from the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation of Canada, Inter American Heart Foundation, Resuscitation Council of Asia, and the Resuscitation Council of Southern Africa); the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Peri-operative, and Critical Care; the Council on Clinical Cardiology; and the Stroke Council. *Circulation* 2008; 118: 2452–2483.
2. Bunch TJ, White RD, Gersh BJ, Meverden RA, Hodge DO, Ballman KV, et al. Long-term outcomes of out-of-hospital cardiac arrest after successful early defibrillation. *N Engl J Med* 2003; 348: 2626–2633.
3. Lundgren-Nilsson A, Rosen H, Hofgren C, Sunnerhagen KS. The first year after successful cardiac resuscitation: function, activity, participation and quality of life. *Resuscitation* 2005; 66: 285–289.
4. Stiell I, Nichol G, Wells G, De Maio V, Nesbitt L, Blackburn J, et al. Health-related quality of life is better for cardiac arrest survivors who received citizen cardiopulmonary resuscitation. *Circulation* 2003; 108: 1939–1944.
5. Wachelder EM, Moulaert VR, van Heugten C, Verbunt JA, Bekkers SC, Wade DT. Life after survival: long-term daily functioning and quality of life after an out-of-hospital cardiac arrest. *Resuscitation* 2009; 80: 517–522.
6. van Alem AP, Waalewijn RA, Koster RW, de Vos R. Assessment of quality of life and cognitive function after out-of-hospital cardiac arrest with successful resuscitation. *Am J Cardiol* 2004; 93: 131–135.
7. Eisenburger P, List M, Schorkhuber W, Walker R, Sterz F, Laggner AN. Long-term cardiac arrest survivors of the Vienna emergency medical service. *Resuscitation* 1998; 38: 137–143.
8. Kamphuis HC, De Leeuw JR, Derksen R, Hauer R, Winnubst JA. A 12-month quality of life assessment of cardiac arrest survivors treated with or without an implantable cardioverter defibrillator. *Europace* 2002; 4: 417–425.
9. Moulaert VR, Verbunt JA, van Heugten CM, Wade DT. Cognitive impairments in survivors of out-of-hospital cardiac arrest: A systematic review. *Resuscitation* 2009; 80: 297–305.
10. Hochstenbach JB, Anderson PG, van Limbeek J, Mulder TT. Is there a relation between neuropsychologic variables and quality of life after stroke? *Arch Phys Med Rehabil* 2001; 82: 1360–1366.
11. Staltnacke BM. Community integration, social support and life satisfaction in relation to symptoms 3 years after mild traumatic brain injury. *Brain Inj* 2007; 21: 933–942.

12. van de Port IG, Kwakkel G, Schepers VP, Heinemans CT, Lindeman E. Is fatigue an independent factor associated with activities of daily living, instrumental activities of daily living and health-related quality of life in chronic stroke? *Cerebrovasc Dis* 2007; 23: 40–45.
13. Cantor JB, Ashman T, Gordon W, Ginsberg A, Engmann C, Egan M, et al. Fatigue after traumatic brain injury and its impact on participation and quality of life. *J Head Trauma Rehabil* 2008; 23: 41–51.
14. Visser-Meily JM, Rhebergen ML, Rinkel GJ, van Zandvoort MJ, Post MW. Long-term health-related quality of life after aneurysmal subarachnoid hemorrhage. Relationship with psychological symptoms and personality characteristics. *Stroke* 2009; 40: 1526–1529.
15. Bunch TJ, White RD, Smith GE, Hodge DO, Gersh BJ, Hammill SC, et al. Long-term subjective memory function in ventricular fibrillation out-of-hospital cardiac arrest survivors resuscitated by early defibrillation. *Resuscitation* 2004; 60: 189–195.
16. Gamper G, Willeit M, Sterz F, Herkner H, Zoufaly A, Hornik K, et al. Life after death: posttraumatic stress disorder in survivors of cardiac arrest—prevalence, associated factors, and the influence of sedation and analgesia. *Crit Care Med* 2004; 32: 378–383.
17. Gorgels AP, Gijsbers C, de Vreede-Swagemakers J, Lousberg A, Wellens HJ. Out-of-hospital cardiac arrest – the relevance of heart failure. The Maastricht Circulatory Arrest Registry. *Eur Heart J* 2003; 24: 1204–1209.
18. Eisenberg MS, Horwood BT, Cummins RO, Reynolds-Haertle R, Hearne TR. Cardiac arrest and resuscitation: a tale of 29 cities. *Ann Emerg Med* 1990; 19: 179–186.
19. de Vreede-Swagemakers JJ, Gorgels AP, Dubois-Arbouw WI, Dalstra J, Daemen MJ, van Ree JW, et al. Circumstances and causes of out-of-hospital cardiac arrest in sudden death survivors. *Heart* 1998; 79: 356–361.
20. Bekkers SC, Eikemans BJ, Tieleman R, Braat SH, Dassen W, Partouns J, et al. Hypothermia for out-of-hospital cardiac arrest survivors: a single-center experience. *Am J Emerg Med* 2007; 25: 1078–1080.
21. Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderman R, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol* 1998; 51: 1055–1068.
22. Sander AM, Fuchs KL, High WM, Jr., Hall KM, Kreutzer JS, Rosenthal M. The Community Integration Questionnaire revisited: an assessment of factor structure and validity. *Arch Phys Med Rehabil* 1999; 80: 1303–1308.
23. de Haan R, Limburg M, Schuling J, Broeshart J, Jonkers L, van Zuylen P. [Clinimetric evaluation of the Barthel Index, a measure of limitations in daily activities]. *Ned Tijdschr Geneesk* 1993; 137: 917–921 (in Dutch).
24. Turnbull JC, Kersten P, Habib M, McLellan L, Mullee MA, George S. Validation of the Frenchay Activities Index in a general population aged 16 years and older. *Arch Phys Med Rehabil* 2000; 81: 1034–1038.
25. Broadbent DE, Cooper PF, FitzGerald P, Parkes KR. The Cognitive Failures Questionnaire (CFQ) and its correlates. *Br J Clin Psychol* 1982; 21: 1–16.
26. Spinhoven P, Ormel J, Sloekers PP, Kempen GI, Speckens AE, Van Hemert AM. A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. *Psychol Med* 1997; 27: 363–370.
27. Sundin EC, Horowitz MJ. Impact of Event Scale: psychometric properties. *Br J Psychiatry* 2002; 180: 205–209.
28. van der Ploeg E, Mooren TT, Kleber RJ, van der Velden PG, Brom D. Construct validation of the Dutch version of the impact of event scale. *Psychol Assess* 2004; 16: 16–26.
29. Kleinman L, Zodet MW, Hakim Z, Aledort J, Barker C, Chan K, et al. Psychometric evaluation of the fatigue severity scale for use in chronic hepatitis C. *Qual Life Res* 2000; 9: 499–508.
30. Horsted TI, Rasmussen LS, Meyhoff CS, Nielsen SL. Long-term prognosis after out-of-hospital cardiac arrest. *Resuscitation* 2007; 72: 214–218.
31. van Alem AP, de Vos R, Schmand B, Koster RW. Cognitive impairment in survivors of out-of-hospital cardiac arrest. *Am Heart J* 2004; 148: 416–421.
32. de Vos R, de Haes HC, Koster RW, de Haan RJ. Quality of survival after cardiopulmonary resuscitation. *Arch Intern Med* 1999; 159: 249–254.
33. Noble AJ, Baisch S, Mendelow AD, Allen L, Kane P, Schenk T. Posttraumatic stress disorder explains reduced quality of life in subarachnoid hemorrhage patients in both the short and long term. *Neurosurgery* 2008; 63: 1095–1104.
34. Moulaert VR, Verbunt JA, van Heugten CM, Bakx WG, Gorgels AP, Bekkers SC, et al. Activity and Life After Survival of a Cardiac Arrest (ALASCA) and the effectiveness of an early intervention service: design of a randomised controlled trial. *BMC Cardiovasc Disord* 2007; 7: 26.