

Influence of beta(2)-adrenoceptor gene polymorphisms on diet-induced thermogenesis

Citation for published version (APA):

Oomen, J. M., Wajers, P. M., van Rossum, C., Hoebee, B., Saris, W. H., & van Baak, M. A. (2005). Influence of beta(2)-adrenoceptor gene polymorphisms on diet-induced thermogenesis. *British Journal of Nutrition*, 94(5), 647-654. <https://doi.org/10.1079/BJN20051516>

Document status and date:

Published: 01/01/2005

DOI:

[10.1079/BJN20051516](https://doi.org/10.1079/BJN20051516)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Influence of β_2 -adrenoceptor gene polymorphisms on diet-induced thermogenesis

J. M. Oomen^{1*}, P. M. C. M. Waijers², C. van Rossum², B. Hoebee³, W. H. M. Saris¹ and M. A. van Baak¹

¹Department of Human Biology/NUTRIM, Maastricht University, Maastricht, The Netherlands

²Centre for Nutrition and Health and

³Laboratory of Toxicology, Pathology and Genetics, National Institute for Public Health and the Environment, Bilthoven, The Netherlands

(Received 21 October 2004 – Revised 31 March 2005 – Accepted 9 May 2005)

The sympathetic nervous system is involved in the control of energy metabolism and expenditure. Diet-induced thermogenesis is mediated partly by the β -adrenergic component of this system. The aim of the present study was to investigate the role of genetic variation in the β_2 -adrenoceptor in diet-induced thermogenesis. Data from twenty-four subjects (fourteen men and ten women; BMI 26.7(SEM 0.8) kg/m²; age 45.2(SEM 1.4) years) with different polymorphisms of the β_2 -adrenoceptor at codon 16 (Gly16Gly, Gly16Arg or Arg16Arg) were recruited for this study. Subjects were given a high-carbohydrate liquid meal, and the energy expenditure, respiratory exchange ratio, and plasma concentrations of NEFA, glycerol, glucose, insulin and catecholamines were measured before and over 4 h after the meal. The AUC of energy expenditure (diet-induced thermogenesis) was not significantly different between polymorphism groups, nor was the response of any of the other measured variables to the meal. In a multiple regression model, the only variable that explained a significant proportion (32 %) of the variation in diet-induced thermogenesis was the increase in plasma adrenaline in response to the meal ($P < 0.05$). The β_2 -adrenoceptor codon16 polymorphisms did not contribute significantly. In conclusion, an independent contribution of the codon 16 polymorphism of the β_2 -adrenoceptor gene to the variation in thermogenic response to a high-carbohydrate meal could not be demonstrated. The interindividual variation in thermogenic response to the meal was correlated with variations in the plasma adrenaline response to the meal.

β_2 -Adrenoceptor polymorphisms: Diet-induced thermogenesis: Catecholamines

Energy expenditure (EE) is an important factor in body-weight regulation. Diet-induced thermogenesis (DIT) is the EE associated with ingestion, absorption and storage of food and accounts for 10–15 % to the total daily EE. DIT shows considerable inter-individual variation (Donahoo *et al.* 2004), and it can be hypothesised that a low DIT contributes to weight gain. Many studies have investigated DIT in obese and lean subjects, but these studies show equivocal results (de Jonge & Bray, 1997, 2002; Granata & Brandon, 2002). Nevertheless, when multiple interfering factors are taken into account simultaneously, the evidence for a reduction in DIT in obesity becomes stronger (de Jonge & Bray, 2002). In addition, several studies show no change in DIT after weight reduction, suggesting that a reduced DIT in obesity is not the consequence of the obese state *per se* (Bessard *et al.* 1983; Schutz *et al.* 1984).

In response to feeding, especially to carbohydrate intake, sympathetic nervous system activity increases (Schwartz *et al.* 1999; Tappy, 2004). The sympathetic nervous system-mediated thermogenic response is also referred to as facultative thermogenesis (Tappy, 2004). The sympathetic nervous system response is biphasic, with an initial increase in noradrenaline and a delayed adrenaline response (Astrup *et al.* 1986, 1989). The sympathetic nervous system-mediated component of DIT can be blocked by

β -adrenoceptor antagonists (Astrup *et al.* 1989; Tappy, 1996), demonstrating the involvement of the β -adrenergic system.

β -Adrenoceptor genes are suggested to be 'candidate genes' for the development of obesity (Chagnon *et al.* 2003). Several studies (Echwald *et al.* 1998; Hellstrom *et al.* 1999; Kortner *et al.* 1999; Meirhaeghe *et al.* 1999; Ukkola *et al.* 2001; van Rossum *et al.* 2002; Garenc *et al.* 2003; Gonzalez Sanchez *et al.* 2003) have shown an association between polymorphisms at codons 16 and/or 27 of the β_2 -adrenoceptor gene and weight gain, obesity or obesity-related phenotypes.

Functional consequences of these polymorphisms with respect to adipocyte lipolysis have also been reported (Large *et al.* 1997; Eriksson *et al.* 2004). Large *et al.* (1997) demonstrated that isolated abdominal subcutaneous fat cells from women homozygous for the arginine 16 (Arg16) polymorphism of the β_2 -adrenoceptor gene had a 5-fold lower sensitivity for lipolysis induced by the β_2 -adrenoceptor agonist terbutaline than did fat cells from women who were heterozygous or homozygous for glycine 16 (Gly16), independent of body fat. Eriksson *et al.* (2004) showed that homozygous haplotypes of the β_2 -adrenoceptor gene differed about 250-fold in their sensitivity to terbutaline-induced lipolysis, the least sensitive haplotype being homozygous for the arginine variant at codon 16. In addition, we have recently

Abbreviations: absDIT, absolute diet-induced thermogenesis; Arg16, arginine 16; AUC, area under the curve; DIT, diet-induced thermogenesis; EE, energy expenditure; FFM, fat-free mass; Gly16, glycine 16; relDIT, relative diet-induced thermogenesis.

* **Corresponding author:** Dr J. M. Oomen, fax +31 (0)43 367 9776, email j.oomen@hb.unimaas.nl

reported that the thermogenic response to infusion of the β_2 -adrenoceptor agonist salbutamol was blunted in carriers of the Arg16Arg variant of the β_2 -adrenoceptor gene compared with Gly16 carriers (Oomen *et al.* 2005).

We therefore hypothesised that carriers of the Arg16Arg variant of the β_2 -adrenoceptor gene might also have a reduced DIT in response to a high-carbohydrate meal compared with carriers of the Gly16 variant. This might, at least in part, explain the association between this polymorphism and obesity. The present study was designed to test this hypothesis.

Methods

Subjects

Twenty-five volunteers (fourteen men and eleven women) participated in the study. They were recruited from an existing cohort in the Maastricht area in The Netherlands that has previously been described (van Rossum *et al.* 2002). The age of the subjects ranged between 32 and 55 years. They did not use medication at the time of the study or the week before. The study protocol was reviewed and approved by the Ethics Committee of Maastricht University, and all subjects gave written informed consent before participating.

Genotyping of the codon 16 polymorphism of the β_2 -adrenoceptor gene

To genotype the codon 16 polymorphism of the β_2 -adrenoceptor gene, genomic DNA was extracted from leucocytes from each individual by digestion with proteinase K followed by phenol–chloroform extraction. Determination of the polymorphism was performed using a PCR–restriction fragment length polymorphism analysis as described before (Large *et al.* 1997).

Experimental design

The day preceding the experimental day, subjects consumed a fixed diet provided by the researchers. This diet (energy content 10.8(SEM 0.3) MJ/d) consisted of 50 % energy from carbohydrates, 15 % from protein and 35 % from fat, which corresponds to the average macronutrient composition of the Dutch diet. The total amount of energy that each subject received was based on an estimation of resting EE by the Harris–Benedict equation (Harris & Benedict, 1919) multiplied by an estimation of the subjects' activity level. The activity level was estimated using a short questionnaire. For most subjects, the activity level was set at 1.4, except when subjects reported to be at least moderately active for more than 3 h/week, for which a figure of 1.6 was used. Subjects were asked to refrain from unusual or strenuous exercise during the day before the experimental day.

On the experimental day, subjects came to the laboratory in the morning after an overnight fast. They came by car or by bus in order to limit their physical activity before the measurements. Subjects were weighed, and body composition was determined by bio-impedance. Thereafter, a venous catheter was inserted into an antecubital vein for blood sampling. Subjects were then positioned under a ventilated hood in a recumbent position for indirect calorimetry. The indirect calorimetry measurements continued throughout the whole experiment and were interrupted only for the consumption of the meal. Subjects were instructed

to limit their movements as much as possible and not to speak during the experiment. They could watch television or a video.

After 30 min rest, baseline measurements were performed: EE was measured over 30 min, and a baseline blood sample was collected at the end. Subjects then received an energy bolus providing 35 % of the energy they had consumed the preceding day (3.71(SEM 0.12) MJ). This test meal consisted of two liquid formulas (total volume 593(SEM 19) ml; Meritene Polvo, Novartis Nutrition and Isostar Long Energy; Novartis Nutrition, Breda, The Netherlands). The macronutrient composition of the test meal was 84 % from carbohydrate, 13 % from protein and 3 % from fat. A high-carbohydrate and high-protein meal is known to stimulate sympathetic nervous system activity (Schwartz *et al.* 1999). The test meal was consumed within 5 min, and measurements were continued for another 4 h. At 10, 30, 60, 120, 180 and 240 min after the meal, a blood sample was drawn.

Anthropometry and body composition

Weight was measured to the nearest 100 g with a digital scale (Seca Delta, Almere, The Netherlands), the subjects wearing clothes but without shoes. Height was known, since it had been measured before in the cohort study (van Rossum *et al.* 2002). Total body water was measured by single-frequency bioimpedance at 50 kHz, using a Xitron 400B bioimpedance analyser (Xitron Technologies Inc., San Diego, CA, USA; van Marken Lichtenbelt *et al.* 1994). Fat-free mass (FFM) was calculated from weight and total body water using a prediction equation (Lukaski *et al.* 1986).

Energy expenditure, respiratory exchange ratio

V_{O_2} and CO_2 production were determined using an open-circuit ventilated-hood system (Omnical, Maastricht, The Netherlands). This system is based on the analysis system for respiration chambers, which has previously been described (Schoffelen *et al.* 1997). EE was calculated from V_{O_2} and CO_2 production according to the Weir formula (Weir, 1949). Respiratory exchange ratio is the ratio of CO_2 production to V_{O_2} .

Analysis of blood samples

Blood samples for the determination of plasma NEFA, glycerol, glucose and insulin concentrations were collected in sodium EDTA tubes, and samples for plasma noradrenaline and adrenaline concentrations in tubes containing heparin and glutathione (1.5 % w/v). Blood samples were immediately centrifuged for 10 min at 800 g at 4°C. Plasma was transferred into test tubes, rapidly frozen in liquid N and stored at -80°C until further analysis.

Glucose (UniKit III, cat. no. 07367204; Roche, Basel, Switzerland) and NEFA (Wako NEFA-C test kit; Wako Chemicals, Neuss, Germany) were analysed with a COBAS FARA semi-automated analyser (Roche Diagnostica, Basel, Switzerland).

Plasma catecholamine levels (time points 0, 120 and 240 min only) were determined by HPLC according to the method of Alberts *et al.* (1992) using a ClinPrep kit (Recipe, Munich, Germany). The plasma insulin level was determined with a double antibody RIA (Insulin RIA 100; Linco Research, St Charles, MO, USA). The homeostasis model assessment index, a measure of insulin sensitivity, was calculated according to Matthews *et al.* (1985) using baseline plasma glucose and insulin levels.

Statistical methods

Data are presented as means with their standard errors. For between-group comparisons, EE was adjusted for FFM. Besides absolute values of EE adjusted for FFM, the change in EE from baseline was also expressed as the change relative to the baseline (relative EE = EE/baseline EE). DIT was calculated as the AUC of relative and absolute EE over the 4 h post-meal period and expressed as the relative DIT (relDIT) and absolute DIT (absDIT). Areas under the curve of post-meal changes from baseline were also calculated for glucose, insulin, NEFA, glycerol, noradrenaline and adrenaline concentrations. Changes over time were analysed for the whole group by repeated-measures ANOVA. ANOVA and independent sample *t* tests were used for comparisons between groups.

Simple regression analysis (Pearson's correlation) was performed with absDIT as the dependent variable and different parameters as the independent variable (energy content of the meal, AUC for glucose, insulin, NEFA, glycerol, noradrenaline and adrenaline, baseline plasma noradrenaline and adrenaline, and homeostasis model assessment index). Multiple stepwise regression analysis, with variables of the simple regression analysis with a value of *P*<0.2 and polymorphism groups as dummy variables, was conducted to estimate the independent contributions of these variables to absDIT. *P*<0.05 was considered to be statistically significant, and all tests were performed as two-tailed tests. Statistical analyses were performed with the SPSS 11.0 statistical package (SPSS Inc. Chicago, IL, USA).

Results

Responses to the meal in the whole group

One subject could only consume half of the test meal and was therefore excluded from the analysis. Data on twenty-four subjects are reported. Their characteristics are shown in Table 1.

After the test meal, EE increased significantly in the whole group. After 120 min, EE was significantly higher than baseline EE (5.61 (SEM 0.10) kJ/min adjusted for FFM *v.* 4.63 (SEM 0.08) kJ/min adjusted for FFM, respectively; *P*<0.001) (Fig. 1). After 240 min, EE was lower (5.34 (SEM 0.09) kJ/min) than at 120 min (*P*<0.01), but it was still significantly higher than the baseline EE (*P*<0.001) (Fig 1). The respiratory exchange ratio

increased after the meal and remained elevated over the 4 h post-meal period (*t* = 0 *v.* *t* = 240 min, *P*<0.001) (Fig. 2).

Changes in plasma concentrations of metabolites and insulin are shown in Fig. 3. Plasma glycerol and NEFA levels were significantly lowered over the whole post-meal period (*t* = 0 *v.* *t* = 240 min, *P*<0.001). Plasma glucose concentration was significantly increased at 60 min compared with the baseline level (*P*<0.001) and was still higher at 240 min compared with baseline (*P*<0.05). Plasma insulin levels were significantly increased at 60 min (*P*<0.001). The insulin concentration increased after the meal and started to decrease 120 min after the meal but was still higher than baseline at 240 min (*P*<0.001).

Plasma noradrenaline levels (Fig. 4) were significantly elevated at 120 min and 240 min after the meal compared with baseline levels (*P*<0.05). Plasma adrenaline levels (Fig. 4) were not significantly different from baseline at 120 min (*P*=0.08). At 240 min, however, plasma adrenaline levels were significantly higher than at 120 min (*P*<0.01).

Baseline values and responses to the meal in the β₂-adrenoceptor codon 16 polymorphism groups

Baseline EE was similar in all polymorphism groups (4.75 (SEM 0.11), 4.55 (SEM 0.10) and 4.42 (SEM 0.21)(kJ/min adjusted for FFM, for Gly16Gly, Gly16Arg and Arg16Arg respectively; NS). There was no significant difference in absDIT (*P*=0.125; Fig. 1) or relDIT (*P*=0.131) between groups (Table 2). In addition, we compared glycine polymorphism carriers (Gly16Arg and Gly16Gly) with Arg16Arg individuals and found a trend (*P*=0.06) towards Arg16Arg having a higher absDIT and a statistically higher relDIT (*P*=0.048) compared with Gly polymorphism carriers. None of the other parameters differed statistically significantly between the three groups, either at baseline or with respect to meal-induced changes (Figs. 2–4).

Explanation of energy expenditure response to the meal

Of all tested variables, the change in EE after the meal (absDIT) was significantly correlated with the change in adrenaline concentration (AUC adrenaline; *P*=0.003) and with the energy content of the meal (*P*=0.024; Table 3). In a multiple-regression model, only the AUC of plasma adrenaline level contributed significantly to absDIT

Table 1. Subject characteristics, for the total group and the β₂-adrenoceptor codon 16 variants Gly16Gly, Gly16Arg and Arg16Arg. (Values are means with their standard errors)

	All		Gly16Gly		Gly16Arg		Arg16Arg	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
Sex (M/F)								
Males (<i>n</i>)	14		7		3		4	
Females (<i>n</i>)	10		6		3		1	
Estimated energy intake (MJ/d)	10.76	0.30	10.79	0.41	10.36	0.19	11.17	0.83
Energy content test meal (MJ)	3.76	0.11	3.7	0.14	3.62	0.19	3.92	0.29
Height (m)	1.75	0.02	1.76	0.03	1.71	0.03	1.75	0.05
Weight (kg)	81.70	3.00	81.40	4.10	80.50	3.80	84.10	9.80
Age (years)	45.20	1.40	44.00	2.10	49.50	1.50	43.30	2.40
BMI (kg/m ²)	26.70	0.80	26.10	1.10	27.70	1.30	27.00	2.10
Body fat (%)	28.90	2.00	30.70	2.70	29.60	5.10	23.40	2.20
Fat-free mass (kg)	58.30	2.90	58.30	2.90	57.10	5.80	64.30	7.60
Resting energy expenditure (kJ/min)	4.63	0.08	4.75	0.11	4.55	0.10	4.42	0.21




Fig. 1. Energy expenditure (EE) adjusted for fat free mass (FFM; kg/min per kg FFM) for the three genetic variations in the β_2 -adrenoceptor (●, Gly16Gly; ○, Gly16Arg; ▼, Arg16Arg) before ($t=0$) and after (30 min intervals), a high carbohydrate meal. The meal was given directly after $t=0$. Values are means with their standard errors represented by vertical bars. The inset shows the results for the whole group.

($P=0.003$, adjusted r^2 0.323). When the polymorphism groups were entered into the regression model, the dummy variables did not contribute significantly to absDIT ($P>0.20$). Similar results were found for relDIT.

Discussion

The aim of the present study was to investigate the influence of the codon 16 variants of the β_2 -adrenoceptor gene on the thermogenic

response to a meal. We hypothesised that subjects with the Arg16Arg polymorphism of the β_2 -adrenoceptor would have a lower DIT after the meal than subjects with the Arg16Gly or Gly16Gly polymorphism. Our data do not support this hypothesis. In contrast, there was a higher relDIT and a trend towards a higher absDIT in the Arg16Arg carriers compared with the Gly polymorphism carriers. absDIT was most strongly associated with the adrenaline response to the meal, and an additional independent contribution of the polymorphism could not be demonstrated.




Fig. 2. Respiratory exchange ratio (RER) for the three genetic variations in the β_2 -adrenoceptor (●, Gly16Gly; ○, Gly16Arg; ▼, Arg16Arg) before ($t=0$) and after (30 min intervals), a high carbohydrate meal. Values are means with their standard errors represented by vertical bars. The inset shows the results for the whole group.




Fig. 3. Plasma glucose, insulin, glycerol and NEFA for the three genetic variations in the β_2 -adrenoceptor (—●—, Gly16Gly; —○—, Gly16Arg; —▼—, Arg16Arg), before ($t = 0$) and after (10, 30, 60, 120, 180 and 240 min) a high carbohydrate meal. Values are means with their standard errors represented by vertical bars.




Fig. 4. Plasma nonadrenaline and adrenaline levels for the three genetic variations in the β_2 -adrenoceptor (—●—, Gly16Gly; —○—, Gly16Arg; —▼—, Arg16Arg), before ($t = 0$) and after (120 min intervals) a high carbohydrate meal. Values are means with their standard errors represented by vertical bars. The insets show the results for each whole group.

Table 2. Absolute diet-induced thermogenesis (absDIT) and relative diet-induced thermogenesis (relDIT) in the polymorphism groups, Gly16Gly, Gly16Arg and Arg16Arg

(Values are means with their standard errors)

	Gly16Gly <i>n</i> 13		Gly16Arg <i>n</i> 6		Arg16Arg <i>n</i> 5	
	Mean	SEM	Mean	SEM	Mean	SEM
Abs DIT (kJ/4 h)	175.1	14.6	147.5	10.8	232.2	14.3
Rel DIT (%/4 h)	15.5	1.5	13.5	1.1	22.7	6.4

Meal-induced thermogenesis is partly mediated by increased sympathoadrenal activity (Valensi *et al.* 1998; Camastra *et al.* 1999; Schwartz *et al.* 1999; Lowell & Bachman, 2003). β -Adrenoceptor blockade blunts meal-induced thermogenesis, especially between 2 and 4 h after the meal (Astrup *et al.* 1989), indicating that the β -adrenergic branch of the sympathoadrenal system is involved in this response during this period. In particular, the β_2 -adrenoceptor seems to be important in catecholamine-induced thermogenesis (Nagase *et al.* 2001). The magnitude of the meal-induced stimulation of the sympathoadrenal system is mainly determined by the size of the meal and its carbohydrate and protein content (Schwartz *et al.* 1999). Because we wanted to study differences in response to meal-induced stimulation of the β_2 -adrenoceptors between subjects with different variants of the β_2 -adrenoceptor gene, we tried to optimise the meal-induced β -adrenergic stimulation by giving a large meal with a high carbohydrate content.

Over the 4 h post-meal period, the expected increases in EE, respiratory exchange ratio, and glucose and insulin concentrations were found, accompanied by reductions in plasma NEFA and glycerol concentrations (Welle & Campbell, 1983b; Astrup *et al.* 1987). Differences in insulin sensitivity have been shown to affect DIT (Camastra *et al.* 1999). In the present study, however, the homeostasis model assessment index, as a measure of insulin sensitivity, was not correlated with absDIT. This could be due to the fact that the variation in homeostasis model assessment index in our study was relatively small. DIT has also been shown to be impaired in obese compared with lean individuals, although results are not consistent (de Jonge & Bray, 2002). We did not find a significant correlation between percentage fat and absDIT ($r = -0.272$, $P = 0.199$), or between BMI and absDIT ($r = 0.141$,

$P = 0.512$), but the ranges of percentage fat and BMI in our group were relatively small.

The plasma noradrenaline concentration was increased at 2 and 4 h after the meal, the plasma adrenaline level only at 4 h. This early response in plasma noradrenaline level and delayed response in plasma adrenaline level has been shown before (Welle & Campbell 1983a; Astrup *et al.* 1987). Astrup *et al.* (1986) suggested that the delayed increase in plasma adrenaline was elicited by the decrease in plasma glucose during this period. The AUC of the post-meal changes in plasma adrenaline was the only parameter that contributed significantly to the variation in DIT ($r^2 = 0.32$). Because adrenaline has a higher affinity for β_2 -adrenoceptors than β_1 - or β_3 -adrenoceptors (Hoffmann *et al.* 2004), the metabolic effects of adrenaline are predominantly mediated by β_2 -adrenoceptors. Our data do not support a modulation of the adrenaline-induced thermogenic effect by the codon 16 polymorphism of the β_2 -adrenoceptor gene, as hypothesised. There was no evidence for a blunted absDIT in subjects with the Arg16Arg variant of the gene compared with Gly16 polymorphism carriers. Instead, we found a trend towards an increased absDIT in the Arg16Arg group compared with the glycine carriers, and relDIT was significantly higher in the Arg16Arg group. Based on the results of the multiple regression analysis, this difference in DIT was partly explained by a variation in adrenaline response, although we were unable to demonstrate statistically significant differences in adrenaline response between the groups.

Our initial hypothesis of a reduced DIT in Arg16Arg homozygotes was based on several previous findings. The Arg16Arg polymorphism has been associated with a reduced sensitivity to β_2 -adrenoceptor agonist-induced lipolysis in isolated human fat cells (Large *et al.* 1997; Eriksson *et al.* 2004). Moreover, in a previous study, we showed that individuals with the Arg16Arg genotype had a blunted thermogenic response to stimulation with the β_2 -adrenoceptor agonist salbutamol compared with Gly16 carriers (Oomen *et al.* 2005). Both factors might contribute to an increased susceptibility to weight gain and obesity in Arg16Arg carriers. The Arg16 polymorphism has indeed been associated with obesity or obesity-related phenotypes in many studies (Echwald *et al.* 1998; Hellstrom *et al.* 1999; Kortner *et al.* 1999; Meirhaeghe *et al.* 1999; Ukkola *et al.* 2001; van Rossum *et al.* 2002; Garenc *et al.* 2003; Gonzalez Sanchez *et al.* 2003), although there are also studies that do not find this association (Ishiyama-Shigemoto *et al.* 1999; Hayakawa *et al.* 2000). The results of this study, however, suggest that even though the Arg16 variant of the β_2 -adrenoceptor may be less sensitive to direct β_2 -adrenoceptor stimulation *in vitro* and *in vivo*, the thermogenic response to a meal is increased rather than reduced in Arg16 homozygotes compared with Gly16 carriers.

Whether the lower sensitivity of the Arg16Arg β_2 -adrenoceptor is compensated by a higher level of sympathetic stimulation, or whether a higher level of stimulation induces a reduced responsiveness of the β_2 -adrenoceptor, cannot be derived from our data. The latter mechanism seems unlikely in view of the fact that, *in vitro*, an enhanced agonist-promoted downregulation of the Gly16 receptor compared with the Arg16 receptor is well established (Liggett, 1997). However, the opposite has been demonstrated *in vivo* for isoprenaline-induced venodilatation, which was insensitive to downregulation in Gly16 homozygotes, whereas those homozygous for Arg16 showed significant downregulation during the 2 h isoprenaline infusion (Dishy *et al.* 2001).

How a reduced sensitivity of the β_2 -adrenoceptor could lead to an increased activation of the sympathoadrenal system is not clear from

Table 3. Pearson correlations (r) of different variables with absolute diet-induced thermogenesis (absDIT), and corresponding P values.

	r	P value
Energy content test meal	0.459	0.024*
AUC glucose	0.116	0.606
AUC insulin	0.292	0.177
AUC NEFA	0.182	0.442
AUC glycerol	-0.142	0.528
Baseline plasma noradrenaline	0.395	0.056
Baseline plasma adrenaline	0.096	0.656
AUC noradrenaline	0.109	0.620
AUC adrenaline	0.595	0.003*
HOMA index	0.260	0.220
Fat percentage	-0.272†	0.199†
BMI	0.141	0.512

AUC, area under the curve; HOMA, homeostasis model assessment.

* $P < 0.05$; † $P < 0.2$.

our data, but assuming that the adrenaline response is indeed linked to changes in plasma glucose (Astrup *et al.* 1986), it is conceivable that a reduced glucose output by the liver to compensate for the fall in blood glucose late after the meal would induce a higher adrenaline response. Glucose output by the liver could be compromised by less responsive β_2 -adrenoceptors involved in liver glycogenolysis or in reduced gluconeogenesis from glycerol owing to compromised β_2 -adrenoceptor-mediated lipolysis.

In conclusion, the present study shows that there is no independent contribution of the codon 16 variant of the β_2 -adrenoceptor gene to the thermogenic response after a high-carbohydrate meal, but that this response is positively correlated with the plasma adrenaline response to the meal.

References

- Alberts G, Boomsma F, Man in't Veld AJ & Schalekamp MA (1992) Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection. *J Chromatogr* **583**, 236–240.
- Astrup A, Andersen T, Henriksen O, Christensen NJ, Bulow J, Madsen J & Quaade F (1987) Impaired glucose-induced thermogenesis in skeletal muscle in obesity. The role of the sympathoadrenal system. *Int J Obes* **11**, 51–66.
- Astrup A, Bulow J, Christensen NJ, Madsen J & Quaade F (1986) Facultative thermogenesis induced by carbohydrate: a skeletal muscle component mediated by epinephrine. *Am J Physiol* **250**, E226–E229.
- Astrup A, Simonsen L, Bulow J, Madsen J & Christensen NJ (1989) Epinephrine mediates facultative carbohydrate-induced thermogenesis in human skeletal muscle. *Am J Physiol* **257**, E340–E345.
- Bessard T, Schutz Y & Jequier E (1983) Energy expenditure and postprandial thermogenesis in obese women before and after weight loss. *Am J Clin Nutr* **38**, 680–693.
- Camastra S, Bonora E, Del Prato S, Rett K, Weck M & Ferrannini E (1999) Effect of obesity and insulin resistance on resting and glucose-induced thermogenesis in man. EGIR (European Group for the Study of Insulin Resistance). *Int J Obes Relat Metab Disord* **23**, 1307–1313.
- Chagnon YC, Rankinen T, Snyder EE, Weisnagel SJ, Perusse L & Bouchard C (2003) The human obesity gene map: the 2002 update. *Obes Res* **11**, 313–367.
- de Jonge L & Bray GA (1997) The thermic effect of food and obesity: a critical review. *Obes Res* **5**, 622–631.
- de Jonge L & Bray GA (2002) The thermic effect of food is reduced in obesity. *Nutr Rev* **60**, 295–297, author reply 299–300.
- Dishy V, Sofowora GG, Xie HG, Kim RB, Byrne DW, Stein CM & Wood AJ (2001) The effect of common polymorphisms of the beta2-adrenergic receptor on agonist-mediated vascular desensitization. *N Engl J Med* **345**, 1030–1035.
- Donahoo WT, Levine JA & Melanson EL (2004) Variability in energy expenditure and its components. *Curr Opin Clin Nutr Metab Care* **7**, 599–605.
- Echwald SM, Sorensen TI, Tybjaerg-Hansen A, Andersen T & Pedersen O (1998) Gln27Glu variant of the human beta2-adrenoreceptor gene is not associated with early-onset obesity in Danish men. *Diabetes* **47**, 1657–1658.
- Eriksson P, Dahlman I, Ryden M, Hoffstedt J & Arner P (2004) Relationship between beta-2 adrenoceptor gene haplotypes and adipocyte lipolysis in women. *Int J Obes Relat Metab Disord* **28**, 185–190.
- Garenc C, Perusse L, Chagnon YC, *et al.* (2003) Effects of beta2-adrenergic receptor gene variants on adiposity: the HERITAGE Family Study. *Obes Res* **11**, 612–618.
- Gonzalez Sanchez JL, Proenza AM, Martinez Larrad MT, Ramis JM, Fernandez Perez C, Palou A & Serrano Rios M (2003) The glutamine 27 glutamic acid polymorphism of the beta2-adrenoceptor gene is associated with abdominal obesity and greater risk of impaired glucose tolerance in men but not in women: a population-based study in Spain. *Clin Endocrinol (Oxf)* **59**, 476–481.
- Granata GP & Brandon LJ (2002) The thermic effect of food and obesity: discrepant results and methodological variations. *Nutr Rev* **60**, 223–233.
- Harris J & Benedict F (1919) *A Biometric Study of Basal Metabolism in Man*. Washington: Carnegie Institute.
- Hayakawa T, Nagai Y, Kahara T, Yamashita H, Takamura T, Abe T, Nomura G & Kobayashi K (2000) Gln27Glu and Arg16Gly polymorphisms of the beta2-adrenergic receptor gene are not associated with obesity in Japanese men. *Metabolism* **49**, 1215–1218.
- Hellstrom L, Large V, Reynisdottir S, Wahrenberg H & Arner P (1999) The different effects of a Gln27Glu beta 2-adrenoceptor gene polymorphism on obesity in males and in females. *J Intern Med* **245**, 253–259.
- Hoffmann C, Leitz MR, Oberdorf-Maass S, Lohse MJ & Klotz KN (2004) Comparative pharmacology of human beta-adrenergic receptor subtypes – characterization of stably transfected receptors in CHO cells. *Naunyn Schmiedebergs Arch Pharmacol* **369**, 151–159.
- Ishiyama-Shigemoto S, Yamada K, Yuan X, Ichikawa F & Nonaka K (1999) Association of polymorphisms in the beta2-adrenergic receptor gene with obesity, hypertriglyceridaemia, and diabetes mellitus. *Diabetologia* **42**, 98–101.
- Kortner B, Wolf A, Wendt D, Beisiegel U & Evans D (1999) Lack of association between a human beta-2 adrenoceptor gene polymorphism (gln27glu) and morbid obesity. *Int J Obes Relat Metab Disord* **23**, 1099–1100.
- Large V, Hellstrom L, Reynisdottir S, Lonnqvist F, Eriksson P, Lannfelt L & Arner P (1997) Human beta-2 adrenoceptor gene polymorphisms are highly frequent in obesity and associated with altered adipocyte beta-2 adrenoceptor function. *J Clin Invest* **100**, 3005–3013.
- Liggett SB (1997) Polymorphisms of the beta2-adrenergic receptor and asthma. *Am J Respir Crit Care Med* **156**, S156–S162.
- Lowell BB & Bachman ES (2003) Beta-adrenergic receptors, diet-induced thermogenesis, and obesity. *J Biol Chem* **278**, 29385–29388.
- Lukaski HC, Bolonchuk WW, Hall CB & Siders WA (1986) Validation of tetrapolar bioelectrical impedance method to assess human body composition. *J Appl Physiol* **60**, 1327–1332.
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF & Turner RC (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* **28**, 412–419.
- Meirhaeghe A, Helbecque N, Cotel D & Amouyel P (1999) Beta2-adrenoceptor gene polymorphism, body weight, and physical activity. *Lancet* **353**, 896.
- Nagase I, Yoshida T & Saito M (2001) Up-regulation of uncoupling proteins by beta-adrenergic stimulation in L6 myotubes. *FEBS Lett* **494**, 175–180.
- Oomen JM, van Rossum CT, Hoebee B, Saris WH & van Baak MA (2005) β_2 -Adrenergic receptor polymorphisms and salbutamol-stimulated energy expenditure. *J Clin Endocrinol Metab* **90**, 2301–230.
- Schoffelen PF, Westerterp KR, Saris WH & Ten Hoor F (1997) A dual-respiration chamber system with automated calibration. *J Appl Physiol* **83**, 2064–2072.
- Schutz Y, Golay A, Felber JP & Jequier E (1984) Decreased glucose-induced thermogenesis after weight loss in obese subjects: a predisposing factor for relapse of obesity? *Am J Clin Nutr* **39**, 380–387.
- Schwartz MW, Baskin DG, Kaiyala KJ & Woods SC (1999) Model for the regulation of energy balance and adiposity by the central nervous system. *Am J Clin Nutr* **69**, 584–596.
- Tappy L (1996) Thermic effect of food and sympathetic nervous system activity in humans. *Reprod Nutr Dev* **36**, 391–397.
- Tappy L (2004) Metabolic consequences of overfeeding in humans. *Curr Opin Clin Nutr Metab Care* **7**, 623–628.

- Ukkola O, Tremblay A & Bouchard C (2001) Beta-2 adrenergic receptor variants are associated with subcutaneous fat accumulation in response to long-term overfeeding. *Int J Obes Relat Metab Disord* **25**, 1604–1608.
- Valensi P, Lormeau B, Dabbech M, Miossec P, Paries J, Dauchy F & Attali JR (1998) Glucose-induced thermogenesis, inhibition of lipid oxidation rate and autonomic dysfunction in non-diabetic obese women. *Int J Obes Relat Metab Disord* **22**, 494–499.
- van Marken Lichtenbelt WD, Westerterp KR, Wouters L & Lijndijk SC (1994) Validation of bioelectrical-impedance measurements as a method to estimate body-water compartments. *Am J Clin Nutr* **60**, 159–166.
- van Rossum CT, Hoebee B, Seidell JC, Bouchard C, van Baak MA, de Groot CP, Chagnon M, de Graaf C & Saris WH (2002) Genetic factors as predictors of weight gain in young adult Dutch men and women. *Int J Obes Relat Metab Disord* **26**, 517–528.
- Weir JB (1949) New methods for calculating metabolic rate with special reference to protein metabolism. *Journal of Physiology* **109**, 1–9.
- Welle S & Campbell RG (1983a) Stimulation of thermogenesis by carbohydrate overfeeding. Evidence against sympathetic nervous system mediation. *J Clin Invest* **71**, 916–925.
- Welle SL & Campbell RG (1983b) Normal thermic effect of glucose in obese women. *Am J Clin Nutr* **37**, 87–92.