

A Licence to Prescribe

Citation for published version (APA):

Kramers, C., Janssen, B. J., Knol, W., Hessel, M. H. M., Mulder, W. M., Dumont, G., van den Brink, A. M., & Tichelaar, J. (2017). A Licence to Prescribe. *British Journal of Clinical Pharmacology*, 83(8), 1860-1861. <https://doi.org/10.1111/bcp.13257>

Document status and date:

Published: 01/08/2017

DOI:

[10.1111/bcp.13257](https://doi.org/10.1111/bcp.13257)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

LETTER TO THE EDITOR

A Licence to Prescribe

Correspondence Jelle Tichelaar PhD, Department of Internal Medicine, Section of Pharmacotherapy, VU University Medical Center, De Boelelaan 1117 (ZH 4A47), 1081 HV AMSTERDAM, The Netherlands. Tel.: +31 20 444 8090; Fax: +31 20 444 8100; E-mail: j.tichelaar@vumc.nl

Received 7 November 2016; **Revised** 9 January 2017; **Accepted** 2 February 2017

Cornelis Kramers¹, Ben. J. Janssen², Wilma Knol³, Marleen H. M. Hessel⁴, Wilhelmina M. Mulder⁴, Glenn Dumont⁵, Antoinette Maassen van den Brink⁶ and Jelle Tichelaar⁷

¹Department of Pharmacology-Toxicology, Radboud University Medical Center and Department of Clinical Pharmacy, Canisius Wilhelmina Ziekenhuis, Nijmegen, The Netherlands, ²Department of Pharmacology & Toxicology, Maastricht University, Maastricht, The Netherlands, ³Department of Geriatric Medicine and Expertise Centre Pharmacotherapy in Old Persons (EPHOR), University Medical Centre Utrecht, Utrecht, The Netherlands, ⁴Department of Clinical Pharmacy and Toxicology, Leiden University Medical Centre, Leiden, The Netherlands, ⁵Department of Pharmacy, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands, ⁶Division of Vascular Pharmacology and Metabolic Diseases, Department of Internal Medicine, Erasmus MC, University Medical Center Rotterdam, The Netherlands, and ⁷Department of Internal Medicine section Pharmacotherapy and Research & Expertise Center In Pharmacotherapy Education (RECIPE), VU University Medical Center, Amsterdam, The Netherlands

Keywords graduation, medication safety, medical students, pharmacotherapy, prescribing

Mistakes in prescribing may have serious consequences, ranging from inefficacy to serious adverse events and even death [1]. However, in the Netherlands in most medical schools, pharmacotherapy is only a minor component of an integrated assessment programme. This means that students can, theoretically, skip most clinical pharmacology teaching during their training yet still qualify and become licensed prescribers of medication [2]. While many prescribers report not feeling confident in prescribing when they start clinical work [3, 4], it is these inexperienced doctors who prescribe the majority of drugs [5]. Not surprisingly, many studies report a high rate of inappropriate prescribing, a potential cause of preventable morbidity and mortality [1, 6].

Appropriate prescribing requires a sufficient level of skills and knowledge. In the Netherlands, the World Health Organization six-step method is now used as the basis for pharmacotherapy education in all eight medical schools [2]. However, although most clinical pharmacology knowledge is readily accessible to prescribers, without a basic knowledge of potentially harmful medication the unconsciously incompetent prescriber may still make serious mistakes.

To tackle this problem, eight Dutch and three Belgian medical schools have joined forces to develop a unique pharmacotherapy test focusing on medication safety, to assess whether future prescribers are able to prescribe safely. It is recognized that certain drug groups cause the majority of preventable serious adverse reactions [7], and based on this literature we generated a list of drugs that cause most harm (Table 1). For each drug listed, medical students are expected to know its mechanism of action, side effects that may cause serious harm, associated risk factors, contributing comedication, and measures to prevent and to treat side effects. In addition, students are expected to know about the proper use of antibiotics and general subjects, such as the six-step method, medicine and driving, prescribing laws and regulations, dosage calculations, and basic pharmacokinetics. The multiple-choice test we developed covers these subjects and assesses whether students have a basic level of specific knowledge about drugs that can cause preventable serious adverse events. Currently it consists of 40 questions, to be answered within 1 h with a caesura of 34 correct answers. Ideally, the test should be taken just before students start prescribing under

Table 1

Subjects covered in the pharmacotherapy end test

Analgesics
Anticoagulants
Cardiovascular drugs
Drugs for diabetes mellitus
Antidepressants
Benzodiazepines
Antibiotics
Laws and regulations
Drug allergy
Proper usage of drugs
Basic pharmacokinetics

supervision. Educational material has been developed including a reader, cases in www.pscribe.nl, a YouTube channel with tutorials, and even a game – *Battle of the meds* – which can be downloaded from the App Store for free. Students must pass this test before they can graduate.

The test was introduced in Nijmegen in 2014, and the other participating medical schools are in the process of introducing it. The Belgian universities are currently working on an additional set of questions for Belgian students, since laws and regulations (a part of the test) differ in Belgium. In Nijmegen, every month 30 students perform an online test, 70% pass the first time and only a single student needs a fourth attempt. A similar initiative has been undertaken in the UK [8]. We think that this test can contribute substantially to patient safety by reducing preventable medication-induced morbidity and mortality, much in the same way that passing a driving test contributes to traffic safety. Studies are in progress to evaluate whether this test has a long-term effect on prescribing skills when these students start prescribing as physicians.

Competing Interests

All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on

request from the corresponding author) and declare no support from any organization for the submitted work, no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years and no other relationships or activities that could appear to have influenced the submitted work.

References

- 1 Leendertse AJ, Egberts AC, Stoker LJ, van den Bemt PM. Frequency of and risk factors for preventable medication-related hospital admissions in the Netherlands. *Arch Intern Med* 2008; 168: 1890–6.
- 2 Keijsers CJ, de Wit JE, Tichelaar J, Brouwers JR, de Wildt DJ, de Vries PG, *et al*. Education on prescribing for older patients in the Netherlands: a curriculum mapping. *Eur J Clin Pharmacol* 2015; 71: 603–9.
- 3 Brinkman DJ, Tichelaar J, van Agtmael MA, de Vries TP, Richir MC. Self-reported confidence in prescribing skills correlates poorly with assessed competence in fourth-year medical students. *J Clin Pharmacol* 2015; 55: 825–30.
- 4 Ryan C, Ross S, Davey P, Duncan EM, Fielding S, Francis JJ, *et al*. Junior doctors' perceptions of their self-efficacy in prescribing, their prescribing errors and the possible causes of errors. *Br J Clin Pharmacol* 2013; 76: 980–7.
- 5 Ryan C, Ross S, Davey P, Duncan EM, Francis JJ, Fielding S, *et al*. Prevalence and causes of prescribing errors: the PRescribing outcomes for trainee doctors engaged in clinical training (PROTECT) study. *PLoS One* 2014; 9: e79802.
- 6 Miguel A, Azevedo LF, Araujo M, Pereira AC. Frequency of adverse drug reactions in hospitalized patients: a systematic review and meta-analysis. *Pharmacoepidemiol Drug Saf* 2012; 21: 1139–54.
- 7 Warlé-van Herwaarden MF, Kramers C, Sturkenboom MC, van den Bemt PM, De Smet PA, Dutch HARM-Wrestling Task Force. Targeting outpatient drug safety: recommendations of the Dutch HARM-wrestling task force. *Drug Saf* 2012; 35: 245–59.
- 8 Maxwell SR, Cameron IT, Webb DJ. Prescribing safety: ensuring that new graduates are prepared. *Lancet* 2015; 385: 579–81.