



Downy Mildew of Alfalfa

Claudia Nischwitz, Associate Professor and Extension Plant Pathologist, Utah State University

What You Should Know

- There are three downy mildew species that can occur on alfalfa. The most common one is *Peronospora trifoliorum*.
- In 2019 we found in Utah the second species that mostly occurs in Europe and Asia - *Peronospora aestivalis*.
- All downy mildew species overwinter in the crown and buds of infected plants.
- Only the first cutting is affected.
- Resistant varieties can help minimize damage.



Fig. 1. Field symptoms of downy mildew in alfalfa

INTRODUCTION

Downy mildew is an obligate parasite. Even though they look like fungi they are not. They belong to the Oomycota and are more closely related to algae than true fungi. The most common alfalfa downy mildew species in the USA is *Peronospora trifoliorum*. In Utah, we found *P. aestivalis*, a species more common in Europe and Asia.



Fig. 2. Close-up of foliar symptoms

SYMPTOMS

The initial symptoms of downy mildew in alfalfa consist of yellow or chlorotic spots on new leaves (Fig. 1 and 2). The spores and fruiting structures (Fig. 3) emerge from the leaves. On the underside the leaves are covered in a white to pale violet colored fungal growth (Fig. 4). Once plants get systemically infected, they can have stems that have a wider diameter and a rosette-like growth at the tip.

plants and infect the leaves. The spores are very fragile and only survive for a few hours to a few days without suitable host tissue to infect. Only young leaf tissue is susceptible to infection. Once the tissue has matured the pathogen can no longer colonize it. Therefore, usually only the first cut of alfalfa is negatively affected. The pathogen is only a problem during cool, wet spring weather. Germination can occur at temperatures between 39-84F. Optimum temperature for germination is 64F. The spores require standing water on leaves for germination (Skinner and Stuteville, 2015).

DISEASE CYCLE

The pathogen overwinters in the cortex of the crown as well as in bud tissue. During the following spring the downy mildew colonizes the infected buds and shoots as they grow and spores develop on the new leaves. The spores are then blown by wind to neighboring alfalfa

MANAGEMENT

Cultural Practices

Cutting

The first cut of alfalfa in the spring should not be delayed because inoculum as well as susceptible host tissue is removed. The downy mildew spores are very short lived and will be dead before new susceptible tissue has grown. Cutting also reduces humidity in the canopy thus reducing the risk of infection.

Resistant varieties

There are resistant varieties available for *P. trifoliorum*. If these varieties are also resistant to *P. aestivalis* is currently unknown. There is no list of varieties that are resistant to downy mildew as this is a characteristic the varieties are not rated for.

Chemical control

Seed treatment

Seed treatments using products containing mefenoxam and metalaxyl can reduce seedling infection of new alfalfa plantings.

REFERENCES

Skinner, D.Z. and Stuteville, D.L. 2015. Downy mildew; in : Compendium of Alfalfa Diseases and Pests. APS Press




Fig. 3. Spores and fruiting structures of downy mildew¹




Fig. 4. Downy mildew growth on underside of leaf

¹Images courtesy of Discoverlife. org

Precautionary Statement: Utah State University Extension and its employees are not responsible for the use, misuse, or damage caused by application or misapplication of products or information mentioned in this document. All pesticides are labeled with active ingredients, directions for use, and hazards, and not all are registered for edible crops. "Restricted use" pesticides may only be applied by a licensed applicator. The pesticide applicator is legally responsible for proper use. USU makes no endorsement of the products listed herein.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Dept. of Ag., Ken White, Vice President for Extension and Agriculture, Utah State University.