

Schiff's Bases Derived from Amino Acids- Imidazole Conjugates as Promising Antioxidant and Antimicrobial Agents

Research Article

BJ Ullas¹, PG Chandrashekar¹, R Suhas², KP Rakesh², P Avinash² and D Channe Gowda^{2*}

¹Department of Chemistry, Yuvaraja's College, University of Mysore, Mysore-570 008, Karnataka, India

²Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore-570 006, Karnataka, India

*Corresponding author: D. Channe Gowda, Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore-570 006, Karnataka, India, Tel.: +91 821 2419664; E-mail: dchannegowda@yahoo.co.in

Copyright: © 2015 Channe G, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article Information: Submission: 29/09/2015; Accepted: 02/11/2015; Published: 10/11/2015

Abstract

A series of amino acids conjugated imidazole derivatives were synthesized and characterized by standard spectroscopical techniques. The title compounds were evaluated for their in vitro antimicrobial activity against bacteria and fungi by zone of inhibition method. The results showed that this skeletal framework exhibited interesting potency as antimicrobial agents. Amongst, compounds 10,11,13,14,16,17 and 18 showed most potent gram negative and gram positive antibacterial activity whereas compounds 11,12,14,15,17 and 26 displayed most potent antifungal activity against *A. niger* and compounds 16,17,18, 21,22 and 24 exhibited good antifungal activity against *F. oxysporum* species. On the other hand, conjugates 11,12,13,14,15,16,17 and 18 showed very good antioxidant activity compared to commercial standard drugs. The results highlight the amino acids conjugated imidazole derivatives as potential leads for designing new future antimicrobial drug candidates possessing good antioxidant properties.

Keywords: Imidazole; Amino acids; Conjugation; Antimicrobial activity.

Introduction

Heterocyclic compounds of nitrogen containing five membered ring systems have been described for their biological activity against various microorganisms [1]. Besides this, the chemistry of imidazoles has also been reviewed in literature and a number of derivatives of imidazoles serve as valuable therapeutic agents [2]. Considerable interest has been created in the chemistry of imidazoles due to their versatile therapeutic activities like antibacterial [3,4], antimalarial [5], antihypertensive [6], antidepressant [7], antitubercular [8], antiviral [9], antiepileptic [10], antirheumatoid arthritis [11], anti-inflammatory [12,13], anticancer activities [14,15] etc. The emergence of powerful and elegant imidazole has stimulated major advances in chemotherapeutic agents of remarkable significance in medicine, biology and pharmacy. Besides this, it is also reported that

imidazole compounds are one of the effective antifungal agents [16]. Considering the importance of imidazole, we planned to incorporate this crucial moiety in the design of our project.

On the other hand, earlier reports have shown that conjugation of different amino acids/ peptides to various biologically active scaffolds has fetched remarkable results which are very promising and even enthusiastic [17-19]. Further, amino acid/ peptide-based drugs have low toxicity, ample bioavailability and permeability, modest potency and good metabolic and pharmacokinetic properties [20]. Prompted by all these observations and with a further interest to develop more biologically active compounds, the present work encompasses the synthesis of novel imidazole-based amino acids analogues as promising antimicrobial and antioxidant agents.

Material and Methods

General

Imidazole 1 was gifted from Jubilant Life Sciences Ltd, all other chemicals and reagents were obtained from Merck (India) and Avra Synthesis (India) which were used directly. Melting point was determined on a Superfit melting point apparatus (India) and is uncorrected. FT-IR was performed using a Jasco spectrometer (Japan) using nujol media. ^1H NMR (400 MHz) and ^{13}C NMR (100 MHz) spectra were recorded on a Agilent Technologies spectrometer (USA) using DMSO (d_6) as solvent. High resolution mass spectroscopic analysis was performed on a Bruker MicroTOF QII mass spectrometer in positive mode. Progress of the reaction was monitored by TLC using silica gel coated on glass plates with the solvent system comprising chloroform/ methanol/ acetic acid in the ratio 98:02:03 (R_f^a) and 95:05:03 (R_f^b) and the compounds on TLC plates were detected by iodine vapors. Chemicals and reagents used for antimicrobial studies were of bacteriological grade.

General procedure for *N*-alkylation

Compound 1 (2.0 g, 0.008 mol) was dissolved in acetone and added base K_2CO_3 (2.3 g, 2 mol) followed by slow addition of 4-chlorobenzyl chloride (1.33 g, 0.008 mol). Reaction was maintained for 10 h at reflux (monitored by TLC), filtered K_2CO_3 , evaporated solvent, cooled and recrystallized *N*-alkylated product 2 from ethanol.

General procedure for hydrolysis

Compound 2 (2.0 g, 0.0054 mol) was dissolved in methanol, added 0.1 N NaOH solution (0.43 g, 0.011 mol) and the reaction was maintained for 5 h at 50 °C. Completion of reaction was monitored by TLC, cooled and pH was adjusted to 1-2 using dil. HCl solution. Compound was extracted into DCM, washed with water, evaporated solvent to obtain compound 3.

General procedure for conjugation of 3 with different amino acids (HCl.NH₂-Xaa-COOMe) where Xaa = Phe, Tyr, Trp, Pro, Met and Lys

To compound 3 (0.002 mol) and HOBT (0.305 g, 0.002 mol) dissolved in DMF (10 mL/g of compound) cooled to 0 °C was added NMM (0.55 mL, 0.002 mol). EDCI (0.383 g, 0.005 mol) was added under stirring while maintaining the temperature at 0 °C. Reaction mixture was stirred for an additional 10 min and a pre-cooled solution of HCl.NH₂-Xaa-COOMe (0.631 g, 0.002 mol) and NMM (0.25 mL, 0.002 mol) in DMF (10 mL) was added slowly. After 20 min, pH of the solution was adjusted to 8 by the addition of NMM and the reaction mixture was stirred over night at room temperature. DMF was removed under reduced pressure and the residue was poured into about 200 mL of ice-cold 90% saturated KHCO_3 solution and stirred for 30 min. The precipitated product was taken into CHCl_3 and washed with 5% NaHCO_3 solution (2×100 mL), water (1×100 mL), HCl (2×100 mL) and brine (1×100 mL). The organic layer was dried over anhydrous Na_2SO_4 and solvent was removed under reduced pressure. The products so obtained were recrystallized from ether/petroleum ether to get conjugates 4 (Phe), 5 (Tyr), 6 (Trp), 7 (Pro), 8 (Met) and 9 (Lys).

General procedure for the synthesis of hydrazides

Compound 4, 5, 6, 7, 8 and 9 (0.3mmol) was dissolved

separately in ethanol (2 mL) and hydrazine hydrate (0.03 mL, 2 mol) was added slowly and the reaction was maintained at reflux for 12 h. Completion of the reaction was monitored by TLC, evaporated solvent, cooled and poured into ice cold water. Solid separated was filtered and dried to obtain 4a, 5a, 6a, 7a, 8a and 9a respectively.

General procedure for the condensation of aldehydes to 4a, 5a, 6a, 7a, 8a and 9a

To compound 4a, 5a, 6a, 7a, 8a and 9a (0.3 mmol) and benzaldehyde (0.3 mmol) dissolved in ethanol (10 mL/g) was added few drops of acetic acid as a catalyst which was maintained at reflux for 15 h. Completion of the reaction was monitored by TLC, solvent was evaporated, cooled and poured into ice cold water precipitate is obtained and then filtered, washed with cold water to obtain compound (10-27).

Antibacterial activity

In vitro antibacterial activity was evaluated against human pathogens of both gram positive organisms namely *X. oryzae* and gram negative organisms namely *E. coli* by agar well diffusion method [21].

The microorganisms were inoculated in to the sterilized nutrient broth and maintained at 37 °C for 24 h. On the day of testing, bacteria were subcultured separately into 25 mL of sterilized nutrient broth. Inoculated subcultured broths were kept at room temperature for the growth of inoculums. Each test compound (10-27) and standard drug of 10 mg was dissolved in 10 mL of DMSO to get a concentration of 1 µg/mL and further diluted to get a final concentration of 50 µg/mL. About 15-20 mL of molten nutrient agar was poured into each of the sterile plates. With the help of cork borer of 6mm diameter, the cups were punched and scooped out of the set agar and the plates were inoculated with the suspension of particular organism by spread plate technique. The cups of inoculated plates were then filled with 0.1 mL of the test solution, Chloromphenicol solution and DMSO (negative control). The plates were allowed to stay for 24 h at 37 °C and zone of inhibition (mm) was then measured.

Antifungal activity

In vitro antifungal activity was evaluated against two fungal species namely *A.niger*, and *F. oxysporum* by agar well diffusion method [22].

The fungal strains were subcultured separately into 25 mL of sterilized nutrient broth and compounds and standard drug (bavistin) of 10 mg was dissolved in 10 mL of DMSO to get a concentration of 1 mg/ml and further diluted to get a final concentration of 50 µg/mL. Molten media of sabouraud agar of 10-15 mL was poured into the petri plates and allowed to solidify. Fungal subculture was inoculated on the solidified media. With the help of 6 mm cork borer, the cups were punched and scooped out of the set agar. The cups of inoculated plates were then filled with 0.1 mL of the test solution, bavistin solution and DMSO (negative control). The plates were allowed to stay for 3 days at room temperature and zone of inhibition (mm) was then measured.

Antioxidant activities

DPPH (1,1-diphenyl-2-picryl-hydrazyl) assay

The radical scavenging activity of DPPH free radicals by synthesized compounds were determined according to the reported method [23]. Briefly, 50 μL of test compounds was mixed at different concentrations (25, 50, 100, 200 and 300 $\mu\text{g}/\text{mL}$) with 1 mL of 0.1 mM DPPH in methanol solution and 450 μL of 50 mM Tris HCl buffer (pH 7.4). Methanol (50 μL) only was used as the experimental control. After 30 min of incubation at room temperature, reduction in the number of DPPH free radicals was measured by reading absorbance at 517 nm. BHT (butylatedhydroxytoluene) was used as control similar to test concentrations. Percent inhibition was calculated from the following equation:

$$\% \text{ Inhibition} = \left[\frac{\text{Absorbance of control} - \text{Absorbance of test sample}}{\text{Absorbance of control}} \right] \times 100$$

ABTS (2,2-azinobis-(3-ethylbenzothiazoline-6-sufonic acid) assay

The ability of the test sample to scavenge ABTS^+ radical cation was determined according to the literature method [24] with slight modification. The ABTS^+ radical cation was pregenerated by mixing 7 mM ABTS^+ stock solution with 2.45 mM potassium persulfate (final concentration) and incubating for 12-16 h in dark at room temperature until the reaction was complete and absorbance was stable. Absorbance of the ABTS^+ solution was equilibrated to 0.70 (± 0.02) by diluting with distilled water at room temperature, then 2 mL was mixed with different concentration of the test sample (25, 50, 100, 200, and 300 $\mu\text{g}/\text{mL}$) and the absorbance was measured at 734 nm after 6 min. The scavenging capability of ABTS^+ radical was calculated using the following equation:

$$\text{ABTS}^+ \text{ scavenging effect (\%)} = [(A_c - A_s) / A_c] \times 100$$

Where, A_c is the initial concentration of the ABTS^+ and A_s is the absorbance of the remaining concentration of ABTS^+ in the presence of compounds.

DMPD (N, N-dimethyl-p-phenylenediamine) assay

The DMPD radical scavenging ability of synthesized compounds was determined by the Fogliano et al., method [25] with slight modifications by Gulcin [26]. This assay is based on the capacity of the extract to inhibit DMPD^+ cation radical formation. Briefly, 105 mg of DMPD was dissolved in 5 mL of distilled water. Then, 1 mL of this solution was added to 100 mL of 0.1 M acetate buffer (pH 5.3). DMPD^+ was produced by adding 0.3 mL ferric chloride (0.05 M) to this solution. Different concentrations of standard antioxidants or synthesized compounds (25-300 $\mu\text{g}/\text{mL}$) were added, and the total volume was adjusted to 1 mL with distilled water. One millilitre of the DMPD^+ solution was directly added to the reaction mixture. The reaction mixtures were incubated in the dark for 15 min. The absorbance was measured at 505 nm and percent inhibition was calculated according to the formula:

$$\% \text{ Inhibition} = \left[\frac{\text{Absorbance of control} - \text{Absorbance of test sample}}{\text{Absorbance of control}} \right] \times 100$$

Results and Discussion**Chemistry**

Starting material 1 was subjected to *N*-alkylation using benzyl chloride that resulted in 2 which was ascertained by the absence of NH proton peak. Product 2 on saponification yielded 3 which exhibited carboxylic acid peak in its PMR spectrum. This compound was conjugated with different amino acids to obtain 4-9 which did not have COOH peak and showed -NH peak at $\delta \sim 8.00$ (1H, s) in ^1H NMR spectra. Compounds 4-9 was separately reacted with hydrazine hydrate to get 4a-9a which was confirmed by disappearance of $-\text{OCH}_3$ peak. The final imines 10-27 were synthesized by condensation of 4a-9a with substituted aldehydes (Scheme). Synthesis were confirmed by the presence of all requisite peaks and absence of extraneous peaks in ^1H NMR and ^{13}C NMR spectra and also mass values were in accordance with the mol. wt. calculated. The analytical and spectroscopic data of the synthesized compounds are given in Table 1.

Biological evaluation**Antimicrobial activity**

The Schiff bases of imidazole-amino acid conjugates were evaluated for their antibacterial studies against both gram-positive and gram-negative bacteria namely *X. oryzae* and *E. coli* respectively and antifungal studies against *A. niger* and *F. oxysporum*. Synthesized compounds exhibited a significant broad spectrum inhibitory effect against all the tested microorganisms. The result obtained as zone of inhibition (mm) is presented in Table 2. The standard drugs chloromphenicol and bavistin were used as standards for antibacterial and antifungal activities respectively.

Previous reports have shown that there is no significant activity with just imidazoles linked amino acids [27,28]. Further, derivatives of amino acids-heterocycle conjugates have exhibited promising results and in continuation it was decided to introduce different aldehydes *viz.* dihydroxy, trihydroxy and trimethoxy benzaldehyde in the present study to obtain imines 10-27. Among the twenty naturally occurring amino acids, selective moieties like Phe, Trp and Tyr (aromatics); Pro (an imine); Met (thio) and Lys (basic) were used in the current investigation. Amongst, compounds 10-18, 21 and 27 appeared to be more active than the standards which may be due to the presence of aromatic side chains in Phe, Trp and Tyr and also could be due to more hydrophobicity of these amino acids along with the presence of more electron donating groups like OH and OCH_3 present on the other end of the molecules. Together these parameters would have contributed to the superior activity. And also it is observed that compounds with trihydroxy benzaldehyde derivatives have shown enhanced activity compared to rest of the analogues which imparts that these hydroxyl functions play a key role in enhancing the activity. On the other hand, compounds containing thio group (22-24) have shown slightly decreased activity than Pro (19-21) which are further less active than Lys (25-27) analogues and

Table 1: Physical and analytical data of the conjugated compounds (10-27)

Entry	R _f value	Yield (%)	M.P. (°C)	Theoretical Mol. Wt.	Actual MS	IR (ν cm ⁻¹) and NMR (CDCl ₃ , δ ppm)
10	0.61	89	103-105	617.10	618.3	IR: 3250 (NH), 1690 (CO), 1590 (-HC=N-) ¹ H NMR: 0.90 (3H,t,-CH ₃ of propyl), 1.57 (6H,s,-CH ₃), 1.68 (2H,m,-CH ₂ of propyl), 2.45 (2H,t,-CH ₂ of propyl), 2.92 (2H,t,β-CH ₂), 4.92 (1H,t,α-CH), 4.97 (2H,s,-CH ₂ of benzyl), 5.0 (2H, s,-OH), 6.66-7.22 (12H,m,Ar-H), 8.1 (1H,s,-HC=N-), 8.8 (2H,d,-NH) ¹³ C NMR: 13.6, 24.4, 28.5, 31.1, 37.8, 40.1, 55.2, 69.7, 115.8, 115.8., 116.2, 117.5, 123.2, 126.4, 128.2, 128.7, 130.9, 131.2, 133.9, 138.7, 143.1, 147.2, 148.6, 150.1, 155.6, 156.5, 160.6, 177.2
11	0.59	91	101	633.30	634.5	IR: 3260 (NH), 1695 (CO), 1620 (-HC=N-) ¹ H NMR: 0.93 (3H,t,-CH ₃ of propyl), 1.54 (6H,s,-CH ₃), 1.70 (2H,m,-CH ₂ of propyl), 2.48 (2H,t,-CH ₂ of propyl), 2.89 (2H,t,β-CH ₂), 4.88 (1H,t,α-CH), 4.90 (2H,s,-CH ₂ of benzyl), 5.6 (3H, s,-OH), 6.70-7.30 (11H,m,Ar-H), 8.1 (1H,s,-HC=N-), 8.6 (2H,d,-NH) ¹³ C NMR: 14.2, 26.2, 27.6, 33.1, 38.4, 41.2, 55.8, 66.6, 116.4, 117.5, 118.8, 119.2, 121.6, 128.6, 128.9, 130.5, 131.4, 131.9, 133.2, 138.4, 140.9, 146.0, 149.5, 151.4, 154.1, 156.0, 161.4, 175.8
12	0.62	92	113-114	675.0	676.1	IR: 3258 (NH), 1680 (CO), 1640 (-HC=N-) ¹ H NMR: 1.0 (3H,t,-CH ₃ of propyl), 1.55 (6H, s, -CH ₃), 1.69 (2H,m,-CH ₂ of propyl), 2.31 (2H,t,-CH ₂), 2.42 (2H,t,-CH ₂ of propyl), 3.09 (1H,m,β-CH ₂), 3.53 (9H,s,-OMe), 4.95 (1H,α,-CH), 4.92 (2H,s,-CH ₂ of benzyl), 6.6-7.22 (11H,m,Ar-H), 8.2 (2H,d,NH), 8.2 (1H,s,-HC=N-) ¹³ C NMR: 14.5, 26.0, 27.9, 30.1, 38.4, 41.2, 53.7, 56.2, 63.6, 105.1, 114.8, 126.1, 126.6, 129.1, 132.2, 133.8, 134.0, 134.4, 136.9, 140.2, 143.2, 150.5, 150.8, 151.8, 156.0, 156.4, 176.6
13	0.55	85	120-122	633.5	634.5	IR: 3264 (NH), 1691 (CO), 1640 (-HC=N-) ¹ H NMR: 0.92 (3H,t,-CH ₃ of propyl), 1.60 (6H,s,-CH ₃), 1.64 (2H,m,-CH ₂ of propyl), 2.41 (2H,t,-CH ₂ of propyl), 2.92 (2H,t,β-CH ₂), 4.92 (1H,t,α-CH), 4.99 (2H,s,-CH ₂ of benzyl), 5.2 (3H, s,-OH), 6.66-7.18 (11H,m,-Ar-H), 8.0 (1H,s,-HC=N-), 8.3 (2H,d,-NH) ¹³ C NMR: 13.7, 24.2, 28.3, 31.0, 37.9, 40.2, 55.3, 69.8, 115.7, 115.8., 116.4, 117.4, 123.2, 128.8, 129.2, 130.5, 131.3, 132.1, 134.4, 138.6, 143.0, 147.4, 149.6, 150.7, 155.7, 156.4, 160.8, 177.0
14	0.58	92	Gum	649.10	650.5	IR: 3245 (NH), 1698 (CO), 1630 (-HC=N-) ¹ H NMR: 0.93 (3H,t,-CH ₃ of propyl), 1.54 (6H,s,-CH ₃), 1.70 (2H,m,-CH ₂ of propyl), 2.48 (2H,t,-CH ₂ of propyl), 2.89 (2H,t,β-CH ₂), 4.88 (1H,t,α-CH), 4.93 (2H,s,-CH ₂ of benzyl), 5.6 (4H, s,-OH), 6.70-7.30 (10H,m,-Ar-H), 8.1 (1H,s,-HC=N-), 8.6 (2H,d,-NH) ¹³ C NMR: 13.9, 24.8, 28.7, 32.0, 37.7, 40.3, 54.9, 69.8, 115.4, 115.5, 115.9, 117.2, 122.8, 128.6, 129.1, 130.6, 131.4, 131.9, 134.3, 138.5, 139.9, 147.2, 149.5, 150.6, 155.7, 156.2, 160.6, 177.2
15	0.70	88	122	691.20	692.2	IR: 3246 (NH), 1686 (CO), 1595 (-HC=N-) ¹ H NMR: 1.0 (3H,t,-CH ₃ of propyl), 1.55 (6H, s, -CH ₃), 1.69 (2H,m,-CH ₂ of propyl), 2.31 (2H,t,-CH ₂), 2.42 (2H,t,-CH ₂ of propyl), 3.09 (1H,m, β-CH ₂), 3.53 (9H,s,-OMe), 4.95 (1H,α,-CH), 4.98 (2H,s,-CH ₂ of benzyl), 5.40 (1H,s,-OH), 6.6-7.22 (11H,m,Ar-H), 8.2 (2H,d,NH), 8.2 (1H,s,-HC=N-) ¹³ C NMR: 13.5, 24.3, 28.5, 31.3, 37.8, 40.1, 55.4, 56.2, 64.8, 106.5, 115.8, 128.0, 128.6, 129.2, 130.4, 131.5, 132.0, 134.4, 138.6, 141.5, 143.0, 150.7, 150.9, 150.9, 155.7, 156.4, 177.5
16	0.66	96	Gum	657.14	658.5	IR: 3266(NH),1670 (CO), 1590 (-HC=N-) ¹ H NMR: 0.82 (3H,t,-CH ₃ of propyl), 1.37 (6H,s,-CH ₃), 1.68 (2H,m,-CH ₂ of propyl), 2.45 (2H,t,-CH ₂ of propyl), 3.06 (2H,d,β-CH ₂), 4.92 (1H, t,δ -CH), 5.40 (2H,s,-OH), 4.99 (2H,s,- CH ₂ of benzyl), 6.6-7.18 (11H,m,Ar-H), 8.4 (1H,s,-HC=N-), 10.1 (2H,t,-NH) ¹³ C NMR: 12.9, 22.2, 28.6, 32.0, 31.6, 40.4, 56.4, 69.8, 110.9, 111.1, 116.4, 116.9, 118.2, 120.6, 121.1, 122.2, 123.1, 127.6, 127.9, 128.6, 130.4, 131.3, 134.4, 136.4, 138.6, 143.0, 147.3, 149.6, 150.4, 156.4, 160.4, 178.0
17	0.57	86	89-90	672.2	673.5	IR: 3261 (NH), 1675 (CO), 1625 (-HC=N-) ¹ H NMR: 0.88 (3H,t,-CH ₃ of propyl), 1.34 (6H,s,-CH ₃), 1.64 (2H,m,-CH ₂ of propyl), 2.45 (2H,t,-CH ₂ of propyl), 3.16 (2H,d,β-CH ₂), 4.90 (1H, t,δ -CH), 5.39 (3H,s,-OH), 4.99 (2H,s,-CH ₂ of benzyl), 6.6-7.18 (10H,m,Ar-H), 8.1 (1H,s,-HC=N-), 10.1 (2H,t,-NH) ¹³ C NMR: 11.7, 23.1, 27.5, 31.6, 31.9, 40.1, 54.2, 66.5, 109.0, 116.4, 118.2, 120.6, 121.1, 122.2, 127.6, 127.9, 128.6, 129.2, 130.4, 131.3, 134.4, 136.4, 143.0, 143.1, 146.7, 148.5, 151.2, 155.1, 159.0, 176.2

18	0.60	90	96	714.29	715.4	IR: 3258 (NH),1660 (CO), 1605 (-HC=N-) ¹ H NMR: 0.82 (3H,t,-CH ₃ of propyl), 1.42 (6H,s,-CH ₃), 1.55 (2H,m,-CH ₂ of propyl), 2.51(2H,t,-CH ₂ of propyl), 3.16 (2H,d,β-CH ₂), 3.23 (2H,d,β-CH ₂), 3.77 (9H,s,-OCH ₃), 4.88 (1H,t,α-CH), 5.0 (2H,s,-CH ₂ of benzyl), 6.3-7.02 (10H,m,Ar-H), 8.2(1H,s,-HC=N-), 9.36 (2H,d,-NH) ¹³ C NMR:13.8,23.6,29.0,30.5,31.3,39.8,56.0,69.8, 106.7,110.9,111.1,120.0,122.2,122.9,127.5,128.1,128.8,130.5,131.3,134.4,136.5,138.6,141.5,143.0,150.7,150.9,160.8,173.0
19	0.55	92	102	567.10	568.4	IR: 3244 (NH), 1685(CO), 1650 (-HC=N-) ¹ H NMR: 0.86 (3H,t,-CH ₃ of propyl), 1.54 (2H,m,β -CH ₂), 1.65 (6H,s,-CH ₃), 1.70 (2H,m,α-CH ₂), 1.72 (2H,m,-CH ₂ of propyl), 2.56 (2H,t,-CH ₂ of propyl), 3.29 (2H,t,α -CH ₂), 4.29 (1H,t,δ-CH ₂), 5.2 (2H,s,-CH ₂ of benzyl), 5.40 (2H,s, -OH), 6.66-7.18 (7H,m,Ar-H), 8.1 (1H,s,-HC=N-) 8.6 (1H,d,-NH) ¹³ C NMR: 12.7, 24.1, 26.0, 31.0, 32.1, 37.0, 55.0, 65.4, 108.6, 110.9, 116.2, 116.8, 119.0, 122.0, 122.3, 122.8, 123.0, 126.5, 127.0, 128.8, 130.4, 131.0, 133.8, 135.5, 137.5, 142.0, 148.2, 149.1, 150.7, 160.6, 169.0
20	0.60	90	109-110	583.20	584.5	IR : 3256 (NH),1669 (CO), 1635 (-HC=N-) ¹ H NMR : 1.02 (3H,t,-CH ₃ of propyl), 1.52 (2H,m,β-CH ₂), 1.56 (6H,s,-CH ₃), 1.68 (2H,m,-CH ₂ of propyl), 1.71 (2H,m,γ-CH ₂), 2.32 (2H,t,-CH ₂ of propyl), 3.30 (2H,s,-CH ₂ of propyl), 4.29 (1H,t,δ-CH), 5.03 (3H,s,-OH), 6.5-7.15 (6H,s,Ar-H), 8.0 (1H,s,-NH), 8.3 (1H,s,-HC=N-) ¹³ C NMR: 15.1, 20.4, 24.2, 25.1, 28.3, 39.6, 55.0, 64.1, 106.3, 110.4, 115.8, 118.5, 121.3, 126.4, 128.3, 129.4, 130.1, 132.1, 134.1, 138.3, 138.6, 140.4, 148.8, 152.2, 154.3, 164.8, 172.7
21	0.68	93	Gm	625.27	626.2	IR: 3244 (NH),1692 (CO), 1625 (-HC=N-) ¹ H NMR: 0.99 (3H,t,-CH ₃ of propyl), 1.49 (2H,m,β-CH ₂), 1.57 (6H,s,-CH ₃), 1.62 (2H,m,-CH ₂ of propyl), 1.72 (2H,m,γ-CH ₂), 2.30 (2H,t,-CH ₂ of propyl), 3.30 (2H,s,-CH ₂ of propyl), 4.31 (1H,t,δ-CH), 5.13 (9H,s,-OMe), 6.5-7.15 (6H,s,Ar-H), 8.1 (1H,s,-NH), 8.4 (1H,s,-HC=N-) ¹³ C NMR: 15.0, 20.2, 30.3, 30.6, 31.2, 39.3, 54.2, 55.0, 68.0, 106.7, 110.8, 113.0, 118.0, 119.1, 122.2, 122.9, 128.8, 128.1, 131.3, 134.4, 138.4, 141.5, 143.0, 150.9, 150.9, 160.8, 168.0
22	0.55	94	Gum	587.2	588.6	IR: 3267 (NH),1700 (CO), 1610 (-HC=N-) ¹ H NMR: 0.99 (3H,t,-CH ₃ of propyl), 1.39 (6H,s,-CH ₃), 1.62 (2H,m,-CH ₂ of propyl), 2.09 (3H,s,δ-CH ₃), 2.53 (2H,t,-CH ₂ of propyl), 2.92 (2H,d,β-CH ₂), 4.92 (1H,t,α-CH), 4.99 (2H,s,-CH ₂ of benzyl), 5.13 (2H,s,-OH), 6.6-7.10 (7H,m,Ar-H), 8.15 (1H,s,-HC=N-) 8.21 (1H,s,-NH) ¹³ C NMR: 14.8, 17.1, 24.7, 27.3, 30.1, 36.5, 40.0, 50.6, 69.8, 114.4, 115.2, 116.4, 121.1, 123.2, 126.8, 127.8, 128.9, 130.3, 131.0, 134.4, 138.6, 147.4, 149.6, 150.7, 156.4, 166.9
23	0.56	88	Gum	603.19	604.3	IR : 3270 (NH),1695 (CO), 1595 (-HC=N-) ¹ H NMR: 1.1 (3H,t,-CH ₃ of propyl), 1.35 (6H,s,-CH ₃), 1.58 (2H,m,-CH ₂ of propyl), 2.16 (3H,s,δ-CH ₃), 2.54 (2H,t,-CH ₂ of propyl), 2.88 (2H,d,β-CH ₂), 4.88 (1H,t,α-CH), 4.96 (2H,s,-CH ₂ of benzyl),5.11 (3H,s,-OH), 6.6-7.10 (6H,m,Ar-H), 8.15 (1H,s,-HC=N-), 8.20 (1H,s,-NH) ¹³ C NMR: 14.5, 21.2, 29.3, 32.2, 32.6, 47.9, 51.9, 71.2, 109.0, 115.2, 118.1, 119.4, 128.8, 130.5, 138.2, 143.0, 147.6, 148.5, 150.1, 151.2, 154.1, 160.8, 171.6
24	0.60	92	108	645.24	646.3	IR: 3240 (NH),1689 (CO), 1645 (-HC=N-) ¹ H NMR: 0.90 (3H,t,-CH ₃ of propyl), 1.35 (6H,s,-CH ₃), 1.58 (2H,m,-CH ₂ of propyl), 2.16 (3H,s,δ-CH ₃), 2.54 (2H,t,-CH ₂ of propyl),3.07 (2H,d,β-CH ₂), 4.88 (1H,t,α-CH), 4.96 (2H,s,-CH ₂ of benzyl), 5.20 (9H,s,-OCH ₃), 6.6-7.10 (6H,m,Ar-H), 8.0 (1H,s,-HC=N-), 8.10 (1H,s,-NH) ¹³ C NMR: 11.6, 17.2, 24.2, 26.1, 31.1, 31.4, 52.8, 56.2, 67.2, 106.7, 114.1, 116.7, 128.1, 131.3, 134.4, 141.0, 144.8, 147.7, 146.0, 149.6, 150.9, 154.0, 161.5, 169.5
25	0.56	91	118	686.27	687.3	IR: 3243 (NH),1695 (CO), 1630 (-HC=N-) ¹ H NMR: 0.88 (3H,t,-CH ₃ of propyl), 1.29 (2H,m,γ-CH ₂), 1.36 (6H,s,-CH ₃), 1.37 (2H,m,-CH ₂ of propyl), 1.41 (2H,m,δ-CH ₂), 1.96 (2H,t,-CH ₂ of propyl), 2.55 (2H,t,ω-CH ₂), 2.46 (2H,s,-CH ₂ of benzyl), 3.81 (2H,d,-CH ₂ of benzyl), 4.51 (1H,t,-CH), 5.12 (2H,s,-OH), 6.6-7.22 (12H,m,Ar-H), 8.0 (3H,s,-NH), 8.1 (1H,s,-HC=N-) ¹³ C NMR: 13.6, 23.1, 26.3, 32.2, 32.6, 37.0, 51.9, 52.8, 56.2, 69.8, 116.3, 117.0, 118.4, 126.0, 127.8, 128.7, 134.3, 138.6, 139.5, 148.7, 149.0, 150.7, 156.4, 162.4, 170.3

26	0.59	86	Gum	702.32	703.3	IR: 3260 (NH), 1698 (CO), 1620 (-HC=N-) ¹ H NMR: 0.96 (3H,t,-CH ₃ of propyl), 1.30 (2H,m,γ-CH ₂), 1.33 (6H,s,-CH ₃), 1.27 (2H,m,-CH ₂ of propyl), 1.42 (2H,m,δ-CH ₂), 1.88 (2H,t,-CH ₂ pf propyl), 2.45 (2H,t,ω,-CH ₂), 2.54 (2H,s,-CH ₂ of benzyl), 3.80 (2H,d,-CH ₂ of benzyl), 4.49 (1H,t,-CH), 5.10 (3H,s,-OH), 6.6-7.22 (12H,m,Ar-H), 8.2 (3H,s,-NH), 8.3 (1H,s,-HC=N-) ¹³ C NMR: 13.7, 22.6, 24.2, 28.3, 28.9, 31.0, 32.6, 45.7, 56.2, 58.7, 68.5, 69.8, 116.3, 117.3, 118.5, 127.2, 127.7, 129.0, 134.3, 138.6, 141.2, 148.7, 149.0, 150.7, 156.4, 165.4, 171.6
27	0.60	90	128	744.37	745.5	IR: 3256 (NH), 1696 (CO), 1610 (-HC=N-) ¹ H NMR: 0.90 (3H,t,-CH ₃ of propyl), 1.33 (2H,m,γ-CH ₂), 1.35 (6H,s,-CH ₃), 1.38 (2H,m,-CH ₂ of propyl), 1.41 (2H,m,δ-CH ₂), 1.96 (2H,t,-CH ₂ of propyl), 2.55 (2H,t,ω,-CH ₂), 2.46 (2H,s,-CH ₂ of benzyl), 3.81 (2H,d,-CH ₂ of benzyl), 4.51 (1H,t,-CH), 5.05 (9H,s,-CH ₃), 6.6-7.22 (12H,m,Ar-H), 8.0 (3H,s,-NH), 8.1 (1H,s,-HC=N-) ¹³ C NMR: 12.5, 24.1, 27.3, 31.8, 33.4, 37.0, 51.9, 52.8, 54.3, 68.6, 115.8, 116.1, 118.2, 118.5, 129.2, 132.1, 135.3, 138.4, 148.7, 149.6, 151.6, 155.7, 156.2, 160.8, 171.2

Table 2: Antimicrobial and Antioxidant activity of synthesized derivatives


Entry	Antibacterial activity		Antifungal activity		Antioxidant activity		
	Zone of inhibition (mm) ± SD (n=3)				IC ₅₀ µg/ml		
	<i>E. coli</i>	<i>X. oryzae</i>	<i>A. niger</i>	<i>F. oxysporum</i>	DPPH	ABTS	DMPD
1	3±0.12	2±0.63	4±0.42	3±0.40	220±2.01	175±2.15	180±1.45
10	16±0.52	14±0.20	15±0.12	5±0.20	40±0.98	40±0.46	45±0.47
11	18±0.20	12±0.52	18±0.22	7±0.52	25±0.66	35±0.79	35±0.52
12	15±0.50	14±0.35	16±0.52	8±0.2	35±1.02	40±0.64	40±0.95
13	17±0.50	14±0.41	14±0.44	5±0.10	35±0.89	35±0.43	40±0.15
14	20±0.41	16±0.50	20±0.36	6±0.40	20±0.49	30±0.63	30±0.65
15	16±0.46	10±0.15	18±0.10	6±0.25	35±0.65	40±0.14	35±0.24
16	20±0.45	14±0.15	15±0.40	10±0.33	30±0.78	35±0.57	35±0.65
17	22±0.47	16±0.25	16±0.25	12±0.40	20±0.61	25±0.69	20±0.54
18	19±0.41	15±0.26	10±0.33	11±0.25	30±0.43	35±0.52	40±0.26
19	12±0.50	10±0.30	14±0.40	9±0.31	60±0.55	45±0.48	40±0.54
20	14±0.20	14±0.20	16±0.25	9±0.10	55±1.04	35±0.33	35±0.25
21	11±0.40	11±0.10	10±0.33	10±0.40	70±1.22	45±0.49	40±0.45
22	10±0.45	8±0.20	9±0.12	11±0.25	95±1.69	50±0.49	80±0.59
23	10±0.36	7±0.10	12±0.22	10±0.33	75±1.64	45±0.43	65±0.65
24	9±0.30	6±0.20	11±0.30	8±0.40	80±1.69	50±0.58	90±0.45
25	13±0.35	9±0.20	13±0.33	9±0.25	65±0.98	60±0.55	80±0.26
26	14±0.30	12±0.20	15±0.35	7±0.32	60±0.24	75±1.02	85±0.47
27	12±0.35	10±0.26	16±0.44	9±0.21	70±0.66	85±1.54	95±1.23
Chloromphenicol	14±0.65	10±0.24	-	-	-	-	-
Bavistin	-	-	13±0.94	8±0.15	-	-	-
BHA	-	-	-	-	55±0.52	45±0.54	50±0.56
AA	-	-	-	-	55±0.69	50±0.66	60±0.42
GA	-	-	-	-	60±0.48	50±0.49	55±0.26

this trend in activity reveals that presence of basic moiety is more helpful in obtaining good results. Further these compounds are found to be more selective in arresting the growth of *A. niger* fungus than *F. oxysporum*.

In vitro antioxidant activities of all the synthesized compounds were evaluated by (i) 1,1-diphenyl-2-picryl-hydrazyl (DPPH) assay, an convenient technique for screening the antioxidant activities of

the antioxidants; (ii) 2,2-azinobis-3-ethylbenzothiazoline-6-sufonic acid (ABTS) cation radical assay which is a conventional and excellent model for assessing the antioxidant activities of hydrogen donating and chain breaking antioxidants [29]; and (iii) *N,N*-dimethyl-*p*-phenylenediamine dihydrochloride (DMPD) cation radical assay which is similar to the DPPH radical scavenging assay. The IC₅₀ values is the effective concentration at which 50% of the

Scheme: Schematic for target compounds



Reagents and Conditions: (i) 4-Chloro benzyl chloride, K_2CO_3 , acetone, reflux, 12 h (ii) 0.1 N NaOH, MeOH, 50-55 °C, 5 h (iii) EDCl, HOBt, NMM, 0-5 °C, 15 h (iv) Hydrazine hydrate, EtOH, reflux, 10 h (v) Substituted aldehydes, EtOH, AcOH, reflux, 14 h

Comp. No.	Xaa	R ¹	R ²	R ³
10		OH	OH	H
11		OH	OH	OH
12		OCH ₃	OCH ₃	OCH ₃
13		OH	OH	H
14		OH	OH	OH
15		OCH ₃	OCH ₃	OCH ₃
16		OH	OH	H
17		OH	OH	OH
18	OCH ₃	OCH ₃	OCH ₃	
19		OH	OH	H
20		OH	OH	OH
21		OCH ₃	OCH ₃	OCH ₃
22		OH	OH	H
23		OH	OH	OH
24		OCH ₃	OCH ₃	OCH ₃
25		OH	OH	H
26		OH	OH	OH
27		OCH ₃	OCH ₃	OCH ₃

radicals were scavenged, were calculated to evaluate the antioxidant activities. A lower IC_{50} value indicate greater antioxidant activity. IC_{50} values of lower than 10 mg/mL usually implied effective activities in antioxidant properties [30]. The IC_{50} of butylatedhydroxylanisole (BHA), ascorbic acid (AA) and gallic acid (GA) was also determined for comparison. The results are presented in Table 2.

Most of the synthesized analogues showed potent antioxidant activities especially 11, 12, 13, 14, 15, 16, 17 and 18 showed excellent radical scavenging activities with IC_{50} values 25, 35, 35, 20, 35, 30, 20 and 30 $\mu\text{g/mL}$ respectively in DPPH assay much better than the standard BHA ($IC_{50} = 55 \mu\text{g/mL}$). In ABTS⁺ radical scavenging assay, the compounds 11, 12, 13, 14, 15, 16, 17 and 18 showed potent antioxidant activity with IC_{50} values 35, 40, 35, 30, 40, 35, 25 and 35 $\mu\text{g/mL}$ respectively which is much better than the commercial standards AA ($IC_{50} = 55 \mu\text{g/mL}$). The compounds 11, 12, 13, 14, 15,

16, 17 and 18 also exhibited striking antioxidant activity with IC_{50} values 35, 40, 40, 30, 35, 35, 20 and 40 $\mu\text{g/mL}$ respectively which is better than the standards BHA ($IC_{50} = 50 \mu\text{g/mL}$), AA ($IC_{50} = 60 \mu\text{g/mL}$) and GA ($IC_{50} = 55 \mu\text{g/mL}$) in DMPD assay. In all the three assays performed, compounds 11, 12, 13, 14, 15, 16, 17 and 18 showed excellent antioxidant activities with IC_{50} values much lower than the standards. The IC_{50} values of these compounds have antioxidant activities lower than 10 mg/mL demonstrating greater capacity in all the three assays. On the basis of the above observation, compounds having -OH (phenolic), -indole and phenyl ring (11, 12, 13, 14, 15, 16, 17 and 18) were found to be the most potent antioxidants. Conjugates with alkyl, imino and thio group showed least antioxidant activity.

Conclusion

A novel series of Schiff base analogues of amino acids-imidazole conjugates has been presented in this work. These compounds were evaluated for antibacterial and antifungal activities by agar well diffusion method. Activity profile revealed that compounds containing aromatic amino acids have shown profound results along with the beneficiary electron loving hydroxyl groups. Presence of thio (Met) and imino (Pro) amino acids have shown moderate activity compared to Lys, a basic amino acid. These target molecules have exhibited selective activity against *A. niger* fungal species and hence these could be used to treat fungal infections. In case of antioxidant activity compounds containing phenyl ring, indole ring and -OH (phenolic) group showed remarkable activity which may be due to electron donating activity and resonating property of phenyl ring. Compounds like 22, 23, 24, 25, 26, and 27 showed least activity which could be due to long alkyl group and also may be due to thio group in methionine.

References

1. Yu YB, Chen HL, Wang LY, Chen XZ, Fu B (2009) A Facile Synthesis of 2,4-Disubstituted Thiazoles Using MnO_2 . *Molecules* 14: 4858-4865.
2. Grimmett MR (1970) *Advances in Heterocyclic Chemistry*. Academic Press, New York, 241-326.
3. Jain AK, Ravichandran V, Sisodiya M, Agrawal RK (2010) Synthesis and antibacterial evaluation of 2-substituted-4,5-diphenyl-N-alkyl imidazole derivatives. *Asian Pac J Trop Med* 3: 471-474.
4. Sharma S, Gangal S, Rauf A (2009) Convenient one-pot synthesis of novel 2-substituted benzimidazoles, tetrahydrobenzimidazoles and imidazoles and evaluation of their in vitro antibacterial and antifungal activities. *Eur J Med Chem* 44: 1751-1757.
5. Boiani M, González M, Mini-Rev (2005) Imidazole and benzimidazole derivatives as chemotherapeutic agents. *Med Chem* 5: 409-424.
6. Villalobos MR, Ibarra BM (2010) Synthesis, vasorelaxant activity and antihypertensive effect of benzo[d]imidazole derivatives. *Bioorg Med Chem* 18: 3985-3991.
7. Hadizadeh F, Hosseinzadeh H, Motamed-Shariaty VS, Seifi M, Kazemi SH (2008) Synthesis and antidepressant activity of N-Substituted imidazole-5-Carboxamides in forced swimming test model. *Iranian J Pharm Res* 7: 29-33.
8. Pandey J, Tiwari VK, Verma SS, Chaturvedi V, Bhatnagar S, et al. (2009) Synthesis and antitubercular screening of imidazole derivatives. *Eur J Med Chem* 44: 3350-3355.
9. Sharma D, Narasimhan B, Kumar P, Judge V, Narang R, et al. (2009) Synthesis, antimicrobial and antiviral evaluation of substituted imidazole

- derivatives. *Eur J Med Chem* 44: 2347-2353.
10. Puratchikody A, GopalaKrishnan S, Nallu M (2005) Synthesis and pharmacological evaluation of some Potent 2-(4-Substitutedphenyl)-4,5-Diphenyl-1H-Imidazoles. *Ind J Pharm Sci* 67: 725-731.
 11. Sisko J, Mellinger M (2009) Development of a general process for the synthesis of highly substituted imidazoles. *Pure Appl Chem* 74: 1349-1357.
 12. Navidpour L, Shadnia H, Shafaroodi H, Amini M, Dehpour AR, et al. (2007) Design, synthesis, and biological evaluation of substituted 2-alkylthio-1,5-diarylimidazoles as selective COX-2 inhibitors. *Bioorg Med Chem* 15: 1976-1982.
 13. Puratchikody A, Doble M (2007) Antinociceptive and anti-inflammatory activities and QSAR studies on 2-substituted-4,5-diphenyl-1H-imidazoles. *Bioorg Med Chem* 15: 1083-1090.
 14. Sharma GK, Kumar S, Pathak D (2010) Synthesis, antibacterial and anticancer activities of some novel imidazoles. *Der Pharmacia Lett* 2: 223-230.
 15. Baroniya S, Anwer Z, Sharma PK, Dudhe R, Kumar N (2010) Recent advancement in imidazole as anti cancer agents: A review. *Der Pharmacia Sinica* 1: 172-182.
 16. Narayana B, Vijaya-Raj KK, Ashalatha BV, Kumari NS, Sarojini BK (2004) Synthesis of some new 5-(2-substituted-1,3-thiazol-5-yl)-2-hydroxy benzamides and their 2-alkoxy derivatives as possible antifungal agents. *Eur J Med Chem* 39: 867-872.
 17. Suhas R, Chandrashekar S, Gowda DC (2011) Synthesis of elastin based peptides conjugated to benzisoxazole as a new class of potent antimicrobials -a novel approach to enhance biocompatibility. *Eur J Med Chem* 46: 704-711.
 18. Suhas R, Chandrashekar S, Gowda DC (2012) A new family of highly potent inhibitors of microbes: synthesis and conjugation of elastin based peptides to piperazine derivative. *Int J Pept Res Thera* 18: 89-98.
 19. Suresha GP, Suhas R, Kapfo W, Gowda DC (2011) Urea/thiourea derivatives of quinazolinone-lysine conjugates: synthesis and structural activity relationships of a new series of antimicrobials. *Eur J Med Chem* 46:2530-2540.
 20. Gadek TR, Nicholas JB (2003) Small molecule antagonists of proteins. *Biochem Pharmacol* 65: 1-8.
 21. Perez C, Paul M, Bazerque P (1990) An antibiotic assay by the agar well diffusion method. *Acta Biol Med Exp* 15:113-115.
 22. Singh I, Singh VP (2000) Antifungal properties of aqueous and organic solution extracts of seed plants against *Aspergillus flavus* and *A. niger*. *Phytomorphology* 50: 151-157.
 23. Blois MS (1958) Antioxidant determinations by the use of a stable free radical. *Nature* 181: 1199-1200.
 24. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, et al. (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Bio Med* 26: 1231-1237.
 25. Fogliano V, Verde V, Randazzo G, Ritieni A (1999) Method for measuring antioxidant activity and its application to monitoring the antioxidant capacity of wines. *J Agric Food Chem* 47: 1035-1040.
 26. Gülçin I (2010) Antioxidant properties of resveratrol: a structure-activity insight. *Food Sci Emerg* 11: 210-218.
 27. Lee Y, Martasek P, Roman LJ, Masters BS, Silverman RB (1999) Imidazole-containing amino acids as selective inhibitors of nitric oxide synthases. *Bioorg Med Chem* 7: 1941-1951.
 28. Eldon EB, Peter BD (1996) Solid phase synthesis of polyamides containing imidazole and pyrrole amino acids. *J Am Chem Soc* 118: 6141-6146.
 29. Leong LP, Shui G (2002) An investigation of antioxidant capacity of fruits in Singapore markets. *Food Chem* 76: 69-75.
 30. Lee YL, Yen MT, Mau JL (2007) Antioxidant properties of various extracts from *Hypsizygus marmoreus*. *Food Chem* 104: 1-9.

Copyright: © 2015 Channe Gowda D, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.