


B. Sridhar,<sup>a</sup> K. Ravikumar,<sup>a\*</sup> H. S. Yathirajan,<sup>b</sup> H. G. Anilkumar<sup>b</sup> and B. Narayana<sup>c</sup><sup>a</sup>Laboratory of X-ray Crystallography, Indian Institute of Chemical Technology, Hyderabad 500 007, India, <sup>b</sup>Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and <sup>c</sup>Department of Chemistry, Mangalore University, Mangalagangotri 574 199, IndiaCorrespondence e-mail:  
ravikumar\_iict@yahoo.co.in

## Key indicators

Single-crystal X-ray study  
 $T = 273$  K  
Mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å  
 $R$  factor = 0.038  
 $wR$  factor = 0.115  
Data-to-parameter ratio = 14.7For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.4-Amino-6-*tert*-butyl-3-thioxo-3,4-dihydro-1,2,4-triazin-5(2*H*)-oneIn the title compound,  $\text{C}_7\text{H}_{12}\text{N}_4\text{OS}$ , the triazine ring is planar. The crystal structure is stabilized by intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds which link the molecules into chains running along the  $c$  axis.Received 13 February 2006  
Accepted 7 March 2006

## Comment

1,2,4-Triazines and the compounds derived from them are found to possess a wide variety of pharmacological activities (Holla *et al.*, 1999). Triazine derivatives include herbicides having a broad spectrum of action that kill many species of weeds, and also herbicides with a narrow selectivity (Gruzdyev *et al.*, 1983). The title compound, (I), is an intermediate in the synthesis of a herbicide, 4-amino-6-(*tert*-butyl)-3-methylthio-1,2,4-triazin-5(4*H*)-one, commonly known as 'Sencor' (Eue & Tietz, 1972). It is used for the control of weeds in potato crops and is a broad-spectrum herbicide used in pre-emergence application in potato crops at a rate of 0.75–1 kg per hectare. In view of the importance of the title compound, (I), its crystal structure is reported here.In all essential details, the molecular geometry (Fig. 1 and Table 1) is in good agreement with related structures (Ghassemzadeh *et al.*, 1998). The triazine ring is planar and exists in the thione form; the  $\text{C}=\text{S}$  bond length of 1.663 (2) Å is slightly longer than the pure double-bond distance (1.61 Å; Pauling, 1960). The  $\text{N}-\text{N}$  [1.341 (2) Å] and  $\text{C}-\text{N}$  [mean value 1.341 (2) Å] bond distances are intermediate between the expected single (1.45 and 1.47 Å, respectively) and double (1.20 and 1.27 Å, respectively) bond distances.The crystal structure is stabilized by intra- and intermolecular hydrogen bonds (Table 2). Atom N1 of the triazine ring links the carbonyl O atom of an adjacent triazine ring, forming a chain running parallel to the  $c$  axis. The tertiary butyl group is oriented such that there are alternate hydrophobic and hydrophilic layers in the crystal packing (Fig. 2).

## Experimental

The title compound, (I), was obtained as a gift sample from Rallies India Ltd, Bangalore. The compound was recrystallized by slow evaporation of an acetone–toluene (9:1) solution (m.p. 488 K).




Figure 1

A view of (I), with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii.




Figure 2

A packing diagram of (I), viewed down the *b* axis, showing the chain formation. Dashed lines indicate N—H...O hydrogen bonds. H atoms not involved in hydrogen bonding have been omitted for clarity.

### Crystal data

$C_7H_{12}N_4OS$   
 $M_r = 200.27$   
 Monoclinic,  $P2_1/c$   
 $a = 12.5196$  (8) Å  
 $b = 6.9535$  (4) Å  
 $c = 12.5987$  (8) Å  
 $\beta = 112.039$  (1)°  
 $V = 1016.64$  (11) Å<sup>3</sup>  
 $Z = 4$

$D_x = 1.308$  Mg m<sup>-3</sup>  
 Mo  $K\alpha$  radiation  
 Cell parameters from 6656 reflections  
 $\theta = 3.0$ – $27.9$ °  
 $\mu = 0.29$  mm<sup>-1</sup>  
 $T = 273$  (2) K  
 Block, colorless  
 $0.21 \times 0.17 \times 0.09$  mm

### Data collection

Bruker SMART APEX CCD area-detector diffractometer  
 $\omega$  scan  
 Absorption correction: none  
 9276 measured reflections  
 1780 independent reflections

1640 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.018$   
 $\theta_{max} = 25.0$ °  
 $h = -14 \rightarrow 14$   
 $k = -8 \rightarrow 8$   
 $l = -14 \rightarrow 14$

### Refinement

Refinement on  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.115$   
 $S = 1.07$   
 1780 reflections  
 121 parameters  
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0662P)^2 + 0.2933P]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{max} < 0.001$   
 $\Delta\rho_{max} = 0.40$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.35$  e Å<sup>-3</sup>

Table 1

Selected geometric parameters (Å, °).

C1—N1	1.336 (3)	C2—N2	1.378 (2)
C1—N2	1.370 (2)	C3—N3	1.288 (2)
C2—O1	1.2244 (19)	N2—N4	1.401 (2)
C1—N1—N3	127.70 (15)		

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 <sup>i</sup>	0.86	1.94	2.790 (2)	168
N4—H4B...O1	0.86	2.31	2.607 (2)	100
N4—H4A...S1	0.86	2.59	2.976 (2)	108

Symmetry code: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

All H atoms were included in calculated positions ( $C-H = 0.93$ – $0.98$  Å and  $N-H = 0.86$  Å) and refined using a riding model, with  $U_{iso}(H) = 1.2U_{eq}(N,C)$  or  $1.5U_{eq}(\text{methyl } C)$ . The methyl groups were allowed to rotate but not to tip.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL/PC* (Sheldrick, 1990) and *MERCURY* (Bruno *et al.*, 2002); software used to prepare material for publication: *SHELXL97*.

BS and KR thank Dr J. S. Yadav, Director, IICT, Hyderabad, for his kind encouragement.

### References

- Bruker (2001). *SAINT* (Version 6.28a) and *SMART* (Version 5.625). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruno, I. J., Cole, J. C., Edgington, P. R., Kessler, M., Macrae, C. F., McCabe, P., Pearson, J. & Taylor, R. (2002). *Acta Cryst.* **B58**, 389–397.
- Eue, L. & Tietz, H. (1972). *Chem. Abstr.* **77**, 30196a.
- Ghassemzadeh, M., Aghapoor, K. & Heravi, M. (1998). *Z. Anorg. Allg. Chem.* **624**, 1969–1972.
- Gruzdyev, G. S., Zinchenko, V. A., Kalinin, V. A. & Slotvov, R. I. (1983). *The Chemical Protection of Plants*, pp. 401–402. Moscow: Mir Publishers.
- Holla, S. B., Sarojini, B. K., Shridhara, K. & Antony, G. (1999). *Il Farmaco*, **54**, 149–151.
- Pauling, L. (1960). *The Nature of the Chemical Bond*, 3rd ed. Ithaca: Cornell University Press.
- Sheldrick, G. M. (1990). *SHELXTL/PC* User's Manual. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.