

Kennesaw State University
DigitalCommons@Kennesaw State University

Faculty Publications

12-2009

A Graph Theoretic Summation of the Cubes of the First n Integers

Joseph DeMaio

Kennesaw State University, jdemai@kennesaw.edu

Andy Lightcap

Georgia State University

Follow this and additional works at: <https://digitalcommons.kennesaw.edu/facpubs>

 Part of the [Applied Mathematics Commons](#), and the [Mathematics Commons](#)

Recommended Citation

DeMaio, J., & Lightcap, A. (2009). A graph theoretic summation of the cubes of the first n integers. *Mathematics Magazine*, 82(5), 363-364. doi:10.4169/002557009X478409

This Article is brought to you for free and open access by DigitalCommons@Kennesaw State University. It has been accepted for inclusion in Faculty Publications by an authorized administrator of DigitalCommons@Kennesaw State University. For more information, please contact digitalcommons@kennesaw.edu.

A Graph Theoretic Summation of the Cubes of the First n Integers

Joe DeMaio
 Kennesaw State University
 Kennesaw, GA 30144
 jdemaio@kennesaw.edu

Andy Lightcap
 Kennesaw State University
 Kennesaw, GA 30144
 andy.lightcap@gmail.com

The complete graph K_{n+1} contains $n+1$ vertices and $\binom{n+1}{2}$ edges. Iteratively building the complete graph K_{n+1} by introducing vertices one at a time and counting the new edges incident to the new vertex provides a combinatorial proof that $\sum_{i=1}^n i = \binom{n+1}{2}$.




Figure 1: $\sum_{i=1}^4 i = \binom{4+1}{2}$

Since $\sum_{i=1}^n i^3 = \binom{n+1}{2}^2$ it seems natural to look for a combinatorial proof that also uses graphs. Consider the complete bipartite graph $K_{\binom{n+1}{2}, \binom{n+1}{2}}$ that contains $2\binom{n+1}{2}$ vertices and $\binom{n+1}{2}^2$ edges. As before, we will count the new edges incident to newly introduced vertices in n stages. At stage i we introduce i new vertices to each side of the graph and count the edges incident to these new vertices. Since $\sum_{i=1}^n i = \binom{n+1}{2}$ this process enumerates all the edges in $K_{\binom{n+1}{2}, \binom{n+1}{2}}$. New vertices on one side are adjacent only to vertices on the other side. When just considering the edges between the new vertices, the subgraph $K_{i,i}$ immediately appears with i^2 edges. It turns out that these i^2 edges along with the additional edges constructed between a new vertex on one side and an old vertex on the other side will always total i^3 new edges. This shows that $\sum_{i=1}^n i^3 = \binom{n+1}{2}^2$.

In order to see that we always introduce i^3 new edges at stage i , we will partition the new edges into complete bipartite graphs. At stage i , there exist

$\binom{i}{2} = \frac{i(i-1)}{2}$ previously introduced vertices on each side of the graph and the new vertices on each side are labeled $\binom{i}{2} + 1, \binom{i}{2} + 2, \dots, \binom{i}{2} + i = \binom{i+1}{2}$. The partition of these edges into complete bipartite graphs depends upon the parity of i . Figure 2 illustrates these stages for $n = 5$. To prevent a deluge of edges in the graph, a complete bipartite graph such as $K_{2,4}$ is represented as $\boxed{1,2} - \boxed{1,2,3,4}$.




Figure 2: $\sum_{i=1}^5 i^3 = \binom{5+1}{2}^2$

When i is odd, the new edges quickly form i disjoint copies of $K_{i,i}$. For odd i we partition the old vertices into $\frac{i-1}{2}$ sets of i vertices for each side. Both sets of i new vertices are adjacent to each of the $\frac{i-1}{2}$ sets of i vertices on the other side. This yields $2\left(\frac{i-1}{2}\right) = i - 1$ additional copies of $K_{i,i}$. Along with the initial copy of $K_{i,i}$ on only the new vertices, we have i copies of $K_{i,i}$ for a total of i^3 new edges.

When i is even, we have to work a bit harder. For even i , we partition the old vertices on each side into $\frac{i}{2} - 1$ sets of i vertices and one set of $\frac{i}{2}$ vertices. This yields $2\left(\frac{i}{2} - 1\right)$ copies of $K_{i,i}$ and two copies of $K_{\frac{i}{2},i}$ for $2\left(\frac{i}{2} - 1\right)i^2 + 2\frac{i}{2}i = i^3 - i^2$ edges. As before, with the original $K_{i,i}$ between the sets of new vertices, the total once again is i^3 new edges.

References

- [1] J. DeMaio and J. Tyson, Proof without words: a graph theoretic summation of the first n integers, *The College Mathematics Journal* **38** (2007) 296.