



University of Dundee

Ways to unwind with HROB, a new player in homologous recombination

Saredi, Giulia; Rouse, John

Published in:
Genes & Development

DOI:
[10.1101/gad.331850.119](https://doi.org/10.1101/gad.331850.119)

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Saredi, G., & Rouse, J. (2019). Ways to unwind with HROB, a new player in homologous recombination. *Genes & Development*, 33(19-20), 1293-1294. <https://doi.org/10.1101/gad.331850.119>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

OUTLOOK

Ways to unwind with HROB, a new player in homologous recombination

Giulia Saredi and John Rouse

MRC Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, University of Dundee, Dundee DD1 5EH, Scotland, United Kingdom

Homologous recombination (HR) is an important route for repairing DNA double-strand breaks (DSBs). The early stages of HR are well understood, but later stages remain mysterious. In this issue of *Genes & Development*, Hustedt and colleagues (pp. 1397–1415) reveal HROB as a new player in HR required for recruitment of the MCM8–9 complex, which is paralogous to the MCM2–7 replicative helicase. HROB functions closely with MCM8–9 to promote postsynaptic DNA repair synthesis. This study sheds valuable light on late events in HR and suggests that HROB may load MCM8–9 onto HR intermediates to facilitate the DNA unwinding required for DNA repair synthesis.

Homologous recombination (HR) is a major route for repairing DNA double-strand breaks (DSBs) (Prakash et al. 2015). HR starts with resection of broken DNA ends, involving nucleases and helicases. The resulting overhangs, coated by RAD51 to yield a nucleoprotein filament, invade the intact sister chromatid, searching for a homologous sequence. The pairing of RAD51-coated single-stranded DNA (ssDNA) with the donor strand (termed synapsis) results in the displacement of the strand complementary to the donor (forming a structure called a “D loop”) and the establishment of a RAD51-bound heteroduplex species that primes DNA repair synthesis (Fig. 1). The postsynaptic stages of HR are poorly understood, but a number of proteins are involved at this stage specifically. For example, the MCM8 and MCM9 proteins, which are related to the subunits of the hexameric MCM2–7 replicative helicase, have been implicated in postsynaptic DNA synthesis. In contrast to MCM2–7, the MCM8–9 complex is dispensable for bulk DNA replication (Griffin and Trakselis 2019). The residual DNA synthesis that is maintained in MCM2-depleted cells requires MCM8–9, but this reflects DNA repair synthesis occurring during HR-mediated repair of the high level of DSBs that occur upon MCM2 depletion (Natsume et al.

2017). Importantly, MCM8–9 was shown to act downstream from RAD51 in the HR-mediated repair of DSBs induced by nuclease overexpression, in meiotic HR, and during the HR stage of DNA interstrand cross-link (ICL) repair (Lutzmann et al. 2012; Nishimura et al. 2012; Natsume et al. 2017). The dominant mode of ICL repair is initiated by collision of replisomes with ICLs, resulting in programmed formation of DSBs that are repaired by HR. Crucially, point mutations in MCM8–9 predicted to abolish helicase activity suppress ICL repair, suggesting that helicase activity is important for HR (Nishimura et al. 2012). One explanation is that MCM8–9 might facilitate postsynaptic DNA repair synthesis by unwinding D loops and enabling extension of the invading RAD51-coated DNA end (Fig. 1).

In this issue of *Genes & Development*, Hustedt et al. (2019) identify a new factor—HROB (HR OB-fold)—that functions closely with MCM8–9 in HR. Rationalizing that HR defects should cause sensitivity to inhibitors of ATR kinase and PARP, the investigators mined published CRISPR screens to identify new genes whose deletion sensitizes to both drug classes. This approach highlighted the uncharacterized *C17ORF53* gene (renamed *HROB*) that, when disrupted in human cells or mice, causes pronounced HR defects. Disruption of mouse *Hrob*, for example, showed defects in meiosis and gametogenesis. Human *HROB*-deleted cells are defective in the HR step of ICL repair, and, whereas RAD51 loading is normal, RAD51 foci persist, hinting that HROB acts at a postsynaptic HR stage downstream from RAD51, similar to MCM8–9. In this light, *HROB* is epistatic with *MCM8* and *MCM9* in ICL repair and in HR-mediated repair of DSBs in a reporter cassette. Furthermore, residual DNA synthesis in MCM2-depleted cells, which depends on MCM8–9 (Natsume et al. 2017), requires HROB, with HROB and MCM8–9 acting epistatically in this context. A breakthrough came with the discovery that HROB recruits MCM8–9 to DNA lesions. HROB was found to physically interact with MCM8–9, and, excitingly, siRNA-mediated depletion of

[**Keywords:** CRISPR screens; cisplatin; DNA damage; DNA repair; DNA synthesis; germ cells; helicase; homologous recombination; infertility]

Corresponding author: j.rouse@dundee.ac.uk

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.331850.119>.

© 2019 Saredi and Rouse This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genesdev.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.




Figure 1. Model for the concerted actions of HROB–MCM8–MCM9 and HELQ in postsynaptic DNA synthesis during HR.

HROB compromised the accumulation of MCM8 at DNA repair foci but not vice versa. These data make a compelling case for HROB as an important HR factor acting with MCM8–9 to promote postsynaptic DNA repair synthesis. The investigators suggest that HROB acts by directly loading MCM8–9 onto HR intermediates, which draws interesting mechanistic parallels with MCM2–7, whose loading involves transient opening of a gate in the MCM2–7 hexamer (Deegan and Diffley 2016). It is not yet clear whether the MCM8–9 complex is hexameric in nature, nor is it known whether gate opening, perhaps mediated by HROB, is required for loading.

Another notable finding is that *HROB* deletion in cells lacking the DNA helicase *HELQ* causes a stronger HR defect than in the single mutants. *HELQ* was known to be involved in HR during ICL repair (Takata et al. 2013) and in HR-mediated meiotic DSB repair (Adelman and Boulton 2010; Ward et al. 2010). In these contexts, *HELQ* deletion causes persistent RAD51 foci (Adelman and Boulton 2010). The new study shows that *HELQ* and *HROB*–MCM8–MCM9 function redundantly, perhaps representing parallel pathways for promoting DNA synthesis downstream from RAD51 (Fig. 1). How might these two pathways converge at the molecular level? Perhaps the simplest model is that unwinding of HR intermediates, such as D loops, requires the helicase activities of both *HELQ* and MCM8–9 to enable extension of the RAD51-coated DNA end. Such a model predicts that helicase-inactivating mutations in *HELQ* would phenocopy, at least partly, the gene deletion. This remains to be tested. It is interesting that *HELQ* can remove RAD51 from dsDNA *in vitro* independently of helicase activity, possibly by direct RAD51 binding (Adelman and Boulton 2010; Ward et al. 2010). Furthermore, *HELQ* interacts with the BCDX2 complex of RAD51 paralogs that may also have a RAD51-stripping activity (Adelman et al. 2013; Takata et al. 2013). It is conceivable, therefore, that the ability of *HELQ* to strip RAD51 off the terminus of the invading end in the D loop, perhaps together with BCDX2, facilitates postsynaptic DNA synthesis through freeing the terminus for

primer extension. Such a model might be suggested if mammalian cells with helicase-inactivating *HELQ* mutations were HR-proficient. Whatever the case, the identification of *HROB*, working together with MCM8–9 and *HELQ* in facilitating postsynaptic DNA repair synthesis, sheds light on a poorly understood aspect of HR.

Acknowledgments

G.S. is supported by an EMBO Long-Term Postdoctoral Fellowship (ALTF 951-2018). Work in the Rouse laboratory is funded by the Medical Research Council (grant no. MC_UU_12016/1).

References

- Adelman CA, Boulton SJ. 2010. Metabolism of postsynaptic recombination intermediates. *FEBS Lett* **584**: 3709–3716. doi:10.1016/j.febslet.2010.05.023
- Adelman CA, Lolo RL, Birkbak NJ, Murina O, Matsuzaki K, Horjési Z, Parmar K, Borel V, Skehel JM, Stamp G, et al. 2013. *HELQ* promotes RAD51 paralogue-dependent repair to avert germ cell loss and tumorigenesis. *Nature* **502**: 381–384. doi:10.1038/nature12565
- Deegan TD, Diffley JF. 2016. MCM: one ring to rule them all. *Curr Opin Struct Biol* **37**: 145–151. doi:10.1016/j.sbi.2016.01.014
- Griffin WC, Trakselis MA. 2019. The MCM8/9 complex: A recent recruit to the roster of helicases involved in genome maintenance. *DNA Repair (Amst)* **76**: 1–10. doi:10.1016/j.dnarep.2019.02.003
- Hustedt N, Saito Y, Zimmermann M, Álvarez-Quilón A, Setiaputra D, Adam S, McEwan A, Yuan JY, Olivieri M, Zhao Y, et al. 2019. Control of homologous recombination by the HROB–MCM8–MCM9 pathway. *Genes Dev* (this issue). doi:10.1101/gad.329508.119
- Lutzmann M, Grey C, Traver S, Ganier O, Maya-Mendoza A, Ranisavljevic N, Bernex F, Nishiyama A, Montel N, Gavois E, et al. 2012. MCM8- and MCM9-deficient mice reveal gametogenesis defects and genome instability due to impaired homologous recombination. *Mol Cell* **47**: 523–534. doi:10.1016/j.molcel.2012.05.048
- Natsume T, Nishimura K, Minocherhomji S, Bhowmick R, Hickson ID, Kanemaki MT. 2017. Acute inactivation of the replicative helicase in human cells triggers MCM8–9-dependent DNA synthesis. *Genes Dev* **31**: 816–829. doi:10.1101/gad.297663.117
- Nishimura K, Ishiai M, Horikawa K, Fukagawa T, Takata M, Takisawa H, Kanemaki MT. 2012. Mcm8 and Mcm9 form a complex that functions in homologous recombination repair induced by DNA interstrand crosslinks. *Mol Cell* **47**: 511–522. doi:10.1016/j.molcel.2012.05.047
- Prakash R, Zhang Y, Feng W, Jasin M. 2015. Homologous recombination and human health: the roles of BRCA1, BRCA2, and associated proteins. *Cold Spring Harb Perspect Biol* **7**: a016600. doi:10.1101/cshperspect.a016600
- Takata K, Reh S, Tomida J, Person MD, Wood RD. 2013. Human DNA helicase *HELQ* participates in DNA interstrand crosslink tolerance with ATR and RAD51 paralogs. *Nat Commun* **4**: 2338. doi:10.1038/ncomms3338
- Ward JD, Muzzini DM, Petalcorin MI, Martinez-Perez E, Martin JS, Plevani P, Cassata G, Marini F, Boulton SJ. 2010. Overlapping mechanisms promote postsynaptic RAD-51 filament disassembly during meiotic double-strand break repair. *Mol Cell* **37**: 259–272. doi:10.1016/j.molcel.2009.12.026



Ways to unwind with HROB, a new player in homologous recombination

Giulia Saredi and John Rouse

Genes Dev. 2019, **33**:

Access the most recent version at doi:[10.1101/gad.331850.119](https://doi.org/10.1101/gad.331850.119)

Related Content	Control of homologous recombination by the HROB MCM8 MCM9 pathway Nicole Hustedt, Yuichiro Saito, Michal Zimmermann, et al. Genes Dev. October , 2019 33: 1397-1415
References	This article cites 11 articles, 3 of which can be accessed free at: http://genesdev.cshlp.org/content/33/19-20/1293.full.html#ref-list-1 Articles cited in: http://genesdev.cshlp.org/content/33/19-20/1293.full.html#related-urls
Creative Commons License	This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see http://genesdev.cshlp.org/site/misc/terms.xhtml). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at http://creativecommons.org/licenses/by-nc/4.0/ .
Email Alerting Service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .

**CRISPR KO, CRISPRa,
CRISPRi libraries.**
Custom or genome-wide.

[VIEW PRODUCTS >](#)

