

MAD/TH/92-8

Jets and jet multiplicities in high energy photon-nucleon interactions

Loyal Durand and Katsuhiko Honjo

*Department of Physics, University of Wisconsin-Madison,
Madison, Wisconsin 53706*

Raj Gandhi

*Department of Physics, Texas A&M University,
College Station, TX 77843*

Hong Pi

*Department of Theoretical Physics, University of Lund,
Lund, Sweden S-22362*

Ina Sarcevic

*Department of Physics, University of Arizona,
Tucson, Arizona 85721*

Abstract

We discuss the theory of jet events in high-energy photon-proton interactions using a model which gives a good description of the data available on total inelastic γp cross sections up to $\sqrt{s}=210$ GeV. We show how to calculate the jet cross sections and jet multiplicities and give predictions for these quantities for energies appropriate for experiments at the HERA ep collider and for very high energy cosmic ray observations.

PACS number(s): 13.60.Hb, 13.87.Ce, 12.40.Lk

The hadronic structure of the photon and its interaction with hadrons are currently subjects of considerable interest. The ZEUS and H1 groups at the ep collider HERA have presented preliminary results [1] on the γp photoproduction cross section at 210 GeV center-of-mass energy which show the rise with energy typical of other hadronic interactions. It was shown by Gandhi and Sarcevic [2] in a simple model that this rise can be used to discriminate between different sets of photon structure functions. The hadronic behavior of the photon at very high energies ($\sim 10^8$ GeV photon energy) is also of interest for the theory of air showers triggered by cosmic ray photons [3].

We have developed an improved QCD-based model of the photon interaction with nucleons [4] which gives predictions for the total inelastic γp cross section which agree well with the HERA data. The model has the interesting feature that the information needed to calculate $\sigma_{inel}^{\gamma p}$ is largely determined by high energy πN scattering. In this paper, we will use the model to discuss the total jet cross section $\sigma_{jet}^{\gamma p}$ and the probabilities for multiple-jet events, and give our predictions for those quantities at HERA energies and the energies relevant for cosmic ray experiments.

We begin by sketching our model for inelastic γp scattering. The theory is developed in more detail in [4]. The incoming physical photon state $|\gamma\rangle_{phys}$ can be expressed as a superposition of the bare photon and the set of virtual hadronic states $|m\rangle$ which can be distinguished in the subsequent interaction, and appear with probabilities \mathcal{P}_m . Accordingly, the total inelastic γp cross section is given in a semiclassical picture as

$$\sigma_{inel}^{\gamma p} = (1 - \mathcal{P}_{had})\sigma_{dir} + \sum_m' \mathcal{P}_m \sigma^{mp}, \quad \mathcal{P}_{had} = \sum_m \mathcal{P}_m \ll 1. \quad (1)$$

Included among the states $|m\rangle$ are low-mass vector meson states such as the ρ, ω and ϕ , and complex non-resonant final states which can be described on the average in a

quark-gluon basis.

Whether the photon appears as a virtual vector meson state or a high-mass non-resonant state depends on the relative transverse momentum $2p_{\perp 0}$ of the quark and antiquark in the virtual transition $\gamma \rightarrow q\bar{q}$ which initiates the hadronic interaction. We note that $p_{\perp 0} \approx 1/r_{\perp}$, where r_{\perp} is the average transverse separation of the virtual quarks during the lifetime of the $q\bar{q}$ system. If r_{\perp} is greater than, or on the order of, the average transverse radius $R_{\perp} \equiv 1/Q_0$ of a vector meson, QCD confinement effects will clearly set in, and the $q\bar{q}$ system will most likely appear in a hadronic collision as a light vector meson. For $r_{\perp} < R_{\perp}$ the $q\bar{q}$ system will be smaller than a vector meson. In this picture, the hadronic system behaves for $p_{\perp 0} < Q_0$ like a vector meson, and for $p_{\perp 0} > Q_0$, like a system of quasi-free quarks and gluons with a transverse area smaller than that of a vector meson by a factor $(Q_0/p_{\perp 0})^2$. Its interaction cross section will be reduced accordingly.

After eikonalizing the hadronic cross sections using an impact parameter representation to account for possible multiple parton scatterings in a single γp collision, the total inelastic γp cross section is given by [4] as

$$\begin{aligned} \sigma_{inel}^{\gamma p} &= \sigma_{dir} + \lambda \mathcal{P}_{\rho} \int d^2b (1 - e^{-2Re\chi^{\rho p}}) \\ &\quad + \sum_q e_q^2 \frac{\alpha_{em}}{\pi} \int_{Q_0^2} \frac{dp_{\perp 0}^2}{p_{\perp 0}^2} \int d^2b (1 - e^{-2Re\chi^{q\bar{q}p}}). \end{aligned} \quad (2)$$

Here \mathcal{P}_{ρ} is the probability that a photon exists in the ρ meson state, $\mathcal{P}_{\rho} = 4\pi\alpha_{em}/f_{\rho}^2$, where f_{ρ}^2 is the $\gamma\rho$ coupling in the vector meson dominance model. We will treat the ρ, ω and ϕ as equivalent, and will use the quark model ratios for the couplings f_V ; then $\lambda = 4/3$ for equal $\rho p, \omega p$, and ϕp cross sections, and $\lambda = 10/9$ for complete suppression of the ϕ contribution at low $p_{\perp 0}$ [5]. The third term in Eq. (2) includes the contributions from the excited hadronic states of the photon, which we collectively call $q\bar{q}$ states. The factor $(e_q^2\alpha_{em}/\pi p_{\perp 0}^2)dp_{\perp 0}^2$ is just the differential probability of

producing a $q\bar{q}$ pair at a relative transverse momentum $2p_{\perp 0}$.

The real part of the eikonal function for the ρp interaction can be written in the form [6]

$$\begin{aligned} \text{Re}\chi^{\rho p}(b, s) &= \text{Re}\chi_{soft}^{\rho p}(s) + \text{Re}\chi_{QCD}^{\rho p}(b, s) \\ &= \frac{1}{2}A^{\rho p}(b)[\sigma_{soft}^{\rho p}(s) + \sigma_{QCD}^{\rho p}(s)], \end{aligned} \quad (3)$$

where $A^{\rho p}(b)$ is the density overlap function,

$$A^{\rho p}(b) = \int d^2b' \rho_{\rho}(b) \rho_p(|\vec{b} - \vec{b}'|), \quad \int d^2b A^{\rho p}(b) = 1, \quad (4)$$

and $\sigma_{soft}^{\rho p}$ and $\sigma_{QCD}^{\rho p}$ are the soft- and hard-scattering parts of the intrinsic cross section. $\sigma_{soft}^{\rho p}$ was parametrized in [4] using a Regge-like form,

$$\sigma_{soft}^{\rho p} = \sigma_0 + \sigma_1(s - m_p^2)^{-1/2} + \sigma_2(s - m_p^2)^{-1}, \quad (5)$$

while $\sigma_{QCD}^{\rho p}$ was identified with the inclusive parton-level cross section for ρp scattering,

$$\sigma_{QCD}^{\rho p} = \sum_{ij} \frac{1}{1 + \delta_{ij}} \int_0^1 dx_1 \int_0^1 dx_2 \int_{p_{\perp, min}^2} dp_{\perp}^2 f_i^p(x_1, p_{\perp}^2) f_j^p(x_2, p_{\perp}^2) \frac{d\hat{\sigma}_{ij}}{dp_{\perp}^2}. \quad (6)$$

Using the equivalence of the ρ and π states in the quark model, we can equate the parton distribution and density profile functions for the ρ meson to the corresponding functions for the pion and take $f_i^{\rho} = f_i^{\pi}$ and $\rho_{\rho}(b) = \rho_{\pi}(b)$. The parton distributions in the pion are known reasonably well [7]. The density profile functions $\rho_{\rho}(b)$ and $\rho_{\pi}(b)$ can be taken as the Fourier transforms of the electromagnetic form factors of the proton and the π meson. The density overlap function $A^{\rho p} \approx A^{\pi p}$ is then given by [8]

$$A^{\rho p}(b) = \frac{1}{4\pi} \frac{\nu^2 \mu^2}{\mu^2 - \nu^2} \left\{ \nu b K_1(\nu b) - \frac{2\nu^2}{\mu^2 - \nu^2} [K_0(\nu b) - K_0(\mu b)] \right\}, \quad (7)$$

where $K_n(x)$ is a hyperbolic Bessel function, and the “size” parameters of the pion and the proton are $\mu^2 = 0.47 \text{ GeV}^2$ and $\nu^2 = 0.71 \text{ GeV}^2$, respectively. The former corresponds to a root-mean-square transverse radius for the pion or ρ meson $R_\perp = 1/Q_0 \approx 2/\mu$.

The eikonal function for the $q\bar{q}p$ scattering terms in Eq. (2) can be expressed similarly as

$$2Re\chi^{q\bar{q}p} = A^{q\bar{q}p}(b, p_\perp) [\sigma_{soft}^{q\bar{q}p}(s, p_\perp) + \sigma_{QCD}^{q\bar{q}p}(s, p_\perp)]. \quad (8)$$

We assume that $\rho_{q\bar{q}}(b)$ has the same form as $\rho_\rho(b)$ except with the size parameter $\mu = 2Q_0$ of the ρ meson replaced by $2p_\perp$ [4]. With this replacement, $A^{q\bar{q}p}(b, p_\perp)$ and $\rho_{q\bar{q}}(b, p_\perp)$ are continuous with the corresponding functions for the ρ meson at $p_\perp = Q_0$, but describe a shrinking system for $p_\perp > Q_0$. Finally, the cross sections $\sigma_{soft}^{q\bar{q}p}(s, p_\perp)$ and $\sigma_{QCD}^{q\bar{q}p}(s, p_\perp)$, were assumed to scale with the physical cross sectional area of the $q\bar{q}$ system so that

$$\sigma_{soft}^{q\bar{q}p}(s, p_\perp) + \sigma_{QCD}^{q\bar{q}p}(s, p_\perp) = (Q_0^2/p_\perp^2) [\sigma_{soft}^{\rho p}(s) + \sigma_{QCD}^{\rho p}(s)]. \quad (9)$$

This model guarantees that σ_{soft} and σ_{QCD} are continuous at $p_\perp = Q_0$, and that the soft contributions from the $q\bar{q}$ states die out as $1/p_\perp^2$ as expected for higher twist contributions.

We have calculated the total inelastic γp cross section using Eq. (2) and the assumptions above. The soft cross section in Eq. (5) was determined [4] using the low-energy γp data [9]. The calculation used the parton distributions of Owens [7] for the pion and those of Eichten *et al.* [10] for the proton. The parameter $p_{\perp, min}$ in Eq. (6) which determines the transverse momentum at which hard parton-level collisions come into play was taken from a similar fit to $\pi^\pm p$ scattering [8]. The results agree with the preliminary HERA data [1] and those at lower energies as shown by the top curve in Fig. 1(a). The calculated rise in $\sigma_{inel}^{\gamma p}$ at higher energies

arises from the hard-scattering contributions, and is a prediction of the model rather than a fit to the HERA data. It is therefore of interest to look for other tests of the model. The jet cross sections predicted by the model provide one such test.

The total jet cross section $\sigma_{jet}^{\gamma p}(s, Q^2)$ is defined to be the part of the inelastic γp cross section which includes events with at least one semihard parton-parton (or γ -parton) scattering with a momentum transfer $p_{\perp}^2 \geq Q^2$, irrespective of any soft processes that may occur. To find an expression for the jet cross section σ_{jet}^{mp} in the interaction between the proton and the hadronic state $|m\rangle$ of the photon, we use the fact that semiclassically $exp[-2Re\chi_{QCD}(b, s, Q^2)]$ can be interpreted as the probability that there is *no* parton-parton scattering with $p_{\perp}^2 \geq Q^2$ in a hadronic collision at impact parameter b . Using this observation, we can rewrite the expression for the total hadronic cross section σ^{mp} to separate out the jet-free part,

$$\begin{aligned}
\sigma_{had}^{mp}(s) &= \sigma_{nojet}^{mp}(s, Q^2) + \sigma_{jet}^{mp}(s, Q^2) \\
&= \int d^2b \left(1 - e^{-2Re\chi_{QCD}^{mp} - 2Re\chi_{soft}^{mp}} \right) \\
&= \int d^2b \left(1 - e^{-2Re\chi_{soft}^{mp}'}(b, s, Q^2) \right) e^{-2Re\chi_{QCD}^{mp}(b, s, Q^2)} \\
&\quad + \int d^2b \left(1 - e^{-2Re\chi_{QCD}^{mp}(b, s, Q^2)} \right). \tag{10}
\end{aligned}$$

Here

$$\sigma_{jet}^{mp}(s, Q^2) = \int d^2b \left(1 - e^{-2Re\chi_{QCD}^{mp}(b, s, Q^2)} \right) \tag{11}$$

is the total cross section for events associated with $|m\rangle$ which contain jets with $p_{\perp} \geq Q$. The factor $\sigma_{QCD}^{\gamma p}$ which appears in the eikonal function in Eq. (11) is now to be evaluated using the expression in Eq. (6) with $p_{\perp, min}^2$ replaced by Q^2 . The remaining ‘‘nojet’’ cross section involves a modified eikonal function

$$Re\chi_{soft}^{mp}{}'(s, b, Q^2) = Re\chi_{soft}^{mp} + Re\chi_{QCD}^{mp} - Re\chi_{QCD}^{mp}(b, s, Q^2), \tag{12}$$

and includes contributions from “soft” jets with $p_{\perp,min}^2 \leq p_{\perp}^2 \leq Q^2$ as well as the usual soft term.

The total jet cross section in γp scattering is the generalized sum of the cross sections $\sigma_{jet}^{mp}(s, Q^2)$, weighted with the probabilities \mathcal{P}_m , over all the possible hadronic states of the photon, plus the very small jet contribution from $\sigma_{dir}(s, Q^2)$,

$$\begin{aligned} \sigma_{jet}^{\gamma p}(s, Q^2) &= \sigma_{dir}(s, Q^2) + \sum_m \mathcal{P}_m \sigma_{jet}^m \\ &= \sigma_{dir}(s, Q^2) + \lambda \mathcal{P}_\rho \sigma_{jet}^{\rho p}(s, Q^2) \\ &\quad + \sum_q e_q^2 \frac{\alpha_{em}}{\pi} \int_{Q_0^2} \frac{dp_{\perp 0}^2}{p_{\perp 0}^2} \sigma_{jet}^{q\bar{q}p}(s, Q^2, p_{\perp 0}^2). \end{aligned} \quad (13)$$

The jet cross sections for specific kinematic cuts, e.g., on jet angle, can be calculated using the same basic formulas with the cuts imposed on the integral for $\sigma_{QCD}^{\gamma p}$ in Eq. (6).

We have calculated the total jet cross section $\sigma_{jet}^{\gamma p}$ in the HERA energy range for $Q = 2, 3, 4,$ and 5 GeV, using the same parameters as were used in [4]. Our predictions are shown in Fig. 1 along with the calculated total inelastic γp cross section and the experimental data [1, 9]. At $\sqrt{s}=200$ GeV, the jet cross section is predicted to be approximately 1.3% of the total inelastic cross section for $Q=5$ GeV in rough agreement with earlier predictions [2], and 25% of the total for $Q=2$ GeV.

The rapid growth of the jet cross section with energy is associated with the growth of the parton distribution functions in the photon and the proton at the small values of x which become accessible at high energies. This growth also leads to an increasing probability of the scattering of more than one parton in a single γp collision. For this reason, the inclusive parton-level γp cross section $\sigma_{QCD}^{\gamma p}$, which counts each parton collision separately, is not the same as $\sigma_{jet}^{\gamma p}$. The latter has the overcounting due to multiple independent scatterings removed in the eikonalization. We show the effect of the eikonalization up to HERA energies in Fig. 2. The effect is not very significant

for $Q = 3, 4$ and 5 GeV at $\sqrt{s} = 200$ GeV, but causes about a 10% reduction in the jet cross section for $Q = 2$ GeV.

In Fig. 2 we show $\sigma_{jet}^{\gamma p}$ for the cosmic ray energy range for the same choices of Q as in Fig. 1, together with the calculated cross sections $\sigma_{inel}^{\gamma p}$ and $\sigma_{QCD}^{\gamma p}$ for $Q = 2$ GeV. At very high energies, the jet cross section clearly comprises a large part of the total inelastic cross section, e.g., at $E_\gamma = 2 \times 10^8$ GeV (or $\sqrt{s} \approx 2 \times 10^4$ GeV) $\sigma_{jet}^{\gamma p}$ constitutes $\approx 75\%$ of $\sigma_{inel}^{\gamma p}$ for $Q = 2$ GeV. Even for $Q = 5$ GeV, $\sigma_{jet}^{\gamma p}$ gives nearly 40% of $\sigma_{inel}^{\gamma p}$. Also, as more of the incident partons scatter, the effect of eikonalization becomes quite large as shown for $Q = 2$ GeV.

To calculate the multiplicity of jets, we note that the average number of parton-parton scatterings with $p_\perp > Q$ in a single γp collision at impact parameter b associated with the hadronic component $|m\rangle$ of the photon is

$$\bar{n}_m(b, s, Q^2) = \sigma_{QCD}^{mp}(s, Q^2) A^{mp}(b) = 2Re\chi_{QCD}^{mp}(b, s, Q^2). \quad (14)$$

Since the parton-parton scatterings in our model are independent, the probability of having n scatterings ($2n$ jets) in such a hadronic collision has a Poisson distribution,

$$P_n^{mp}(b, s, Q^2) = \frac{1}{n!} [\bar{n}_m(b, s, Q^2)]^n e^{-\bar{n}_m(b, s, Q^2)}. \quad (15)$$

The probability $P_n(s, Q^2)$ of having n scatterings relative to *all* the inelastic events at an impact parameter b is then

$$\begin{aligned} P_n(s, Q^2) &= (\sigma_{inel}^{\gamma p})^{-1} \sum_m \mathcal{P}_m \sigma_{jet,n}^{mp} \\ &= (\sigma_{inel}^{\gamma p})^{-1} \sum_m \mathcal{P}_m \int d^2b P_n^{mp}(b, s, Q^2) \\ &= (\sigma_{inel}^{\gamma p})^{-1} \sum_m \mathcal{P}_m \frac{1}{n!} \int d^2b [\bar{n}_m(b, s, Q^2)]^n e^{-\bar{n}_m(b, s, Q^2)} \end{aligned} \quad (16)$$

for $n \geq 1$. $P_0(s, Q^2)$ is defined to be the probability of having a soft interaction with

no associated jets,

$$P_0(s, Q^2) = (\sigma_{inel}^{\gamma p})^{-1} \sum_m \mathcal{P}_m \int d^2b \left(1 - e^{-2Re\chi_{soft}^{mp'}(b, s, Q^2)} \right) e^{-\bar{n}_m(b, s, Q^2)}. \quad (17)$$

Note that we have neglected the very small two-jet contribution to these expressions from the direct QCD interaction of the photon with the partons in the proton.

We have calculated P_n for $n = 0, 1, 2$ and 3 at $\sqrt{s} = 200$ GeV using $Q = 2, 3, 4,$ and 5 GeV. The results are shown in Table I. For $Q = 5$ GeV the probability of having a 2-jet event (a single parton scattering) is about 1%; the probability for 2n-jet events with n larger than 1 is essentially zero. For $Q = 2$ GeV the *total* probability of having any hadronic jet event is approximately 27%, with the 2-jet events comprising more than 90 % of the total.

We also investigated jet production in γp scattering at very high energies. In Fig. 3 we show the distribution of P_n at $\sqrt{s} = 2 \times 10^4$ GeV with the choice of $Q = 3$ GeV. It is more likely at this energy to have jets in an event than not, with an approximately 70% jet probability. While the 2-jet production is still the most probable, events with more than 2 jets occur with about 20% probability.

It will be quite interesting to see if the predictions above are verified in future experiments since they connect the hadronic properties of the photon quite directly to those of the pion. In particular, the jet structure of pion and photon-induced reactions on protons should be quite similar.

ACKNOWLEDGMENTS

This work was supported in part through U. S. Department of Energy Grants Nos. DE-AC02-76ER00881 and DE-FG02-85ER40213. One of the authors (RG) would like to thank the World Laboratory for support.

References

- [1] The preliminary values of $\sigma_{inel}^{\gamma p}$ are: ZEUS Collaboration, Phys. Lett. B **293**, 465 (1992), $\sigma_{inel}^{\gamma p}=154\pm 16\pm 32 \mu\text{b}$ at a average energy of 210 GeV; H1 Collaboration (reported at the XXVI International Conference on High Energy Physics by F. Eisele), $\sigma_{inel}^{\gamma p}= 165\pm 28 \mu\text{b}$ or $150\pm 22 \mu\text{b}$ depending on the model used in the analysis. We show a value $160\pm 30 \mu\text{b}$ for H1 in our figures to span the experimental and theoretical uncertainties.
- [2] R. Gandhi and I. Sarcevic, Phys. Rev. D **44**, R10 (1991).
- [3] See, for example, M. Drees and F. Halzen, Phys. Rev. Lett. **61**, 275 (1988); M. Drees, F. Halzen, and K. Hikasa, Phys. Rev. D **39**, 1310 (1989); T.K. Gaisser, F. Halzen, T. Stanev, and E. Zas, Phys. Lett. B **243**, 444 (1990).
- [4] K. Honjo, L. Durand, R. Gandhi, I. Sarcevic, and H. Pi, University of Wisconsin Report No. MAD/TH/92-6.
- [5] There is evidence for considerable suppression of the cross section expected for photoproduction of the ϕ relative to that for the ρ for $\sqrt{s} \approx 10$ GeV. See R.M. Eglyoff *et al.*, Phys. Rev. Lett. **43**, 657 (1979). We therefore used $\lambda=10/9$ in the calculations reported here. This has rather little effect on the high energy results [4].
- [6] L. Durand and H. Pi, Phys. Rev. Lett. **58**, 303 (1987); Phys. Rev. D **40**, 1436 (1989).
- [7] J.F. Owens, Phys. Rev. D **30**, 943 (1984).
- [8] L. Durand and H. Pi, Phys. Rev. D **43**, 2125 (1991).

- [9] The lower energy data shown in Fig. 1 are from D.O. Caldwell *et al.*, Phys. Rev. Lett. **25**, 609 (1970); *ibid.* **40**, 1222 (1978).
- [10] E. Eichten, K. Lane, I. Hinchliffe, and C. Quigg, Rev. Mod. Phys. **56**, 579 (1984).

FIGURE CAPTIONS

FIG. 1. (a) The calculated total jet cross sections as functions of \sqrt{s} and the transverse momentum cutoff Q . The total γp inelastic cross section calculated in [4] is compared to the low-energy [9] and HERA [1] data in the upper curve. (b) The eikonalized and inclusive jet cross sections for the same values of Q .

FIG. 2. The total inelastic and jet cross sections at cosmic ray energies. $\sigma_{QCD}^{\gamma p}$ is the inclusive jet cross section.

FIG. 3. The probability distributions for n parton-parton collisions or $2n$ jets in a γp collision.

Q (GeV)	P_0	P_1	P_2	P_3
2	0.734	0.228	0.020	0.002
3	0.904	0.078	0.002	0.000
4	0.954	0.030	0.000	0.000
5	0.971	0.013	0.000	0.000

Table 1: Probabilities P_n of having n parton-parton collisions or $2n$ jets in a single inelastic γp event at $\sqrt{s}=200$ GeV.