



**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang^{1,2}, D. B. A. Jones^{1,3}, J. Worden², H. M. Worden⁴, D. K. Henze⁵, and Y. Wang^{6,7}

¹Department of Physics, University of Toronto, Toronto, ON, Canada

²Jet Propulsion Laboratory, California Institute of Technology, Pasadena CA, USA

³JIFRESSE, University of California, Los Angeles, Los Angeles, CA, USA

⁴National Center for Atmospheric Research, Boulder, CO, USA

⁵University of Colorado Boulder, CO, USA

⁶Department of Marine Sciences, Texas A&M University at Galveston, Galveston, TX, USA


⁷Ministry of Education Key Laboratory for Earth System Modeling, Center for Earth System Science, Institute for Global Change Studies, Tsinghua University, Beijing, China

Received: 26 January 2015 – Accepted: 9 February 2015 – Published: 25 February 2015

Correspondence to: Z. Jiang (zhe.jiang@jpl.nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Abstract

Chemical transport models (CTMs) driven with high-resolution meteorological fields can better resolve small-scale processes, such as frontal lifting or deep convection, and thus improve the simulation and emission estimates of tropospheric trace gases. In this work, we explore the use of the GEOS-Chem four-dimensional variational (4-D-Var) data assimilation system with the nested high-resolution version of the model ($0.5^\circ \times 0.67^\circ$) to quantify North American CO emissions during the period of June 2004 – May 2005. With optimized lateral boundary conditions, regional inversion analyses can reduce the sensitivity of the CO source estimates to errors in long-range transport and in the distributions of the hydroxyl radical (OH), the main sink for CO. To further limit the potential impact of discrepancies in chemical aging of air in the free troposphere, associated with errors in OH, we use surface level multispectral MOPITT CO retrievals, which have greater sensitivity to CO near the surface and reduced sensitivity in the free troposphere, compared to previous versions of the retrievals. We estimate that the annual total anthropogenic CO emission from the contiguous US 48 states was 97 Tg CO, a 14 % increase from the 85 Tg CO in the a priori. This increase is mainly due to enhanced emissions around the Great Lakes region and along the west coast, relative to the a priori. Sensitivity analyses using different OH fields and lateral boundary conditions suggest a possible error, associated with local North America OH distribution, in these emission estimates of 20 % during summer 2004, when the CO lifetime is short. This 20 % OH-related error is 50 % smaller than the OH-related error previously estimated for North American CO emissions using a global inversion analysis. We believe that reducing this OH-related error further will require integrating additional observations to provide a strong constraint on the CO distribution across the domain. Despite these limitations, our results show the potential advantages of combining high-resolution regional inversion analyses with global analyses to better quantify regional CO source estimates.

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Inverse modeling is a powerful tool to improve our understanding of emissions of greenhouse gases and pollutant tracers, by combining observations of atmospheric composition with models. Despite more than a decade of inverse modeling work to better quantify emissions of atmospheric CO (e.g., Palmer et al., 2003; Pétron et al., 2004; Heald et al., 2004; Arellano et al., 2006; Jones et al., 2009; Kopacz et al., 2010; Gonzi et al., 2011; Fortems-Cheiney et al., 2012), there is significant uncertainty in regional CO source estimates, reflecting varying source estimates from the inverse modeling analyses. As noted in previous studies, the discrepancies between the estimated CO emissions from different inversion analyses are due, in part, to errors in the atmospheric models used in the inversions. Model errors in long-range transport, vertical convective transport, diffusion, and chemistry (e.g. Arellano et al., 2006; Fortems-Cheiney et al., 2011; Jiang et al., 2013; Locatelli et al., 2013; Worden et al., 2013; Jiang et al., 2011, 2015) all adversely impact the inverse modeling of CO and other trace constituents (such as methane), and mitigating these errors in global models is challenging.

One way to reduce the effects of some model errors is to carry out the model simulations at high spatial resolution, which allows an improved description of small-scale processes, particularly those associated with vertical convection and diffusion. There have been several studies using high-resolution mesoscale models for inversion analyses (e.g. Stroud et al., 2011; Valin et al., 2011; Kilch et al., 2014; Stock et al., 2014) with the lateral boundary conditions provided from global, coarse resolution models (e.g. Curci et al., 2010; Peylin et al., 2011). However, the consistency of boundary conditions becomes a critical issue in these regional analyses (e.g. Gockede et al., 2010). The boundary conditions have also been imposed based on independent data, such as aircraft in-situ measurements (e.g. Brioude et al., 2012; Lauvaux et al., 2012; Wecht et al., 2014).

Regional inverse modeling of CO emissions with adequate boundary condition optimization will also reduce the impact of discrepancies in long-range transport and in the

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

boundary conditions is also problematic because observational coverage is often limited in space and time. A better approach for imposing the boundary conditions is to assimilate satellite observations that can provide a strong constraint on the distribution of CO throughout the free troposphere. Here we explore the use of the MOPITT data to constrain the lateral boundary conditions as well as the surface CO emissions. We also examine the potential impact of discrepancies in the abundance of the hydroxyl radical (OH), the main CO sink, on the estimates CO sources in a regional inverse modeling context.

This paper is organized as follows: in Sect. 2 we describe the MOPITT instruments and the GEOS-Chem model. In Sect. 3 we outline the inversion framework used in this work. In Sect. 4, we describe our approach for initial and boundary condition optimization, and present the estimated monthly mean North American emissions. The sensitivity of the source estimates to the chemical sink is examined by comparing the inversion results obtained with two different OH fields. Our conclusions are then provided in Sect. 5.

2 Observations and model

2.1 MOPITT

The MOPITT instrument was launched on 18 December 1999, on NASA's Terra spacecraft. We employ the multispectral version 5 (V5J) retrievals, in which the thermal infrared (TIR) radiances at $4.7\ \mu\text{m}$ are combined the near infrared (NIR) radiances at $2.3\ \mu\text{m}$ to provide greater sensitivity to lower tropospheric CO over land (Worden et al., 2010; Deeter et al., 2011). The retrievals are conducted with respect to the logarithm of the volume mixing ratio (VMR), and are reported on a 10-level pressure grid (surface, 900, 800, 700, 600, 500, 400, 300, 200, and 100 hPa). Although we use only the surface level MOPITT retrievals in our analysis, it is necessary to transform the modeled CO profile to account for the vertical resolution of the MOPITT retrieval. This

transformation is carried out using the following observation operator

$$F(\mathbf{x}) = \mathbf{y}_a + \mathbf{A}(H(\mathbf{x}) - \mathbf{y}_a) \quad (1)$$

where \mathbf{A} is the MOPITT averaging kernel, $H(\mathbf{x})$ is the GEOS-Chem profile of CO (interpolated onto the MOPITT retrieval grid), and \mathbf{y}_a is the MOPITT a priori profile. After transforming the modeled profile, the modeled CO at the surface is compared to the surface level MOPITT CO, as described in Eq. (2) in Sect. 3. Deeter et al. (2012, 2013) evaluated the multispectral MOPITT data and reported a small positive bias of 2.7 % at the surface and a larger positive bias of 14 % at 200 hPa for the V5J data. The large bias in the upper troposphere is not an issue here since we focus on the surface level data. Further details for the MOPITT instrument and the multispectral retrievals are given in Jiang et al. (2015).

2.2 GEOS-Chem

The GEOS-Chem global chemical transport model (CTM) (<http://www.geos-chem.org>) is driven by assimilated meteorological fields from the NASA Goddard Earth Observing System (GEOS-5) at the Global Modeling and data Assimilation Office. The standard GEOS-Chem chemical mechanism includes 43 tracers, and simulates a detailed description of tropospheric O_3 - NO_x -hydrocarbon chemistry, including the radiative and heterogeneous effects of aerosols. The native horizontal resolution of GEOS-5 is $0.5^\circ \times 0.667^\circ$, but the meteorological fields are usually degraded to $4^\circ \times 5^\circ$ or $2^\circ \times 2.5^\circ$ for global-scale simulations.

Our analysis is based on the CO-only simulation in GEOS-Chem v8-02-01, with relevant updates through v9-01-01, using archived monthly OH fields from the full chemistry run. The standard OH field used in this work is from GEOS-Chem version v5-07-08 (Evans et al., 2005). In order to study the influence of the OH distribution on the inversion analyses, we also archive the OH fields from a v8-02-01 GEOS-Chem full chemistry simulation. Additional details about OH fields and emission inventories can be found in Jiang et al. (2015). Briefly, the annual North America sources of CO

are 134 Tg CO from fossil fuel and biofuel combustion and biomass burning, 61 Tg CO from the oxidation of biogenic volatile organic compounds (VOCs), and 71 Tg CO from the oxidation of CH₄. Figure 1 shows the distribution of the annual mean CO emissions for June 2004 to May 2005.

The inversion analyses here are carried out using the GEOS-Chem four-dimensional variational (4-D-Var) data assimilation system, which was first described by Henze et al. (2007) and has been widely used in the chemical assimilation of CO and other tracer gases (e.g. Kopacz et al., 2009, 2010; Singh et al., 2011, Wells et al., 2014, Deng et al., 2014). Previous GEOS-Chem CO inversion analyses were conducted with the global version of the model. Here we extend the 4-D-Var system to enable regional inverse modeling of CO using the nested version of GEOS-Chem.

The nested capability was first implemented in GEOS-Chem by Wang et al. (2004) for the GEOS-3 version of the meteorological fields. The model was then updated by Chen et al. (2009) to support the GEOS-5 meteorological fields, with 0.5° × 0.667° resolution, which are used here. In the nested simulation the boundary conditions are based on fields archived from a global simulation (at 4° × 5° or 2° × 2.5°) with a 3 h temporal resolution, which are used to rewrite the tracer concentrations in a buffer zone around the nested domain before every transport step. Along the boundary of the nested domain, the direction of the wind field is used to identify whether the flow is directed into or out of the domain, and the mixing ratios of the tracers in the buffer zone are used to provide the necessary upstream information. A key benefit of using the nested model was shown by Wang et al. (2004), who found that the CO mixing ratios in the high-resolution nested simulation were lower than in the coarse resolution global model, which they attributed to the failure of the coarse global model to capture subgrid vertical motions. Figure 2 shows the simulated CO mixing ratio on 1 May 2006, obtained with the 4° × 5° global simulation and with the nested North America simulation. The yellow box in Fig. 2 indicates the buffer zone in which the boundary conditions are applied. As shown in the figure, the high-resolution CO distribution better reflects the

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



influence of the mid-latitude cyclone present over central North America, and the urban emission centers can be more clearly identified in the high-resolution simulation.

3 Inversion approach

The inverse method seeks an optimal estimate of the CO sources that is consistent with both the observed atmospheric concentrations and the a priori constraints on the sources by minimizing the cost function $J(\mathbf{x})$

$$J(\mathbf{x}) = \sum_{i=1}^N (F_i(\mathbf{x}) - \mathbf{y}_i)^T \mathbf{S}_\Sigma^{-1} (F_i(\mathbf{x}) - \mathbf{y}_i) + (\mathbf{x} - \mathbf{x}_a)^T \mathbf{S}_a^{-1} (\mathbf{x} - \mathbf{x}_a) \quad (2)$$

where \mathbf{x} is the state vector of emissions, N is the total number of observations assimilated over the assimilation window (which is one month), \mathbf{y}_i is the i th vector of observed concentrations (the MOPITT surface level retrievals), and $F(\mathbf{x})$ is the forward model, which accounts for the vertical smoothing of the MOPITT retrieval and is described in Eq. (1). Here \mathbf{x}_a is the a priori estimate and \mathbf{S}_Σ and \mathbf{S}_a are the observational and a priori error covariance matrices, respectively. The first term on the right in Eq. (2) represents the mismatch between the simulated and observed concentrations weighted by the observation error covariance. The second term represents the departure of the estimate from the a priori. The cost function is iteratively minimized using the L-BFGS algorithm (Liu and Nocedal, 1989). The inversion approach is exactly the same as described in Jiang et al. (2015). We, therefore, refer the reader to Jiang et al. (2015) for details of the optimizing scheme, the MOPITT data selection criteria, and the specification of the error covariance matrices. We employed an Observing System Simulation Experiments (OSSE) to evaluate our high-resolution 4-D-Var system in the Appendix, which suggested the nested inversion has similar reliability as the global scale assimilation system.

4 Results and discussion

4.1 Optimization on the initial and boundary conditions

We produce initial conditions following the approach of Jiang et al. (2015), by assimilating MOPITT V5J tropospheric profile data using the sequential sub-optimal Kalman filter (Parrington et al., 2008) from 1 January 2004 to 1 May 2005. Because of the bias in the V5J data at 200 hPa, we assimilate the profile data only below 200 hPa. The optimized CO distribution from the Kalman filter is archived at the beginning of each month, providing the initial conditions at the beginning of each month for the 4-D-Var inversion analyses.

As mentioned above, the lateral boundary conditions for the nested simulation could be specified from the global model. However, a better approach would be to constrain the global-scale and regional-scale emissions within the same inversion framework, so that the optimized emissions on the global-scale will provide less biased boundary conditions for the regional inversion. Such an approach has been used to constrain CH₄ and N₂O emissions over South America and Europe (Meirink et al., 2008; Bergamaschi et al., 2010; Corazza et al., 2011) with the nested TM5 model. An issue with this approach is that the adjustment in the emissions on the global scale will have to be projected through long-range transport to the nested domain. If there are any biases in the model transport, those biases will also be projected onto the nested inversion.

Because the GEOS-Chem nesting is one-way, we cannot implement the same approach that is used in TM5. Instead, we conduct a global-scale inversion analysis and use the a posteriori CO fields as boundary conditions for the regional inversion. The a posteriori simulation from the global scale inversion should provide less biased boundary conditions for the regional scale inversion than the free running model (without assimilation). However, as mentioned above, the boundary conditions could potentially be problematic if there are biases in the model transport. Alternatively, one could optimize the model CO distribution (using the sequential sub-optimal Kalman filter, for example), over writing any potential discrepancy in the tracer distribution associated

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with errors in the model transport (or chemistry). Here we compare the CO distribution obtained from optimizing the CO distribution using the Kalman filter and from optimizing the CO sources using the 4-D-Var system.

The relative differences in the CO distribution obtained from these two assimilation approaches are shown in Fig. 3. The assimilation of the MOPITT tropospheric profiles with the sequential sub-optimal Kalman filter from 1 January 2004 to 1 June 2005 is referred as CO_KF. The a posteriori CO distribution obtained from optimizing the monthly mean CO emissions using the 4-D-Var scheme is referred as CO_EMS. Shown in Fig. 3 are the relative differences of the lower tropospheric (surface–500 hPa) partial columns, calculated as $(CO_EMS - CO_KF)/CO_KF$. Since both approaches used the same initial conditions, archived from the Kalman Filter assimilation, at the beginning of each month, the relative differences shown in Fig. 3 can be considered as the residual bias in the a posteriori simulation that cannot be effectively removed within the one-month assimilation period by adjusting only the surface emissions. The most significant feature is the positive residual bias along the Intertropical Convergence Zone (ITCZ), suggesting errors in convective transport in the model (e.g., Jiang et al., 2013; Worden et al., 2013). There are also large regional differences (positive and negative) in both hemispheres that vary from month to month that are likely due to errors in chemistry and transport. For example, there are significant differences over Central America and the Gulf of Mexico in June–September 2004.

Since the objective of this work is to constrain the North American CO emissions, using CO_EMS for the boundary conditions may lead to biases in Mexico, the southern US, and along the North American west coast. Consequently, we decided to use the a posteriori fields from the Kalman filter as our optimized boundary conditions. The impact of the initial and boundary conditions from the Kalman filter assimilation is shown in Fig. 4. Driven with original initial and boundary conditions (without assimilation), the modeled CO columns (Fig. 4b) are obviously lower than that of the MOPITT observations (Fig. 4a), whereas the difference is much smaller when the initial and boundary conditions are optimized (Fig. 4c) with the assimilation of the MOPITT profiles. It is clear

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

that a significant bias will be introduced in the a posteriori regional emission estimates if the original initial and boundary conditions were used in the inversion analyses. Note, that as described in Jiang et al. (2015), we do not assimilate MOPITT data at high latitudes due to a potential positive bias in the CO retrievals at high latitudes; we only assimilate MOPITT data equator-ward of 40° over oceans and 52° over land, as shown in Fig. 4.

The distribution of the relative differences between the modeled and observed CO fields is shown in Fig. 4d. With both the free running model and the optimized initial and boundary conditions, the distribution of the differences with respect to the MOPITT data are approximately Gaussian. The free running model has a low bias of -13.3% . Assimilating the MOPITT profile data to optimize the initial and boundary conditions reduced the mean bias to 3.5% , which should produce a better constraint on local North American emissions.

4.2 CO source estimates for June 2004–May 2005

Figure 5 shows the monthly scaling factors, which are the ratio of the a posteriori to a priori emissions, for June 2004–May 2005. The figure shows enhancement of anthropogenic CO emissions in the Great Lakes region and along the US west coast. The annual total anthropogenic emission for the contiguous US 48 states is increased by 14% , from 85 to 97 Tg. This estimate is consistent with the results of our global inversion analysis presented in Jiang et al. (2015). The annual total North America CO emission from the oxidation of biogenic VOCs is reduced by 17% , from 61 to 51 Tg with the largest reduction around the Gulf of Mexico in July–September 2004. A possible reason for this reduction is the overestimation of isoprene emission in the MEGAN 2.0 inventory used in this work. As discussed in Jiang et al. (2015) several previous studies have suggested that the MEGAN 2.0 isoprene emissions are biased high over North America.

The time series for the a priori and a posteriori estimates for different emission categories are shown in Fig. 6 for all of North America ($15\text{--}65^\circ\text{N}$) and for the contiguous

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

US 48 states. For the whole continent, generally, the “bottom-up” inventory shows high CO emissions in summer and lower values in winter. This seasonal variation is driven by the oxidation of biogenic VOCs, which is significant in May–September and peaks in July, and by biomass burning, which is at a maximum in April in Mexico and in August in boreal Canada. The uncertainty of the anthropogenic emissions is assumed to be small. The analysis, however, does suggest greater anthropogenic emissions in January–May 2005, which accounts for the larger total CO emissions for this period, as shown in Fig. 6.

The United States is the largest CO source in North America, contributing 63% to the total North America source. The monthly a priori source in July is 16 Tg, twice as large as the source in winter. The distinct seasonal variation is driven by the strong biogenic VOCs source in summer. During January–April 2005, the total a posteriori CO source in this region is 59 Tg, 36% higher than the a priori value. On the contrary, during June–August 2004, the total a posteriori CO source in this region is 31 Tg, 29% lower than the a priori value. This significant discrepancy between summer and winter was also observed by Kopacz et al. (2010). The estimated winter emissions of Kopacz et al. (2010) are about 50% larger than the summer emissions. Kopacz et al. (2010) and Stein et al. (2014) attributed the low bias of Northern Hemisphere CO in winter to an underestimation of road traffic emission.

The monthly total CO emissions for the contiguous US 48 states agrees well with the results from the global $4^\circ \times 5^\circ$ resolution inversion of Jiang et al. (2015). The largest difference is observed in December 2004, when the a posteriori emission estimate from the coarse-resolution inversion is 17% higher. The smallest difference is observed in October 2004, when the a posteriori emission estimate of the coarse-resolution inversion is 2% higher.

Although there is good agreement at continental scales between our high-resolution inverse analysis and the course-resolution inversion of Jiang et al. (2015), we do observe significant differences at regional scales. In June 2004, the high-resolution inversion shows large changes, along the west coast, whereas the spatial variation is much

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

smaller in the coarse-resolution inversion, due to averaging on the model grid. More pronounced discrepancies are observed in December 2004. The coarse-resolution inversion shows the a priori CO emission estimates over Eastern US should be increased by 20–50 %. However, the high-resolution inversion shows much more variation on regional or urban scales. For example, there is a significant CO emission decrease in the Toronto area, whereas there are large increases to the east of Lake Ontario. There is also significant CO emission reduction in West Virginia, and large increases in North and South Carolina. The greater spatial structure in the regional emission estimates is somewhat expected because of the higher spatial resolution, but it is unclear as to how reliable these features are, given the information content of the MOPITT data.

The Intercontinental Transport Experiment-North America, Phase A (INTEX-A) campaign was conducted during 1 July–15 August 2004, over North America (Singh et al., 2006). A DC-8 aircraft was used to measure gas and aerosol abundances, including CO, NO₂, formaldehyde (HCHO), and H₂O, over an altitude range from 0.2 to 12.5 km. In this work, the aircraft measurements from the INTEX-A DC-8 aircraft in July 2004 are used to evaluate the inversion results obtained from MOPITT data. Figure 7 shows the difference between the GEOS-Chem simulation and the INTEX-A DC-8 aircraft observation in free troposphere. The inverse model significantly reduces the positive bias in the model bias relative to the aircraft measurements, from 7.2 to 0.5 ppb, suggesting that the a posteriori CO does indeed provide a better regional fit to the independent aircraft data.

4.3 Sensitivity of regional source estimates to OH

Following Jiang et al. (2015) we assess the sensitivity of the source estimates to the OH fields by repeating the inversion for June–August 2004 with the OH fields from version v8-02-01 of GEOS-Chem. We focus on just the summer months for comparison with Jiang et al. (2015). Also, the OH impact is expected to be greater in summer, when the CO lifetime is short. The initial and boundary conditions for this inversion were obtained by assimilating the MOPITT V5J profiles into the model with the Kalman filter and the

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

v8-02-01 OH fields from 1 January 2004 to 1 September 2004. The inversion based on these initial and boundary conditions is referred to as v8OH_BCv8. Our standard inversion with initial and boundary conditions based on the v5-07-08 OH is referred to as v5OH_BCv5. As discussed in Jiang et al. (2015), the OH concentrations in v8-02-01 are significantly higher than the v5-07-08 version in the Northern Hemisphere, and consequently, the CO lifetime is about 30 % shorter.

Figure 8a–c shows the scaling factors for the v8OH_BCv8 inversion. The differences relative to the standard inversion (v5OH_BCv5) is shown in Fig. 8d–f. The relative difference in the a posteriori CO emission estimates in the contiguous US 48 states inferred from the two OH fields is 32 %, suggesting that the OH fields still have a significant impact on the a posteriori estimates, even with the optimized boundary conditions. The relative difference of 32 % is 20 % smaller than the relative difference (40 %) obtained by Jiang et al. (2015) in their $4^\circ \times 5^\circ$ resolution global-scale inversion. Although the OH-related error is smaller than in the global, coarse-resolution inversion, it is still surprisingly large. Figure 8j–l shows the relative differences between the boundary conditions obtained from the v5-07-08 and the v8-02-01 OH fields, which were used in the v5OH_BCv5 and v8OH_BCv8 inversions, respectively. Under ideal conditions, the difference between two boundary conditions should be small because both were optimized with the same MOPITT data using the same approach. However, we do see large differences of more than 15 % along the northern and northeastern boundaries, suggesting our optimization of the boundary conditions is inadequate. We believe that the main reason for this is that we neglected MOPITT data at high latitudes (see Fig. 4) to avoid a potential positive bias in the data. Assimilating data from multi-instruments, such as was done by Kopacz et al. (2010), could provide better data coverage at high-latitudes, and thus a better constraint on the northern boundary conditions. Moving the northern boundary to lower latitude would be also helpful.

To quantify the contribution of the differences in the boundary conditions to the differences in the source estimates, we repeated the inversion using the v8-02-01 OH fields, but with the initial and boundary conditions obtained with v5-07-08 OH. This inversion is

referred to as v8OH_BCv5. Since the initial and boundary conditions in the v8OH_BCv5 and v5OH_BCv5 inversions are identical, the differences in the source estimates obtained from these will reflect only the influence of the OH differences over North America. Figure 8g–i shows the differences between v8OH_BCv5 and v5OH_BCv5 (our standard inversion). The relative difference in the a posteriori CO emission estimates for the contiguous US 48 states is only 20 %, which is 50 % smaller than the relative difference obtained by Jiang et al. (2015) in their 4° × 5° global inversion. The large reduction in the impact of OH on the source estimates compared to the global-scale inversion is encouraging, and demonstrates the potential advantages of high-resolution regional inversion analyses.

5 Summary

High-resolution CTM simulations have obvious accuracy advantages over coarse resolution simulations, particularly for small-scale processes (e.g., Stroud et al., 2011; Kilch et al., 2014; Stock et al., 2014). In this work, we used the adjoint of the nested GEOS-Chem model, at a resolution of 0.5° × 0.67°, to constrain North American CO emissions during the period of June 2004–May 2005. To reduce the potential impact of discrepancies in the modeled OH field on the source estimates, we used the V5J surface level MOPITT retrievals. Our results show that the annual total anthropogenic CO emissions for the contiguous US 48 states should be increased by 14 %, from 85 to 97 Tg for 2004–2005. The adjustment was mainly caused by an increase in emissions near the Great Lakes and along the west coast. The inversion analysis also suggested that the total CO emissions should be increased by 36 % during January–April 2005, and decreased by 29 % during June–August 2004. This seasonal variation was also observed by Kopacz et al. (2010), and could be associated with an underestimation of road traffic emission in winter (Stein et al., 2014). The inversion results were evaluated with in-situ measurements from the DC-8 aircraft during the INTEX-A campaign in July 2004. The

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mean bias between the model and the aircraft data in the free troposphere was reduced from 7.2 ppb with the a priori emissions to 0.5 ppb with the a posteriori emissions.

Reliable initial and boundary conditions are critical for regional inversion analyses. We used a sequential sub-optimal Kalman filter (Parrington et al., 2008) to assimilate MOPITT CO profiles to optimize the distribution of CO (rather than the emissions) to produce improved initial and lateral boundary conditions for the regional inversion analyses. Because of the restricted domain of the regional analyses, the optimized boundary conditions should significantly reduce the sensitivity of the estimated CO sources to errors in long-range transport and in the OH distribution. Comparison of the inversion results driven with two different OH fields, from version v5-07-08 and v8-02-01 of GEOS-Chem, produced relative differences in the North American source estimates of 32 % for June–August 2004. This OH-related difference in the source estimates is about 20 % smaller than the differences obtained by Jiang et al. (2015) for the same period in their global-scale inversion analysis.

Examination of the differences in the boundary conditions based on the two OH fields showed large relative differences (greater than 15 %) in the northern and northeastern boundaries, suggesting that our optimization of the boundary conditions was inadequate. In our assimilation of the MOPITT data we neglected data pole-ward of 52 and 40° over land and oceans, respectively, to avoid the influence of a potential high-latitude bias in the data, and we believe that this accounted for the weaker constraint on the northern boundary conditions in the analysis. To assess the influence of the boundary conditions we repeated the inversion with the v8-02-01 OH fields using the same initial and boundary conditions from the standard inversion using the v5-07-08 OH. In this case, we estimated a relative difference between the source estimates based on the v8-02-01 and v5-07-08 OH fields of 20 %, which is 50 % smaller than that reported by Jiang et al. (2010). Thus, our best estimate for North American CO emissions for 2004–2005 is 97 Tg, with a potential error of 20 %, associated with discrepancies in local North America OH.

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Our results demonstrate that high-resolution, regional inversion analyses can reduce the sensitivity of the inferred CO source estimates to errors in long-range transport and in the OH distributions. However, the 20 % OH-related discrepancy that we estimated is still large, and could indicate that more stringent constraints on the regional boundary conditions are needed. This may be achieved by integrating data from multiple sources. The OH-related discrepancies could also reflect that fact that in summer, air in the middle and upper troposphere over North America is trapped by a semipermanent anticyclone, which allows greater chemical aging than direct lateral export from the continent (Li et al., 2005; Cooper et al., 2007). Although the MOPITT surface level retrievals have peak sensitivity to CO near the boundary layer, their sensitivity extend up to the middle troposphere (see Fig. 1a of Jiang et al., 2013). Consequently, the inversion analyses could be sensitivite to chemical aging of air in the North American anticyclone. Work is needed to determine the residence time for air in the anticyclone compared to the spatio-temporal variability of the constraints on the North American source estimates provided by the MOPITT surface level retrievals. Improving the source estimates will likely require assimilating sufficient information to obtain a strong constraint on the CO distribution on a timescales shorter than the timescale for chemical aging in the domain. Despite these limitations, we believe that our results show the potential advantages of combining high-resolution regional inversion analyses with global analyses to better quantify regional CO source estimates.

Appendix: Observing System Simulation Experiments (OSSE)

The reliability of the nested inversion is examined with an OSSE for the period 1–15 June 2004. In the OSSE, we firstly create pseudo-observations, by archiving model output with CO emission unchanged. In the pseudo-inversion, we reduced the CO emission by 50 % and the objective of the OSSE is to observe whether the scaling factors can return to true state (1.0). Figure A1a shows the result of the reference global scale inversion with $4^\circ \times 5^\circ$ resolution. The a posteriori estimation converges to

the true state in all major emission regions. In the nested inversion (Fig. A1c), the model converges to the true state in Eastern US, whereas the result is noisy in Western US and Canada, which is consistent with the global scale inversion, as shown in Fig. A1b. The OSSE demonstrates the nested inversion has similar reliability as the global scale
5 assimilation system.

Acknowledgements. This work was supported by funding from the Natural Science and Engineering Research Council of Canada, the Canadian Space Agency, and NASA grants NNX10AT42G and NNX11AI54G.

References

- 10 Arellano Jr., A. F., Kasibhatla, P. S., Giglio, L., van der Werf, G. R., Randerson, J. T., and Collatz, G. J.: Time dependent inversion estimates of global biomass-burning CO emissions using Measurement of Pollution in the Troposphere (MOPITT) measurements, *J. Geophys. Res.*, 111, D09303, doi:10.1029/2005JD006613, 2006.
- 15 Bergamaschi, P., Krol, M., Meirink, J. F., Dentener, F., Segers, A., van Aardenne, J., Monni, S., Vermeulen, A. T., Schmidt, M., Ramonet, M., C., Yver, Meinhardt, F., Nisbet, E. G., R. E., Fisher, O'Doherty, S., and Dlugokencky, E. J.: Inverse modeling of European CH₄ emissions 2001–2006, *J. Geophys. Res.*, 115, D22309, doi:10.1029/2010JD014180, 2010.
- 20 Brioude, J., Petron, G., Frost, G. J., Ahmadov, R., Angevine, W. M., Hsie, E.-Y., Kim, S.-W., Lee, S.-H., McKeen, S. A., Trainer, M., Fehsenfeld, F. C., Holloway, J. S., Peischl, J., Ryerson, T. B., and Gurney, K. R.: A new inversion method to calculate emission inventories without a prior at mesoscale: application to the anthropogenic CO₂ emission from Houston, Texas, *J. Geophys. Res.*, 117, D05312, doi:10.1029/2011JD016918, 2012.
- 25 Chen, D., Wang, Y., McElroy, M. B., He, K., Yantosca, R. M., and Le Sager, P.: Regional CO pollution and export in China simulated by the high-resolution nested-grid GEOS-Chem model, *Atmos. Chem. Phys.*, 9, 3825–3839, doi:10.5194/acp-9-3825-2009, 2009.
- Cooper, O., Trainer, M., Thompson, A., Oltmans, S., Tarasick, D., Witte, J., Stohl, A., Eckhardt, S., Lelieveld, J., Newchurch, M., Johnson, B., Portmann, R., Kalnajs, L., Dubey, M., Leblanc, T., McDermid, I., Forbes, G., Wolfe, D., Carey-Smith, T., Morris, G., Lefer, B., Rappenglück, B., Joseph, E., Schmidlin, F., Meagher, J., Fehsenfeld, F., Keating, T., Curen, R. and Min-

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

schwamer, K.: Evidence for a recurring eastern North America upper tropospheric ozone maximum during summer, *J. Geophys. Res.*, 112, D23304, doi:10.1029/2007JD008710, 2007.

Corazza, M., Bergamaschi, P., Vermeulen, A. T., Aalto, T., Haszpra, L., Meinhardt, F., O'Doherty, S., Thompson, R., Moncrieff, J., Popa, E., Steinbacher, M., Jordan, A., Dlugokencky, E., Brühl, C., Krol, M., and Dentener, F.: Inverse modelling of European N₂O emissions: assimilating observations from different networks, *Atmos. Chem. Phys.*, 11, 2381–2398, doi:10.5194/acp-11-2381-2011, 2011.

Curci, G., Palmer, P. I., Kurosu, T. P., Chance, K., and Visconti, G.: Estimating European volatile organic compound emissions using satellite observations of formaldehyde from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 10, 11501–11517, doi:10.5194/acp-10-11501-2010, 2010.

Deeter, M. N., Worden, H. M., Gille, J. C., Edwards, D. P., Mao, D., and Drummond, J. R.: MOPITT multispectral CO retrievals: origins and effects of geophysical radiance errors, *J. Geophys. Res.*, 116, D15303, doi:10.1029/2011JD015703, 2011.

Deeter, M. N., Worden, H. M., Edwards, D. P., Gille, J. C., and Andrews, A. E.: Evaluation of MOPITT retrievals of lower-tropospheric carbon monoxide over the United States, *J. Geophys. Res.*, 117, D13306, doi:10.1029/2012JD017553, 2012.

Deeter, M. N., Martínez-Alonso, S., Edwards, D. P., Emmons, L. K., Gille, J. C., Worden, H. M., Pittman, J. V., Daube, B. C., and Wofsy, S. C.: Validation of MOPITT Version 5 thermal-infrared, near-infrared, and multispectral carbon monoxide profile retrievals for 2000–2011, *J. Geophys. Res. Atmos.*, 118, 6710–6725, doi:10.1002/jgrd.50272, 2013.

Deng, F., Jones, D. B. A., Henze, D. K., Bousseres, N., Bowman, K. W., Fisher, J. B., Nasar, R., O'Dell, C., Wunch, D., Wennberg, P. O., Kort, E. A., Wofsy, S. C., Blumenstock, T., Deutscher, N. M., Griffith, D. W. T., Hase, F., Heikkinen, P., Sherlock, V., Strong, K., Sussmann, R., and Warneke, T.: Inferring regional sources and sinks of atmospheric CO₂ from GOSAT XCO₂ data, *Atmos. Chem. Phys.*, 14, 3703–3727, doi:10.5194/acp-14-3703-2014, 2014.

Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32, L09813, doi:10.1029/2005GL022469, 2005.

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Szopa, S., Deeter, M. N., and Clerbaux, C.: Ten years of CO emissions as seen from Measurements of Pollution in the Troposphere (MOPITT), *J. Geophys. Res.*, 116, D05304, doi:10.1029/2010JD014416, 2011.

Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Saunio, M., Szopa, S., Cressot, C., Kurosu, T. P., Chance, K., and Fried, A.: The formaldehyde budget as seen by a global-scale multi-constraint and multi-species inversion system, *Atmos. Chem. Phys.*, 12, 6699–6721, doi:10.5194/acp-12-6699-2012, 2012.

Göckede, M., Turner, D. P., Michalak, A. M., Vickers, D., and Law, B. E.: Sensitivity of a subregional scale atmospheric inverse CO₂ modeling framework to boundary conditions, *J. Geophys. Res.*, 115, D24112, doi:10.1029/2010JD014443, 2010.

Gonzi, S., Feng, L., and Palmer, P. I.: Seasonal cycle of emissions of CO inferred from MOPITT profiles of CO: sensitivity to pyroconvection and profile retrieval assumptions, *Geophys. Res. Lett.*, 38, L08813, doi:10.1029/2011GL046789, 2011.

Heald, C. L., Jacob, D. J., Jones, D. B. A., Palmer, P. I., Logan, J. A., Streets, D. G., Sachse, G. W., Gille, J. C., Hoffman, R. N., and Nehrkorn, T.: Comparative inverse analysis of satellite (MOPITT) and aircraft (TRACE-P) observations to estimate Asian sources of carbon monoxide, *J. Geophys. Res.* 109, D23306, doi:10.1029/2004JD005185, 2004.

Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, *Atmos. Chem. Phys.*, 7, 2413–2433, doi:10.5194/acp-7-2413-2007, 2007.

Jiang, Z., Jones, D. B. A., Kopacz, M., Liu, J., Henze, D. K., and Heald, C.: Quantifying the impact of model errors on top-down estimates of carbon monoxide emissions using satellite observations, *J. Geophys. Res.*, 116, D15306, doi:10.1029/2010JD015282, 2011.

Jiang, Z., Jones, D. B. A., Worden, H. M., Deeter, M. N., Henze, D. K., Worden, J., Bowman, K. W., Brenninkmeijer, C. A. M., and Schuck, T. J.: Impact of model errors in convective transport on CO source estimates inferred from MOPITT CO retrievals, *J. Geophys. Res. Atmos.*, 118, 2073–2083, doi:10.1002/jgrd.50216, 2013.

Jiang, Z., Jones, D. B. A., Worden, H. M., and Henze, D. K.: Sensitivity of top-down CO source estimates to the modeled vertical structure in atmospheric CO, *Atmos. Chem. Phys.*, 15, 1521–1537, doi:10.5194/acp-15-1521-2015, 2015.

Jones, D. B. A., Bowman, K. W., Logan, J. A., Heald, C. L., Liu, J., Luo, M., Worden, J., and Drummond, J.: The zonal structure of tropical O₃ and CO as observed by the Tropospheric Emission Spectrometer in November 2004 – Part 1: Inverse modeling of CO emissions, *Atmos. Chem. Phys.*, 9, 3547–3562, doi:10.5194/acp-9-3547-2009, 2009.

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Junhua Liu, Logan, J. A., Jones, D. B. A., Livesey, N. J., Megretskaia, I., Carouge, C., and Nedelec, P.: Analysis of CO in the tropical troposphere using Aura satellite data and the GEOS-Chem model: insights into transport characteristics of the GEOS meteorological products, *Atmos. Chem. Phys.*, 10, 12207–12232, doi:10.5194/acp-10-12207-2010, 2010.
- 5 Klich, C. A. and Fuelberg, H. E.: The role of horizontal model resolution in assessing the transport of CO in a middle latitude cyclone using WRF-Chem, *Atmos. Chem. Phys.*, 14, 609–627, doi:10.5194/acp-14-609-2014, 2014.
- Kopacz, M., Jacob, D. J., Fisher, J. A., Logan, J. A., Zhang, L., Megretskaia, I. A., Yantosca, R. M., Singh, K., Henze, D. K., Burrows, J. P., Buchwitz, M., Khlystova, I., McMillan, W. W., Gille, J. C., Edwards, D. P., Eldering, A., Thouret, V., and Nedelec, P.: Global estimates of CO sources with high resolution by adjoint inversion of multiple satellite datasets (MOPITT, AIRS, SCIAMACHY, TES), *Atmos. Chem. Phys.*, 10, 855–876, doi:10.5194/acp-10-855-2010, 2010.
- 10 Lauvaux, T., Schuh, A. E., Uliasz, M., Richardson, S., Miles, N., Andrews, A. E., Sweeney, C., Diaz, L. I., Martins, D., Shepson, P. B., and Davis, K. J.: Constraining the CO₂ budget of the corn belt: exploring uncertainties from the assumptions in a mesoscale inverse system, *Atmos. Chem. Phys.*, 12, 337–354, doi:10.5194/acp-12-337-2012, 2012.
- Liu, D. C. and Nocedal, J.: On the limited memory method for large scale optimization, *Math. Program.*, 45, 503–528, 2010.
- 20 Locatelli, R., Bousquet, P., Chevallier, F., Fortems-Cheney, A., Szopa, S., Saunois, M., Agustí-Panareda, A., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Gloor, E., Houweling, S., Kawa, S. R., Krol, M., Patra, P. K., Prinn, R. G., Rigby, M., Saito, R., and Wilson, C.: Impact of transport model errors on the global and regional methane emissions estimated by inverse modelling, *Atmos. Chem. Phys.*, 13, 9917–9937, doi:10.5194/acp-13-9917-2013, 2013.
- 25 Meirink, J. F., Bergamaschi, P., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modelling of atmospheric methane emissions: method and comparison with synthesis inversion, *Atmos. Chem. Phys.*, 8, 6341–6353, doi:10.5194/acp-8-6341-2008, 2008.
- Parrington, M., Jones, D. B. A., Bowman, K. W., Horowitz, L. W., Thompson, A. M., Tarasick, D. W., and Witte, J. C.: Estimating the summertime tropospheric ozone distribution over North America through assimilation of observations from the Tropospheric Emission Spectrometer, *J. Geophys. Res.*, 113, D18307, doi:10.1029/2007JD009341, 2008.
- 30

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Parrington, M., Palmer, P. I., Henze, D. K., Tarasick, D. W., Hyer, E. J., Owen, R. C., Helmig, D., Clerbaux, C., Bowman, K. W., Deeter, M. N., Barratt, E. M., Coheur, P.-F., Hurtmans, D., Jiang, Z., George, M., and Worden, J. R.: The influence of boreal biomass burning emissions on the distribution of tropospheric ozone over North America and the North Atlantic during 2010, *Atmos. Chem. Phys.*, 12, 2077–2098, doi:10.5194/acp-12-2077-2012, 2012.
- Pétron, G., Granier, C., Khattatov, B., Yudin, V., Lamarque, J.-F., Emmons, L., Gille, J., and Edwards, D. P.: Monthly CO surface sources inventory based on the 2000–2001 MOPITT satellite data, *Geophys. Res. Lett.*, 31, L21107, doi:10.1029/2004GL020560, 2004.
- Peylin, P., Houweling, S., Krol, M. C., Karstens, U., Rödenbeck, C., Geels, C., Vermeulen, A., Badawy, B., Aulagnier, C., Pregger, T., Delage, F., Pieterse, G., Ciais, P., and Heimann, M.: Importance of fossil fuel emission uncertainties over Europe for CO₂ modeling: model inter-comparison, *Atmos. Chem. Phys.*, 11, 6607–6622, doi:10.5194/acp-11-6607-2011, 2011.
- Singh, H. B., Brune, W. H., Crawford, J. H., Jacob, D. J., and Russell, P. B.: Overview of the summer 2004 Intercontinental Chemical Transport Experiment–North America (INTEX-A), *J. Geophys. Res.*, 111, D24S01, doi:10.1029/2006JD007905, 2006.
- Singh, K., Jardak, M., Sandu, A., Bowman, K., Lee, M., and Jones, D.: Construction of non-diagonal background error covariance matrices for global chemical data assimilation, *Geosci. Model Dev.*, 4, 299–316, doi:10.5194/gmd-4-299-2011, 2011.
- Stein, O., Schultz, M. G., Bouarar, I., Clark, H., Huijnen, V., Gaudel, A., George, M., and Clerbaux, C.: On the wintertime low bias of Northern Hemisphere carbon monoxide found in global model simulations, *Atmos. Chem. Phys.*, 14, 9295–9316, doi:10.5194/acp-14-9295-2014, 2014.
- Stock, Z. S., Russo, M. R., and Pyle, J. A.: Representing ozone extremes in European megacities: the importance of resolution in a global chemistry climate model, *Atmos. Chem. Phys.*, 14, 3899–3912, doi:10.5194/acp-14-3899-2014, 2014.
- Stroud, C. A., Makar, P. A., Moran, M. D., Gong, W., Gong, S., Zhang, J., Hayden, K., Mihele, C., Brook, J. R., Abbatt, J. P. D., and Slowik, J. G.: Impact of model grid spacing on regional- and urban-scale air quality predictions of organic aerosol, *Atmos. Chem. Phys.*, 11, 3107–3118, doi:10.5194/acp-11-3107-2011, 2011.
- Valin, L. C., Russell, A. R., Hudman, R. C., and Cohen, R. C.: Effects of model resolution on the interpretation of satellite NO₂ observations, *Atmos. Chem. Phys.*, 11, 11647–11655, doi:10.5194/acp-11-11647-2011, 2011.

**Regional data
assimilation of
multi-spectral
MOPITT observations
of CO over North
America**

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Wang, Y. X., McElroy, M. B., Jacob, D. J., and Yantosca, R. M.: A nested grid formulation for chemical transport over Asia: applications to CO, *J. Geophys. Res.*, 109, D22307, doi:10.1029/2004JD005237, 2004.

5 Wecht, K. J., Jacob, D. J., Sulprizio, M. P., Santoni, G. W., Wofsy, S. C., Parker, R., Bösch, H., and Worden, J.: Spatially resolving methane emissions in California: constraints from the CalNex aircraft campaign and from present (GOSAT, TES) and future (TROPOMI, geostationary) satellite observations, *Atmos. Chem. Phys.*, 14, 8173–8184, doi:10.5194/acp-14-8173-2014, 2014.

10 Wells, K. C., Millet, D. B., Cady-Pereira, K. E., Shephard, M. W., Henze, D. K., Bousseres, N., Apel, E. C., de Gouw, J., Warneke, C., and Singh, H. B.: Quantifying global terrestrial methanol emissions using observations from the TES satellite sensor, *Atmos. Chem. Phys.*, 14, 2555–2570, doi:10.5194/acp-14-2555-2014, 2014.

15 Worden, H. M., Deeter, M. N., Edwards, D. P., Gille, J. C., Drummond, J. R., and Nédélec, P.: Observations of near-surface carbon monoxide from space using MOPITT multispectral retrievals, *J. Geophys. Res.*, 115, D18314, doi:10.1029/2010JD014242, 2010.

Worden, J., Jiang, Z., Jones, D., Alvarado, M., Bowman, K., Frankenberg, C., Kort, E., Kulawik, S., Lee, M., Liu, J., Payne, V., Wecht, K., and Worden, H.: El Niño, the 2006 Indonesian peat fires, and the distribution of atmospheric methane, *Geophys. Res. Lett.*, 40, 4938–4943, doi:10.1002/grl.50937, 2013.

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 1. Annual mean CO emissions from fossil fuel (FF), biofuel (BF), biomass burning (BB) and the oxidation of biogenic NMVOC and CH₄, averaged for June 2004 to May 2005. The unit is 10^{12} molec cm⁻² s⁻¹.

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 2. CO mixing ratio in GEOS-Chem on level 10 (about 850 hPa) on 1 May 2006. The influence of a mid-latitude cyclone is clearly shown in the high-resolution ($0.5^\circ \times 0.667^\circ$) simulation (right), whereas it is not obvious in that with coarse resolution ($4^\circ \times 5^\circ$) simulation (left). The light yellow line on the $0.5^\circ \times 0.667^\circ$ plot demarcates the buffer zone in which the coarse resolution boundary conditions are imposed.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 3. Relative difference between the optimized lower tropospheric partial columns (surface–500 hPa) between Kalman Filter assimilation, referred as CO_KF and a posteriori simulation of global scale inversion, referred as CO_EMS. The value is calculated as $(CO_EMS - CO_KF)/CO_KF$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 4. Mean tropospheric CO columns ($10^{18} \text{ molec cm}^{-2}$) in the GEOS-Chem North America nested domain in June 2004 from (a) MOPITT version 5; (b) GEOS-Chem model with the original initial and boundary conditions; (c) with the optimized initial and boundary conditions. (d) Relative difference, $(\text{Mod} - \text{MOP})/\text{MOP}$, using original (red), and optimized (blue) initial and boundary conditions. The numbers in panel (d) are mean bias. Note that MOPITT data poleward of 40° and 52° over oceans and land, respectively, are not used in this work to reduce the influence of potential positive bias in MOPITT CO retrievals, as described in Jiang et al. (2014).

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 5. Monthly scaling factor for total CO emissions (from combustion sources and the oxidation of biogenic NMVOC) during June 2004–May 2005.

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 6. Monthly CO emissions during June 2004–May 2005 for different emission categories: total emissions (black), anthropogenic emissions (blue), biomass burning (red), and the oxidation of biogenic NMVOCs (green). The a priori estimates are shown with the solid line and a posteriori values are indicated with the dashed line. The unit is $\text{Tg}(\text{month})^{-1}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 7. Difference between the GEOS-Chem simulation with INTEX-A DC-8 aircraft observation in free troposphere in July 2004. **(a)** A priori model simulation. **(b)** A posteriori model simulation. The model is sampled at the aircraft measurements time, location and altitude.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure 8. (a–c): Scaling factors of v8OH_BCv8 inversion, based on the v8-02-01 OH; (d–f): difference between scaling factors of the v5OH_BCv5 and v8OH_BCv8 inversions; (g–i): similar to panels (d–f), but for the v8OH_BCv5 inversion, with the initial and boundary conditions from the standard inversion with the v5-07-08 OH fields; (j–l) relative difference between the lower tropospheric CO partial columns (surface–500 hPa) of the boundary conditions for the v5OH_BCv5 (v5) and the v8OH_BCv8 (v8) inversions, calculated as: $2 \cdot (v8 - v5) / (v8 + v5)$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regional data assimilation of multi-spectral MOPITT observations of CO over North America

Z. Jiang et al.




Figure A1. OSSE scaling factors for 1–15 June 2004. **(a)** Global reference inversion with $4^{\circ} \times 5^{\circ}$ resolution. **(b)** Zoomed in North America region of the global inversion. **(c)** Nested inversion results. The scaling factor for the first guess is 0.5 and for the true state is 1.0.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)