

AMENABLE GROUPS WITHOUT FINITELY PRESENTED AMENABLE COVERS

MUSTAFA GÖKHAN BENLİ, ROSTISLAV GRIGORCHUK
AND PIERRE DE LA HARPE
WITH A POST-SCRIPTUM BY RALPH STREBEL

ABSTRACT. The goal of this article is to study results and examples concerning finitely presented covers of finitely generated amenable groups. We collect examples of groups G with the following properties: (i) G is finitely generated, (ii) G is amenable, e.g. of intermediate growth, (iii) any finitely presented group with a quotient isomorphic to G contains non-abelian free subgroups, or the stronger (iii') any finitely presented group with a quotient isomorphic to G is large.

After submission of the final version of this paper to the *Bulletin of Mathematical Sciences* <http://www.springer.com/birkhauser/mathematics/journal/13373>, we have received a note of Ralph Strebél on an construction of Yves de Cornulier, providing a strenghtening of Corollary C.6. This note, slightly modified, is added as a post-scriptum to the present file.

1. Introduction

1.A. Motivation. In the study of finiteness conditions on groups, the following type of question is natural:

Question 1.1. *Given a Property (\mathcal{P}) of groups, does any finitely generated group with (\mathcal{P}) have a finitely presented cover with (\mathcal{P}) ?*

A **cover** of a group G is a group E given together with an epimorphism $E \twoheadrightarrow G$. A *cover* fits into an *extension* $1 \rightarrow N \rightarrow E \twoheadrightarrow G \rightarrow 1$.

Date: March 20, 2013, and April 27 for the Post Scriptum.

2000 Mathematics Subject Classification. 20E08, 20E22, 20F05 43A07.

Key words and phrases. Finitely generated groups, finitely presented covers, self-similar groups, metabelian groups, soluble groups, groups of intermediate growth, amenable groups, groups with non-abelian free subgroups, large groups.

We are grateful to the Swiss National Science Foundation for supporting a visit of the first author to Geneva. The first author is grateful to the department of mathematics, University of Geneva, for their hospitality during his stay. The first and the second authors are partially supported by NSF grant DMS – 1207699.

The answer can be positive for trivial reasons, for example when Property (\mathcal{P}) holds for free groups (such as exponential growth) or when Property (\mathcal{P}) implies finite presentation (such as nilpotency, or polynomial growth). The answer is also positive when (\mathcal{P}) is Kazhdan's Property (\mathcal{T}) , by a non-trivial result of Shalom [Shal-00].

Here, we concentrate on Question 1.1 for amenability, a case in which the answer is negative. Indeed, the goal of this article is to study examples and results concerning *finitely generated amenable groups that do not have finitely presented amenable covers*.

Our motivation is better to understand the class of *finitely presented amenable groups*, and related groups. To describe the situation, it is convenient to introduce the class \mathcal{AG} of amenable groups, and the subclass \mathcal{EG} of elementary amenable groups, which is easier to understand (Appendix D is a reminder on these classes, and on growth).

On the one hand, many examples are known in \mathcal{EG} of finitely presented soluble groups, including the standard examples described in the exposition [Stre-84], the Baumslag-Remeslennikov group, which is a finitely presented metabelian group with derived group free abelian of infinite rank (references in our Remark A.4.c), and the Kharlampovich group, which is a finitely presented soluble group with unsolvable word problem (references in the proof of Proposition A.5). There are also finitely presented groups in \mathcal{EG} which are not virtually soluble (Example D.1).

In the complement $\mathcal{AG} \setminus \mathcal{EG}$, a reasonable number of *finitely generated* examples are known, in particular the *groups of intermediate growth*. Also known are *finitely presented* examples, all closely related to self-similar groups. But not enough examples are known to allow one to guess at a general picture; this is illustrated by the following old and basic problem in the theory of growth of groups (see, e.g., [Kour-06], Problems 4.5.b, 9.8, and 10.11):

Basic problem 1.2. *Does there exist a finitely presented group of intermediate growth?*

The answer does not seem to be at hand. Here is a possibly easier problem:

Problem 1.3. *Does there exist a finitely generated group of intermediate growth that is the quotient of a finitely presented group without non-abelian free subgroups? or the quotient of a finitely presented amenable group?*

Finitely generated elementary amenable groups are never of intermediate growth [Chou-80], so that Problems 1.2 and 1.3 do not arise

in \mathcal{EG} . Problem 1.3 involves *two* questions, because there exist finitely presented groups that are non-amenable and do not contain non-abelian free subgroups [OlSa-02].

Strictly speaking, the solution of the analogue of Problem 1.2 is known for the class $\mathcal{AG} \setminus \mathcal{EG}$. Indeed, one knows a few sporadic examples of finitely presented groups in this class, such as the HNN-extension $\widehat{\mathfrak{G}}$ of the Grigorchuk group [Grig-98], and the HNN-extension of the Brunner-Sidki-Vieira group that appears in [GrZ-02b, Proposition 6] (the latter group is amenable by [BaKN-10] and not elementary amenable by [GrZ-02b]). The finitely presented HNN-extension of the basilica group that appears in [GrZ-02b, Theorem 1.7] is not even in the class $\mathcal{AG} \setminus \mathcal{SG}$, defined in Appendix D; see also [BaVi-05, Theorem 12]. But, despite these examples, very little is known about finitely presented groups in $\mathcal{AG} \setminus \mathcal{EG}$, and it would be interesting to find methods providing new examples; a priori, one could hope and try finitely presented covers, but our paper shows that this does not look very promising.

Before stating the next problem, here is a definition: the **elementary amenable radical** $\text{Rad}_{\text{ea}}(G)$ of a group G is its unique maximal normal elementary amenable subgroup. Note that $\text{Rad}_{\text{ea}}(G)$ is contained in the amenable radical of G , that appears (but for its name) in [Day-57, Lemma 1 of Section 4].

Problem 1.4. *Is there a finitely presented group G in $\mathcal{AG} \setminus \mathcal{EG}$, with $\text{Rad}_{\text{ea}}(G) = \{1\}$, that has infinitely presented¹ quotients? or that has uncountably many pairwise non-isomorphic quotients?*

1.B. First examples. When (\mathcal{P}) is amenability, we know two approaches to Question 1.1: One is based on the theory of Σ -invariants, developed in a series of papers by Bieri, Strebel, and Neumann (see Appendix C). The other involves self-similar groups and approximation methods (see Sections 2 and 3).

The first result we quote is due to Bieri and Strebel [BiSt-80, Theorem 5.5 and Corollary 5.6]. On the one hand, we reformulate it in a slightly weaker version, assuming that E is finitely presented (instead of assuming that E has Property FP_2 , as in the original paper); on the other hand, we formulate it for *virtually* metabelian groups, because this follows immediately from the case of metabelian groups.

Theorem 1.5 (Bieri-Strebel). *Let G be a virtually metabelian group that is finitely generated and infinitely presented.*

¹A finitely generated group is “infinitely presented” if it is not finitely presented.

Any finitely presented cover E of G has non-abelian free subgroups. In particular, E is non-soluble, indeed non-amenable.

The proof of Theorem 1.5 can now be understood in a much simpler way than in [BiSt–80], as we indicate in Corollary C.4. The reason is that we can use better invariants, namely those defined in [BiNS–87] and their later reformulations (see Strebel’s exposition in progress [Stre]).

Examples of metabelian groups that are finitely generated and infinitely presented include matrix groups like $\left(\begin{pmatrix} \frac{\ell}{m} & \mathbf{Z} \\ 0 & \mathbf{Z} \left[\frac{1}{\ell m} \right] \end{pmatrix} \right)$ where $\ell, m \geq 2$ are coprime integers, wreath products $A \wr \mathbf{Z}$ where $A \neq \{1\}$ is a finitely generated abelian group, and free metabelian groups $\text{FSol}(k, 2) := F_k / [[F_k, F_k], [F_k, F_k]]$, where F_k is the free group on $k \geq 2$ generators.

Appendix A includes a reminder on metabelian groups. Appendix B is a reminder on wreath products and lamplighter groups. Appendix C is a reminder on Bieri-Neumann-Strebel invariants. Proposition B.2, Corollary B.3, Proposition B.9, Corollary C.4, and Corollary C.6 give examples of finitely generated groups of which all finitely presented covers have non-abelian free subgroups.

We denote by \mathfrak{G} the *Grigorchuk group*², introduced in [Grig–80]; see Example 2.16. Recall here that \mathfrak{G} is finitely generated, indeed generated by four involutions traditionally written a, b, c, d . This group has many remarkable properties, including that of being an infinite 2-group of intermediate growth; in particular it is amenable. The group \mathfrak{G} has a presentation, due to Lysenok, with four generators and infinitely many relators. Appropriate finite subsets of these relators naturally define a sequence $(\mathfrak{G}_n)_{n \geq 0}$ of four-generated finitely presented groups converging to \mathfrak{G} (Definition 5.2).

There is a reminder on convergence of groups in Section 3.

Theorem 1.6 ([GrHa–01]). *Any finitely presented cover of the group \mathfrak{G} is large.*

Recall that a group is **large** if it contains a subgroup of finite index that has non-abelian free quotients. Note that large groups have non-abelian free subgroups.

In view of Proposition 3.3 below, Theorem 1.6 is a straightforward consequence of the main result of [GrHa–01]; see also Theorem 2.14 and Example 2.16. More precisely, it was shown in [GrHa–01] that

²Two of the authors insist that we mention this terminology, often used today. The third author submits.

each \mathfrak{G}_n is virtually a direct product of finitely generated non-abelian free groups; this has been sharpened in [BaCo–06], and can be further improved:

Theorem 1.7 (Section 5). *Let \mathfrak{G} and \mathfrak{G}_n be as above. For each $n \geq 0$, the group \mathfrak{G}_n has a normal subgroup H_n of index $2^{2^{n+1}+2}$ that is isomorphic to the direct product of 2^n free groups of rank 3.*

Another way of sharpening Theorem 1.6 is given in Proposition 5.11, where the condition of finite presentability for the cover of \mathfrak{G} is replaced by the weaker Condition FP_2 .

We denote by \mathfrak{B} the Basilica group. Recall here that \mathfrak{B} can be generated by two elements, and is an amenable torsion-free group of exponential growth. See Example 2.17 and Appendix D for some other properties of \mathfrak{B} .

Theorem 1.8 (Erschler). *Any finitely presented cover of the Basilica group \mathfrak{B} has non-abelian free subgroups.*

Given an invertible automaton (A, τ) over a finite alphabet X , Erschler introduces a notion of “automatically presented group $G^*(A, \tau)$ generated by the automaton” (see [Ersc–07] for details). She shows that, if $G^*(A, \tau)$ is not virtually abelian, then any finitely presented cover of it has non-abelian free subgroups. For the Basilica automaton, the group $G^*(A, \tau)$ coincides with \mathfrak{B} , and Theorem 1.8 follows.

As noted in [Ersc–07], these arguments do not apply to the group \mathfrak{G} of Theorem 1.6; this is due to the fact that, for (A, τ) the automaton of \mathfrak{G} , the cover $G^*(A, \tau) \twoheadrightarrow \mathfrak{G}$ has a non-trivial kernel.

In our setting, for $G = \mathfrak{B}$, the universal contracting cover G_0 of Definition 2.5 is free of rank 2 (see Example 2.17), and Theorem 1.8 follows from our Theorem 2.14. Indeed, our argument shows that

any finitely presented cover of \mathfrak{B} is large.

Notation: In this paper, the symbols \mathfrak{G} and \mathfrak{B} will be used *only* for the two groups introduced above. In Examples 2.18 to 2.21, we introduce four other groups, with their usual notation in this subject: the iterated monodromy group of $z^2 + i$ denoted by \mathfrak{J} , the Gupta-Sidki group \mathfrak{GS} , the Fabrykowski-Gupta group \mathfrak{FG} , and the Hanoi Towers group \mathfrak{H} .

We show in Section 2 that Theorem 2.14 implies Theorems 1.6 on \mathfrak{G} , 1.8 on \mathfrak{B} , and 2.22 on \mathfrak{J} , \mathfrak{GS} , \mathfrak{FG} , and \mathfrak{H} . We state now a shorthand version of 2.14. Notation and technical terms are defined in Section 2.

Theorem 1.9. *Let G be an infinite finitely generated self-similar group. Assume that G is contracting faithful self-replicating. Let G_0 denote a standard contracting cover of G , as in Definition 2.8.*

If G_0 has non-abelian free subgroups, then so does any finitely presented cover of G .

If G_0 is large, any finitely presented cover of G is large.

1.C. Infinitely more examples. The group \mathfrak{G} has uncountably many³ relatives G_ω introduced in [Gri–84a]. Each of these groups is generated by a set S_ω of four involutions. They are parameterized by the space $\Omega = \{0, 1, 2\}^{\mathbb{N}}$. We have $\mathfrak{G} = G_\omega$ when ω is the 3-periodic sequence $012012012\cdots$. Let Ω_+ denote the complement in Ω of the space of eventually constant sequences. Section 4 contains a description of the family $(G_\omega)_{\omega \in \Omega_+}$.

Theorem 1.10. *For $\omega \in \Omega_+$, the group G_ω is of intermediate growth, and any finitely presented cover of G_ω is large.*

Note also the following straightforward consequence of Theorems 1.6, 1.8, and 1.10:

Corollary 1.11. *Let H be a finitely generated cover of one of \mathfrak{G} (as in Theorem 1.6), \mathfrak{B} (as in Theorem 1.8), or G_ω , $\omega \in \Omega_+$ (as in Theorem 1.10). Any finitely presented cover of H is large.*

There are several interesting classes of groups that qualify to be the H of Corollary 1.11:

(i) The uncountably many groups of [Ersc–04], which are finitely generated, of intermediate growth, and not residually finite, each one being a *central* cover of \mathfrak{G} .

(ii) The groups of [BaEr–a], which are finitely generated groups of intermediate growth, with exactly known growth functions, each one being a cover of \mathfrak{G} .

(iii) Permutational wreath products of the form $A \wr_X G_\omega$, where $A \neq \{1\}$ is a finite group and G_ω is as in Theorem 1.10 [BaEr–b].

There is an uncountable family of finitely generated amenable simple groups, which are topological full groups of minimal homeomorphisms of the Cantor space [JuMo]. None of these groups is finitely presented (see [Matu–06, Theorem 5.7], as well as [GrMe]). In our context, it is natural to formulate:

³Here and elsewhere, “uncountably many” groups means “uncountably many pairwise non-isomorphic” groups.

Problem 1.12. *Let G be one of the finitely generated amenable simple groups that appears in [JuMo]. Does G have an amenable finitely presented cover?*

Remark 1.13. We state two kinds of results concerning appropriate covers: some establish that the covers are large (see Theorem 1.6), others, weaker, that they contain non-abelian free groups (see Theorem 1.5). Strong statements do not hold in all cases.

For example, let $G = \text{Met}(k, \ell) \simeq \mathbf{Z} \left[\frac{1}{\ell m} \right] \rtimes_{\ell/m} \mathbf{Z}$, for two coprime integers $k, \ell \geq 2$ (Definition B.7). Then G satisfies the hypothesis of Theorem 1.5. The Baumslag-Solitar group $\text{BS}(\ell, m)$ is a finitely presented cover of G ; it is known that $\text{BS}(\ell, m)$ has non-abelian free subgroups, but is not large (Proposition B.6.iv).

1.D. A dual to Question 1.1. Given a Property (\mathcal{P}) of groups, it is standard to ask whether any countable group with (\mathcal{P}) is a *subgroup* of some finitely generated group with (\mathcal{P}) . For example, the answer is known to be positive if (\mathcal{P}) is solubility [NeNe–59], or amenability, or elementary amenability [OlOs, Corollary 1.3]. An earlier result of [HiNN–49] also gives a positive answer to a similar question: for a given integer n , if G is a group that has a presentation with a countable number (possibly infinite) of generators and n relators, then G can be embedded in a finitely presented group with 2 generators and n relators. But the answer is negative if (\mathcal{P}) is nilpotency, because any subgroup of a finitely generated nilpotent group is finitely generated. It is also negative if (\mathcal{P}) is metabelianity: indeed, for any prime p , the abelian quasi-cyclic group $\mathbf{Z}(p^\infty)$ cannot be embedded in a finitely generated metabelian group [NeNe–59, Lemma 5.3].

As a digression from our main theme, and since recursively presented groups are mentioned in Appendix A, we formulate one more question, which is in some sense dual to Question 1.1.

Question 1.14. *Given a Property (\mathcal{P}) of groups, is any finitely generated recursively presented group with (\mathcal{P}) a subgroup of some finitely presented group with (\mathcal{P}) ?*

The answer to Question 1.14 is known to be positive if (\mathcal{P}) is metabelianity (Proposition A.2), or solubility of the word problem ([Clap–67], and also [Mill–89, Theorem 2.8]). The answer is not known if (\mathcal{P}) is amenability [OlOs, Problem 1.7].

1.E. Plan of the paper. Section 2: Non-abelian free subgroups of finitely presented covers of contracting self-similar groups. Proofs of Theorems 1.9 and 2.14; proofs of Theorems 1.6, 1.8, and their analogues for \mathfrak{J} , \mathfrak{CS} , \mathfrak{FS} , and \mathfrak{H} .

Section 3: Marked groups and the Chabauty topology, a reminder.

Section 4: The analogue of Theorem 1.6 for the family $(G_\omega)_{\omega \in \Omega}$ of [Gri–84a], Theorem 1.10 and its proof.

Section 5: The group of intermediate growth \mathfrak{G} , and the proof of Theorem 1.7; this is a quantitative sharpening of Theorem 1.6. Proposition 5.11 on FP_2 -covers of \mathfrak{G} .

Appendix A: On soluble groups, metabelian groups, and finite presentations.

Appendix B: On wreath products and lamplighter groups.

Appendix C: On Bieri-Neumann-Strebel invariants.

Appendix D: On growth and amenability.

Post-scriptum: *A construction of large groups, due to Yves de Cornulier*, by Ralph Strebel.

1.F. Acknowledgements. We are most grateful to Yves de Cornulier and Ralph Strebel, for generous comments on preliminary versions of this paper, as well as to Laurent Bartholdi, Gilbert Baumslag, Marc Burger, Tullio Ceccherini-Silberstein, Anna Erschler, Sergei Ivanov, Olga Kharlampovich, Volodymyr Nekrashevych, Alexander Ol’shanskii, Vitalii Roman’kov, and Mark Sapir, for useful discussions and mail exchanges.

2. Non-abelian free subgroups of finitely presented covers of contracting self-similar groups

Let G, H be two groups and X a H -set; here, this means that H acts on X *from the right*. The corresponding **permutational wreath product** is the semi-direct product

$$G \wr_X H := G^{(X)} \rtimes H.$$

We denote by $G^{(X)}$ the group of functions $(g_x)_{x \in X} : X \rightarrow G, x \mapsto g_x$ with finite support, and we consider the action *from the left* of H on $G^{(X)}$, for which the action of h on $(g_x)_{x \in X}$ is $(g_{xh})_{x \in X}$. Hence the product of two elements in $G \wr_X H$ is given by

$$((g_x)_{x \in X}, h) ((g'_x)_{x \in X}, h') = ((g_x g'_{xh})_{x \in X}, hh').$$

In case G acts *from the right* on some set W , the group $G \wr_X H$ acts naturally *from the right* on $W \times X$, by

$$(w, y) ((g_x)_{x \in X}, h) = (wg_y, yh).$$

In expositions of wreath products and self-similar groups, choices of which groups actions are from the left and which from the right vary from one paper to the other, but a (sometimes hidden) mixture of left actions and right actions seems unavoidable.

If G, G' are two groups and $\alpha : G \rightarrow G'$ a homomorphism, we have a natural homomorphism

$$(1) \quad \alpha \wr_X 1_H : G \wr_X H \rightarrow G' \wr_X H, \quad ((g_x)_{x \in X}, h) \mapsto ((\alpha(g_x))_{x \in X}, h),$$

where 1_H stands for the identity automorphism of H .

If X is clear from the context, we write “ \wr ” for “ \wr_X ”. In particular, with $X = \{0, 1, \dots, d-1\}$ and S_d the full symmetric group of X , we write $G \wr S_d$ for $G \wr_X S_d$. Also, we write 1_d for 1_{S_d} , and

$$((g_x)_{x \in X}, \tau) = (g_0, \dots, g_{d-1}, \tau) \quad \text{with } g_0, \dots, g_{d-1} \in G \text{ and } \tau \in S_d$$

for a typical element of $G \wr S_d = G^X \rtimes S_d$.

The **iterated wreath products** with S_d are defined inductively, for each integer $n \geq 0$, by

$$G \wr^n S_d = \begin{cases} G & \text{for } n = 0 \\ (G \wr^{n-1} S_d) \wr S_d & \text{for } n \geq 1. \end{cases}$$

We have the following **associativity of permutational wreath products**: for a H -set X and a K -set Y , the canonical mapping

$$\begin{cases} (G \wr_X H) \wr_Y K & \longrightarrow & G \wr_{X \times Y} (H \wr_Y K) \\ \left(((g_{x,y})_{x \in X}, h_y)_{y \in Y}, k \right) & \longmapsto & \left((g_{x,y})_{(x,y) \in X \times Y}, ((h_y)_{y \in Y}, k) \right) \end{cases}$$

is an isomorphism of groups (this is standard, see e.g. [Meld–95, Chapter 1, Theorem 3.2]). In particular, for $n \geq 1$, we have

$$G \wr^n S_d = (G \wr^{n-1} S_d) \wr S_d = (G \wr S_d) \wr^{n-1} S_d = G^{X^n} \rtimes S_d^{(n)},$$

where $S_d^{(n)} = (\dots (S_d \wr S_d) \wr \dots) \wr S_d = S_d^{(n-1)} \wr S_d$ is the appropriate permutation group of X^n , acting here *from the right*. We write

$$((g_v)_{v \in X^n}, \tau) \quad \text{with } g_v \in G \text{ for all } v \in X^n \text{ and } \tau \in S_d^{(n)}$$

for a typical element of $G \wr^n S_d$.

Definition 2.1. Let G be a group and $d \geq 2$ an integer. A **self-similar structure of degree d** on G is a homomorphism

$$(2) \quad \Phi : G \rightarrow G \wr S_d.$$

A **self-similar group** is such a pair (G, Φ) ; when Φ is clear from the context, we write also “ G is a self-similar group”.

If (G, Φ) be a self-similar group, the construction (1) gives rise to a sequence of homomorphisms

$$(3) \quad \Phi_n : G \xrightarrow{\Phi_{n-1}} G \wr^{n-1} S_d \xrightarrow{\Phi^{1_{d^{n-1}}}} (G \wr S_d) \wr^{n-1} S_d = G \wr^n S_d$$

for $n \geq 2$; we write $\Phi_0 = \text{id}_G$ and $\Phi_1 = \Phi$. Note that, if Φ is injective, so is Φ_n for all $n \geq 0$. It is routine to check that Φ_{m+n} is the composition

$$(4) \quad \Phi_{m+n} : G \xrightarrow{\Phi_n} G \wr^n S_d \xrightarrow{\Phi_m \wr 1_d^n} (G \wr^m S_d) \wr^n S_d = G \wr^{m+n} S_d$$

for all $m, n \geq 0$.

The composition of Φ_n and the quotient map $G \wr^n S_d \longrightarrow S_d^{(n)}$ is a homomorphism

$$(5) \quad G \longrightarrow S_d^{(n)}, \quad g \longmapsto \tau_g^{(n)}.$$

Thus, introducing the v -coordinates of $\Phi_n(\cdot)$, we have

$$\Phi_n(g) = ((g_v)_{v \in X^n}, \tau_g^{(n)}) \in G \wr_{X^n} S_d^{(n)} = G \wr^n S_d$$

for all $g \in G$. Note that

$$(6) \quad \tau_g^{(n)} = ((\tau_{g_x}^{(n-1)})_{x \in X}, \tau_g^{(1)}) \in S_d^{(n)}$$

for all $g \in G$ and $n \geq 1$.

Let $X^* = \bigsqcup_{n \geq 0} X^n$ be the free monoid over $X = \{0, 1, \dots, d-1\}$. The **length** of $v \in X^*$ is the integer $n = |v|$ such that $v \in X^n$. The **d -regular rooted tree** is the tree with vertex set X^* , and with edges connecting pairs of vertices of the form $(x_1 \cdots x_n, x_1 \cdots x_n x_{n+1})$, with $n \geq 0$ and $x_1, \dots, x_{n+1} \in X$; abusively, we denote this tree also by X^* . The homomorphisms of (5) define an action *from the right* of G on the tree X^* .

For $n \geq 1$ and $v \in X^n$, we define the **stabilizer of v** to be the subgroup

$$(7) \quad \text{Stab}_G(v) = \{g \in G \mid v\tau_g^{(n)} = v\},$$

and we have a homomorphism

$$(8) \quad \Phi_v : \text{Stab}_G(v) \longrightarrow G, \quad g \longmapsto g_v$$

where $g_v = \Phi_v(g)$ is the v -coordinate of $\Phi_n(g)$.

Lemma 2.2. *With the notation above,*

$$g_{vw} = (g_v)_w, \quad (gh)_v = g_v h_{v\tau_g^{(n)}}, \quad \text{and} \quad (h^{-1})_v = \left(h_{v\tau_{h^{-1}}^{(n)}} \right)^{-1},$$

for all $g, h \in G$, $n \geq 1$, $v \in X^n$, and $w \in X^*$.

Proof. To illustrate the fact that G acts on X^* *from the right*, we spell out the proof of the second equality, writing vg for $v\tau_g^{(n)}$. We have on the one hand

$$((vw)g)h = ((vg)(wg_v))h = (vgh)(wg_v h_{vg})$$

and on the other hand

$$(vw)(gh) = (vgh)(w(gh)_v).$$

Hence $(gh)_v = g_v h_{vg}$. \square

Definition 2.3. A self-similar group (G, Φ) is **faithful** if its action on the tree X^* described above is faithful; this implies that the homomorphism Φ is injective (but the converse does not hold⁴).

A self-similar group (G, Φ) is **contracting** if there is a finite subset $\mathcal{M} \subset G$ with the following property:

for all $g \in G$, there exists an integer $k \geq 0$
such that $g_v \in \mathcal{M}$ for all $v \in X^*$ with $|v| \geq k$.

The smallest such \mathcal{M} , namely

$$\mathcal{N} := \bigcup_{g \in G} \bigcap_{k \geq 0} \{g \in G \mid \exists h \in G, \ell \geq k, v \in X^\ell \text{ with } h_v = g\}$$

is called the **nucleus** of (G, Φ) .

A self-similar group (G, Φ) is **self-replicating**⁵ if, for all $g \in G$ and $x \in X$, there exists $h \in \text{Stab}_G(x)$ such that $h_x = g$, namely if, for all $x \in X$, the homomorphism Φ_x of (8) is onto. When this is so, it is easy to check by induction on the level that, for all $g \in G$, $n \geq 1$, and $v \in X^n$, there exists $h \in \text{Stab}_G(v)$ such that $\Phi_v(h) = g$, namely Φ_v is onto.

Observe that, by definition, we have $1 \in \mathcal{N}$. Moreover for $g \in G$, we have $g \in \mathcal{N}$ if and only if $g^{-1} \in \mathcal{N}$, by Lemma 2.2.

The following proposition records basic facts about the nucleus of a contracting group.

Proposition 2.4. *Let (G, Φ) be a contracting self-similar group with nucleus \mathcal{N} , as above.*

(i) *For $g \in \mathcal{N}$ and $x \in X$, we have $g_x \in \mathcal{N}$.*

(ii) *If (G, Φ) is self-replicating and G is finitely generated, then \mathcal{N} generates G .*

Proof. For $g \in \mathcal{N}$, there exist $h \in G$, $k \geq 0$, and $v \in X^k$ such that $h_v = g$ and $h_w \in \mathcal{N}$ for all $w \in X^*$ with $|w| \geq k$ (otherwise, \mathcal{N} would not be minimal). Hence $g_x = (h_v)_x = h_{vx} \in \mathcal{N}$ for all $x \in X$. This proves (i).

⁴Set $H = \Phi^{-1}(G^X)$. The kernel of the action of G on X^* is the largest normal subgroup N of G that is contained in H and such that $\Phi(N) \subset N^X$.

⁵Or **recurrent**, or **fractal**, as in [Nekr-05, Definition 2.8.1]. See [BaGN-03] for relations of such groups with fractal sets.

For (ii), we paraphrase [Nekr–05, Lemma 2.11.3]. Denote by $\langle \mathcal{N} \rangle$ the subgroup of G generated by \mathcal{N} . Let S be a symmetric finite generating set of G . For all $s \in S$, there exists $k_s \geq 0$ such that $s_v \in \mathcal{N}$ for all $v \in X^*$ with $|v| \geq k_s$. Set $k = \max\{k_s \mid s \in S\}$.

Let $g \in G$ and $v \in X^*$ with $|v| \geq k$. There exist $s_1, \dots, s_m \in S$ with $g = s_1 \cdots s_m$, so that

$$\begin{aligned} g_v &= (s_1)_v (s_2 \cdots s_m)_{vs_1} = \cdots \\ &= (s_1)_v (s_2)_{vs_1} (s_3)_{vs_1 s_2} \cdots (s_m)_{vs_1 \cdots s_{m-1}} \in \langle \mathcal{N} \rangle, \end{aligned}$$

where the last inclusion follows from $|v| = |vs_1| = \cdots |vs_1 \cdots s_{m-1}| \geq k$. In particular, the image of Φ_v , as defined in (8), is contained in $\langle \mathcal{N} \rangle$.

If (G, Φ) is self replicating, then Φ_v is onto for all $v \in X^*$ with $|v| \geq 1$. The conclusion follows. \square

The next proposition was inspired to us by [Nekr–05, Lemma 2.13.4]. We need some notation and a definition; our exposition borrows from [Bart].

Definition 2.5. Let (G, Φ) be a self-replicating contracting self-similar group, with nucleus $\mathcal{N} = \{n_1, \dots, n_\ell\}$. Let $S = \{s_1, \dots, s_\ell\}$ be a finite set given with a bijection $s_j \leftrightarrow n_j$ with \mathcal{N} . Let R be the set of relators in the letters of S of one the forms

$$\begin{aligned} s_i &= 1 & \text{if } n_i &= 1 \in G, \\ s_i s_j &= 1 & \text{if } n_i n_j &= 1 \in G, \\ s_i s_j s_k &= 1 & \text{if } n_i n_j n_k &= 1 \in G. \end{aligned}$$

Note that these relators are of length at most 3; they are indexed by a subset of $\mathcal{N} \sqcup \mathcal{N}^2 \sqcup \mathcal{N}^3$.

Assume furthermore that G is finitely generated. The **universal contracting cover** of G is the finitely presented group G_0^{un} defined by the presentation with S as set of generators and R as set of relators. The assignment $\pi^{\text{un}}(s_i) = n_i$ extends to a group homomorphism

$$(9) \quad \pi^{\text{un}} : G_0^{\text{un}} = \langle S \mid R \rangle \longrightarrow G,$$

because $\pi^{\text{un}}(r) = 1$ for any $r \in R$.

Note that π^{un} is onto, by Proposition 2.4. We define finally

$$(10) \quad \widehat{\pi}^{\text{un}} = \pi^{\text{un}} \wr 1_d : G_0^{\text{un}} \wr S_d \longrightarrow G \wr S_d.$$

Remark 2.6. In particular examples, and for simplicity, it is often convenient to delete from S the generator corresponding to $1 \in \mathcal{N}$, to delete s_k if there exist $i, j \in \{1, \dots, \ell\}$ with $n_k = n_i n_j$, and to delete one generator of every pair corresponding to $\{n, n^{-1}\} \subset \mathcal{N}$. For example, in Example 2.17, we have $\mathcal{N} = \{1, a^{\pm 1}, b^{\pm 1}, c^{\pm 1}\}$ with 7 elements, and $c = a^{-1}b$, but $S = \{a, b\}$ with 2 elements.

Note however that, in Example 2.16, we keep d in the generating set $\{a, b, c, d\}$ of G_0 , even though $d = bc$.

Proposition 2.7. *Let (G, Φ) be a self-replicating contracting self-similar group of degree d , with nucleus \mathcal{N} . Assume that G is finitely generated. Let $G_0^{\text{un}} = \langle S \mid R \rangle$ and $\pi^{\text{un}} : G_0^{\text{un}} \rightarrow G$ be the universal contracting cover and the projection of Definition 2.5.*

Then there exists a homomorphism

$$\varphi_1^{\text{un}} : G_0^{\text{un}} \longrightarrow G_0^{\text{un}} \wr S_d$$

such that the self-similar group $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ is contracting, with nucleus S . Moreover the diagram

$$(11) \quad \begin{array}{ccc} G_0^{\text{un}} & \xrightarrow{\varphi_1^{\text{un}}} & G_0^{\text{un}} \wr S_d \\ \pi^{\text{un}} \downarrow & & \downarrow \widehat{\pi}^{\text{un}} \\ G & \xrightarrow{\Phi} & G \wr S_d \end{array}$$

commutes.

Proof. Step 1, definition of φ_1^{un} . Denote by ℓ the order of \mathcal{N} , and write $\mathcal{N} = \{n_1, \dots, n_\ell\}$, as above. Let $i \in \{1, \dots, \ell\}$. By Proposition 2.4, there exist $i_0, \dots, i_{d-1} \in \{1, \dots, \ell\}$ and $\tau_i \in S_d$ such that

$$\Phi(n_i) = (n_{i_0}, \dots, n_{i_{d-1}}, \tau_i).$$

We set

$$\varphi_1^{\text{un}}(s_i) = (s_{i_0}, \dots, s_{i_{d-1}}, \tau_i) \in G_0^{\text{un}} \wr S_d,$$

and we claim that this extends to a group homomorphism φ_1^{un} as in (11).

Consider a relator as in Definition 2.5, say $s_i s_j s_k = 1$ (shorter relators are dealt with similarly); hence $n_i n_j n_k = 1 \in G$. Choose $x \in X$; recall that X stands for $\{0, \dots, d-1\}$. There exist $p, q, r \in \{1, \dots, \ell\}$ and $\tau_p, \tau_q, \tau_r \in S_d$ such that the x -coordinate and the last coordinate of $\Phi(n_i n_j n_k)$ can be written as

$$(n_i n_j n_k)_x = n_p n_q n_r \quad \text{and} \quad \tau_{n_i n_j n_k} = \tau_p \tau_q \tau_r.$$

Since $n_i n_j n_k = 1 \in G$, we have

$$n_p n_q n_r = 1 \in G \quad \forall x \in X \quad \text{and} \quad \tau_p \tau_q \tau_r = 1 \in S_d.$$

Hence $\varphi_1^{\text{un}}(s_i) \varphi_1^{\text{un}}(s_j) \varphi_1^{\text{un}}(s_k) = 1 \in G_0^{\text{un}}$. The claim is proven.

Step 2: $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ is a contracting group with nucleus S . For any word w in the letters of S , we have to show that there exists a vertex

$v \in X^*$ such that $(w)_v \in S$. By induction on the word length, and by Lemma 2.2, it is enough to show this for a word of length 2.

Let $s_i, s_j \in S$ and $v \in X^*$ be such that $(n_i n_j)_v \in \mathcal{N}$, say $(n_i n_j)_v = n_k$. We have

$$(n_i)_v (n_j)_{v\tau_{n_i}^{(|v|)}} = n_k \quad \text{in } G,$$

which is a relator of length at most 3. Hence the corresponding relator $(s_i s_j)_v = s_k$ holds in S .

It follows that S is the nucleus of the group $(G, \varphi_1^{\text{un}})$.

Step 3, commutativity of the diagram. This can be checked on the set S of generators of G_0^{un} . \square

The universal contracting cover $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ of (G, Φ) is uniquely defined by (G, Φ) , and contracting. But we believe it need not be self-replicating (even though we do not know of any specific example). In all cases, $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ has quotients by finite sets of relations that are self-replicating contracting covers of (G, Φ) , as described in the Definition 2.8 and Proposition 2.9. Note however that these quotients are no more uniquely defined by (G, Φ) , since choices are involved.

In each of Examples 2.16 to 2.21 below, the universal contracting cover is self-replicating.

Definition 2.8. Let (G, Φ) be a self-replicating contracting self-similar group, with nucleus $\mathcal{N} = \{n_1, \dots, n_\ell\}$; assume that G is finitely generated. Let $S = \{s_1, \dots, s_\ell\}$ be in bijection with \mathcal{N} , and $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ the universal contracting cover of (G, Φ) , as in Definition 2.5. Let $\pi^{\text{un}} : G_0^{\text{un}} \rightarrow G$ be as in (9).

Let $x \in X$ and $n_i \in \mathcal{N}$. Since the pair (G, Φ) is self-replicating, there exists⁶ $g(x, n_i) \in \text{Stab}_G(x)$ such that $(g(x, n_i))_x = n_i$. Since π^{un} is onto, there exists $h(x, n_i) \in G_0^{\text{un}}$ such that $\pi^{\text{un}}(h(x, n_i)) = g(x, n_i)$; moreover, since $\widehat{\pi}^{\text{un}}$ is the identity on the permutations of the wreath product, we have $h(x, n_i) \in \text{Stab}_{G_0^{\text{un}}}(x)$. By commutativity of Diagram (11), we have $\pi^{\text{un}}((h(x, n_i))_x) = n_i$. Set $w(x, n_i) = (h(x, n_i))_x s_i^{-1}$; then $w(x, n_i)$ belongs to the kernel of π^{un} .

Again, by commutativity of (11), we have $(w(x, n_i))_v \in \ker(\pi^{\text{un}})$ for all $v \in X^*$. Since $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ is contracting, the subset

$$E(x, n_i) = \{g \in G_0^{\text{un}} \mid g = (w(x, n_i))_v \text{ for some } v \in X^*\}$$

of G_0^{un} is finite. Define

$$E = \bigcup_{x \in X, n \in \mathcal{N}} E(x, n) \quad \text{and} \quad H = \langle\langle E \rangle\rangle \subset G_0^{\text{un}},$$

⁶Note that a choice is involved here.

where $\langle\langle E \rangle\rangle$ denote the *normal* subgroup of G_0^{un} generated by E .

A **standard contracting cover** of G is a quotient group of the form $G_0 = G_0^{\text{un}}/H$, with H as above; the image of S in G is a generating set, that we denote again (abusively) by S . Note that E is a finite subset of G_0^{un} , and consequently that G_0 is a finitely presented group.

The epimorphism π^{un} factors through a homomorphism $\pi : G_0 \rightarrow G$, because E is a subset of $\ker \pi^{\text{un}}$. It follows from the definition that the restriction of π to the generating set S of G_0 is injective.

The following proposition is the analogue of proposition 2.7 for G_0 .

Proposition 2.9. *Let (G, Φ) be a self-replicating contracting self-similar group of degree d , with nucleus \mathcal{N} . Assume that G is finitely generated. Let G_0 and $\pi : G_0 \rightarrow G$ be a standard contracting cover of (G, Φ) and its projection to G , as in Definition 2.8.*

Then there exists a homomorphism

$$\varphi_1 : G_0 \rightarrow G_0 \wr S_d$$

such that the self-similar group (G_0, φ_1) is contracting and self-replicating with nucleus S . Moreover the diagram

$$(12) \quad \begin{array}{ccc} G_0 & \xrightarrow{\varphi_1} & G_0 \wr S_d \\ \pi \downarrow & & \downarrow \widehat{\pi} \\ G & \xrightarrow{\Phi} & G \wr S_d \end{array}$$

commutes.

Proof. By construction of the set E , for any element $g \in E$ and any $x \in X$, we have $g_x \in E$. Hence the homomorphism $\varphi_1^{\text{un}} : G_0^{\text{un}} \rightarrow G_0^{\text{un}} \wr S_d$ induces a homomorphism $\varphi_1 : G_0 \rightarrow G_0 \wr S_d$. Since $(G_0^{\text{un}}, \varphi_1^{\text{un}})$ is contracting with nucleus S , the self-similar group (G_0, φ_1) is contracting with nucleus S .

Let $x \in X$, and $n_i \in \mathcal{N}$. We continue with the notation of Definition 2.8. By construction of G_0 , the relation $h(x, n_i) = s_i$ holds in G_0 ; moreover $h(x, n_i)$ is an element of $\text{Stab}_{G_0}(x)$. This shows that the pair (G_0, φ_1) is self-replicating.

The commutativity of diagram (12) can be checked on the generators of G_0 . \square

From here to Corollary 2.15, we keep the same notation as in Definition 2.8 and Proposition 2.9 for G_0 , π , and φ_1 , in relation with a given contracting self-replicating self-similar group (G, Φ) , with G finitely generated.

Definition 2.10. For an integer $n \geq 0$, define

- (i) the homomorphism $\varphi_n : G_0 \rightarrow G_0 \wr^n S_d$ as in (3),
- (ii) its kernel $N_n = \ker(\varphi_n)$ and the quotient $G_n = G_0/N_n$,
- (iii) the homomorphism

$$(13) \quad \widehat{\pi}_n = \pi \wr 1_{d^n} : G_0 \wr^n S_d \rightarrow G \wr^n S_d$$

as in (1); note that $\widehat{\pi}_1$ is the $\widehat{\pi}$ of (10).

We have $\Phi_n \pi = \widehat{\pi}_n \varphi_n$, i.e. the diagram

$$(14) \quad \begin{array}{ccc} G_0 & \xrightarrow{\varphi_n} & G_0 \wr^n S_d \\ \pi \downarrow & & \downarrow \widehat{\pi}_n \\ G & \xrightarrow{\Phi_n} & G \wr^n S_d \end{array}$$

commutes. Observe that $N_0 \subset \dots \subset N_n \subset N_{n+1} \subset \dots$ and define

$$N = \bigcup_{n=0}^{\infty} N_n.$$

Remark 2.11. As noted in Definition 2.8, the restriction of π to S is injective. More generally, in Definition 2.10, the restriction of $\widehat{\pi}_n$ to the subset $(S^{X^n}, 1)$ of $G_0 \wr^n S_d = G_0^{X^n} \rtimes S_d^{(n)}$ is injective.

Lemma 2.12. *Let (G, Φ) be a self-similar group; assume that G is finitely generated and that (G, Φ) is faithful contracting self-replicating. With the notation above, we have*

$$N = \ker \pi, \text{ namely } G_0/N = G,$$

so that

$$\lim_{n \rightarrow \infty} G_n = G$$

in the space of marked groups on $|S|$ generators (in the sense of Section 3).

Proof. Let $g \in N$. Let $n \geq 1$ be such that $g \in \ker(\varphi_n)$. Then $\Phi_n \pi(g) = \widehat{\pi}_n \varphi_n(g) = 1$, hence $g \in \ker(\pi)$ by the faithfulness assumption.

Conversely, let $k \in \ker(\pi)$. On the one hand, since (G_0, φ) is contracting, there exists $n \geq 0$ such that $k_v \in S$ for all $v \in X^n$. On the other hand, $\pi(k) = 1$ implies $\pi(k_v) = 1$ for all $v \in X^n$; moreover, the $S_d^{(n)}$ -coordinate of $\varphi_n(k)$ is 1, by commutativity of Diagram (14). Hence, by Remark (2.11), we have $k_v = 1$ for all $v \in X^n$, and therefore $k \in N_n = \ker(\varphi_n)$, a fortiori $k \in N$. \square

Lemma 2.13. *In the situation of the previous lemma, for all $n \geq 1$, we have*

$$N_n = \varphi_1^{-1}(N_{n-1}^d)$$

so that $\varphi_1 : G_0 \rightarrow G_0 \wr S_d$ induces a homomorphism

$$\psi_n : \begin{cases} G_n \longrightarrow & G_{n-1} \wr S_d \\ gN_n \longmapsto & \left((\varphi_1(g)_x N_{n-1})_{x \in X}, \tau_g^{(1)} \right). \end{cases}$$

Moreover ψ_n is injective.

Proof. For $g \in G$, write

$$(15) \quad \varphi(g) = ((g_x)_{x \in X}, \tau_g^{(1)}) \quad \text{and} \quad \varphi_n(g) = ((g_v)_{v \in X^n}, \tau_g^{(n)}).$$

Assume first that $g \in N_n$. Thus $(g_x)_{v'} = 1$ and $\tau_{g_x}^{(n-1)} = 1$ for all $x \in X$ and $v' \in X^{n-1}$. This can be written

$$\varphi_{n-1}(g_x) = (((g_x)_{v'})_{v' \in X^{n-1}}, \tau_{g_x}^{(n-1)}) = 1 \quad \forall x \in X,$$

namely $g_x \in N_{n-1} \forall x \in X$. We have checked that $\varphi_1(N_n) \subset N_{n-1}^d$, and $N_n \subset \varphi_1^{-1}(N_{n-1}^d)$ follows.

Assume now that $g \in \varphi_1^{-1}(N_{n-1}^d)$, namely that $(g_x)_{v'} = 1$ and $\tau_{g_x}^{(n-1)} = 1$ for all $x \in X$ and $v' \in X^{n-1}$. This can be written $g_v = 1$ for all $v \in X^n$ and $\tau_g^{(n)} = 1$, namely $g \in N_n$. Hence $\varphi_1^{-1}(N_{n-1}^d) \subset N_n$. \square

The next theorem is a detailed version of Theorem 1.9.

Theorem 2.14. *Let (G, Φ) be a self-similar group; assume that G is finitely generated and that (G, Φ) is faithful contracting self-replicating. Let G_0 be a standard contracting cover, as in Definition 2.8.*

Assume that G_0 contains non-abelian free subgroups. Then, for each $n \geq 0$, the group G_n of Definition 2.10 contains non-abelian free subgroups. More generally, every finitely presented cover of G contains non-abelian free subgroups.

Assume moreover that G_0 is large. Then every finitely presented cover of G is large.

Proof. Let $S_d^{(0)}$ be the subgroup of S_d of permutations fixing the letter $x = 0$. For $n \geq 1$, let H_n be the finite index subgroup of G_n defined by

$$H_n = \psi_n^{-1}(G_{n-1}^{\{0,1,\dots,d-1\}} \rtimes S_d^{(0)}),$$

where ψ_n is as in lemma 2.13. Projection onto the first coordinate (i.e. the coordinate $x = 0$)

$$p_n^{(0)} : G_n^{\{0,1,\dots,d-1\}} \rtimes S_d^{(0)} \longrightarrow G_n$$

defined by

$$p_n^{(0)}((g_x N_n)_{x \in X}, \tau) = g_0 N_n$$

is a group homomorphism. It turns out that the composition

$$q_n^{(0)} : H_n \xrightarrow{\psi_n} \psi_n(H_n) \xrightarrow{p_{n-1}^{(0)}} G_{n-1}$$

defines a group homomorphism from H_n to G_{n-1} .

Given a generator sN_{n-1} of G_{n-1} (where s is a generator of G_0), using the self-replicating property of (G_0, φ_1) let $h \in St_{G_0}(0)$ be such that $\varphi_1(h)_0 = s$. It turns out that $q^{(0)}(hN_n) = sN_{n-1}$ which shows that $q_n^{(0)}$ is onto G_{n-1} . The conclusion is that for each $n \geq 1$, G_n contains a finite index subgroup H_n which maps onto G_{n-1} .

Therefore, if G_0 contains non-abelian free subgroups (respectively is large), by induction on n , each G_n will contain non-abelian free subgroups (respectively will be large). Then by lemma 2.12 and corollary 3.4 below, every finitely presented cover of G will contain non-abelian free subgroups (respectively will be large). \square

Corollary 2.15. *Let G be as in Theorem 2.14. If G_0 contains non-abelian free subgroups, then G is infinitely presented.*

Proof. Since G_0 does contain non-abelian free subgroups, by assumption, and G does not, by [Nekr–10, Theorem 4.2], G cannot be finitely presented, by the previous theorem. \square

Consider integers $d, \ell \geq 1$, and a system of relations

$$(16) \quad \begin{cases} s_1 = ((s_1)_0, \dots, (s_1)_{d-1}) \tau_1 \\ \dots\dots\dots \\ s_\ell = ((s_\ell)_0, \dots, (s_\ell)_{d-1}) \tau_\ell \end{cases}$$

with

$$(s_j)_x \in \{s_1, \dots, s_\ell\} \text{ and } \tau_j \in S_d \\ \text{for } j \in \{1, \dots, \ell\} \text{ and } x \in X = \{0, \dots, d-1\}.$$

By induction on $n \geq 0$, the system (16) defines a set S of automorphisms of the d -regular rooted tree X^* , again denoted by s_1, \dots, s_ℓ . These generate a group $G = \langle S \rangle$ of automorphisms of the tree X^* , and (16) define a self-similarity structure Φ on G ; thus G is a self-similar group as in Definition 2.1, and moreover G is faithful. The system (16) is often denoted by Φ again.

We review below some classical examples of self-similar groups defined this way.

Example 2.16. The 4-generated group $\mathfrak{G} = \langle a, b, c, d \rangle$ of Theorem 1.6 and [Grig–80, Grig–83] is a self-similar group of degree 2, with $\Phi : \mathfrak{G} \rightarrow \mathfrak{G} \wr C_2$ defined by

$$\Phi(a) = (1, 1)\tau, \quad \Phi(b) = (a, c), \quad \Phi(c) = (a, d), \quad \Phi(d) = (1, b).$$

Here, $C_2 = \{1, \tau\}$ denotes the cyclic group of order 2 (written S_2 in Definition 2.1). The self-similar group (\mathfrak{G}, Φ) is faithful, contracting, and self-replicating. The group \mathfrak{G} is of intermediate growth.

The nucleus is

$$\mathcal{N} = \{1, a, b, c, d\}.$$

The universal contracting cover of Definition 2.5 has the presentation

$$G_0 = \langle a, b, c, d \mid a^2, b^2, c^2, d^2, bcd \rangle \simeq C_2 * V,$$

where C_2 is now the group $\{1, a\}$ and V the Klein Vierergruppe $\{1, b, c, d\}$, isomorphic to $C_2 \times C_2$. The sign \simeq indicates an isomorphism of groups. It can easily be checked that this cover is self-replicating, so that a cover as in Definition 2.8 is not needed here.

Proofs of these facts, and of other properties of \mathfrak{G} , can be found in [Grig–80], [Harp–00, Chapter VIII], or [Nekr–05, Section 1.6], to quote *some* of the existing expositions only; see also our Section 5. The group \mathfrak{G} can be viewed as the IMG of an orbifold version of the *tent map* $T : [0, 1] \rightarrow [0, 1]$, defined by $T(x) = 2x$ for $x \leq 1/2$ and $T(x) = 2 - 2x$ for $x \geq 1/2$ [Nekr–11, Section 5.3]. Here and below, “IMG” stands for “**I**terated **M**onodromy **G**roup” (see [Nekr–05]).

Observe that $C_2 * V$ is virtually a non-abelian free group, because it is a free product of finite groups, distinct from $C_2 * C_2$ (see for example [Serr–77, Proposition 4 in Number I.1.3, Page 16]). It contains a free subgroup F_3 of index 8; one easy way to check this involves computing virtual Euler-Poincaré characteristics, as in [Serr–71, Section 1.8]: if

$$F_x = \ker(p : C_2 * V \xrightarrow{\text{canonical}} C_2 \times V),$$

where F_x stands for the free group of rank x , then

$$\chi(C_2 * V) = \frac{1}{2} + \frac{1}{4} - 1 = \frac{1}{[C_2 * V : \ker p]} \chi(F_x) = \frac{1}{8}(1 - x),$$

and therefore $x = 3$.

Hence Theorem 1.6 is a particular case of Theorem 2.14.

Example 2.17. The 2-generated **Basilica group** $\mathfrak{B} = \langle a, b \rangle$ is a self-similar group of degree 2, with homomorphism $\Phi : \mathfrak{B} \rightarrow \mathfrak{B} \wr C_2$ defined by

$$\Phi(a) = (b, 1)\tau, \quad \Phi(b) = (a, 1).$$

The self-similar group (\mathfrak{B}, Φ) is faithful, contracting, and self-replicating. The group \mathfrak{B} is of exponential growth.

The nucleus is

$$\mathcal{N} = \{1, a^{\pm 1}, b^{\pm 1}, c^{\pm 1}\}, \text{ where } c = a^{-1}b.$$

The universal contracting cover has the presentation

$$G_0 = \langle a, b \mid \emptyset \rangle \simeq F_2.$$

It is self-replicating.

The group \mathfrak{B} has been introduced in [GrZ-02a, GrZ-02b]. The name ‘‘Basilica’’ was given by Mandelbrot to the Julia set of the quadratic transformation $z \mapsto z^2 - 1$ of the complex plane, in honour of the *Basilica Cattedrale Patriarcale di San Marco*, and its reflection in Venetian waters [Mand-80, Page 254]. The group \mathfrak{B} was identified as $\text{IMG}(z^2 - 1)$ in [BaGN-03, Theorem 5.8]⁷, and the group was named ‘‘Basilica’’ in [BaVi-05], [Kaim-05], and [Nekr-05].

Our notation for $\Phi(a)$ and $\Phi(b)$ is essentially that of [Nekr-05, Page 208]; the roles of a and b are exchanged in [GrZ-02a].

Again, Theorem 1.8 is a particular case of Theorem 2.14.

Incidentally, since \mathfrak{B} is amenable (references in Appendix D), Theorem 1.8 shows that \mathfrak{B} is *not* finitely presented. Since we could not find references for a direct proof the latter statement in the literature, let us allude to two other simple ways to show that \mathfrak{B} is not finitely presented. One, suggested by Julia Bartsch (private communication), uses the infinite presentation of \mathfrak{B} given in [GrZ-02b] and obtained together with Laurent Bartholdi; then a nice argument concludes that this presentation is minimal (erasing any of its relators would change the group). The other uses the contracting property of the Basilica group established in [GrZ-02a] and follows the idea indicated in [Grig-80] for \mathfrak{G} .

Example 2.18. The **IMG of $z^2 + i$**

$$\mathfrak{J} = \text{IMG}(z^2 + i) = \langle a, b, c \rangle$$

is defined by

$$\Phi(a) = (1, 1)\tau, \quad \Phi(b) = (a, c), \quad \Phi(c) = (b, 1).$$

It was studied in detail in [GrSS-07], and was shown to be of intermediate growth in [BuPe-06].

Its nucleus is

$$\mathcal{N} = \{1, a, b, c\},$$

⁷As acknowledged in [BaGN-03], part of the credit for this is due to Richard Pink.

and the only non-trivial relators of length ≤ 3 among elements of \mathcal{N} are $a^2 = b^2 = c^2 = 1$ (this can best be checked with the GAP package (<http://www.gap-system.org/Packages/automgrp.html>). Thus the universal contracting cover

$$G_0 = \langle a, b, c \mid a^2, b^2, c^2 \rangle \simeq C_2 * C_2 * C_2$$

has a free subgroup of finite index (indeed a subgroup F_5 of index 8). It is self-replicating.

Example 2.19. The **Gupta-Sidki group** $\mathfrak{GS} = \langle a, b \rangle$ is the 2-generated group of automorphisms of the ternary rooted tree defined by

$$\Phi(a) = (1, 1, 1)\tau, \quad \Phi(b) = (a, a^{-1}, b),$$

where $\tau \in S_3$ is the cyclic permutation $(0, 1, 2)$. It is the infinite 3-group introduced in [GuSi-83]; it is just infinite [BaGr-02, Proposition 8.3]; it can be viewed as an IMG [Nekr-11, Section 4.5].

Its nucleus is

$$\mathcal{N} = \{1, a, a^{-1}, b, b^{-1}\}.$$

The universal contracting cover is

$$G_0 = \langle a, b \mid a^3, b^3 \rangle \simeq C_3 * C_3,$$

and contains a free subgroup F_4 of index 9. It is self-replicating.

The growth type of \mathfrak{GS} is not known.

Example 2.20. The **Fabrykowski-Gupta group** $\mathfrak{FG} = \langle a, b \rangle$ is the 2-generated group of automorphisms of the ternary rooted tree defined by

$$\Phi(a) = (1, 1, 1)\tau, \quad \Phi(b) = (a, 1, b),$$

with τ as in Example 2.19 [FaGu-85, FaGu-91]. It is of intermediate growth (see the original papers, and an exposition with improved estimates of growth in [BaPo-09]), it is just infinite [BaGr-02, Proposition 6.2], and it is the IMG of the cubic polynomial $z^3(-\frac{3}{2} + i\frac{\sqrt{3}}{2}) + 1$ [Nekr-11, Section 5.4].

As in the previous example, the nucleus is

$$\mathcal{N} = \{1, a, a^{-1}, b, b^{-1}\}.$$

The universal contracting cover is

$$G_0 = \langle a, b \mid a^3, b^3 \rangle \simeq C_3 * C_3.$$

It is self-replicating.

Example 2.21. The ternary **Hanoi Towers group** $\mathfrak{H} = \langle a, b, c \rangle$ is the 3-generated group of automorphisms of the ternary rooted tree defined by

$$\Phi(a) = (a, 1, 1)\tau_{1,2}, \quad \Phi(b) = (1, b, 1)\tau_{0,2}, \quad \Phi(c) = (1, 1, c)\tau_{0,1}$$

where $\tau_{1,2}$ is the transposition of S_3 exchanging 1 and 2, and similarly for $\tau_{0,2}, \tau_{0,1}$. It was introduced in [GrSu-06] as a model for the well-known Hanoi Towers problem; it is known to be of exponential growth ([GrSu-07, Subsection 6.1] and [Grig-06]), and isomorphic to $\text{IMG}(z^2 - \frac{16}{27z})$ [GrSu-07, Example 8].

The nucleus is

$$\mathcal{N} = \{1, a, b, c\}.$$

The universal contracting cover is

$$G_0 = \langle a, b, c \mid a^2, b^2, c^2 \rangle \simeq C_2 * C_2 * C_2.$$

It is self-replicating.

Theorem 2.22. *Any finitely presented cover of one of the groups \mathfrak{J} , \mathfrak{GS} , \mathfrak{FG} , \mathfrak{H} of the four previous examples is large.*

This is a straightforward consequence of Theorem 2.14. In Section 4, we will show how to modify 2.14 to cover uncountably many examples.

Remark 2.23. Groups of interest here are often known to have rather few quotients, of special kinds. Let us illustrate this as follows.

(i) A group is **just infinite** if all its proper quotients are finite. The group \mathfrak{G} is just infinite. More generally, with the notation of section 4, the group G_ω is just infinite for all $\omega \in \Omega_0$ (as we repeat below in Proposition 4.2.ii).

(ii) Without recalling here the technical definitions, let us mention the following property of a finitely generated group G assumed to be branch, or even weakly branch: for any normal subgroup $N \neq \{1\}$ of G , there exists an integer $n \geq 1$ such that N contains the derived group of the rigid stabilizer $\text{Rist}_G(n)$; this follows from the proof of [Grig-00, Theorem 4].

As a consequence, if G is branch, then any proper quotient of G is virtually abelian.

In particular, any proper quotient of one of the groups \mathfrak{J} , \mathfrak{GS} , \mathfrak{FG} , and \mathfrak{H} , is virtually abelian.

(This does *not* apply to \mathfrak{B} , which is weakly branch but *not* branch group. This applies to \mathfrak{G} , but it is of little interest in this case since the property of (i) is strictly stronger.)

(iii) It is shown in [Grig–98] that \mathfrak{G} has a finitely presented HNN-extension $\widehat{\mathfrak{G}}$ which is in $\mathcal{SG} \setminus \mathcal{EG}$. Any proper quotient of $\widehat{\mathfrak{G}}$ is metabelian and virtually abelian [SaWi–02, Theorem 2.3].

(iv) The Basilica group \mathfrak{B} is **just non-soluble**, which means that all its proper quotients are soluble [GrZ–02a, Proposition 6].

(v) Recall however that there exist groups of intermediate growth with uncountably many quotients: see [Gri–84b] and Definition 4.13.

3. Marked groups and the Chabauty topology

For k a positive integer, let F_k denote the free group of rank k , given together with an ordered free basis (s_1, \dots, s_k) of generators. A **marked group of rank k** is a pair (G, S) where G is a group and S an ordered set of k generators (for distinct $s, t \in S$, equalities $s = 1$ and $s = t \in G$ are allowed). To such a pair corresponds a free cover $\pi_G : F_k \twoheadrightarrow G$, with $\pi_G(s_j)$ being the j th generator of S ($1 \leq j \leq k$). We denote by \mathcal{M}_k the set of marked groups on k generators, identified here with the set of normal subgroups of F_k via the bijection $(G, S) \longleftrightarrow \ker \pi_G$.

The idea to furnish a space of (sub)groups with a topology goes back at least to Chabauty [Chab–50], and has been revisited on many occasions, among others by Bourbaki [Bour–63, chapitre VIII, § 5], Gromov [Grom–81, final remarks], one of us [Gri–84a], Stepin [Step–96], Champetier [Cham–00], Champetier-Guirardel [ChGu–05], and Ceccherini-Silberstein & Coornaert [CeCo–10].

The **Chabauty topology** on \mathcal{M}_k , also called the **Cayley topology**, is that defined by the basis⁸

$$(17) \quad \mathcal{O}_{K, K'} = \{N \triangleleft F_k : N \cap K = \emptyset \text{ and } K' \subset N\},$$

with K, K' finite subsets in F_k . This topology makes \mathcal{M}_k a totally disconnected compact space. It is also completely metrisable, as we now recall. For two subsets A, A' in F_k , let $v(A, A')$ denote the largest integer n such that $A \cap B_S^{F_k}(n) = A' \cap B_S^{F_k}(n)$, where $S = (s_1, \dots, s_k)$ in F_k is as above, and where balls $B_S^{F_k}(n)$ are as in Appendix D. Set $d(A, A') = \exp(-v(A, A'))$. Then d is a metric (indeed an ultrametric) and makes the set 2^{F_k} of subsets of F_k a totally discontinuous compact

⁸There is an equivalent definition in terms of the subbasis

$$\mathcal{O}_{K, V} = \{N \triangleleft F_k : N \cap K = \emptyset \text{ and } N \cap V \neq \emptyset\},$$

indexed by pairs (K, V) where K is a finite subset of F_k and V a subset of F_k . With K compact and V open, it has the advantage to carry over to the space of closed subgroups of a locally compact group G .

metric space, in which the space \mathcal{M}_k of marked groups on k generators (namely the space of normal subgroups of F_k) is closed. The topology induced by d on \mathcal{M}_k coincides with that defined by (17).

Here is an elementary and basic fact about this topology. The earliest written reference we know for it is [CoGP–07, Lemma 1.3 and Lemma 1].

Proposition 3.1. *Let k, ℓ be two positive integers. Let $(G, S) \in \mathcal{M}_k$ and $(G, T) \in \mathcal{M}_\ell$ be two marked groups with the same underlying group. Then there exist neighbourhoods $U \subset \mathcal{M}_k$ of (G, S) and $V \subset \mathcal{M}_\ell$ of (G, T) that are homeomorphic.*

In loose words, local properties of (G, S) are properties of G itself.

This proposition justifies the following definitions: a property (\mathcal{P}) of finitely generated groups is **open** [respectively **closed**] if, for any positive integer k , the subset of \mathcal{M}_k of marked groups (G, S) such that G has Property (\mathcal{P}) is open [respectively closed]. A finitely generated group G is **isolated** if, for any ordered generating set $S = (s_1, \dots, s_k)$ of G , the point (G, S) is isolated in \mathcal{M}_k . We collect a few examples as follows:

Proposition 3.2. *For $k \geq 2$, in the space \mathcal{M}_k of marked groups of rank k :*

- (i) *“Being abelian” is both an open and a closed property; more generally, for $d \geq 1$, “being nilpotent of nilpotent class at most d ” is both open and closed. “Being nilpotent” is open and non-closed.*
- (ii) *“Being soluble of solubility class at most k ” is closed and non-open. “Being soluble” is neither open nor closed.*
- (iii) *“Being finite” and “having torsion” are open and non-closed.*
- (iv) *If $(G, S) \in \mathcal{M}_k$ is a marking of a finitely presented group G , there exists a neighbourhood of (G, S) in \mathcal{M}_k containing only marked quotients of (G, S) .*
- (v) *A necessary condition for (G, S) to be an isolated point in \mathcal{M}_k is that G is finitely presented. Finite groups and finitely presented simple groups are isolated.*
- (vi) *There exists an isolated group that is 3-soluble and non-Hopfian; the group $\widehat{\mathfrak{G}}$ mentioned in Remark 2.23.iii is isolated.*
- (vii) *Amenability is neither open nor closed.*
- (viii) *Kazhdan Property (T) is open in \mathcal{M}_k .*
- (ix) *Serre Property (FA) is not open in \mathcal{M}_k .*

On the proof. Claims (i) to (v) are elementary; most of them appear explicitly in [ChGu–05, Section 2.6 and Lemma 2.3]. For (i), note moreover that “being nilpotent” is open by (iv), because nilpotent

groups are finitely presented. For (ii), note that “being soluble” is non-open, because metabelian groups like $\mathbf{Z} \wr \mathbf{Z}$ are limits of non-soluble groups (see Example B.2, say).

“Being nilpotent”, “being soluble”, “being finite”, “being amenable” and “having torsion” are non-closed properties, because non-abelian free groups are residually finite p -groups, for any prime p (due to [Hall–50], see also [Vale–93]).

For (v), observe that a finitely generated infinitely presented group G is always a limit of finitely presented groups G_n ; more precisely

$$G = \langle s_1, \dots, s_k \mid (r_i)_{i \geq 1} \rangle = \lim_{n \rightarrow \infty} G_n$$

with $G_n = \langle s_1, \dots, s_k \mid r_1, \dots, r_n \rangle$.

Necessary *and sufficient* conditions for isolated points are known in terms of the existence of “finite discriminating subsets”; we refer to [CoGP–07, Proposition 2]; see also [Grig–05, Theorem 2.1]. The class of isolated groups contains considerably more groups than the finite groups and the finitely presented simple groups [CoGP–07].

The first part of Claim (vi) is [CoGP–07, Proposition 10]; the second part is implicit in [SaWi–02], and explicit in [CoGP–07, Proposition 5.18]). “Being amenable” is non-open, again because $\mathbf{Z} \wr \mathbf{Z}$ is a limit of groups with non-abelian free subgroups (Example B.2). Claim (viii) is a result of [Shal–00], and (ix) of [DuMi]. \square

The Chabauty topology on \mathcal{M}_k plays an important role in connection with many group properties including “Property LEF” and “soficity”. The two latter properties define subspaces in \mathcal{M}_k that are closed [CeCo–10, Propositions 7.3.7 and 7.5.13].

Note the contraposition of (v): for $(G, S) \in \mathcal{M}_k$ with G infinitely presented, there exists a sequence $((G_n, S_n))_{n \geq 1}$ of pairwise distinct points in \mathcal{M}_k such that $\lim_{n \rightarrow \infty} (G_n, S_n) = (G, \bar{S})$.

The simplest examples of converging sequences in \mathcal{M}_k are of the following kind. Let

$$N_1 \subset \dots \subset N_n \subset N_{n+1} \subset \dots \subset N := \bigcup_{n \geq 1} N_n$$

be a nested sequence of normal subgroups in F_k . Let S_0 be a free basis of F_k . Denote by $p_n : F_k \rightarrow G_n := F_k/N_n$ ($n \geq 1$) and $p : F_k \rightarrow G := F_k/N$ the canonical projections. Set $S_n = p_n(S_0)$ and $S = p(S_0)$. Then $((G_n, S_n))_{n \geq 1}$ is a sequence in \mathcal{M}_k converging to (G, S) . In this case we often suppress the emphasis on the generating sets and write simply that the sequence $(G_n)_{n \geq 1}$ converges to G in \mathcal{M}_k .

Converging sequences in \mathcal{M}_k need not be of this special kind, with G a quotient of G_n for all n large enough. See below, Proposition 4.2.vi.

The following observation about \mathcal{M}_k and covers, basic for us, is well-known; see e.g. [CoKa–11, Lemma 3.1]. We provide a proof for the convenience of the reader.

Proposition 3.3. *Let $((G_n, S_n))_{n \geq 1}$ be a converging sequence in \mathcal{M}_k ; set $(G, S) = \lim_{n \rightarrow \infty} (G_n, S_n)$. Let Γ be a finitely presented group; assume there exists a cover $\pi : \Gamma \twoheadrightarrow G$.*

Then Γ is a cover of G_n for n large enough.

Note. In case G itself is finitely presented, this lemma is an immediate consequence of Proposition 3.2.iv.

Proof. Denote as above by (s_1, \dots, s_k) an ordered free basis of F_k . Let $p_n : F_k \twoheadrightarrow G_n$ and $p : F_k \twoheadrightarrow G$ be the free covers corresponding to (G_n, S_n) and (G, S) respectively. Set $N_n = \ker(p_n)$ and $N = \ker(p)$. Let (t_1, \dots, t_ℓ) an ordered generating set of Γ . Consider the free group F_ℓ on an ordered basis $U = (u_1, \dots, u_\ell)$ and the free cover $q : F_\ell \twoheadrightarrow \Gamma$ defined by $q(u_j) = t_j$ for $j = 1, \dots, \ell$.

Since Γ is finitely presented, there exists a finite subset $R \subset F_\ell$ of words v_1, \dots, v_m in the letters of $U \cup U^{-1}$ such that $\ker(q)$ is the normal subgroup of F_ℓ generated by R , namely such that $\langle U \mid R \rangle$ is a presentation of Γ . For $j \in \{1, \dots, \ell\}$, choose a word w_j in the letters $p(s_1), \dots, p(s_k)$ and their inverses such that $\pi(t_j) = w_j$. Let \tilde{w}_j be the word in $\{s_1, s_1^{-1}, \dots, s_k, s_k^{-1}\}$ obtained by substitution of $s_i^{\pm 1}$ in place of $p(s_i)^{\pm 1}$; observe that $p(\tilde{w}_j) = w_j = \pi(t_j)$. Consider the homomorphism

$$h : F_\ell \longrightarrow F_k \quad \text{defined by} \quad h(u_j) = \tilde{w}_j \quad (1 \leq j \leq \ell).$$

Then $ph(u_j) = p(\tilde{w}_j) = w_j = \pi(t_j) = \pi q(u_j)$ for all j , so that $ph = \pi q$, and therefore $h(R) \subset N$.

The last inclusion means that the open subset

$$\mathcal{O}' := \{M \triangleleft F_k : h(R) \subset M\} = \bigcap_{i=1}^m \mathcal{O}_{\emptyset, \{h(r_i)\}}$$

is a neighbourhood of N in \mathcal{M}_k . Hence, for n large enough, we have $N_n \in \mathcal{O}'$ and therefore $h(R) \subset N_n$.

Denote by $\langle\langle T \rangle\rangle$ the *normal* subgroup of a group H generated by a subset $T \subset H$. Let

$$h_1 : \Gamma = F_\ell / \langle\langle R \rangle\rangle \longrightarrow F_k / \langle\langle h(R) \rangle\rangle$$

be the cover induced by h , and

$$h_2 : F_k / \langle\langle h(R) \rangle\rangle \longrightarrow F_k / N_n = G_n$$

that defined by the inclusion $\langle\langle h(R) \rangle\rangle \subset N_n$ (for $n \gg 1$). The composition $h_2 h_1$ is a cover $\Gamma \twoheadrightarrow G_n$, and this concludes the proof. \square

An immediate consequence of the previous proposition is the following corollary, of very frequent use in our work.

Corollary 3.4. *Consider the three following group properties:*

- (NA) *non-amenability,*
- (Fr) *containing non-abelian free groups,*
- (La) *being large.*

Let $k \geq 2$ and $((G_n, S_n))_{n \geq 1}$ be a converging sequence in \mathcal{M}_k , with limit (G, S) .

If, for all n large enough, G_n has one of the three properties above, then any finitely presented cover of G has the same property.

Proof. The point is that a group that has a quotient with one of the properties (NA), (Fr), (La) has itself the same property. \square

4. The analogue of Theorem 1.6 for the family $(G_\omega)_{\omega \in \Omega}$ of [Gri-84a]

Let Ω be the Cantor space $\{0, 1, 2\}^{\mathbb{N}}$ of all sequences of 0's, 1's and 2's, with the product topology. Denote by Ω_- the countable subspace of eventually constant sequences, by Ω_+ its complement, and by Ω_0 the subspace of sequences with infinitely many occurrences of each of 0, 1, 2; thus

$$\Omega_0 \subset \Omega_+ \subset \Omega = \Omega_+ \sqcup \Omega_-.$$

We denote by σ the shift on Ω , defined by $(\sigma(\omega))_n = \omega_{n+1}$ for all $n \geq 1$.

We will recall the construction of [Gri-84a], which is a generalisation of that of Section 2. It associates with each point $\omega \in \Omega$ a marked group (G_ω, S_ω) with S_ω consisting of 4 generators of order 2; for example, $\mathfrak{G} = G_{\overline{012}}$, where $\overline{012}$ stands for the 3-periodic sequence $012012012 \dots$. In this section, set

$$X = \{0, 1\}$$

and identify X^* with the 2-regular rooted tree. We proceed to define for all $\omega \in \Omega$ a marked group $(G_\omega, S_\omega) \in \mathcal{M}_4$ of automorphisms of X^* .

Definition 4.1. The flip $a \in \text{Aut}(X^*)$ is defined by

$$a(0v) = 1v \text{ and } a(1v) = 0v \text{ for all } v \in X^*.$$

Set

$$\begin{array}{lll} a_{\beta(0)} = a & a_{\beta(1)} = a & a_{\beta(2)} = 1 \\ a_{\gamma(0)} = a & a_{\gamma(1)} = 1 & a_{\gamma(2)} = a \\ a_{\delta(0)} = 1 & a_{\delta(1)} = a & a_{\delta(2)} = a. \end{array}$$

Define for each $\omega = (\omega_n)_{n \geq 1} \in \Omega$ a set $S_\omega = \{a, b_\omega, c_\omega, d_\omega\}$ of four automorphisms of X^* by

$$\begin{aligned} b_\omega &= (a_{\beta(\omega_1)}, b_{\sigma(\omega)}) \\ c_\omega &= (a_{\gamma(\omega_1)}, c_{\sigma(\omega)}) \\ d_\omega &= (a_{\delta(\omega_1)}, d_{\sigma(\omega)}). \end{aligned}$$

It is easy to check that

$$(18) \quad \begin{aligned} ac_\omega a &= (b_{\sigma(\omega)}, a_{\beta(\omega_1)}) \\ ad_\omega a &= (c_{\sigma(\omega)}, a_{\gamma(\omega_1)}) \\ ab_\omega a &= (d_{\sigma(\omega)}, a_{\delta(\omega_1)}) \\ a^2 &= b_\omega^2 = c_\omega^2 = d_\omega^2 = b_\omega c_\omega d_\omega = 1. \end{aligned}$$

We define the group

$$G_\omega = \langle a, b_\omega, c_\omega, d_\omega \rangle$$

generated by S_ω ; it is a subgroup of $\text{Aut}(X^*)$. It follows from the last line of (18) that any element of G_ω can be written as

$$(19) \quad (*) a * a * \cdots * a (*)$$

with $*$ $\in \{b_\omega, c_\omega, d_\omega\}$, $(*) \in \{1, b_\omega, c_\omega, d_\omega\}$, and $n \geq 0$ occurrences of a .

Observe that any permutation τ of $\{0, 1, 2\}$ induces a permutation of Ω , again denoted by τ ; the groups $G_{\tau(\omega)}$ and G_ω are isomorphic.

In [Gri–84a, Section 6], there is moreover a modified construction providing a marked group $(\tilde{G}_\omega, \tilde{S}_\omega)$; we refer to the original paper. Note that (v) below holds for the modified groups, but not for the groups G_ω .

Proposition 4.2. *Let $\Omega = \{0, 1, 2\}^{\mathbb{N}}$. For $\omega \in \Omega$, let G_ω and \tilde{G}_ω be as above.*

- (i) *For $\omega \in \Omega$, the groups G_ω and \tilde{G}_ω are both infinite, and \tilde{G}_ω is infinitely presented.*
- (ii) *For $\omega \in \Omega_+$, the marked groups (G_ω, S_ω) and $(\tilde{G}_\omega, \tilde{S}_\omega)$ are isomorphic; the group G_ω is of intermediate growth.
For $\omega \in \Omega_0$, the group G_ω is an infinite 2-group, and is just infinite.⁹*
- (iii) *For $\omega \in \Omega_-$, the group G_ω is virtually free abelian, and consequently finitely presented of polynomial growth, while the group \tilde{G}_ω is virtually metabelian and of exponential growth.*

⁹Let Ω_1 be the subset of Ω_+ of sequences containing infinitely many occurrences of two of 0, 1, 2, and finitely many occurrences of the third, so that $\Omega_+ = \Omega_0 \sqcup \Omega_1$. For $\omega \in \Omega_1$, the group G_ω is not a 2-group, indeed it has elements of infinite order.

- (iv) For $\omega, \omega' \in \Omega_+$, the groups G_ω and $G_{\omega'}$ are isomorphic if and only if $\omega' = \tau(\omega)$ for some permutation τ of $\{0, 1, 2\}$.
- (v) The mapping $\Omega \longrightarrow \mathcal{M}_4$, $\omega \longmapsto (\tilde{G}_\omega, \tilde{S}_\omega)$ is a homeomorphism onto its image.
- (vi) For a converging sequence $(\omega_{(n)})_{n \geq 1}$ of points in Ω_+ with a limit ω in Ω_+ , we have $\lim_{n \rightarrow \infty} (G_{\omega_{(n)}}, S_{\omega_{(n)}}) = (G_\omega, S_\omega)$ in \mathcal{M}_4 . If, moreover, $\omega_n \in \Omega_0$ for all $n \geq 1$ and $\omega \notin \Omega_0$, then, for all $n \geq 1$, the group G_ω is not a quotient of $G_{\omega_{(n)}}$.

On the proof. Most of this is proved in [Gri–84a]; more precisely:

(i) G_ω is infinite [Gri–84a, Theorem 2.1] and \tilde{G}_ω is infinitely presented [Gri–84a, Theorem 6.2].

(ii) For $\omega \in \Omega_+$, we have $(G_\omega, S_\omega) = (\tilde{G}_\omega, \tilde{S}_\omega)$ [Gri–84a, observation just before Theorem 6.1], and G_ω is of intermediate growth [Gri–84a, Corollary 3.2]. For $\omega \in \Omega_0$, G_ω is an infinite 2-group that is just infinite [Gri–84a, Theorems 2.1 and 8.1].

(iii) G_ω is virtually free abelian [Gri–84a, Theorem 2.1.(3)], while \tilde{G}_ω is virtually metabelian and of exponential growth [Gri–84a, Theorem 6.1].

About (iv), see [Nekr–05, Theorem 2.10.13]. A weaker statement is proved in [Gri–84a, Section 5].

For (v), see [Gri–84a, Proposition 6.2].

For (vi), given any $n \geq 1$, note that G_ω is neither isomorphic to $G_{\omega_{(n)}}$, by (iv), nor a non-trivial quotient of $G_{\omega_{(n)}}$, by (ii). \square

For the main result of this section (Theorem 4.5), we will need an analogue in the present context of the homomorphisms (2) and (3) of Section 2. Recall that we have a natural *isomorphism*

$$\Phi_X : \text{Aut}(X^*) \xrightarrow{\cong} \text{Aut}(X^*) \wr S_2.$$

We keep the notation of Definition 4.1.

Definition 4.3. Let $\omega \in \Omega$. The restriction to G_ω of the isomorphism Φ_X provides an injective homomorphism

$$\Phi_\omega^{(1)} = \Phi_\omega : G_\omega \longrightarrow G_{\sigma(\omega)} \wr S_2.$$

On the generators, we have

$$\begin{aligned} \Phi_\omega(a) &= (1, 1)\tau \\ \Phi_\omega(b_\omega) &= (a_{\beta(\omega_2)}, b_{\sigma(\omega)}) \\ \Phi_\omega(c_\omega) &= (a_{\gamma(\omega_2)}, c_{\sigma(\omega)}) \\ \Phi_\omega(d_\omega) &= (a_{\delta(\omega_2)}, d_{\sigma(\omega)}) \end{aligned}$$

(recall that $S_2 = \{1, \tau\}$). The sequence of homomorphisms $(\Phi_\omega^{(n)})_{n \geq 1}$ is defined inductively by

$$\Phi_\omega^{(n)} : G_\omega \xrightarrow{\Phi_\omega^{(n-1)}} G_{\sigma^{n-1}(\omega)} \wr^{n-1} S_2 \xrightarrow{\Phi_{\sigma^{n-1}(\omega)}^{(1)} \wr^{1_{dn-1}}} G_{\sigma^n(\omega)} \wr^n S_2.$$

Lemma 4.4 (contraction in G_ω). *Let $\omega \in \Omega$. We keep the notation above.*

- (i) *For each $n \geq 1$, the homomorphism $\Phi_\omega^{(n)}$ is injective.*
- (ii) *For all $g \in G_\omega$, there exists an integer $n \geq 1$ such that*

$$\Phi_\omega^{(n)}(g) = ((g_v)_{v \in X^n}, \tau_g^{(n)})$$

with $g_v \in \{1, a, b_{\sigma^n(\omega)}, c_{\sigma^n(\omega)}, d_{\sigma^n(\omega)}\} \forall v \in X^n$ and $\tau_g^{(n)} \in S_2$.

Proof. By induction on the length of g , in the sense of (19). □

In Theorem 1.10 of the Introduction, the claim on intermediate growth is a repetition of part of Proposition 4.2, and the claim on covers is the theorem below.

Theorem 4.5. *For $\omega \in \Omega_+$, any finitely presented cover of G_ω is large.*

Remark 4.6. (1) Let $\omega \in \Omega_-$. Any finitely presented cover of the infinitely presented group \tilde{G}_ω contains non-abelian free groups, by Theorem 1.5. As recorded in Proposition 4.2.iii, the group G_ω is virtually free abelian, and finitely presented. For example, if ω is the constant sequence $000 \dots$, then G_ω is the infinite dihedral group.

(2) If we replace “is large” by “contains non-abelian free subgroups” in Theorem 4.5, the resulting statement has a short proof. More precisely:

For any $\omega \in \Omega$, any finitely presented cover of \tilde{G}_ω has non-abelian free subgroups.

Indeed, let $(\omega_n)_{n \geq 1}$ be a sequence of eventually constant sequences converging to ω in Ω . Then \tilde{G}_{ω_n} is virtually metabelian and infinitely presented for all $n \geq 1$ (Claims (i) and (iii) in Proposition 4.2), and $(\tilde{G}_{\omega_n})_{n \geq 1}$ converges to \tilde{G}_ω (Proposition 4.2.v). Let E be a finitely presented cover of \tilde{G}_ω . Then E is a cover of G_{ω_n} for n large enough (Proposition 3.3). Hence Bieri-Strebel Theorem 1.5 shows that E contains non-abelian free subgroups.

From now on, we assume that

$$\omega \in \Omega_+.$$

Our strategy for the proof of Theorem 4.5 is to adapt to the present context the steps of Section 2.

The following definition should be compared with Definition 2.10. Note however that G_0 has not quite the same meaning here and there.

Definition 4.7. Set again

$$G_0 = \langle a, b, c, d \mid a^2, b^2, c^2, d^2, bcd \rangle \simeq C_2 * V,$$

as in Example 2.16. Observe that any element of G_0 can be written as

$$(20) \quad (*)a * a * \cdots * a(*)$$

with $* \in \{b, c, d\}$, $(*) \in \{1, b, c, d\}$, and $n \geq 0$ occurrences of a (compare with Equation (19)).

For $i \in \{0, 1, 2\}$, set

$$\varphi_i(a) = (1, 1)\tau \text{ for all } i \in \{0, 1, 2\}$$

and

$$\begin{array}{lll} \varphi_0(b) = (a, b) & \varphi_1(b) = (a, b) & \varphi_2(b) = (1, b) \\ \varphi_0(c) = (a, c) & \varphi_1(c) = (1, c) & \varphi_2(c) = (a, c) \\ \varphi_0(d) = (1, d) & \varphi_1(d) = (a, d) & \varphi_2(d) = (a, d). \end{array}$$

It is easy to check that these formulas define homomorphisms

$$\varphi_i : G_0 \longrightarrow G_0 \wr S_2 \quad (i = 0, 1, 2).$$

Set $\varphi_\omega^{(1)} = \varphi_{\omega_1}$ and define, inductively for $n \geq 2$, homomorphisms

$$\varphi_\omega^{(n)} : G_0 \xrightarrow{\varphi_\omega^{(n-1)}} G_0 \wr^{n-1} S_2 \xrightarrow{\varphi_{\omega_n} \wr 1_2^n} G_0 \wr^n S_2.$$

For $n \geq 1$, set

$$N_{\omega, n} = \ker(\varphi_\omega^{(n)}) \text{ and } G_{\omega, n} = G_0 / N_{\omega, n}.$$

We have natural homomorphisms

$$\begin{aligned} \pi_\omega &: G_0 \longrightarrow G_\omega, \\ \widehat{\pi}_\omega = \widehat{\pi}_{\omega, 1} &: G_0 \wr S_2 \longrightarrow G_{\sigma(\omega)} \wr S_2, \\ \widehat{\pi}_{\omega, n} &: G_0 \wr^n S_2 \longrightarrow G_{\sigma^n(\omega)} \wr^n S_2. \end{aligned}$$

(Compare with (9), (10), and (13), but note that $\widehat{\pi}_{\omega, 1} = \pi_\omega \wr 1_2$ does not hold here.)

The next lemma is about diagrams analogous to (11) and (14). Its proof uses an argument similar to one in the proof of Proposition 2.7, and will be omitted.

Lemma 4.8. *The diagram*

$$(21) \quad \begin{array}{ccc} G_0 & \xrightarrow{\varphi_\omega^{(n)}} & G_0 \wr^n S_2 \\ \pi_\omega \downarrow & & \downarrow \widehat{\pi}_{\omega,n} \\ G & \xrightarrow{\Phi_\omega^{(n)}} & G_{\sigma^n(\omega)} \wr^n S_2 \end{array}$$

commutes for each $n \geq 1$.

The next lemma is analogous to Step 2 in the proof of Proposition 2.7.

Lemma 4.9 (contraction in G_0). *For all $k \in G_0$, there exists an integer $n \geq 1$ such that*

$$\varphi_\omega^{(n)}(k) = ((k_v)_{v \in X^n}, \tau_k^{(n)})$$

with $k_v \in \{1, a, b, c, d\} \forall v \in X^n$ and $\tau_k^{(n)} \in S_2$.

Proof: by induction on the length of k , in the sense of (20). □

Define now

$$N_\omega = \bigcup_{n \geq 1} N_{\omega,n}$$

(compare with Definition 2.10). The two following lemmas are appropriate modifications of Lemmas 2.12 and 2.13; we repeat the proof for the first one, and not for the second one.

Lemma 4.10. *We have*

$$N_\omega = \ker(\pi_\omega : G_0 \longrightarrow G_\omega), \text{ namely } G_\omega \simeq G_0/N_\omega,$$

so that

$$\lim_{n \rightarrow \infty} G_{\omega,n} = G_\omega$$

in the space of marked groups on 4 generators.

Proof. Let $g \in N$. Let $n \geq 1$ be such that $g \in \ker(\varphi_\omega^{(n)})$. Since $\Phi_\omega^{(n)} \pi_\omega(g) = \widehat{\pi}_{\omega,n} \varphi_\omega^{(n)}(g)$, we have $\pi_\omega(g) = 1$ by Lemma 4.4.i.

Conversely, let $k \in G_0$. There exists $n \geq 0$ such that $(\varphi_\omega^{(n)}(k))_v \in \{1, a, b, c, d\}$ for all $v \in X^n$, by Lemma 4.9. Assume that $k \in \ker(\pi_\omega)$. Then $\widehat{\pi}_{\omega,n}(\varphi_\omega^{(n)}(k)) = 1$. As $\widehat{\pi}_{\omega,n}$ is injective “on generators” (in a sense similar to that of Remark 2.11), we have $\varphi_\omega^{(n)}(k) = 1$, and therefore $k \in N_{\omega,n} \subset N_\omega$.

(Note that the hypothesis “ $\omega \in \Omega_+$ ” is necessary for the previous argument. If ω_n were eventually constant, one of $b_{\sigma^n(\omega)}, c_{\sigma^n(\omega)}, d_{\sigma^n(\omega)}$ would be the identity of $G_{\sigma^n(\omega)}$ for n large enough). □

Lemma 4.11. *In the situation of the previous lemma, we have for all $n \geq 1$*

$$\varphi_\omega^{(1)}(N_{\omega,n}) \subset N_{\omega,n-1}^2 \subset G_0 \wr S_2 \text{ and } (\varphi_\omega^{(1)})^{-1}(N_{\omega,n-1}^2) \subset N_{\omega,n}.$$

It follows that $\varphi_\omega^{(1)} : G_0 \rightarrow G_0 \wr S_2$ induces a homomorphism

$$\psi_\omega^{(n)} : \begin{cases} G_{\omega,n} \longrightarrow G_{\omega,n-1} \wr S_2 \\ gN_{\omega,n} \longmapsto \left(((\varphi_{\omega_n}(g))_x N_{\omega,n-1})_{x \in X}, \tau_g^{(1)} \right) \end{cases}$$

which is injective.

Proposition 4.12. *For each $\omega \in \Omega_+$ and $n \geq 0$, the group $G_{\omega,n}$ is large.*

Proof. The group $G_0 = C_2 * V$ has a free subgroup of finite index, indeed a subgroup isomorphic to F_3 of index 8. For $n \geq 1$, because of the previous lemma and as in the proof of Theorem 2.14, there exists a subgroup of index 2 in $G_{\omega,n}$ and a homomorphism from this subgroup onto $G_{\omega,n-1}$. It follows by induction on n that $G_{\omega,n}$ is large. \square

End of proof of Theorem 4.5. Since $G_{\omega,n}$ is large for n large enough, it follows from Lemma 4.10 and Corollary 3.4 that any cover of G_ω is large. \square

Definition 4.13. For $\omega \in \Omega$, let M_ω denote the kernel of the defining cover $F_4 \rightarrow G_\omega$; in other terms, M_ω is the inverse image of N_ω by the epimorphism $F_4 \rightarrow G_0$ mapping the four generators of F_4 onto $a, b, c, d \in G_0$. For a subset Ψ of Ω , the Ψ -**universal group** is the group

$$\mathcal{U}_\Psi = F_4 / \bigcap_{\omega \in \Psi} M_\omega.$$

For example, $\mathcal{U}_\emptyset = \{1\}$, and $\mathcal{U}_{\{\omega\}} = G_\omega$ for all $\omega \in \Omega$.

The terminology is justified by cases that have appeared in the literature, with Ψ large. For example, let Λ denote the subset of Ω_0 of sequences that are concatenations of blocks 012, 120, 201. Then \mathcal{U}_Λ has uncountably many quotients (a consequence of Proposition 4.2.iv); it has intermediate growth, and therefore is amenable (established in [Grig, Theorem 9.7]).

Suppose that Ψ contains some $\omega \in \Omega_+$. Then any cover of \mathcal{U}_Ψ is a cover of G_ω . Theorem 4.5 implies:

Corollary 4.14. *For any $\Psi \subset \Omega$ such that $\Psi \cap \Omega_+ \neq \emptyset$, the Ψ -universal group \mathcal{U}_Ψ is infinitely presented, and any finitely presented cover of it is large.*

In particular, this corollary solves the first part of Problem 9.5 in [Grig–05], by showing that \mathcal{U}_Ω is infinitely presented.

5. The group of intermediate growth \mathfrak{G}

Let \mathfrak{G} be the self-similar group of degree 2 of Example 2.16. On the one hand, \mathfrak{G} is a group of the family studied in the previous section: $\mathfrak{G} = G_{\overline{012}}$; thus Theorem 4.5 “contains” Theorem 1.6. On the other hand, in this particular case, we can describe much more precisely a sequence of finitely presented covers converging to \mathfrak{G} , and this is the subject of the present section. Note however that, even if the cover \mathfrak{G}_{-1} below is the same as G_0 in Example 2.16, the sequence $(\mathfrak{G}_n)_{n \geq 0}$ is *not* the particular case for \mathfrak{G} of the sequence $(G_n)_{n \geq 1}$ (even shifted) of Section 2.

Immediately after its discovery it was observed that \mathfrak{G} is infinitely presented. Then, Lysenok found a presentation that we recall below.

Set

$$\mathfrak{G}_{-1} = \langle a, b, c, d \mid a^2 = b^2 = c^2 = d^2 = bcd = 1 \rangle \simeq C_2 * V,$$

and denote by S the system of four involutions $\{a, b, c, d\}$ generating \mathfrak{G}_{-1} . Elements in \mathfrak{G}_{-1} are in natural bijection with “reduced words” of the form

$$t_0 a t_1 a \cdots a t_{k-1} a t_k$$

with $k \geq 0$, $t_1, \dots, t_{k-1} \in \{b, c, d\}$, and $t_0, t_k \in \{\emptyset, b, c, d\}$. Throughout the remainder of this section, we use the same symbol to denote an element of \mathfrak{G}_{-1} and its image in any quotient of \mathfrak{G}_{-1} , in particular in \mathfrak{G} ; thus, $S = \{a, b, c, d\}$ denotes a set of generators in \mathfrak{G}_{-1} and in any quotient of \mathfrak{G}_{-1} .

The substitution σ defined by

$$\sigma(a) = aca, \quad \sigma(b) = d, \quad \sigma(c) = b, \quad \sigma(d) = c$$

extends to reduced words, for example $\sigma(abac) = acadacab$, and the resulting map

$$\sigma : \mathfrak{G}_{-1} \longrightarrow \mathfrak{G}_{-1}$$

is a group endomorphism. Define

$$\begin{aligned} u_0 &= (ad)^4 & u_n &= \sigma^n(u_0) \quad \forall n \geq 0 \\ v_0 &= (adacac)^4 & v_n &= \sigma^n(v_0) \quad \forall n \geq 0 \end{aligned}$$

Theorem 5.1 ([Lyse–85]). *The group \mathfrak{G} has a presentation*

$$\langle a, b, c, d \mid a^2 = b^2 = c^2 = d^2 = bcd = 1, \quad u_n = v_n = 1 \quad \forall n \geq 0 \rangle.$$

Note. It is moreover known that this presentation is minimal [Grig–99]. Lysenok’s presentation is the prototype of what is now called an *L-presentation* [Bart–03].

Definition 5.2. For $n \geq 0$, define a pair $(\mathfrak{G}_n, S) \in \mathcal{M}_4$ by

$$\mathfrak{G}_n = \left\langle a, b, c, d \left| \begin{array}{l} a^2 = b^2 = c^2 = d^2 = bcd = 1 \\ u_0 = \cdots = u_n = v_0 = \cdots = v_{n-1} = 1 \end{array} \right. \right\rangle$$

$$S = \{a, b, c, d\} \subset \mathfrak{G}_n.$$

Observe that $\lim_{n \rightarrow \infty} (\mathfrak{G}_n, S) = (\mathfrak{G}, S)$ in \mathcal{M}_4 , and that there are natural surjections $\mathfrak{G}_{-1} \twoheadrightarrow \mathfrak{G}_n \twoheadrightarrow \mathfrak{G}$ for all $n \geq 0$.

Theorem 5.3. *For each $n \geq 0$, the group \mathfrak{G}_n has a normal subgroup H_n of index $2^{2^{n+1}+2}$ which is isomorphic to the direct product of 2^n free groups of rank 3.*

Remark 5.4. (i) A weaker result was first established in [GrHa–01]: For each $n \geq 0$, \mathfrak{G}_n contains a subgroup of finite index isomorphic to the direct product of 2^n copies of finitely generated non-abelian free groups. This by itself implies that any finitely presented cover of \mathfrak{G} contains non-abelian free subgroups.

(ii) The result of [GrHa–01] was improved in [BaCo–06]: For each $n \geq 0$, the group \mathfrak{G}_n has a normal subgroup H_n of index 2^{α_n} , where $\alpha_n \leq (11 \cdot 4^n + 1)/3$, and H_n is a subgroup of index 2^{β_n} in a finite direct product of 2^n non-abelian free groups of rank 3, where $\beta_n \leq (11 \cdot 4^n - 8)/3 - 2^n$.

(iii) Our proof of Theorem 5.3 is split in several lemmas, until 5.10.

If x, \dots, y are elements of a group H , we denote by $\langle x, \dots, y \rangle_H$ the subgroup of H they generate, and by $\langle\langle x, \dots, y \rangle\rangle_H$ the *normal* subgroup of H they generate. Define first

$$\begin{aligned} B_0 &= \langle\langle b \rangle\rangle_{\mathfrak{G}_0}, \\ \Xi_0 &= \langle b, c, d, aba, aca, ada \rangle_{\mathfrak{G}_0}, \\ D_0 &= \langle a, d \rangle_{\mathfrak{G}_0}, \\ D_0^{\text{diag}} &= \langle (a, d), (d, a) \rangle_{\mathfrak{G}_0}. \end{aligned}$$

It is easy to check that $D_0^{\text{diag}} \cap (B_0 \times B_0) = \{1\}$, and that D_0^{diag} normalizes $B_0 \times B_0$. The assignment

$$\begin{array}{ll} b & \mapsto (a, c) & aba & \mapsto (c, a) \\ c & \mapsto (a, d) & aba & \mapsto (d, a) \\ d & \mapsto (1, b) & aba & \mapsto (b, 1) \end{array}$$

extends to a group homomorphism $\psi_0 : \Xi_0 \longrightarrow \mathfrak{G}_0 \times \mathfrak{G}_0$ [GrHa–01, Proposition 1]. For each $n \geq 0$, define now

$$\begin{aligned} N_n &= \langle\langle u_0, \dots, u_n, v_0, \dots, v_{n-1} \rangle\rangle_{\mathfrak{G}_0}; \text{ observe that } N_n \subset \Xi_0; \\ \mathfrak{G}_n &= \mathfrak{G}_0/N_n \text{ and } \pi_n : \mathfrak{G}_0 \twoheadrightarrow \mathfrak{G}_n \text{ the canonical projection;} \\ B_n &= \langle\langle b \rangle\rangle_{\mathfrak{G}_n} = \pi_n(B_0); \\ \Xi_n &= \langle b, c, d, aba, aca, ada \rangle_{\mathfrak{G}_n} = \pi_n(\Xi_0); \\ D_n^{\text{diag}} &= \langle\langle (a, d), (d, a) \rangle\rangle_{\mathfrak{G}_n \times \mathfrak{G}_n}; \\ \sigma_n : \mathfrak{G}_{n-1} &\longrightarrow \mathfrak{G}_n, \quad gN_{n-1} \longmapsto \sigma(g)N_n \quad (\text{for } n \geq 1 \text{ only}). \end{aligned}$$

For the definition of the homomorphism σ_n , note that $\sigma(N_{n-1}) \subset N_n$.

Lemma 5.5 ([GrHa–01], Lemma 3). *Let B_0 denote the normal subgroup of \mathfrak{G}_0 generated by b . Then:*

- (i) B_0 is of index 8 in \mathfrak{G}_0 ;
- (ii) B_0 is generated by the four elements
 $\xi_1 := b, \xi_2 := aba, \xi_3 := dabad, \xi_4 = adabada$;
- (iii) B_0 has the presentation $\langle \xi_1, \xi_2, \xi_3, \xi_4 \mid \xi_1^2 = \xi_2^2 = \xi_3^2 = \xi_4^2 = 1 \rangle$;
- (iv) B_0 contains N_n for all $n \geq 1$.

Lemma 5.6 ([GrHa–01], mostly Proposition 10). *(i) The kernel and the image of the homomorphism ψ_0 are given by*

$$\begin{aligned} \ker(\psi_0) &= \langle\langle u_1, v_0 \rangle\rangle_{\Xi_0}, \\ \text{Im}(\psi_0) &= (B_0 \times B_0) \rtimes D_n^{\text{diag}} \text{ of index 8 in } \mathfrak{G}_0 \times \mathfrak{G}_0. \end{aligned}$$

(ii) For $n \geq 1$, the homomorphism ψ_0 induces an isomorphism

$$\psi_n : \Xi_n \xrightarrow{\cong} (B_{n-1} \times B_{n-1}) \rtimes D_{n-1}^{\text{diag}} <_8 \mathfrak{G}_{n-1} \times \mathfrak{G}_{n-1}$$

where $<_8$ indicates that the left-hand side is a subgroup of index 8 in the right-hand side.

Set $K_0 = \langle\langle (ab)^2 \rangle\rangle_{\mathfrak{G}_0}$; observe that $K_0 \subset B_0$.

Lemma 5.7. *(i) The subgroup K_0 is of index 2 in B_0 . It is generated by*

$$t = (ab)^2 \quad v = (bada)^2 \quad w = (abad)^2$$

Moreover K_0 contains N_n for $n \geq 1$.

(ii) The group K_0 is a free group of rank 3.

Proof. (i) This follows from [Harp–00, Page 230]. Since B_0 contains N_n and each u_n, v_n is a fourth power, necessarily N_n is contained in K_0 .

For (ii), see [BaCo–06, Proposition 4], where the proof uses Kurosh’s theorem. Alternatively one can use the Reidemeister-Schreier method to find a presentation for K_0 and see that it is indeed free of rank 3. \square

Lemma 5.8. *If g is an element of B_{n-1} then*

$$\psi_n(\sigma_n(g)) = (1, g) \quad \text{and} \quad \psi_n(a\sigma_n(g)a) = (g, 1).$$

Proof. For the generators of B_{n-1} that are images of those of Lemma 5.5 for B_0 , we have

$$\begin{aligned} \psi_n(\sigma_n(b)) &= \psi_n(d) = (1, b), \\ \psi_n(\sigma_n(aba)) &= \psi_n(acadaca) = (d^2, aba) = (1, aba), \\ \psi_n(\sigma_n(dabad)) &= \psi_n(cacadacac) = (ad^2a, dabad) = (1, dabad), \\ \psi_n(\sigma_n(adabada)) &= \psi_n(acacacacacacaca) = (1, adabada), \end{aligned}$$

and this shows the first equality. The second follows because, if $\psi_n(h) = (h_0, h_1)$, then $\psi_n(aha) = (h_1, h_0)$. \square

Let $K_n = K_0/N_n$. It is a normal subgroup of \mathfrak{G}_n contained in B_n .

Lemma 5.9. *Let $n \geq 1$.*

- (i) *We have $\sigma_n(K_{n-1}) \subset K_n \subset B_n$.*
- (ii) *If H_{n-1} is a subgroup of K_{n-1} , then $\psi_n^{-1}(H_{n-1} \times H_{n-1}) \subset K_n$.*

Proof. (i) Let t, v, w be now the canonical images in K_n of the elements of K_0 denoted by the same symbols in Lemma 5.7. On the one hand, we have $\psi_n(\sigma_n(t)) = (1, t)$ by Lemma 5.8. On the other hand, we have

$$\psi_n(w) = \psi_n(aba)\psi_n(d)\psi_n(aba)\psi_n(d) = (cc, abab) = (1, t)$$

by the definitions of ψ_n and w . Hence $\sigma_n(t) = w \in K_n$ by Lemma 5.6.ii.

Let $g_1 \in \mathfrak{G}_{n-1}$. From the definition of ψ_n , we see that the composition $\Xi_n \rightarrow \mathfrak{G}_{n-1}$ of ψ_n with a projection onto one of the factors is onto. Hence there exists $g \in \Xi_n$ and $g_0 \in \mathfrak{G}_{n-1}$ such that $\psi_n(g) = (g_0, g_1)$. We have as above¹⁰ $\psi_n(\sigma_n(t^{g_1})) = (1, t^{g_1})$ and

$$\psi_n(w^g) = \psi_n(w)^{\psi_n(g)} = (1, t)^{\psi_n(g)} = (1, t^{g_1}),$$

and therefore $\sigma_n(t^{g_1}) = w^g$. Since K_n is a normal subgroup of \mathfrak{G}_n containing w , we have $\sigma_n(t^{g_1}) \in K_n$ for all $g_1 \in \mathfrak{G}_{n-1}$. The inclusion $\sigma_n(K_{n-1}) \subset K_n$ follows, because K_{n-1} is generated by t as a normal subgroup of \mathfrak{G}_{n-1} .

- (ii) Let $(h_0, h_1) \in H_{n-1} \times H_{n-1}$. We have

$$\psi_n^{-1}(h_0, h_1) = a\sigma_n(h_0)a\sigma_n(h_1)$$

by Lemma 5.8, and the right-hand side is in K_n by (i). \square

¹⁰Remember that $t^h = h^{-1}th$.

Set $H_0 = K_0$. For $n \geq 1$, define inductively

$$H_n = \psi_n^{-1}(H_{n-1} \times H_{n-1}).$$

The definition makes sense by Lemma 5.9.ii. The following lemma finishes the proof of Theorem 5.3.

Lemma 5.10. *Let $n \geq 0$, and the notation be as above.*

- (i) H_n is a normal subgroup of \mathfrak{G}_n contained in K_n .
- (ii) The group H_n is a direct product of 2^n free groups of rank 3.
- (iii) Its index is given by $[\mathfrak{G}_n : H_n] = 2^{(2^{n+1}+2)}$.

Proof. For $n = 0$, the three claims follow from Lemmas 5.5 and 5.7. We suppose now that $n \geq 1$ and that the lemma holds for $n - 1$.

(i) The group H_n is clearly normal in Ξ_n , by Lemma 5.6.ii. To show that H_n is normal in \mathfrak{G}_n , it suffices to check that $aH_na \subset H_n$, because \mathfrak{G}_n is generated by Ξ_n (of index 2 in \mathfrak{G}_n) and a . Let $h \in H_n$. Let $h_0, h_1 \in H_{n-1}$ be defined by $\psi_n(h) = (h_0, h_1)$. Then $\psi_n(aha) = (h_1, h_0) \in H_{n-1} \times H_{n-1}$, and therefore $aha \in H_n$.

(ii) This is a straightforward consequence of the isomorphism $H_n \simeq H_{n-1} \times H_{n-1}$, see again Lemma 5.6.

(iii) By the induction hypothesis, we have

$$\begin{aligned} & [(B_{n-1} \times B_{n-1}) \rtimes D_{n-1}^{\text{diag}} : H_{n-1} \times H_{n-1}] \\ &= \frac{[\mathfrak{G}_{n-1} \times \mathfrak{G}_{n-1} : H_{n-1} \times H_{n-1}]}{[\mathfrak{G}_{n-1} \times \mathfrak{G}_{n-1} : (B_{n-1} \times B_{n-1}) \rtimes D_{n-1}^{\text{diag}}]} \\ &= \frac{2^{2^n+2} \times 2^{2^n+1}}{2^3} = 2^{2^{n+1}+1}. \end{aligned}$$

Thus the commutative diagram

$$\begin{array}{ccc} \mathfrak{G}_n & & \mathfrak{G}_{n-1} \times \mathfrak{G}_{n-1} \\ \downarrow 2 & & \downarrow 2^3 \\ \psi_n : \Xi_n & \xrightarrow{\cong} & (B_{n-1} \times B_{n-1}) \rtimes D_{n-1}^{\text{diag}} \\ \downarrow 2^{(2^{n+1}+1)} & & \downarrow 2^{(2^{n+1}+1)} \\ H_n & \longrightarrow & H_{n-1} \times H_{n-1} \end{array}$$

shows that H_n has index $2^{(2^{n+1}+2)}$ in \mathfrak{G}_n . □

The proof of Theorem 5.3 is now complete.

Recall that a group G is called of type FP_n if the trivial $\mathbf{Z}[G]$ -module \mathbf{Z} has a projective resolution, namely if there exists an exact sequence

$$\cdots \longrightarrow P_{j+1} \longrightarrow P_j \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow \mathbf{Z}$$

where the P_j 's are projective $\mathbf{Z}[G]$ -modules for all $j \geq 0$, and finitely generated projective $\mathbf{Z}[G]$ -modules for all $j \leq n$. It is known that

- (i) a group is of type FP_1 if and only if it is finitely generated;
- (ii) finitely presented groups are of type FP_2 ;
- (iii) Condition FP_2 is *strictly* weaker than finite presentability;
- (iv) a group is of type FP_2 if and only if it is the quotient of some finitely presented group by a *perfect* normal subgroup.

For (i) and (ii), see for example the notes in which “type FP_n ” was first defined [Bier–76]; see [BeBr–97] for (iii) and [Brow–82, Section VII.5, Exercise 3] for (iv). The following question is natural:

Does \mathfrak{G} have an amenable cover of type FP_2 ?

The answer is due to Yves de Cornulier (unpublished). We reproduce it here, with our thanks to him.

Proposition 5.11 (de Cornulier). *Any cover of type FP_2 of the group \mathfrak{G} is large.*

Proof. Let $E \twoheadrightarrow \mathfrak{G}$ be a cover, with E of type FP_2 . By (iv) above, there exists a finitely presented group F and a perfect normal subgroup P of F such that F/P is isomorphic to E . For n large enough and \mathfrak{G}_n as in Theorem 5.3, there exists by Proposition 3.3 a normal subgroup K_n of F such that F/K_n is isomorphic to \mathfrak{G}_n . Observe that the group

$$F/PK_n \simeq (F/K_n)/(P/P \cap K_n) \simeq \mathfrak{G}_n/(P/P \cap K_n)$$

is a quotient of \mathfrak{G}_n by a perfect normal subgroup.

Since \mathfrak{G}_n has a subgroup of index a power of 2 which is a direct product of free groups (Theorem 5.3, or [BaCo–06]), \mathfrak{G}_n is residually soluble. It follows that the only perfect subgroup of \mathfrak{G}_n is $\{e\}$, so that $F/PK_n \simeq \mathfrak{G}_n$. Hence F/PK_n is large, and so is its cover $F/P \simeq E$. \square

APPENDIX A. On soluble groups, metabelian groups, and finite presentations

The existence of groups that are finitely generated and infinitely presented was established by B.H. Neumann in 1937. More precisely, he constructed *uncountably many 2-generator groups* [Neum–37, Theorem 14]; at most countably many of them are finitely presented. Later it was checked that none of them is finitely presented (see the last proof of Appendix C, as well as [BaMi–09, Theorem C]).

Infinitely generated *soluble* groups are equally abundant, as we recall below after having fixed some notation.

The groups of the derived series of a group G are defined inductively by $D^0G = G$ and $D^{\ell+1}G = [D^\ell G, D^\ell G]$. The **free soluble group of rank k and solubility class ℓ** is the quotient $\text{FSol}(k, \ell) = F_k/D^\ell F_k$, where F_k stands for the free group of rank k . Any k -generated soluble group of solubility class at most ℓ is a quotient of $\text{FSol}(k, \ell)$. A group G is **metabelian** if $D^2G = 1$, namely if it is a cover of an abelian group with abelian kernel. The group $\text{FSol}(k, 2)$ is the **free metabelian group of rank k** .

Philip Hall established the existence of *uncountably many finitely generated soluble groups*. His result is much more precise [Hall–54, Theorem 6]: given any countable abelian group $A \neq 1$, there exist uncountably many groups G such that

$$d(G) = 2, \quad Z(G) \simeq A, \quad [G, D^2G] = 1.$$

Here $d(G)$ stands for the minimal number of generators of G , and $Z(G)$ for its centre. The condition $[G, D^2G] = 1$ can be translated in words: G is a *centre-by-metabelian group*. It is moreover known that there are uncountably many finitely generated soluble groups which are not quasi-isometric to each other [CoTe, Corollary 1.8].

On the contrary, there are only countably many finitely generated metabelian groups (this is repeated as Proposition A.3 below), and more generally¹¹ abelian-by-polycyclic groups ([Hall–54, Corollary 2 to Theorem 3], see also [LeRo–04, Corollary 4.2.5]). Before comparing soluble groups in general with metabelian groups in particular, we collect some well-known facts in the following lemma.

Recall that a group G satisfies **Max-n, the maximal condition for normal subgroups**, if any increasing sequence of normal subgroups of G is ultimately stationary, or equivalently if any normal subgroup of G is finitely generated *as normal subgroup*.

Lemma A.1. *Let G be a finitely generated group, N a normal subgroup, and Z a central subgroup.*

(i) *If G/N is finitely presented, there exists a finite subset $S \subset N$ such that N is the smallest normal subgroup of G containing S .*

(ii) *If G/Z is finitely presented, then Z is finitely generated.*

(iii) *If G has uncountably many normal subgroups, then G has uncountably many pairwise non-isomorphic quotients.*

¹¹Let \mathcal{P} and \mathcal{Q} be group properties. A group G is \mathcal{P} -by- \mathcal{Q} if G has a normal subgroup N with Property \mathcal{P} such that G/N has Property \mathcal{Q} .

(iv) Suppose that G is finitely presented and satisfies Max-n. Then G/N is finitely presented.

Proof. Claim (i) is [Robi–96, Lemma 14.1.3]. It is a simple consequence of the following fundamental observation of B.H. Neumann: let S, S' be two finite generating sets of a group G ; assume that G has a finite presentation $\langle S \mid R \rangle$ involving S and a finite set R of relators; then there exists a finite set R' of relators in the letters of S' such that $\langle S' \mid R' \rangle$ is also a finite presentation of G [Neum–37, Lemma 8].

Claim (ii) is the special case of (i) for a central subgroup.

For Claim (iii), consider an uncountable family $(N_\alpha)_{\alpha \in A}$ of distinct normal subgroups of G . Fix $\alpha \in A$. Let B be a subset of A such that, for each $\beta \in B$, there exists an isomorphism $\phi_\beta : G/N_\beta \rightarrow G/N_\alpha$. It suffices to show that B is countable.

For $\beta \in B$, let π_β denote the composition of the canonical projection $G \rightarrow G/N_\beta$ with ϕ_β . Since G is finitely generated and G/N_α countable, there are only countably many homomorphisms from G to G/N_α . As $N_\beta = \ker(\pi_\beta)$, the set B is countable.

For Claim (iv), consider a finite presentation of G , namely a free group F on a finite set S and a normal subgroup M of F generated as normal subgroup by a finite subset R of F , such that $G = F/M$. Since G satisfies Max-n, there exists a finite subset R' of F of which the image in G generates N as a normal subgroup. Then $\langle S \mid R \cup R' \rangle$ is a finite presentation of G/N .

Note that Claim (ii) is a special case of [Hall–54, Lemma 2]. Our argument for Claim (iii) can be found in [Hall–54, Page 433], and that for Claim (iv) is “a well-known principle” cited in [Hall–54, Page 420]. \square

Finitely generated metabelian groups are “well-behaved” in many ways:

Proposition A.2 (Hall, Baumslag, Remeslennikov). *Let G be a finitely generated metabelian group.*

- (i) G satisfies Max-n. In particular, the centre of G is finitely generated.
- (ii) If G is finitely presented, so is any quotient of G .
- (iii) G is residually finite.
- (iv) G has a soluble word problem.
- (v) G can be embedded into a finitely presented metabelian group.
- (vi) G is recursively presented.

References. Claim (i) is [Hall–54, Theorem 3], Claim (ii) follows by Lemma A.1.iv, and Claim (iii) is [Hall–59, Theorem 1].

For the particular case of free metabelian groups, Claim (iii) follows from linearity: it is known that $\text{FSol}(k, 2)$ is a subgroup of $\text{GL}_2(\mathbf{C})$. This is a form of the “Magnus embedding theorem” ; see [Magn–39], and also [Wehr–73, Theorem 2.11]. (On the contrary, $\text{FSol}(k, \ell)$ is not linear when $\ell \geq 3$; see Remark A.6.)

Claim (iv) can be found in [Baum–74]. It is also a consequence of (a particular case of) a result of Wehrfritz: any finitely generated metabelian group is quasi-linear, namely is a subgroup of a group of the form $\prod_{i=1}^r \text{GL}_n(F_i)$, where F_1, \dots, F_r are fields [Wehr–80]. More generally, several algorithmic problems are known to be soluble in finitely generated metabelian groups [BaCR–94].

Claim (v) was proved by Baumslag [Baum–73] and Remeslennikov [Reme–73], independently. See also [LeRo–04, Proposition 11.3.2].

Claim (vi) is [Baum–74, Corollary A1]; it also follows from Claim (v). Note that Claim (vi) is contained in Claim (iv), but we add it for comparison with Proposition A.5. \square

Proposition A.3 (P. Hall). *There are countably many finitely generated metabelian groups.*

Remark A.4. (a) Let G be a finitely generated metabelian group G ; let $k \geq 0$ be such that G can be generated by k elements, so that G is a quotient of the free metabelian group $\text{FSol}(k, 2)$. Though it need not be finitely presented (examples are shown below), the group G is finitely presented *as a metabelian group*, because $\text{FSol}(k, 2)$ satisfies Max-n; in other terms, we can write

$$G = \text{FSol}(k, 2) / \langle\langle r_1, \dots, r_n \rangle\rangle,$$

where the notation $\langle\langle \dots \rangle\rangle$ indicates a normal subgroup generated *as such* by elements r_1, \dots, r_n in $\text{FSol}(k, 2)$. Proposition A.3 follows.

Note that Proposition A.3 is also a straightforward consequence of Claim (vi) in the previous proposition.

(b) Some of the claims in Proposition A.2 can be improved. For example, (i) holds for finitely generated abelian-by-polycyclic groups, and (iii) holds for finitely generated abelian-by-nilpotent groups (Hall). Moreover (iii) holds for abelian-by-polycyclic groups, as shown by Roseblade and Jategaonkar in 1973 and 1974 (see [Sega–75], or Chapter 7 and in particular Theorem 7.2.1 in [LeRo–04]).

(c) Until the early 70’s, there were rather few known examples of finitely presented metabelian groups. The 3-generator 3-relator group

$$H = \langle a, s, t \mid a^t = aa^s, [s, t] = 1 = [a, a^s] \rangle,$$

appeared independently in papers by Baumslag [Baum–72] and Remeslennikov [Reme–73]; see also [Stre–84, Theorem A]. It is metabelian,

its derived group is free abelian of infinite rank, and it contains the wreath product $\mathbf{Z} \wr \mathbf{Z}$ as a subgroup [Baum–74, Pages 72–73]. It was quite a surprise at this time [Baum–72, first lines] to find a finitely presented group containing a normal abelian subgroup of infinite rank. More recently, the quotient group $H/(a^2 = 1)$ was the main character in [GLSZ–00].

For groups of higher solubility degrees, the picture is substantially different, even under the stronger hypothesis of finite presentability. Each of the claims of the next proposition is meant to be compared with the corresponding claim of Proposition A.2.

Proposition A.5. *Let G be a finitely presented soluble group.*

- (i) *G need not satisfy Max-n. Indeed, the centre of G need not be finitely generated.*
- (ii) *G may have uncountably many quotients, and in particular infinitely presented quotients.*
- (ii') *Any metabelian quotient of G is finitely presented.*
- (iii) *G need not be residually finite.*
- (iv) *G need not have a soluble word problem.*

Let G be now a finitely generated soluble group.

- (v) *G need not be recursively presented.*
- (vi) *G need not embed into any finitely presented group.*

On the proof. Let p be a prime. For $n \geq 2$, consider the group A_n of n -by- n triangular matrices of the form

$$\begin{pmatrix} 1 & * & \cdots & * & * \\ 0 & * & \cdots & * & * \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & * & * \\ 0 & 0 & \cdots & 0 & 1 \end{pmatrix}$$

with upper-triangular entries $a_{i,j}$, $1 \leq i < j \leq n$, in $\mathbf{Z}[\frac{1}{p}]$ and diagonal entries $a_{2,2}, \dots, a_{n-1,n-1}$ in $p^{\mathbf{Z}}$. This group is soluble. Its center $Z(A_n)$ is isomorphic to $\mathbf{Z}[\frac{1}{p}]$, that is to A_2 , and therefore is not finitely generated; it follows that A_n does not satisfy Max-n. It is easy to check that A_n is finitely generated when $n \geq 3$.

For $n \geq 4$, the group A_n is finitely presented ([Abel–77] for $n = 4$ and [AbBr–87] for $n \geq 4$). This justifies Claim (i). The existence of a finitely presented soluble group without Max-n solves a problem of P. Hall; Remeslennikov had an earlier claim for this [Reme–72] which was apparently unjustified [Thom–77].

Note that A_3 is infinitely presented. This was most likely known to P. Hall, and can be found in [AbBr–87]. But it is also a consequence of Bieri-Strebel Theorem 1.5; indeed, since $Z(A_3)$ is not finitely generated, $A_3/Z(A_3)$ is infinitely presented (Lemma A.1), and the soluble group A_3 cannot be a finitely presented cover of the metabelian group $A_3/Z(A_3)$.

Since $Z(A_n) \simeq \mathbf{Z}[\frac{1}{p}]$ is not finitely generated, the quotient $A_n/Z(A_n)$, with $n \geq 3$, is finitely generated non-finitely presented, by Lemma A.1.ii. When $n \geq 4$, this justifies the second part of Claim (ii).

Claim (ii') follows from Theorem 1.5.

For $n \geq 3$, the quotient of A_n by the central subgroup

$$\left\{ (z_{i,j})_{1 \leq i,j \leq n} \in A_n \mid \begin{array}{l} z_{i,j} = \delta_{i,j} \text{ for } (i,j) \neq (1,n) \\ z_{1,n} \in \mathbf{Z} \end{array} \right\} \simeq \mathbf{Z}$$

(where $(\delta_{i,j})_{1 \leq i,j \leq n}$ denotes the unit matrix) is finitely generated non-Hopfian¹² (the argument of [Hall–61] for $n = 3$ carries over to all $n \geq 3$), and therefore non-residually finite. When $n \geq 4$, this justifies Claim (iii).

Still for $n \geq 4$, it is known that the quotient $A_n/Z(A_n)$ does satisfy Max-n and does not have any minimal presentation [BCGS, Lemma 3.2 and Corollary 3.6]. The last statement means that any presentation of $A_n/Z(A_n)$ contains redundant relators; in particular, the finitely related group A_n has a quotient that is not finitely related. The group A_n itself has only countably many quotients; see [Lyul–84, Theorem 1] and [BCGS, Corollary 3.4].

Concerning Claim (iv), finitely presented soluble groups with unsolvable word problems have been constructed by Kharlampovich in [Khar–81] and by Baumslag, Gildenhuys and Strebel in [BaGS–86]. Groups in [BaGS–86] have centers that are not finitely generated, and therefore have infinitely presented quotients (see again Claim (ii)).

Earlier, Meskin had constructed a finitely generated recursively presented residually finite soluble group with unsolvable word problem [Mesk–74].

For any prime p , there exists a finitely presented soluble group G with centre $(C_p)^{(\infty)}$, an infinite direct sum of cyclic groups of order p (see [Khar–90], as well as [KhMS, Lemma 4.14]). Hence G has uncountably many quotients, by Lemma A.1.iii. This justifies the first part of Claim (ii).

¹² A group G is **non-Hopfian** if there exists a surjective endomorphism of G onto itself that is not injective. A residually finite finitely generated group is Hopfian [Mal'c–40].

Claim (v) follows from the existence of uncountably many finitely generated soluble groups, because there are only countably many recursively presented groups.

(Digression: Let G be a finitely generated soluble group. Assume that G is of finite Prüfer rank, namely that there exists an integer $d \geq 1$ such that any finitely generated subgroup of G can be generated by d elements. Then G is recursively presented if and only if G has a soluble word problem. For this, and for examples of G with and without soluble word problem, see [CaRo–84].)

Claim (vi) follows from Claim (v) because a finitely generated subgroup of a finitely presented group is recursively presented. (This is straightforward; see the first page of [Higm–61], where Higman establishes the famous non-trivial converse; alternatively, see [Mill–89, Lemma 2.1].) \square

Remark A.6. As noted parenthetically just after Proposition A.2, free solvable groups $\text{FSol}(k, \ell)$ are not linear when $k \geq 2$ and $\ell \geq 3$ [Smir–64, Smir–65]. Here is a proof.

For $k \geq 3$, here is first a short reduction to a more standard result. Let \mathfrak{C} denote the class of (nilpotent-by-abelian)-by-finite groups; observe that quotients of groups in \mathfrak{C} are in \mathfrak{C} . The Lie-Kolchin-Mal'cev theorem (see for example [Robi–96, Section 15.1]) establishes that soluble linear groups are in \mathfrak{C} . If $\text{FSol}(k, \ell)$ were linear, hence in \mathfrak{C} , so would be any quotient, in particular the iterated wreath product $(\mathbf{Z} \wr \mathbf{Z}) \wr \mathbf{Z}$; but this is not [Robi–96, 15.1.5].

The following argument, shown to us by Ralph Strebels, holds for any $k \geq 2$. Denote by x_1, \dots, x_k a set of free generators of $\text{FSol}(k, \ell)$. Let H be a subgroup of finite index in $\text{FSol}(k, \ell)$; there exists an integer $m \geq 1$ such that $x_1^m, x_2^m \in H$; let U denote the subgroup of $\text{FSol}(k, \ell)$ generated by x_1^m and x_2^m , so that

$$U \subset H \subset_{\text{finite index}} \text{FSol}(k, \ell).$$

The group U is isomorphic to $\text{FSol}(k, \ell)$, by a result due independently to Gilbert Baumslag [Baum–63, Theorem 2] and Shmel'kin [Shme–64, Theorem 5.2].

If $\text{FSol}(k, \ell)$ was linear, it would have a nilpotent-by-abelian subgroup of finite index, say H . By the lines above, $\text{FSol}(2, \ell)$ would be nilpotent-by-abelian. Hence any 2-generated soluble group of solubility class at least 3 would be nilpotent-by-abelian. But this is not true.

Indeed, consider the symmetric group S_4 on four letters. It is generated by two elements, a transposition and a 4-cycle. It is soluble of class 3, with $D^1 S_4 = A_4$ (the alternating group on four letters), $D^2 S_4 = V$

(the Klein Vierergruppe), and $D^3S_4 = \{1\}$. It is not nilpotent-by-abelian, namely A_4 is not nilpotent: $[A_4, V] = V$.

The argument of Smirnov is different. It relies on the bi-orderability of the groups $\text{FSol}(k, \ell)$.

APPENDIX B. On wreath products and lamplighter groups

Permutational wreath products have been defined in the beginning of Section 2. The **standard wreath product** $G \wr_H H$ refers to the action of H on itself by left multiplications.

Proposition B.1. *Consider two groups G, H , a non-empty H -set X , and the permutational wreath product $G \wr_X H$. We assume that $G \neq \{1\}$.*

(i) *$G \wr_X H$ is finitely generated if and only if G, H are finitely generated and H has finitely many orbits on X .*

(ii) *$G \wr_X H$ is finitely presented if and only if G, H are finitely presented, the H_x 's ($x \in X$) are finitely generated, and there are finitely many orbits in $X \times X$ for the diagonal action of H (where $H_x = \{h \in H \mid h(x) = x\}$).*

(iii) *In particular, as soon as H is infinite, the standard wreath product $G \wr_H H$ is not finitely presented.*

References. Claim (i) is standard, and easy; if necessary, see [Corn–06, Proposition 2.1]. For Claim (ii), see [Corn–06, Theorem 1.1]. Claim (iii) is the main result of [Baum–61]. \square

As a particular case of Proposition B.1, if G is finitely-generated abelian and $G \neq 1$, the group $G \wr \mathbf{Z}$ is metabelian, finitely generated, and infinitely presented. When G is finite abelian and $H \simeq \mathbf{Z}$ infinite cyclic, we will call $G \wr \mathbf{Z}$ the **lamplighter group** for G . (For this terminology, precise assumptions on G and H vary from one author to the other; some ask that $G = \mathbf{Z}/2\mathbf{Z}$.)

Proposition B.2. *Consider two finitely presented groups G, H , an H -set X such that H has finitely many orbits on X and infinitely many orbits on $X \times X$, and the permutational wreath product $G \wr_X H$. We assume that $G \neq \{1\}$.*

(i) *$G \wr_X H$ is finitely generated and is infinitely presented.*

(ii) *For any finitely presented cover $\pi : E \twoheadrightarrow G \wr_X H$, the group E has non-abelian free subgroups.*

Proof. Claim (i) is a particular case of Proposition B.1. In Proposition 2.10 of [Corn–06], it is shown that the kernel of π contains non-abelian free subgroups. \square

Note that, in the particular case of two abelian groups G and H , Claim (ii) is also a consequence of Theorem 1.5.

On other proofs of Proposition B.2 in the case of $W := \mathbf{Z} \wr_{\mathbf{Z}} \mathbf{Z}$. (We have $G = H = \mathbf{Z}$ and $X = \mathbf{Z}$.) We have a presentation

$$W = \langle s, t \mid [s^{t^i}, s^{t^j}] \ \forall i, j \in \mathbf{Z} \rangle;$$

indeed, any element in the right-hand side can be written as

$$\begin{aligned} & s^{m_1} t^{n_1} s^{m_2} t^{n_2} s^{m_3} t^{n_3} \dots s^{m_\ell} t^{n_\ell} \\ &= s^{m_1} (s^{t^{-n_1}})^{m_2} (s^{t^{-n_1-n_2}})^{m_3} \dots (s^{t^{-n_1-\dots-n_{\ell-1}}})^{m_\ell} t^{n_1+\dots+n_\ell} \end{aligned}$$

for some $m_1, n_1, \dots, m_\ell, n_\ell \in \mathbf{Z}$, and therefore as

$$(s^{t^{j_1}})^{i_1} (s^{t^{j_2}})^{i_2} \dots (s^{t^{j_k}})^{i_k} t^N$$

for appropriate $i_1, j_1, \dots, i_k, j_k, N \in \mathbf{Z}$ with $i_1 < i_2 < \dots < i_k$. It follows that the natural homomorphism

$$\langle s, t \mid [s^{t^i}, s^{t^j}] \ \forall i, j \in \mathbf{Z} \rangle \longrightarrow W$$

is an isomorphism.

Since $t^i [s, s^{t^k}] t^{-i} = [s^{t^i}, s^{t^{i+k}}]$, we have a second presentation

$$W = \langle s, t \mid [s, s^{t^i}] \ \forall i \in \mathbf{N} \rangle.$$

For a positive integer n , define

$$W_n = \langle s, t \mid [s, s^{t^i}], \ i = 0, \dots, n \rangle.$$

Note that $\lim_{n \rightarrow \infty} W_n = W$ in \mathcal{M}_2 . We have a third presentation

$$W_n = \left\langle s_0, \dots, s_n, t \mid \begin{array}{l} [s_i, s_j], \ 0 \leq i, j \leq n, \\ s_k^t = s_{k+1}, \ 0 \leq k \leq n-1 \end{array} \right\rangle.$$

Indeed, it can be checked that the assignments

$$\begin{aligned} \varphi_1 &: s \longmapsto s_0, \ t \longmapsto t \\ \varphi_2 &: s_i \longmapsto s^{t^i}, \ t \longmapsto t \quad (0 \leq i \leq n) \end{aligned}$$

define, between the groups of the two previous presentations, isomorphisms that are inverse to each other.

Let H_n be the free abelian subgroup of W_n generated by s_0, \dots, s_n . Denote by K_n the subgroup of H_n generated by s_0, \dots, s_{n-1} , and by L_n that generated by s_1, \dots, s_n ; observe that $K_n \simeq L_n \simeq \mathbf{Z}^n$. Let $\psi_n : K_n \longrightarrow L_n$ be the isomorphism defined by $\psi(s_{i-1}) = s_i$ for $i = 1, \dots, n$. Then W_n is clearly the HNN-extension corresponding to the data $(H_n, \psi_n : K_n \xrightarrow{\sim} L_n)$. By Britton's lemma, W_n contains non-abelian free subgroups.

It follows from Corollary 3.4 that any finitely presented cover of W contains non-abelian free groups.

Let us finally allude to another argument showing that W is infinitely presented. Let H denote the subgroup of $\mathrm{GL}_3(\mathbf{C})$ generated by the three matrices

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & t & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{and} \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix},$$

where t is some transcendental number. The centre $Z(H)$ of H is free abelian of infinite rank, and the quotient $H/Z(H)$ is isomorphic to $M := \mathbf{Z}[t, t^{-1}]^2 \rtimes \mathbf{Z}$, where \rtimes refers to the action of the generator $1 \in \mathbf{Z}$ by $\begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}$. There is an isomorphism of M onto a subgroup of index 2 in $\mathbf{Z}[t, t^{-1}] \rtimes_t \mathbf{Z} \simeq \mathbf{Z} \wr \mathbf{Z}$, given by $(P(t), Q(t), n) \mapsto (P(t^2) + tQ(t^{-2}), 2n)$. It follows from Lemma A.1.ii that $\mathbf{Z} \wr \mathbf{Z}$ is not finitely presented. We are grateful to Adrien Le Boudec for correcting an earlier version of our argument at this point.

The isomorphism of $H/Z(H)$ with a subgroup of index 2 in $\mathbf{Z} \wr \mathbf{Z}$ is essentially due to P. Hall. See [Hall–54, Theorem 7] and [CaCo, Lemma 3.1]. \square

Concerning Proposition B.2, let us add one more remark about the particular case $W = (\mathbf{Z}/h\mathbf{Z}) \wr \mathbf{Z}$, with $h \geq 2$: it is known that any finitely presented cover of W is large [Baum–93, Section IV.3, Theorem 7].

Recall that the free soluble group $\mathrm{FSol}(k, \ell) = F_k/D^\ell F_k$ of rank k and solubility class ℓ has been defined in Appendix A.

Corollary B.3. *For $k, \ell \geq 2$, the group $\mathrm{FSol}(k, \ell)$ is infinitely presented, and any finitely presented cover of it contains non-abelian free subgroups.*

Note. (i) That $\mathrm{FSol}(k, \ell)$ is infinitely presented is a result due to [Shme–65]. See also [Corn–06, Proposition 2.10 and Corollary 2.14].

(ii) Corollary B.3 and our proof carry over to free polynilpotent groups

$$\mathrm{FPolynilp}(k, \ell_1, \dots, \ell_n) := F_k/C^{\ell_k}(C^{\ell_{k-1}}(\dots C^{\ell_1}(F_k)\dots))$$

for any $k \geq 2$, $n \geq 2$, and $\ell_1, \dots, \ell_n \geq 2$ (where $C^\ell G$ denotes the ℓ th group of the lower central series of a group G , defined by $C^1 G = G$, and $C^{\ell+1} G = [G, C^\ell G]$). These groups are the subject of [Shme–64].

Proof. Since $\mathbf{Z} \wr \mathbf{Z}$ is a two-generator metabelian group, we have an epimorphism $\text{FSol}(2, 2) \twoheadrightarrow \mathbf{Z} \wr \mathbf{Z}$. Indeed, we have a sequence of natural epimorphisms

$$\text{FSol}(k, \ell) \twoheadrightarrow \text{FSol}(k, 2) \twoheadrightarrow \text{FSol}(2, 2) \twoheadrightarrow \mathbf{Z} \wr \mathbf{Z}.$$

Hence any finitely presented cover of $\text{FSol}(k, \ell)$ is also one of $\mathbf{Z} \wr \mathbf{Z}$. If $\text{FSol}(k, \ell)$ was finitely presented, it would contain non-abelian free subgroup by Proposition B.2, but this cannot be in a soluble group. \square

In the situation of Proposition B.2, suppose moreover that H is infinite residually finite, G has at least one non-trivial finite quotient, and consider the standard wreath product ($X = H$). The following strengthening of Claim (ii) is shown in [CoKa–11, Theorem 1.5]: any finitely presented cover of $G \wr H$ is *large*. In particular:

any finitely presented cover of $\mathbf{Z} \wr \mathbf{Z}$ is large.

By the proof of Corollary B.3, it follows that,

for $k, \ell \geq 2$, any finitely presented cover of $\text{FSol}(k, \ell)$ is large.

The following notion provides interesting examples of metabelian groups, as we will illustrate with Baumslag-Solitar groups.

Definition B.4. The **metabelianization** of a group G is the metabelian quotient group G/D^2G .

Definition B.5 ([BaSo–62]). For $\ell, m \in \mathbf{Z} \setminus \{0\}$, the **Baumslag-Solitar group** is defined by the two-generators one-relator presentation

$$\text{BS}(\ell, m) = \langle s, t \mid t^{-1}s^\ell t = s^m \rangle.$$

We collect three well-known properties of these groups as follows.

Proposition B.6. *Let $\ell, m \in \mathbf{Z} \setminus \{0\}$ and $\text{BS}(\ell, m)$ be as above.*

- (i) *$\text{BS}(\ell, m)$ is abelian if and only if $\text{BS}(\ell, m)$ is nilpotent, if and only if $\ell = m = \pm 1$.*
- (ii) *$\text{BS}(\ell, m)$ is metabelian if and only if $\text{BS}(\ell, m)$ is soluble, if and only if $\text{BS}(\ell, m)$ does not contain non-abelian free subgroups, if and only if $|\ell| = 1$ or $|m| = 1$.*
- (iii) *If ℓ, m satisfy $\ell, m \geq 2$ and are coprime, then $\text{BS}(\ell, m)$ is non-Hopfian.*
- (iv) *For ℓ, m as in (iii), the group $\text{BS}(\ell, m)$ contains non-abelian free groups, but is not large.*

On the proof. It is easy to check that the four groups $\text{BS}(\ell, m)$, $\text{BS}(m, \ell)$, $\text{BS}(-\ell, -m)$, $\text{BS}(-m, -\ell)$ are isomorphic. For simplicity, let us assume

from now on that ℓ and m are positive. (For the general case, with all details, we refer to [Souc–01]).

It is an exercise to check that $\text{BS}(1, m) \simeq \mathbf{Z} \left[\frac{1}{m} \right] \rtimes_m \mathbf{Z}$ for any $m \geq 1$. It follows that $\text{BS}(\ell, m)$ is metabelian if $\ell = 1$ or $m = 1$, and abelian if and only if $\ell = m = 1$. If $m \geq 2$, note that $\text{BS}(1, m)$ is not nilpotent, because its subgroup $\mathbf{Z} \left[\frac{1}{m} \right]$ is not finitely generated. If $\ell \geq 2$ and $m \geq 2$, the subgroup generated by $s^{-1}ts$ and t is free of rank 2, by Britton’s Lemma.

Claim (iii) is the main reason for the celebrity of these groups. It is straightforward to check that the assignments $\varphi(s) = s^\ell$ and $\varphi(t) = t$ define an endomorphism φ of $\text{BS}(\ell, m)$. The image of φ contains t and s^ℓ , hence $t^{-1}s^\ell t^{-1} = s^m$, and therefore s ; hence φ is onto. On the one hand, $\varphi([t^{-1}st, s]) = [t^{-1}s^\ell t, s^\ell] = [s^m, s^\ell] = 1$; on the other hand, $[t^{-1}st, s] = t^{-1}st s t^{-1}s^{-1}t s^{-1} \neq 1$, where the last inequality holds by Britton’s Lemma; hence φ is not one-to-one.

More generally, we know necessary and sufficient conditions on ℓ, m for $\text{BS}(\ell, m)$ to be non-Hopfian; see [BaSo–62, Coll–78, CoLe–83].

The first part of Claim (iv) is standard (it also follows from Theorem 1.5, see B.8.ii below). For the second part, see Example 3.2 and Theorem 6 in [EdPr–84]. \square

Definition B.7. For two coprime positive integers ℓ, m , not both 1, let

$$\text{Met}(\ell, m) = \left(\begin{array}{cc} \left(\frac{\ell}{m} \right)^{\mathbf{Z}} & \mathbf{Z} \left[\frac{1}{\ell m} \right] \\ 0 & 1 \end{array} \right) \simeq \mathbf{Z} \left[\frac{1}{\ell m} \right] \rtimes_{\ell/m} \mathbf{Z}$$

be the group of triangular matrices generated by

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} \frac{\ell}{m} & 0 \\ 0 & 1 \end{pmatrix}.$$

Let

$$\mu_{\ell, m} : \text{BS}(\ell, m) \rightarrow \text{Met}(\ell, m)$$

denote the epimorphism defined by

$$\mu_{\ell, m}(s) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \text{ and } \mu_{\ell, m}(t) = \begin{pmatrix} \frac{\ell}{m} & 0 \\ 0 & 1 \end{pmatrix}.$$

The following proposition collects facts on $\text{BS}(\ell, m)$ and $\text{Met}(\ell, m)$. Claims (iii) to (v) constitute a digression from our theme.

Proposition B.8. *Let the notation be as just above, and ℓ, m be two coprime positive integers, not both 1.*

(i) $\mu_{\ell, m}$ is an isomorphism if and only if $\ell = 1$ or $m = 1$.

We assume furthermore that $\min\{\ell, m\} \geq 2$.

(ii) $\text{Met}(\ell, m)$ is infinitely presented.

(iii) The multiplier group $H_2(\text{Met}(\ell, m), \mathbf{Z})$ is trivial.

(iv) $\text{Met}(\ell, m)$ is of cohomological dimension 3.

(v) For $x \in \mathbf{C}$ transcendental, the matrices $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$

generate a group isomorphic to $\mathbf{Z} \wr \mathbf{Z}$.

On proofs. Claim (i) has already been given as an exercise, in the proof of Proposition B.6. Claim (ii) is a consequence of a particular case of the main result of [BaSt–76], or a consequence of [BiSt–78, Theorem C]; see also [LeRo–04, Proposition 11.4.3]. Claim (iv) is [Gild–79, Theorem 4]. Claim (v) is [LeRo–04, Proposition 3.1.4].

For (iii), see [BaSt–76, No 1.8]. Recall that, if a group G is finitely presented, then its multiplier group $H_2(G, \mathbf{Z})$ is finitely generated; this is a simple consequence of the so-called *Schur-Hopf Formula*, for a group $G = F/R$ presented as a quotient of a free group F , which reads “ $H_2(G, \mathbf{Z}) = (R \cap [F, F])/[R, F]$ ”. Claim (iii) is one of the standard examples showing that the converse *does not* hold. \square

Let ℓ, m be coprime positive integers, with $\ell, m \geq 2$. We denote by $p_{\ell, m} : F_2 \rightarrow \text{BS}(\ell, m)$ the defining cover of the corresponding Baumslag-Solitar group, namely the cover mapping a basis of the free group of rank 2 onto $\{s, t\}$. Let $\varphi : \text{BS}(\ell, m) \rightarrow \text{BS}(\ell, m)$ be the usual non-injective surjective endomorphism, as in Proposition B.6. For $n \geq 1$, set

$$M_n = \ker(\varphi^n), \quad N_n = \ker(\varphi^n p_{m, \ell}), \quad G_n = \text{BS}(\ell, m)/M_n = F_2/N_n.$$

Observe that the sequence $(M_n)_{n \geq 1}$ is strictly increasing, yet G_n is isomorphic to $\text{BS}(\ell, m)$ for each $n \geq 1$.

Proposition B.9. *Let ℓ, m be coprime integers, with $\ell, m \geq 2$, and $\text{Met}(\ell, m)$ as in Definition B.7. Let $(G_n)_{n \geq 1}$ be as above.*

(i) $\text{Met}(\ell, m)$ is isomorphic to $\text{BS}(\ell, m)_{\text{metab}}$.

(ii) With the notation above, we have an isomorphism

$$\text{Met}(\ell, m) \simeq F_2 / \left(\bigcup_{n \geq 1} N_n \right),$$

so that $\lim_{n \rightarrow \infty} G_n = \text{Met}(\ell, m)$ in \mathcal{M}_2 .

(iii) Any finitely presented cover of $\text{Met}(\ell, m)$ contains non-abelian free subgroups.

Proof. Claim (i) is part of [Baum–74, Theorem G].

Claim (ii) is [BaSt–76, see 1.8] or [GrMa–97, Theorem 3], there for $(\ell, m) = (2, 3)$, but the argument carries over to the case stated

here. (When working on [GrMa–97], the authors were not aware of [BaSt–76].)

Since $\text{BS}(\ell, m)$ contains non-abelian free subgroups by Proposition B.6.ii, and since $\lim_{n \rightarrow \infty} G_n = \text{Met}(\ell, m)$, Claim (iii) follows by Corollary 3.4. \square

APPENDIX C. On Bieri-Neumann-Strebel invariants

Example C.1. Consider the two metabelian groups

$$\begin{aligned} \text{Met}(1, 6) &= \text{BS}(1, 6) = \mathbf{Z}[1/6] \rtimes_{1/6} \mathbf{Z} \\ \text{Met}(2, 3) &= \text{BS}(2, 3)_{\text{metab}} = \mathbf{Z}[1/6] \rtimes_{2/3} \mathbf{Z}. \end{aligned}$$

The first group is finitely presented. By contrast, the second group is infinitely presented (Propositions B.8). This spectacularly different behavior of two superficially similar-looking groups was the initial motivation of Bieri and Strebel (later joined by Neumann and others) for the work that lead to the Σ -invariants; see [Bier–79], as well as [Baum–74, Problem 1]. We come back to these two examples in Example C.3.

Let G be a finitely generated group, S a finite generating set, and $\text{Cay}(G, S)$ the corresponding **Cayley graph**, with vertex set G and edge set $\{\{g, h\} \mid g^{-1}h \in S \cup S^{-1}\}$. The set of **characters**, namely of group homomorphisms from G to \mathbf{R} , is a real vector space $\text{Hom}(G, \mathbf{R})$ isomorphic to \mathbf{R}^n , where n is the torsion-free rank of the abelian group G/D^1G . Let $S(G)$ denote the **character sphere** of G , quotient of the set of non-zero characters by the natural action of the group of positive reals; we have $S(G) \approx \mathbf{S}^{n-1}$, and we write $[\chi]$ the class in $S(G)$ of a character $\chi \neq 0$ in $\text{Hom}(G, \mathbf{R})$. (The sign \approx indicates a homeomorphism; \mathbf{S}^k denotes the k -sphere, and $\mathbf{S}^{-1} = \emptyset$.) For such a $[\chi]$, set

$$G_{[\chi]} = \{g \in G \mid \chi(g) \geq 0\},$$

and let $\text{Cay}_{[\chi]}(G, S)$ be the subgraph of $\text{Cay}(G, S)$ spanned by $G_{[\chi]}$. The **Bieri-Neumann-Strebel invariant**, or shortly **BNS-invariant**, of G is defined to be

$$\Sigma^1(G) = \{[\chi] \in S(G) \mid \text{Cay}_{[\chi]}(G, S) \text{ is connected}\} \subset S(G).$$

The superscript indicates that Σ^1 is just one out of many “geometric invariants” (see [BiRe–88] and [Stre]).

Observe that there is an antipodal map $[\chi] \mapsto -[\chi] := [-\chi]$ defined on $S(G)$. Let G_0 be a finitely generated group and $p : G_0 \twoheadrightarrow G$ an epimorphism; then p induces a map $p^* : S(G) \longrightarrow S(G_0)$ which a

homeomorphism onto its image and which intertwines the antipodal maps.

On the one hand, the invariant of [BiNS–87] involves one more group A on which D^1G acts in an appropriate way; here, we particularize to $A = D^1G$. On the other hand, the definition above is a reformulation of the original definition, see for example [Stre].

Theorem C.2. *Let G be a finitely generated group, and $\Sigma^1(G) \subset S(G)$ as above.*

(i) $\Sigma^1(G)$ is independent of the choice of S .

(ii) $\Sigma^1(G)$ is open in $S(G)$.

(iii) $\Sigma^1(G) = S(G)$ if and only if the derived group D^1G is finitely generated.

(iv) Suppose that G is metabelian. Then G is finitely presented if and only if $\Sigma^1(G) \cup (-\Sigma^1(G)) = S(G)$, and G is polycyclic if and only if $\Sigma^1(G) = S(G)$.

(v) If G is finitely presented and has no non-abelian free subgroup, then $\Sigma^1(G) \cup (-\Sigma^1(G)) = S(G)$.

(vi) If G_0 is a finitely generated group and $p : G_0 \twoheadrightarrow G$ an epimorphism, then $\Sigma^1(G_0) \cap p^*(S(G)) \subset p^*(\Sigma^1(G))$. In particular:

$$\Sigma^1(G) \cup (-\Sigma^1(G)) \subsetneq S(G) \implies \Sigma^1(G_0) \cup (-\Sigma^1(G_0)) \subsetneq S(G_0).$$

On the proof. For (i) see [Stre, Theorem A2.3].

For (ii), see [BiNS–87, Theorem A] and [Stre, Theorem A3.3].

(iii) If D^1G is finitely generated, then $\Sigma^1(G) = S(G)$, [Stre, Proposition A2.6]. For the “iff”, see [BiNS–87, Theorem B1].

For (iv), see [BiSt–80, Theorem A] and [Stre, Subsection B3.2c].

For (v), see [BiNS–87, Theorem C] and [Stre, Theorem A5.1].

Claim (vi) is rather straightforward from the definitions [Stre, Proposition A4.5]. \square

Example C.3. (i) $\Sigma^1(G) = S(G) \approx \mathbf{S}^{n-1}$ for G a finitely generated abelian group of torsion-free rank n .

(ii) $\Sigma^1(F_n) = \emptyset \subset S(F_n) \approx \mathbf{S}^{n-1}$ for the non-abelian free group F_n of rank $n \geq 2$; see [Stre, Item A2.1a].

(iii) For the soluble Baumslag-Solitar group $\text{BS}(1, n)$, the invariant $\Sigma^1(\text{BS}(1, n))$ is one of the two points of the sphere $S(\text{BS}(1, n)) \approx \mathbf{S}^0$; the argument of [Stre, Item A2.1a] for $n = 2$ carries over to $n \geq 2$.

(iv) For two coprime integers $\ell, m \geq 2$, we have $\Sigma^1(\text{Met}(\ell, m)) = \emptyset \subset \mathbf{S}^0$ [Stre, Item A3.6].

(v) If G is a semi-direct product $H \rtimes \mathbf{Z}$ of an infinite locally finite group H by an infinite cyclic group, then $\Sigma^1(G) = \emptyset \subset \mathbf{S}^0$ [Stre, Lemma B3.1].

(vi) For any $n \geq 1$ and any rational polyhedral subset P of \mathbf{S}^{n-1} , there exists a finitely presented group G with $S(G) \approx \mathbf{S}^{n-1}$ and a homeomorphism $p^* : \mathbf{S}^{n-1} \rightarrow S(G)$ such that $\Sigma^1(G) = p^*(S(G) \setminus P)$.

Some comment is in order for (vi), cited here to show that the invariant $\Sigma^1(G)$ can be more complicated than those of Examples (i) to (v). For a finitely generated group G , a non-zero character $\chi \in \text{Hom}(G, \mathbf{R})$ is *rational* if $\chi(G)$ is an infinite cyclic subgroup of \mathbf{R} . Denote by $\text{Hom}_{\mathbf{Q}}^*(G, \mathbf{R})$ the set of non-zero rational characters on G , and by $S_{\mathbf{Q}}(G)$ its image in $S(G)$; then $S_{\mathbf{Q}}(G)$ is a dense subset of $S(G)$ [Stre, Lemma B3.3].

Consider a positive integer n and the sphere $\mathbf{S}^{n-1} = S(\mathbf{Z}^n)$. A *rational hemisphere* of \mathbf{S}^{n-1} is the closure of the image of the half-space $\{\chi \in \text{Hom}_{\mathbf{Q}}^*(G, \mathbf{R}) \mid \chi(z) \geq 0\}$, for some $z \in \mathbf{Z}^n \setminus \{0\}$. A *rational convex polyhedral subset* of \mathbf{S}^{n-1} is a finite intersection of rational hemispheres. A *rational polyhedral subset* of \mathbf{S}^{n-1} is a finite union of rational convex polyhedral subsets.

Given an integer $n \geq 1$ and a rational polyhedral subset $P \subset \mathbf{S}^{n-1}$, there is a finitely presented group G and an epimorphism $p : G \rightarrow \mathbf{Z}^n$ such that $p^* : S(\mathbf{Z}^n) \rightarrow S(G)$ is a homeomorphism, and $\Sigma^1(G) = p^*(S(\mathbf{Z}^n) \setminus P)$. See [BiNS–87, Corollary 7.6] and [BiSt, Chapter IV, Section 1.1].

Corollary C.4. *Let G be a finitely generated group and E a finitely presented cover of G . If $\Sigma^1(G) \cup (-\Sigma^1(G)) \subsetneq S(G)$, then E contains non-abelian free subgroups. This holds in particular:*

- when G is metabelian and infinitely presented,*
- when $G = H \rtimes \mathbf{Z}$ with H infinite locally finite, as in Example C.3.v.*

Proof. This is straightforward from Claims (iii) to (v) of Theorem C.2 and from Example C.3.v. \square

In Corollary C.4, the claim concerning metabelian groups is precisely Theorem 1.5.

Example C.5 (B.H. Neumann). Let $\mathcal{V} = \{2 \leq v_1 < v_2 < v_3 < \dots\}$ be an infinite increasing sequence of integers. Set

$$X_{\mathcal{V}} = \{(i, j) \in \mathbf{Z}^2 \mid i \geq 1, -v_i \leq j \leq v_i\}.$$

For each $i \geq 1$, denote by $X_{\mathcal{V},i}$ the subset $\{(i, j) \in X_{\mathcal{V}} \mid -v_i \leq j \leq v_i\}$, of cardinal $2v_i + 1$. Define two permutations $\alpha_{\mathcal{V}}, \beta_{\mathcal{V}}$ of the set $X_{\mathcal{V}}$ as follows: for each $i \geq 1$, they preserve $X_{\mathcal{V},i}$, and

- ($\alpha_{\mathcal{V}}$) the restriction of $\alpha_{\mathcal{V}}$ to $X_{\mathcal{V},i}$ is the $(2v_i + 1)$ -cycle $((i, -v_i), (i, -v_i + 1), \dots, (i, v_i))$,

$(\beta_{\mathcal{V}})$ the restriction of $\beta_{\mathcal{V}}$ to $X_{\mathcal{V},i}$ is the 3-cycle $((i, -1), (i, 0), (i, 1))$.

The **Neumann group** corresponding to \mathcal{V} is the group $G_{\mathcal{V}}$ of permutations of $X_{\mathcal{V}}$ generated by $\alpha_{\mathcal{V}}$ and $\beta_{\mathcal{V}}$.

Let $\text{Alt}_f(\mathbf{Z})$ denote the group of permutations of finite supports of \mathbf{Z} that are even on their support. Let $\text{Alt}_f(\mathbf{Z}) \rtimes_{\text{shift}} \mathbf{Z}$ denote its semi-direct product with \mathbf{Z} , where \mathbf{Z} acts on itself by shifts, the generator 1 acting by $\alpha : j \mapsto j + 1$. Observe that $\text{Alt}_f(\mathbf{Z}) \rtimes_{\text{shift}} \mathbf{Z}$ is generated by α and the 3-cycle $\beta = (-1, 0, 1)$. It is easy to check that the assignment $\alpha_{\mathcal{V}} \mapsto \alpha$, $\beta_{\mathcal{V}} \mapsto \beta$ extends to an epimorphism $\pi_{\mathcal{V}} : G_{\mathcal{V}} \twoheadrightarrow \text{Alt}_f(\mathbf{Z}) \rtimes_{\text{shift}} \mathbf{Z}$. Neumann has shown that the kernel of $\pi_{\mathcal{V}}$ is the restricted product $\prod'_{i=1}^{\infty} \text{Alt}(2v_i + 1)$; see [Neum–37], as well as [Harp–00, Complement III.35].

This has two straightforward consequences. On the one hand, any minimal finite normal subgroup of $G_{\mathcal{V}}$ is one of the $\text{Alt}(2v_i + 1)$. Thus, for two distinct sequences \mathcal{V} and \mathcal{V}' , the groups $G_{\mathcal{V}}$ and $G_{\mathcal{V}'}$ are not isomorphic. In particular, there are uncountably many pairwise non-isomorphic 2-generator groups; hence these are infinitely presented, except possibly for a countable number of them (but see below). On the other hand, the groups $G_{\mathcal{V}}$ are elementary amenable.

Corollary C.6. *Let \mathcal{V} be a sequence of integers as in Example C.5. Any finitely presented cover of $G_{\mathcal{V}}$ contains non-abelian free subgroups. In particular $G_{\mathcal{V}}$ is not finitely presented.*

Proof. The claim of Corollary C.4 concerning $H \rtimes \mathbf{Z}$ applies to the group $\text{Alt}_f(\mathbf{Z}) \rtimes_{\text{shift}} \mathbf{Z}$. Since any cover of $G_{\mathcal{V}}$ is a cover of $\text{Alt}_f(\mathbf{Z}) \rtimes_{\text{shift}} \mathbf{Z}$, Corollary C.6 follows from Corollary C.4. \square

Let us also indicate how the last statement of Corollary C.6 is a straightforward consequence of the first paper [BiSt–78] on BNS-invariants.

Proof that $G_{\mathcal{V}}$ is not finitely presented. It follows from the definition of $G_{\mathcal{V}}$ that the kernel $N_{\mathcal{V}}$ of the composition $G_{\mathcal{V}} \twoheadrightarrow \text{Alt}_f(\mathbf{Z}) \rtimes_{\text{shift}} \mathbf{Z} \twoheadrightarrow \mathbf{Z}$ is locally finite. Suppose (ab absurdo) that $G_{\mathcal{V}}$ is finitely presented. By [BiSt–78, Theorem A], we have $G_{\mathcal{V}} = \text{HNN}(H, \varphi : K \xrightarrow{\sim} L)$ for a finitely generated subgroup H of $N_{\mathcal{V}}$ and an isomorphism φ between two subgroups K, L of H . Since non-ascending extensions contain non-abelian free subgroups (by Britton’s lemma), $G_{\mathcal{V}}$ is an ascending HNN-extension; we may assume that $K = H$. Since $N_{\mathcal{V}}$ is locally finite, the finitely generated subgroup H of $N_{\mathcal{V}}$ is finite. It follows that $K = H = L$. Hence $G_{\mathcal{V}} = H \rtimes_{\varphi} \mathbf{Z}$ and the kernel of $G_{\mathcal{V}} \twoheadrightarrow \mathbf{Z}$ is finite. This is preposterous, and the proof is complete. \square

APPENDIX D. **On growth and amenability**

Let G be a group generated by a finite set S . For an integer $n \geq 0$, let $B_S^G(n)$ denote the “ball of radius n around the origin”, namely the set of those elements $g \in G$ that can be written as words $g = s_1 \cdots s_n$, with $s_1, \dots, s_n \in S \cup S^{-1} \cup \{1\}$. Let $\gamma_S^G(n)$ denote the cardinality of $B_S^G(n)$. Then G is said to be

- (pol) of **polynomial growth** if there exist constants $a, d > 0$ such that $\gamma_S^G(n) \leq an^d$ for all $n \geq 1$,
- (exp) of **exponential growth** if there exist a constant $c > 1$ such that $\gamma_S^G(n) \geq c^n$ for all $n \geq 0$,
- (int) of **intermediate growth** in other cases.

It is easy to check that this trichotomy depends only on G , not on the finite generating set S . For information on the growth of groups, we refer to the books [Harp–00] and [Mann–11].

A group G is **amenable** if there exists a left-invariant finitely additive probability measure defined on all subsets of G (there are many other equivalent definitions). Two basic results are important here: (i) amenability of groups is preserved by the four operations of taking subgroups, quotients, direct limits, and extensions with amenable kernels (already in [vNeu–29]), and (ii) groups of intermediate growth are amenable (this goes back to [AdSr–57], and is also a straightforward consequence of Følner’s Criterion [Foln–55]). These results make it natural to define three classes of groups:

\mathcal{AG} is the **class of amenable groups**, defined in [vNeu–29].

\mathcal{EG} is the **class of elementary amenable groups**, defined in [Day–57]; it is the smallest class of groups containing the easiest examples, that are finite groups and abelian groups, and stable by the four operations listed above.

\mathcal{SG} is the **class of subexponentially amenable groups** (see below for an historical comment on this definition); it is the smallest class of groups containing \mathcal{EG} and the next easiest examples, that are the groups of intermediate growth.

We have a partition

$$\mathcal{AG} = (\mathcal{AG} \setminus \mathcal{SG}) \sqcup (\mathcal{SG} \setminus \mathcal{EG}) \sqcup \mathcal{EG}.$$

Let us mention a few groups in each of these three parts.

The class \mathcal{EG} contains all virtually soluble groups and all locally finite groups; other examples are cited in Subsection 1.A.

There are finitely generated groups in \mathcal{EG} which are not virtually soluble: for example all Neumann groups G_ν discussed in Example

C.5, or an example in [Hill–91]. As already mentioned in Subsection 1.D, any countable elementary amenable group embeds in a finitely generated elementary amenable group; the same hold for “amenable” instead of “elementary amenable” [OlOs, Corollary 1.3].

Let us describe a family of finitely presented elementary amenable groups that are not virtually soluble.

Example D.1. Consider an integer $n \geq 1$, the set

$$S_n = \{(j, k) \in \mathbf{Z}^2 \mid j \geq 0, 1 \leq k \leq n\}$$

of n parallel half-intervals in the square lattice, and the **Houghton group** H_n of all permutations h of S_n such that, for each $k \in \{1, \dots, n\}$, there exists a translation $t_k \in \mathbf{Z}$ such that $h(j, k) = (j + t_k, k)$ for all j large enough [Houg–78]. Denote by $\text{Sym}_f(S_n)$ the group of permutations of S_n with finite supports, clearly a normal subgroup of H_n , and set

$$A = \{(t_1, \dots, t_n) \in \mathbf{Z}^n \mid \sum_{k=1}^n t_k = 0\}.$$

We have a short exact sequence

$$1 \longrightarrow \text{Sym}_f(S_n) \longrightarrow H_n \xrightarrow{\pi} A \longrightarrow 1$$

where, with the notation above, $\pi(h) = (t_1, \dots, t_n)$. Since $\text{Sym}_f(S_n)$ is locally finite and A abelian, H_n is elementary amenable.

It is known that the group H_n is of type FP_{n-1} but not FP_n [Brow–87, Theorem 5.1]. In particular, for $n \geq 3$, the group H_n is finitely presented.

Finitely generated groups in the class \mathcal{EG} are either of polynomial growth or of exponential growth [Chou–80]; this has been sharpened: a finitely generated group in the class \mathcal{EG} has either polynomial growth or *uniform* exponential growth [Osin–04] (see also [Breu–07]). By a famous theorem of Gromov [Grom–81], a finitely generated group of polynomial growth is virtually nilpotent, and in particular finitely presented.

The class $\mathcal{SG} \setminus \mathcal{EG}$ contains the class of finitely generated groups of intermediate growth. Historically, the group \mathfrak{G} of Theorem 1.6 and Example 2.16 was the first group shown to be of intermediate growth [Grig–83]. This class also contains finitely presented groups, such as the group with 5 generators and 11 relators of [Grig–98], later shown to have another presentation with 2 generators and 4 relators (due to Bartholdi, see [CeGH–99, Number 12]).

The history of early papers on the class \mathcal{SG} is worth a few lines. This class was first implicitly introduced in a paper on 4-manifold topology, more precisely on 4-manifold surgery and 5-dimensional s-cobordism theorems, [FrTe–95] (see also [KrQu–00]), and then explicitly in [Grig–98]. Freedman and Teichner introduce a class of groups that they call “good”, defined as the groups for which the “ π_1 -Null Disk Lemma” holds; this lemma establishes the existence of 2-discs bounding some closed curves in 4-manifolds of a certain kind. Good groups include finitely generated groups in the class \mathcal{SG} [FrTe–95, Theorem 0.1 and Lemma 1.2].

The class $\mathcal{AG} \setminus \mathcal{SG}$ contains the **Basilica group** \mathfrak{B} of Example 2.17, which was first shown to be not in \mathcal{SG} [GrZ–02a], and later shown to be amenable [BaVi–05]. The method of Bartholdi and Virag was streamlined and generalized by Kaimanovich in [Kaim–05], in terms of entropy and of the legendary “Münchhausen’s trick”. This and later papers show the amenability of \mathfrak{B} and of many other non elementary amenable groups (see [BaKN–10, AmAV], building among other things on [Sidk–00]). The class $\mathcal{AG} \setminus \mathcal{SG}$ contains also the finitely generated amenable simple groups that appear in [JuMo], and in Problem 1.12.

Non-amenable groups include non-abelian free groups, more generally groups containing non-abelian free subgroups [vNeu–29].

We conclude this report by Question D.2, due to Tullio Ceccherini-Silberstein. Before this, we recall some background.

A group G has a **paradoxical decomposition** if there exist integers $p, q \geq 2$, subsets $X_1, \dots, X_p, Y_1, \dots, Y_q$ of G , and elements $g_1, \dots, g_p, h_1, \dots, h_q$ in G such that

$$\begin{aligned} G &= X_1 \sqcup \dots \sqcup X_p \sqcup Y_1 \sqcup \dots \sqcup Y_q \\ &= g_1 X_1 \sqcup \dots \sqcup g_p X_p = h_1 Y_1 \sqcup \dots \sqcup h_q Y_q, \end{aligned}$$

where \sqcup denotes disjoint union. Tarski has shown that G is non-amenable if and only if G has a paradoxical decomposition [Tars–38]; see also [Wago–85] and [HaSk–86]. For G non-amenable, the **Tarski number** $\mathcal{T}(G)$ of G is the minimum of the sum $p+q$, over all paradoxical decompositions of G . When G has non-abelian free subgroups, it is easy to show that $\mathcal{T}(G) = 4$; in particular, if G is a non-elementary Gromov-hyperbolic group, then $\mathcal{T}(G) = 4$. As a student of Tarski in the 1940’s, Jonsson has shown that, conversely, $\mathcal{T}(G) = 4$ implies that G has non-abelian free subgroups. It is also easy to show that $\mathcal{T}(G) \geq 6$ for a non-amenable torsion group G . See [Wago–85] or [CeGH–99, Propositions 20 and 21]. We do not know any example of

a non-amenable group G without non-abelian free subgroups for which the exact value of $\mathcal{T}(G)$ has been computed.

Let $\mathcal{M}_m^{\text{na}}$ be the subspace of \mathcal{M}_m of those pairs (G, S) with G non-amenable. For $m \geq 2$, the Tarski number function

$$\mathcal{M}_m^{\text{na}} \longrightarrow \mathbf{N}, \quad (G, S) \longmapsto \mathcal{T}(G)$$

is not continuous. Indeed, there are sequences of non-elementary Gromov-hyperbolic groups (with Tarski number 4) converging in \mathcal{M}_m to non-amenable torsion groups (with Tarski number at least 6). See [Cham–00, Théorème 1.3], or one of the following classes of examples.

(1) For integers $m, n \geq 2$, let $B(m, n)$ denote the free Burnside group of rank m and exponent n , that is the quotient of the free group F_m by the set of relators $(w^n = 1)_{w \in F_m}$. For n odd and $n \geq 665$, Adyan has shown that $B(m, n)$ is non-amenable [Adya–82]; moreover, it is known that $6 \leq \mathcal{T}(B(m, n)) \leq 14$ [CeGH–99, Theorem 61].

Such groups are limits in \mathcal{M}_m of non-elementary Gromov-hyperbolic groups. More precisely, for n odd and large enough, the group $B(m, n)$ is a limit in \mathcal{M}_m of a sequence $(B(m, n, i))_{i \geq 1}$ of non-elementary Gromov-hyperbolic groups. A similar fact is shown by Ivanov in [Ivan–94, see Lemma 21.1], in the much more difficult case of n even, with $n \geq 2^{48}$ and n divisible by 2^9 ; his proof adapts to the case needed here, with important simplifications (compare with [GrIv–09], in particular Theorems 1.10 and 1.7, and recall that a finitely presented group with a subquadratic Dehn function is hyperbolic). In particular, for $m \geq 2$ and n odd large enough, the free Burnside group $B(m, n)$, of Tarski number between 6 and 14, is a limit in \mathcal{M}_m (indeed in $\mathcal{M}_m^{\text{na}}$) of groups of Tarski number 4.

(2) Ol’shanskii has worked out several “Tarski monster groups”. In particular, for any prime p large enough, he has constructed a 2-generated non-amenable torsion group $\text{TM}(p)$ in which any proper non-trivial subgroup is of order p . Moreover $\text{TM}(p)$, of Tarski number ≥ 6 , is a limit in \mathcal{M}_2 of non-elementary Gromov hyperbolic groups, of Tarski number 4 [Ol’sh–80, Lemma 10.7a].

In several other papers by Ol’shanskii and co-authors, there are several other classes of examples of such Burnside type limits of hyperbolic groups. Let us only quote one paper [OlOS–09], and the book [Ol’sh–91].

Question D.2. *For an integer $k \geq 4$ and a finitely generated non-amenable group G with $\mathcal{T}(G) = k$, does there exist a finitely presented cover E of G with $\mathcal{T}(E) = k$?*

The answer is clearly positive when $\mathcal{T}(G) = 4$ (with E free). If we define $\mathcal{T}(G) = \infty$ when G is amenable, the previous question for $k = \infty$ coincides with Question 1.1, and the answer is negative.

REFERENCES

- [Abel–77] Herbert ABELS, *An example of a finitely presented solvable group*, in “Homological group theory, Proc. Sympos. Durham, 1977”, London math. Soc. Lecture Note Ser. **36** (Cambridge Univ. Press 1979), 205–211.
- [AbBr–87] Herbert ABELS and Kenneth S. BROWN, *Finiteness properties of solvable S -arithmetic groups: an example*, J. Pure Appl. Algebra **44** (1987), 77–83.
- [AdSr–57] Georgy M. ADEL’SON-VEL’SII and Yulii A. SREIDER, *The Banach mean on groups*, Usphei Math. Nauk. (N.S.) **12**(6) (1957), 131–136.
- [Adya–82] Sergei Ivanovich ADYAN, *Random walks on free periodic groups*, Math. USSR Izv. **21**³ (1983), 425–434. [Russian: Izv. Adak. Nauk SSSR, Ser. Math. **46**⁶ (1982), 1139–1149.]
- [AmAV] Gideon AMIR, Omer ANGEL, and Bálint VIRÁG, *Amenability of linear-activity automaton groups*, Preprint, arXiv:0905.2007v2, 21 June 2011, J. European Math. Soc. to appear.
- [Bart–03] Laurent BARTHOLDI, *L -presentations and branch groups*, J. Algebra **268** (2003), 419–443.
- [Bart] Laurent BARTHOLDI, *(Self-)similar groups and the Farrell-Jones conjectures*, Preprint, arXiv:1107.5339v2, 28 Jul 2011.
- [BaGr–02] Laurent BARTHOLDI and Rostislav GRIGORCHUK *On parabolic subgroups and Hecke algebras of some fractal groups*, Serdica Math. J. **28** (2002), 47–90.
- [BaGN–03] Laurent BARTHOLDI, Rostislav GRIGORCHUK and Volodymyr NEKRASHEVYCH, *From fractal groups to fractal sets*, in “Fractal in Graz 2001”, Birkhäuser (2003), 25–118.
- [BaVi–05] Laurent BARTHOLDI and Bálint VIRÁG, *Amenability via random walks*, Duke Math. J. **130** (2005), 39–56.
- [BaCo–06] Laurent BARTHOLDI and Yves DE CORNULIER, *Infinite groups with large balls of torsion elements and small entropy*, Arch. Math. Basel **87** (2006), 104–112.
- [BaPo–09] Laurent BARTHOLDI and Floriane POCHON, *On growth of torsion groups*, Groups Geom. Dyn. **3** (2009), 525–539.
- [BaKN–10] Laurent BARTHOLDI, Vadim KAIMANOVICH and Volodymyr NEKRASHEVYCH, *On amenability of automata groups*, Duke Math. J. **154** (2010), 575–598.
- [BaEr–a] Laurent BARTHOLDI and Anna ERSCHLER, *Growth of permutational extensions*, Inventiones Math. (to appear).
- [BaEr–b] Laurent BARTHOLDI and Anna ERSCHLER, *Groups of given intermediate word growth*, Preprint, arXiv:1110.3650v2, 15 Nov 2011.
- [Baum–61] Gilbert BAUMSLAG, *Wreath products and finitely presented groups*, Math. Z. **75** (1961), 22–28.

- [Baum–63] Gilbert BAUMSLAG *Some subgroup theorems for free v -groups*, Trans. AMS **108** (1963), 516–525.
- [Baum–72] Gilbert BAUMSLAG, *A finitely presented metabelian group with a free abelian derived group of infinite rank*, Proc. Amer. Math. Soc. **35** (1972), 61–62.
- [Baum–73] Gilbert BAUMSLAG, *Subgroups of finitely presented metabelian groups*, J. Australian Math. Soc. **16** (1973), 98–110.
- [Baum–74] Gilbert BAUMSLAG, *Finitely presented metabelian groups*, in “Proc. Conf. Theory of Groups, Canberra, 1973”, Springer Lecture Notes in Math. **372** (1974), 65–74.
- [Baum–93] Gilbert BAUMSLAG, *Topics in combinatorial group theory*, Lectures in Mathematics, ETH Zürich, Birkhäuser, 1993.
- [BaSo–62] Gilbert BAUMSLAG and Donald SOLITAR, *Some two-generator one-relator non-Hopfian groups*, Bull. Amer. Math. Soc. **68** (1962), 199–201.
- [BaSt–76] Gilbert BAUMSLAG and Ralph STREBEL, *Some finitely generated, infinitely related metabelian groups with trivial multiplier*, J. of Algebra **40** (1976), 46–62.
- [BaGS–86] Gilbert BAUMSLAG, Dion GILDENHUYS, and Ralph STREBEL, *Algorithmically insoluble problems about finitely presented solvable groups, Lie and associative algebras. I*, J. Pure Appl. Algebra **39** (1986), 53–94.
- [BaCR–94] Gilbert BAUMSLAG, Frank B. CANNONITO, and Derek J.S. ROBINSON, *The algorithmic theory of finitely generated metabelian groups*, Trans. Amer. Math. Soc. **344** (1994), 629–648.
- [BaMi–09] Gilbert BAUMSLAG and Charles F. MILLER, III, *Reflections on some groups of B.H. Neumann*, J. Group Theory **12** (2009), 771–781.
- [BeBr–97] Mladen BESTVINA and Noel BRADY, *Morse theory and finiteness properties of groups*, Invent. Math. **129** (1997), 445–470.
- [Bier–76] Robert BIERI, *Homological dimension of discrete groups*, Queen Mary College Mathematical Notes, 1976.
- [Bier–79] Robert BIERI, *Finitely presented soluble groups*, Séminaire d’algèbre Paul Dubreuil, 1977–78, Springer Lecture Notes in Math. **740** (1979), 1–8.
- [BiSt–78] Robert BIERI and Ralph STREBEL, *Almost finitely presented soluble groups*, Comment. Math. Helv. **53** (1978), 258–278.
- [BiSt–80] Robert BIERI and Ralph STREBEL, *Valuations and finitely presented metabelian groups*, Proc. London Math. Soc. (3) **41** (1980), 439–464.
- [BiNS–87] Robert BIERI, Walter NEUMANN, and Ralph STREBEL, *A geometric invariant of discrete groups*, Invent. Math. **90** (1987), 451–477.
- [BiRe–88] Robert BIERI and Burkhardt RENZ, *Valuations on free resolutions and higher geometric invariants of groups*, Comment. Math. Helv. **63** (1988), 464–497.
- [BiSt] Robert BIERI and Ralph STREBEL, *Geometric invariants for discrete groups*, unpublished book, 1992 (see [Stre]).
- [BCGS] Robert BIERI, Yves DE CORNULIER, Luc GUYOT, and Ralph STREBEL, *Infinite presentability of groups and condensations*, Preprint, arXiv:1010.0271v2, 17 May 2012.
- [Bour–63] Nicolas BOURBAKI, *Intégration, chapitres 7 et 8*, Hermann, 1963.

- [Breu–07] Emmanuel BREUILLARD, *On uniform exponential growth for solvable groups*, Pure Appl. Math. Q. **3** (2007), 949–967.
- [Brow–82] Kenneth S. BROWN, *Cohomology of groups*, Springer, 1982.
- [Brow–87] Kenneth S. BROWN, *Finiteness properties of groups*, J. Pure Appl. Algebra **44** (1987), 45–75.
- [BuPe–06] Kai-Uwe BUX and Rodrigo PÉREZ, *On the growth of iterated monodromy groups*, in “Topological and asymptotic aspects of group theory”, Contemp. Math. **394** (Amer. Math. Soc. 2006), 61–76.
- [CaRo–84] Frank B. CANNONITO and Derek J.S. ROBINSON, *The word problem for finitely generated soluble groups of finite rank*, Bull. London Math. Soc. **16** (1984), 43–46.
- [CaCo] Pierre-Emmanuel CAPRACE and Yves DE CORNULIER, *On embeddings into compactly generated groups*, Preprint, 17 September 2012.
- [CeCo–10] Tullio CECCHERINI-SILBERSTEIN and Michel COORNAERT, *Cellular automata and groups*, Springer, 2010.
- [CeGH–99] Tullio CECCHERINI-SILBERSTEIN, Rostislav GRIGORCHUK, and Pierre DE LA HARPE, *Amenability and paradoxical decompositions for pseudogroups and for discrete metric spaces*, Proc. Steklov Inst. Math. **224** (1999), 57–97.
- [Chab–50] Claude CHABAUTY, *Limite d’ensembles et géométrie des nombres*, Bull. Soc. Math. France **78** (1950), 143–151.
- [Cham–00] Christophe CHAMPETIER, *L’espace des groupes de type fini*, Topology **39** (2000), 657–680.
- [ChGu–05] Christophe CHAMPETIER and Vincent GUIRADEL, *Limit groups as limits of free groups: compactifying the set of free groups*, Isarel J. Math. **146** (2005), 1–75.
- [Chou–80] Christophe CHOU, *Elementary amenable groups*, Illinois J. Math. **1** (1980), 396–407.
- [Clap–67] Christopher Robert Jasper CLAPHAM, *Finitely presented groups with word problems of arbitrary degrees of insolubility*, Proc. London Math. Soc. **14** (1964), 633–676.
- [Coll–78] Donald J. COLLINS, *Some one-relator Hopfian groups*, Trans. Amer. Math. Soc. **235** (1978), 363–374.
- [CoLe–83] Donald J. COLLINS and Frank LEVIN, *Automorphisms and Hopficity of certain Baumslag-Solitar groups*, Arch. Math. **40** (1983), 385–400.
- [Corn–06] Yves DE CORNULIER, *Finitely presented wreath products and double coset decompositions*, Geom. Dedicata **122** (2006), 89–108.
- [CoGP–07] Yves DE CORNULIER, Luc GUYOT, and Wolfgang PITSCH, *On the isolated points in the space of groups*, Groups Geom. Dyn. **1** (2007), 47–59.
- [CoKa–11] Yves DE CORNULIER and Aditi KAR, *On Property (FA) for wreath products*, J. Group Theory **14** (2011), 165–174.
- [CoTe] Yves DE CORNULIER and Romain TESSERA, *Dehn function and asymptotic cones of Abels’ group*, Preprint, arXiv:1203.4696v1, 21 Mar 2012.
- [Day–57] Mahlon M. DAY, *Amenable semigroups*, Illinois J. Math. **1** (1957), 509–544.
- [DuMi] Martin J. DUNWOODY and Ashot MINASYAN, *An (FA)-group that is not (FR)*, Preprint, arXiv:1203.3317v1, 15 Mar 2012.

- [EdPr-84] Martin EDJVET and Stephen J. PRIDE, *The concept of "largeness" in group theory II*, in A.C. Kim and B.H. Neumann (Eds.), "Groups-Korea 1983", Lecture Notes in Math. **1098** (Springer 1984) 29–54.
- [Ersc-04] Anna ERSCHLER, *Non residually finite groups of intermediate growth, commensurability and non-geometricity*, J. of Algebra **222** (2004), 154–172.
- [Ersc-07] Anna ERSCHLER, *Automatically presented groups*, Groups Geom. Dyn. **1** (2007), 47–59.
- [FaGu-85] Jacek FABRYKOWSKI and Narain GUPTA, *On groups with sub-exponential growth functions*, J. Indian Math. Soc. **49** (1985), 249–256.
- [FaGu-91] Jacek FABRYKOWSKI and Narain GUPTA, *On groups with sub-exponential growth functions. II*, J. Indian Math. Soc. **56** (1991), 217–228.
- [Foln-55] Erling FØLNER, *On groups with full Banach mean value*, Math. Scand. **3** (1955), 243–254.
- [FrTe-95] Michael H. FREEDMAN and Peter TEICHNER, *4-manifold topology I: subexponential groups*, Inventiones Math. **122** (1995), 500–529. See the Appendix in [KrQu-00].
- [Gild-79] Don GILDENHUYS, *Classification of soluble groups of cohomological dimension two*, Math. Z. **166** (1979), 21–25.
- [Grig-80] Rostislav GRIGORCHUK, *On Burnside's problem on periodic groups*, Funktsional. Anal. i Prilozhen, **14** (1980), 53–54.
- [Grig-83] Rostislav GRIGORCHUK, *On the Milnor problem of growth*, Dokl. Acad. Nauk SSSR **271** (1983), 30–33.
- [Gri-84a] Rostislav GRIGORCHUK, *Degrees of growth of finitely generated groups and the theory of invariant means*, Izv. Akad. Nauk SSSR Ser. Mat. **48**(5) (1984), 939–985. [In English Math. USSR Izv. **85** (1985) 259–300.]
- [Gri-84b] Rostislav GRIGORCHUK, *Construction of p -groups of intermediate growth that have a continuum of quotient groups*, Algebra Logic **23** (1984), 265–273.
- [Grig-98] Rostislav GRIGORCHUK, *An example of a finitely presented amenable group that does not belong to the class EG*, Mat. Sb. **189** (1998), 79–100.
- [Grig-99] Rostislav GRIGORCHUK, *On the system of defining relations and the Schur multiplier of periodic groups generated by finite automata*, in "Groups St Andrews 1997 in Bath, 1", C.M. Campbell et al. Editors, London Math. Soc. Lecture Note Ser. **260** (1999), 290–317.
- [Grig-00] Rostislav GRIGORCHUK, *Just infinite branch groups*, in "New horizons in pro- p -groups", M. DU SAUTOY et al. Editors, Birkhäuser (2000), 121–179.
- [Grig-05] Rostislav GRIGORCHUK, *Solved and unsolved problems around one group*, in "Infinite groups: geometric, combinatorial and dynamical aspects", L. Bartholdi, T. Ceccherini-Silberstein, T. Smirnova-Nagnibeda, and A. Żuk Editors, Progr. in Math **248**, Birkhäuser (2005), 117–218.

- [Grig-06] Rostislav GRIGORCHUK, *Hanoi Towers group on 3 pegs and its pro-finite closure, joint work with Volodymyr NEKRASHEVYCH and Zoran ŠUNIĆ*, Mathematisches Forschungsinstitut Oberwolfach Report 25/2006, Pages 1477–1479 [see also Report 19/2006, Pages 1179–1182].
- [Grig] Rostislav GRIGORCHUK, *Milnor’s Problem on the growth of groups and its consequences*, Preprint, arXiv:1111.0512v2, 31 Jan 2012.
- [GrMa-97] Rostislav GRIGORCHUK and Mohammad J. MAMAGHANI, *On use of iterates of endomorphisms for constructing groups with specific properties*, Mat. Stud. **8** (1997), 198–206, 238.
- [GLSZ-00] Rostislav GRIGORCHUK, Peter LINNELL, Thomas SCHICK, and Andrzej ŻUK, *On a question of Atiyah*, C.R. Acad. Sci., Paris, Sér. I, Math. **331**⁹ (2000), 663–668.
- [GrHa-01] Rostislav GRIGORCHUK and Pierre DE LA HARPE, *Limit behaviour of exponential growth rates for finitely generated groups*, in “Essays on geometry and related topics, Mémoires dédiés à André Haefliger, Vol. 2”, Monogr. Enseign. Math. **38** (2001), 351–370.
- [GrZu-01] Rostislav GRIGORCHUK and Andrzej ŻUK, *The lamplighter group as a group generated by a 2-state automaton, and its spectrum*, Geom. Dedicata **87** (2001), 209–244.
- [GrZ-02a] Rostislav GRIGORCHUK and Andrzej ŻUK, *On a torsion-free weakly branch group defined by a three state automaton*, Internat. J. Algebra Comput. **12** (2002), 223–246.
- [GrZ-02b] Rostislav GRIGORCHUK and Andrzej ŻUK, *Spectral properties of a torsion-free weakly branch group defined by a three state automaton*, in “Computational and statistical group theory (Las Vegas, 2001)”, Contemp. Math. **298** (Amer. Math. Soc. 2002), 57–82.
- [GrSu-06] Rostislav GRIGORCHUK and Zoran ŠUNIĆ, *Asymptotic aspects of Schreier graphs and Hanoi Towers groups*, C. R. Math. Acad. Sci. Paris **342**⁸ (2006), 545–550.
- [GrSu-07] Rostislav GRIGORCHUK and Zoran ŠUNIĆ, *Self-similarity and branching in group theory*, in “Groups St. Andrews 2005, Vol. 1”, C.M. Campbell et al. Editors, London Math. Soc. Lecture Note Ser. **339** (2007), 36–95.
- [GrSS-07] Rostislav GRIGORCHUK, Dmytro SAVCHUK, and Zoran ŠUNIĆ, *The spectral problem, substitutions and iterated monodromy*, in “Probability and mathematical physics”, CRM Proc. Lecture Notes **42** (Amer. Math. Soc. 2007), 225–248.
- [GrIv-09] Rostislav GRIGORCHUK and Sergei V. IVANOV, *On Dehn functions of infinite presentations of groups*, Geom. Funct. Anal. **18** (2009), 1841–1874.
- [GrMe] Rostislav GRIGORCHUK and Konstantin MEDYNETS, *Topological full groups are locally embeddable into finite groups*, Preprint, arXiv:1105.0719v3, 23Jan2012.
- [Grom-81] Michael GROMOV, *Groups of polynomial growth and expanding maps (with an appendix by Jacques Tits)*, Publ. math. I.H.É.S. **53** (1981) 53–78.
- [GuSi-83] Narain GUPTA and Said SIDKI, *On the Burnside problem for periodic groups*, Math. Z. **182** (1983), 385–388.

- [Hall-50] Marshall HALL, *A topology for free groups and related groups*, Annals of Math. **52** (1950), 127–139.
- [Hall-54] Philip HALL, *Finiteness conditions for soluble groups*, Proc. London Math. Soc. **4** (1954), 419–436. Collected Works 305–324.
- [Hall-59] Philip HALL, *On the finiteness of certain soluble groups*, Proc. London Math. Soc. **9** (1959), 595–622. Collected Works 515–544.
- [Hall-61] Philip HALL, *The Frattini subgroups of finitely generated groups*, Proc. London Math. Soc. **11** (1961), 327–352. Collected Works 581–608.
- [Harp-00] Pierre de la HARPE, *Topics in geometric group theory*, Chicago Univ. Press, 2000.
- [HaSk-86] Pierre DE LA HARPE and Georges SKANDALIS, *Un résultat de Tarski sur les actions moyennables de groupes et les partitions paradoxales*, Enseign. Math. **32** (1986), 121–138.
- [Higm-61] Graham HIGMAN, *Subgroups of finitely presented groups*, Proc. Roy. Soc. London Ser. A **262** (1961), 455–475.
- [HiNN-49] Graham HIGMAN, Bernhard Hermann NEUMANN, and Hanna NEUMANN, *Embedding theorems for groups*, J. London Math. Soc. **24** (1949), 247–254.
- [Hill-91] Jonathan A. HILLMAN, *Elementary amenable groups and 4-manifolds with Euler characteristic 0*, J. Austral. Math. Soc. (Series A) **50** (1991), 160–170.
- [Houg-78] Chris H. HOUGHTON, *The first cohomology of a group with permutation module coefficients*, Arch. Math **31** (1978) 254–258.
- [Ivan-94] Sergei V. IVANOV, *The free Burnside groups of sufficiently large exponents*, Int. J. Algebra Comput. **4**, No.1-2 (1994), 1–308.
- [JuMo] Kate JUSCHENKO and Nicolas MONOD, *Cantor systems, piecewise translations and simple amenable groups*, Preprint, arXiv:1204.2132v3, 29 Apr 2012.
- [Kaim-05] Vadim KAIMANOVICH, *“Münchhausen trick” and amenability of self-similar groups*, Int. J. Algebra Comput. **15** (2005), 907–937.
- [Khar-81] Olga KHARLAMPOVICH, *A finitely presented solvable group with unsolvable word problem*, Izv. Akad. Nauk SSSR Ser. Math. **45** (1981), 852–873, 928. [English translation: Math. USSR Izvestija **19** (1982), 151–169.]
- [Khar-90] Olga KHARLAMPOVICH, *The word problem for groups and Lie algebras*, Doctor’s Thesis (Russian), Moscow Steklov Mathematical Institute, 1990.
- [KhMS] Olga KHARLAMPOVICH, Alexei MYASNIKOV, and Mark SAPIR, *Algorithmically complex residually finite groups*, Preprint, arXiv:1204.6506v4, 4 Oct 2012.
- [Kour-06] *The Kourovka notebook: unsolved problems in group theory*, 16th Edition. Russian Academy of Sciences, Siberian Division, 2006.
- [KrQu-00] Vyacheslav S. KRUSKHAL and Frank QUINN, *Subexponential groups in 4-manifold topology*, Geometry & Topology **4** (2000), 407–430. With an appendix by Michael H. FREEDMAN and Peter TEICHNER.
- [LeRo-04] John C. LENNOX and Derek J.S. ROBINSON, *The theory of infinite soluble groups*, Clarendon Press, 2004.

- [Lyse–85] Igor LYSËNOK, *A set of defining relations for the Grigorchuk group*, Math. Zametki **38** (1985), 503–516, 636. English translation: Math. Notes Acad. Sc. USSR **38**:4 (1985), 784–792.
- [Lyul–84] Natalia Albertovna LYUL’KO, *Normal subgroups of Abels’ groups*, Math. Notes **36** (1984), 639–641.
- [Magn–39] Wilhelm MAGNUS, *On a theorem of Marshall Hall*, Annals of Math. **40** (1939), 764–768.
- [Mal’c–40] Anatoly Ivanovich MAL’CEV, *On the faithful representation of infinite groups by matrices*, Amer. Math. Soc. Transl. Ser. 2 **45** (1962), 1–18. [Russian: Rec. Math. [Mat. Sbornik] N.S., **8(50)**:3 (1940), 405–422.]
- [Mand–80] Benoît B. MANDELBROT, *Fractal aspects of the iteration of $z \rightarrow \Lambda z(1-z)$ for complex Λ and z* , Annals New York Acad. Sciences **357** (1980), 249–259.
- [Mann–11] Avinoam MANN, *How groups grow*, Cambridge Univ. Press, 2011.
- [Matu–06] Hiroki MATUI, *Some remarks on topological full groups of Cantor minimal systems*, Internat. J. Math. **17** (2006), 231–251.
- [Meld–95] John D.P. MELDRUM, *Wreath products of groups and semigroups*, Pitman Monograph, Longman, 2005.
- [Mesk–74] Stephen MESKIN, *A finitely generated residually finite group with an unsolvable word problem*, Proc. Amer. Math. Soc. **43** (1974), 8–10.
- [Mill–89] Charles F. MILLER, III, *Decision problems for groups – survey and reflections*, in “Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989)” (Springer 1991), 1–59.
- [Nekr–05] Volodymyr NEKRASHEVYCH, *Self-similar groups*, Mathematical Surveys and Monographs **117**, American Mathematical Society 2005.
- [Nekr–10] Volodymyr NEKRASHEVYCH, *Free subgroups in groups acting on rooted trees*, Groups, Geometry, and Dynamics **4** (2010), 847–862.
- [Nekr–11] Volodymyr NEKRASHEVYCH, *Iterated monodromy groups*, in “Groups St Andrews 2009 in Bath”, Volume 1, C.M. Campbell et al. Editors, London Math. Soc. Lecture Note Ser. **387** (2011), 41–93.
- [Neum–37] Bernhard Hermann NEUMANN, *Some remarks on infinite groups*, J. London Math. Soc. **12** (1937), 120–127.
- [NeNe–59] Bernhard Hermann NEUMANN and Hanna NEUMANN, *Embedding theorems for groups*, J. London Math. Soc. **34** (1959), 465–479.
- [vNeu–29] John VON NEUMANN, *Zur allgemeinen Theorie des Masses*, Fund. Math. **13** (1929) 73–116, 333 (= Collected works, Vol. I, 599–643).
- [Ol’sh–80] Alexander Yu. OL’SHANSKII, *An infinite group with subgroups of prime orders* Math. USSR, Izv. **16** (1981) 279–289. Izv. Akad. Nauk SSSR, Ser. Mat. **44** (1980), 309–321.]
- [Ol’sh–91] Alexander Yu. OL’SHANSKII, *Geometry of defining relations in groups*, Kluwer, 1991.
- [OlSa–02] Alexander Yu. OL’SHANSKII and Mark SAPIR, *Non-amenable finitely presented torsion-by-cyclic groups*, Publ. Math. I.H.É.S. **96** (2002), 43–169.
- [OlOS–09] Alexander Yu. OL’SHANSKII, Denis OSIN, and Mark SAPIR, with an appendix by Michael KAPOVICH and Bruce KLEINER, *Lacunary hyperbolic groups*, Geom. Topol. **13** (2009), 2052–2140.

- [OlOs] Alexander Yu. OL'SHANSKII and Denis OSIN, *A quasi-isometric embedding theorem for groups*, Preprint, arXiv:1202.6437v2, 2 Mar 2012.
- [Osin-04] Denis OSIN, *Algebraic entropy of elementary amenable groups*, *Geom. Dedicata* **104** (2004), 133–151.
- [Reme-72] Vladimir Nikanorovich REMESLENNIKOV, *A finitely presented soluble group without maximum condition for normal subgroups*, *Mat. Zametki* **12** (1972), 287–293 [Russian].
- [Reme-73] Vladimir Nikanorovich REMESLENNIKOV, *On finitely presented soluble groups*, *Proc. Fourth All-Union Symposium on the Theory of Groups (Novosibirsk, 1973)*, 164–169 [Russian].
- [Robi-96] Derek J.S. ROBINSON, *A course in the theory of groups*, Second Edition, *Graduate Texts in Math.* **80**, Springer 1996.
- [SaWi-02] Mark SAPIR and Daniel T. WISE, *Ascending HNN extensions of residually finite groups can be non-Hopfian and can have very few finite quotients*, *J. Pure Appl. Algebra* **166** (2002), 191–202.
- [Sega-75] Dan SEGAL, *On abelian-by-polycyclic groups*, *J. London Math. Soc.* **11** (1975), 445–452.
- [Serr-71] Jean-Pierre SERRE, *Cohomologie des groupes discrets*, in “Prospects in mathematics”, *Annals of Math. Studies* **70** (Princeton Univ. Press 1971), 77–169.
- [Serr-77] Jean-Pierre SERRE, *Arbres, amalgames, SL_2* , *Astérisque* **46**, Soc. Math. France, 1977.
- [Shal-00] Yehuda SHALOM, *Rigidity of commensurators and irreducible lattices*, *Invent. Math.* **141** (2000), 1–54.
- [Shme-64] Alfret L. SHMELKIN, *Free polynilpotent groups*, *Amer. Math. Transl. II* **55** (1966), 270–304. [Russian: *Izv. Adad. Nauk. SSSR* **28** (1964), 91–122.]
- [Shme-65] Alfret L. SHMELKIN, *Über auflösbare Produkte von Gruppen*, *Sib. Mat. Sz.* **6** (1965), 212–220.
- [Sidk-00] Said SIDKI, *Automorphisms of one-rooted trees: growth, circuit structure and acyclicity*, *J. of Math. Sci. New York*, **100** (2000), 1925–1943.
- [Smir-64] Dmitrii Matveevich SMIRNOV, *Generalized solvable groups and their group rings*, *Soviet Math. Doklady* **5** (1964), 465–467.
- [Smir-65] Dmitrii Matveevich SMIRNOV, *On generalized solvable groups and their group rings*, *Mat. Sb. (N.S.)* **67 (109)** 1965, 366–383.
- [Souc-01] Estelle SOUCHE, *Quasi-isométries et quasi-plans dans l'étude des groupes discrets*, Ph.D. Thesis, Université de Provence, 17 décembre 2001. *Appendice A: groupes de Baumslag-Solitar*, Pages 69–89.
- [Step-96] Anatolii Mikhailovich STEPIN, *Approximation of groups and group actions, the Cayley topology*, in “Ergodic theory of Z_d -actions: Proceedings of the Warwick symposium, 1993–94”, *Lect. Note Ser.* **228** (Cambridge University Press 1996), 475–484.
- [Stre-84] Ralph STREBEL, *Finitely presented soluble groups*, in “Group theory, Essays for Philip Hall” (Academic Press, 1984), 257–314.
- [Stre] Ralph STREBEL, *Notes on the Sigma-invariants, Part 1*, Preprint, arXiv:1204.0214v1, 1 Apr 2012.
- [Tars-38] Alfred TARSKI, *Algebraische Fassung des Massproblems*, *Fund. Math.* **31** (1938), 47–66 [Collected papers, Vol. 1, pp. 599–643.]

- [Thom–77] Michael W. THOMSON, *Subgroups of finitely presented solvable linear groups*, Trans. Amer. Math. Soc. **231** (1977), 133–142.
- [Vale–93] Alain VALETTE, *A short proof that free groups are residually p -groups*, Expositiones Math. **11** (1993), 65–71.
- [Wago–85] Stan WAGON, *The Banach-Tarski paradox*, Cambridge University Press, 1985.
- [Wehr–73] Bertram A.F. WEHRFRITZ, *Infinite linear groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete **76**, Springer 1973.
- [Wehr–80] Bertram A.F. WEHRFRITZ, *On finitely generated soluble linear groups*, Math. Zeit. **170** (1980), 155–167.

Post-scriptum

A CONSTRUCTION OF LARGE GROUPS, DUE TO YVES DE CORNULIER

By Ralph Strebel

The following proposition implies that any finitely presented cover of one of the groups $G_{\mathcal{V}}$ of Corollary C.6 is large. The proof uses the basic idea of the proof of Theorem A in [BiSt–78], refined for Theorem 6.1 in [BCGS].

Proposition. *Let G be a finitely generated group that is an extension of an infinite, locally finite group by an infinite cyclic group.*

Any finitely presented cover of G is large.

Proof. The beginning of our proof holds for any finitely generated group G given with a projection $\chi : G \rightarrow \mathbf{Z}$; let N denote the kernel of χ .

We can choose a finite subset A of N and an element $t \in G$ such that $\chi(t) = 1$ and $A \cup \{t\}$ generates G . Let \tilde{A} be a finite set given with a bijection $\tilde{a} \leftrightarrow a$ with A ; we denote by F the free group on a set $\tilde{A} \sqcup \{\tilde{t}\}$. Let

$$\rho : F \longrightarrow G$$

be the epimorphism defined by $\rho(\tilde{a}) = a$ for all $\tilde{a} \in \tilde{A}$, and $\rho(\tilde{t}) = t$.

We construct as follows a sequence $(G_m)_{m \geq 1}$ of finitely generated groups. The kernel N is the union of subgroups

$$B_m := \text{gp}\{t^\ell a t^{-\ell} \mid -m \leq \ell \leq m \text{ and } a \in A\}, \quad m \geq 1;$$

the notation indicates that B_m is the subgroup $\text{gp}\{\dots\}$ of G generated by $\{\dots\}$. For $m \geq 1$, consider the two subgroups

$$\begin{aligned} H_m &:= \text{gp}\{t^\ell a t^{-\ell} \mid -m \leq \ell \leq m-1 \text{ and } a \in A\} \\ K_m &:= \text{gp}\{t^\ell a t^{-\ell} \mid -m+1 \leq \ell \leq m \text{ and } a \in A\} \end{aligned}$$

of B_m , the isomorphism

$$\tau_m : H_m \xrightarrow{\cong} K_m, \quad h \mapsto tht^{-1},$$

and the HNN-extension

$$G_m := \langle B_m, u_m \mid u_m h u_m^{-1} = \tau_m(h) \text{ for all } h \in H_m \rangle.$$

By Britton's lemma, we can and do view B_m as a subgroup of G_m . Observe that G_m is actually generated by $A \subset H_m$ and u_m ; indeed, the defining relations in G_m imply that

$$u_m^\ell a u_m^{-\ell} = (\tau_m)^\ell(a) = t^\ell a t^{-\ell}$$

for all ℓ with $-m \leq \ell \leq m$ and $a \in A$.

Let

$$\lambda_{m,m+1} : G_m \longrightarrow G_{m+1}$$

be the homomorphism of which the restriction to B_m is the inclusion $B_m \subset B_{m+1} \subset G_{m+1}$, and such that $\lambda_{m,m+1}(u_m) = u_{m+1}$. Let

$$\lambda_m : G_m \longrightarrow G$$

be the homomorphism of which the restriction to B_m is the inclusion $B_m \subset G$, and such that $\lambda_m(u_m) = t$. Observe that $\lambda_{m,m+1}$ and λ_m are epimorphisms, and that $\lambda_m = \lambda_{m+1} \lambda_{m,m+1}$. Let

$$\rho_m : F \longrightarrow G_m$$

be the homomorphism defined by $\rho_m(\tilde{a}) = a$ for all $\tilde{a} \in \tilde{A}$ and $\rho_m(\tilde{t}) = u_m$. Observe that ρ_m is an epimorphism, and that $\lambda_m \rho_m = \rho$.

We have clearly

$$\ker(\rho_1) \subset \ker(\rho_2) \subset \cdots \subset \ker(\rho_m) \subset \cdots \subset \ker(\rho).$$

We claim that, moreover, $\bigcup_{m=1}^{\infty} \ker(\rho_m) = \ker(\rho)$. Indeed, let $w \in \ker(\rho)$. Since $\chi(\rho(w)) = 0 \in \mathbf{Z}$, the element w is a product of conjugates by powers of \tilde{t} of elements in $\tilde{A} \cup \tilde{A}^{-1}$. Let m be large enough for w to be a product of elements of the form $\tilde{t}^\ell \tilde{a} \tilde{t}^{-\ell}$, with $|\ell| \leq m$ and $\tilde{a} \in \tilde{A} \cup \tilde{A}^{-1}$. Then $\rho(w) = \lambda_m(\rho_m(w)) = 0$, with $\rho_m(w) \in B_m \subset G_m$. Since the restriction of λ_m to B_m is injective, this implies $w \in \ker(\rho_m)$, and the claim holds.

We use now one more hypothesis: N is locally finite. Since B_m is a finitely generated subgroup in the locally finite group N , it is a finite group. It follows from Proposition 11 and its corollary in [Serr-77, Pages 160–161] that the HNN-extension G_m has a finitely generated free subgroup F_m of finite index. We use now the last hypothesis: N is infinite. It implies that F_m cannot be cyclic, i.e. that G_m has a finite

index subgroup F_m that is non-abelian and free. In other words, G_m is large.

Consider finally a finitely presented group E and an epimorphism $\pi : E \twoheadrightarrow G$. Upon replacing A at the beginning of the proof by a larger finite subset of N , we can assume that there exists an epimorphism $\sigma : F \twoheadrightarrow E$ of which the kernel is generated by a finite set of relators, and that ρ factors as the composition $F \xrightarrow{\sigma} E \xrightarrow{\pi} G$. Hence $\ker(\sigma) \subset \ker(\rho)$. Since $\ker(\sigma)$ is the normal closure of a finite set, there exists $m \geq 1$ with $\ker(\sigma) \subset \ker(\rho_m)$. It follows that $E = F/K$ has $G_m = F/\ker(\rho_m)$ as a quotient group. Since G_m is large, we conclude that E is large. \square

As a consequence of the proof, we have:

Corollary. *Let G be a (locally finite)-by-(infinite cyclic) group, as in the previous proposition. Assume that G is finitely generated, say that G has a generating set S of some size $k \geq 1$.*

Then, in the space of marked group \mathcal{M}_k , see Section 3, the pair (G, S) is the limit of finitely generated virtually free groups. More precisely, there exists in \mathcal{M}_k a sequence $(G_m, S_m)_{m \geq 1}$ of virtually free marked groups converging to (G, S) , and for each $m \geq 1$ an epimorphism $G_m \twoheadrightarrow G$ mapping S_m onto S .

M.B. AND R.G.: DEPARTMENT OF MATHEMATICS, MAILSTOP 3368,
TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA.

E-mail address: mbenli@math.tamu.edu, grigorch@math.tamu.edu

P.H.: SECTION DE MATHÉMATIQUES, UNIVERSITÉ DE GENÈVE, C.P. 64,
CH-1211 GENÈVE 4, SUISSE.

E-mail address: Pierre.deLaHarpe@unige.ch

R.S.: UNIVERSITÉ DE FRIBOURG, DÉPARTEMENT DE MATHÉMATIQUES,
CHEMIN DU MUSÉE 23, CH-1700 FRIBOURG, SUISSE.

E-mail address: ralph.strebel@unifr.ch