

Measurement of the azimuthal angle distribution of leptons from W boson decays as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV

D. Acosta,¹⁴ T. Affolder,⁷ M.G. Albrow,¹³ D. Ambrose,³⁶ D. Amidei,²⁷ K. Anikeev,²⁶ J. Antos,¹ G. Apollinari,¹³ T. Arisawa,⁵⁰ A. Artikov,¹¹ W. Ashmanskas,² F. Azfar,³⁴ P. Azzi-Bacchetta,³⁵ N. Bacchetta,³⁵ H. Bachacou,²⁴ W. Badgett,¹³ A. Barbaro-Galtieri,²⁴ V.E. Barnes,³⁹ B.A. Barnett,²¹ S. Baroiant,⁵ M. Barone,¹⁵ G. Bauer,²⁶ F. Bedeschi,³⁷ S. Behari,²¹ S. Belforte,⁴⁷ W.H. Bell,¹⁷ G. Bellettini,³⁷ J. Bellinger,⁵¹ D. Benjamin,¹² A. Beretvas,¹³ A. Bhatti,⁴¹ M. Binkley,¹³ D. Bisello,³⁵ M. Bishai,¹³ R.E. Blair,² C. Blocker,⁴ K. Bloom,²⁷ B. Blumenfeld,²¹ A. Bocci,⁴¹ A. Bodek,⁴⁰ G. Bolla,³⁹ A. Bolshov,²⁶ D. Bortoletto,³⁹ J. Boudreau,³⁸ C. Bromberg,²⁸ E. Brubaker,²⁴ J. Budagov,¹¹ H.S. Budd,⁴⁰ K. Burkett,¹³ G. Busetto,³⁵ K.L. Byrum,² S. Cabrera,¹² M. Campbell,²⁷ W. Carithers,²⁴ D. Carlsmith,⁵¹ A. Castro,³ D. Cauz,⁴⁷ A. Cerri,²⁴ L. Cerrito,²⁰ J. Chapman,²⁷ C. Chen,³⁶ Y.C. Chen,¹ M. Chertok,⁵ G. Chiarelli,³⁷ G. Chlachidze,¹³ F. Chlebana,¹³ M.L. Chu,¹ J.Y. Chung,³² W.-H. Chung,⁵¹ Y.S. Chung,⁴⁰ C.I. Ciobanu,²⁰ A.G. Clark,¹⁶ M. Coca,⁴⁰ A. Connolly,²⁴ M. Convery,⁴¹ J. Conway,⁴³ M. Cordelli,¹⁵ J. Cranshaw,⁴⁵ R. Culbertson,¹³ D. Dagenhart,⁴ S. D'Auria,¹⁷ P. de Barbaro,⁴⁰ S. De Cecco,⁴² S. Dell'Agnello,¹⁵ M. Dell'Orso,³⁷ S. Demers,⁴⁰ L. Demortier,⁴¹ M. Deninno,³ D. De Pedis,⁴² P.F. Derwent,¹³ C. Dionisi,⁴² J.R. Dittmann,¹³ A. Dominguez,²⁴ S. Donati,³⁷ M. D'Onofrio,¹⁶ T. Dorigo,³⁵ N. Eddy,²⁰ R. Erbacher,¹³ D. Errede,²⁰ S. Errede,²⁰ R. Eusebi,⁴⁰ S. Farrington,¹⁷ R.G. Feild,⁵² J.P. Fernandez,³⁹ C. Ferretti,²⁷ R.D. Field,¹⁴ I. Fiori,³⁷ B. Flaughner,¹³ L.R. Flores-Castillo,³⁸ G.W. Foster,¹³ M. Franklin,¹⁸ J. Friedman,²⁶ I. Furic,²⁶ M. Gallinaro,⁴¹ M. Garcia-Sciveres,²⁴ A.F. Garfinkel,³⁹ C. Gay,⁵² D.W. Gerdes,²⁷ E. Gerstein,⁹ S. Giagu,⁴² P. Giannetti,³⁷ K. Giolo,³⁹ M. Giordani,⁴⁷ P. Giromini,¹⁵ V. Glagolev,¹¹ D. Glenzinski,¹³ M. Gold,³⁰ N. Goldschmidt,²⁷ J. Goldstein,³⁴ G. Gomez,⁸ M. Goncharov,⁴⁴ I. Gorelov,³⁰ A.T. Goshaw,¹² Y. Gotra,³⁸ K. Goulios,⁴¹ A. Gresele,³ C. Grosso-Pilcher,¹⁰ M. Guenther,³⁹ J. Guimaraes da Costa,¹⁸ C. Haber,²⁴ S.R. Hahn,¹³ E. Halkiadakis,⁴⁰ R. Handler,⁵¹ F. Happacher,¹⁵ K. Hara,⁴⁸ R.M. Harris,¹³ F. Hartmann,²² K. Hatakeyama,⁴¹ J. Hauser,⁶ J. Heinrich,³⁶ M. Hennecke,²² M. Herndon,²¹ C. Hill,⁷ A. Hocker,⁴⁰ K.D. Hoffman,¹⁰ S. Hou,¹ B.T. Huffman,³⁴ R. Hughes,³² J. Huston,²⁸ J. Incandela,⁷ G. Introzzi,³⁷ M. Iori,⁴² C. Issever,⁷ A. Ivanov,⁴⁰ Y. Iwata,¹⁹ B. Iyutin,²⁶ E. James,¹³ M. Jones,³⁹ T. Kamon,⁴⁴ J. Kang,²⁷ M. Karagoz Unel,³¹ S. Kartal,¹³ H. Kasha,⁵² Y. Kato,³³ R.D. Kennedy,¹³ R. Kephart,¹³ B. Kilminster,⁴⁰ D.H. Kim,²³ H.S. Kim,²⁰ M.J. Kim,⁹ S.B. Kim,²³ S.H. Kim,⁴⁸ T.H. Kim,²⁶ Y.K. Kim,¹⁰ M. Kirby,¹² L. Kirsch,⁴ S. Klimenko,¹⁴ P. Koehn,³² K. Kondo,⁵⁰ J. Konigsberg,¹⁴ A. Korn,²⁶ A. Korytov,¹⁴ J. Kroll,³⁶ M. Kruse,¹² V. Krutelyov,⁴⁴ S.E. Kuhlmann,² N. Kuznetsova,¹³ A.T. Laasanen,³⁹ S. Lami,⁴¹ S. Lammel,¹³ J. Lancaster,¹² M. Lancaster,²⁵ R. Lander,⁵ K. Lannon,³² A. Lath,⁴³ G. Latino,³⁰ T. LeCompte,² Y. Le,²¹ J. Lee,⁴⁰ S.W. Lee,⁴⁴ N. Leonardo,²⁶ S. Leone,³⁷ J.D. Lewis,¹³ K. Li,⁵² C.S. Lin,¹³ M. Lindgren,⁶ T.M. Liss,²⁰ D.O. Litvintsev,¹³ T. Liu,¹³ N.S. Lockyer,³⁶ A. Loginov,²⁹ M. Loreti,³⁵ D. Lucchesi,³⁵ P. Lukens,¹³ L. Lyons,³⁴ J. Lys,²⁴ R. Madrak,¹⁸ K. Maeshima,¹³ P. Maksimovic,²¹ L. Malferrari,³ M. Mangano,³⁷ G. Manca,³⁴ M. Mariotti,³⁵ M. Martin,²¹ A. Martin,⁵² V. Martin,³¹ M. Martinez,¹³ P. Mazzanti,³ K.S. McFarland,⁴⁰ P. McIntyre,⁴⁴ M. Menguzzato,³⁵ A. Menzione,³⁷ P. Merkel,¹³ C. Mesropian,⁴¹ A. Meyer,¹³ T. Miao,¹³ J.S. Miller,²⁷ R. Miller,²⁸ S. Miscetti,¹⁵ G. Mitselmakher,¹⁴ N. Moggi,³ R. Moore,¹³ T. Moulik,³⁹ A. Mukherjee,^M Mulhearn,²⁶ T. Muller,²² A. Munar,³⁶ P. Murat,¹³ J. Nachtman,¹³ S. Nahn,⁵² I. Nakano,¹⁹ R. Napora,²¹ C. Nelson,¹³ T. Nelson,¹³ C. Neu,³² M.S. Neubauer,²⁶ C. Newman-Holmes,¹³ F. Niell,²⁷ T. Nigmanov,³⁸ L. Nodulman,² S.H. Oh,¹² Y.D. Oh,²³ T. Ohsugi,¹⁹ T. Okusawa,³³ W. Orejudos,²⁴ C. Pagliarone,³⁷ F. Palmonari,³⁷ R. Paoletti,³⁷ V. Papadimitriou,⁴⁵ J. Patrick,¹³ G. Pauletta,⁴⁷ M. Paulini,⁹ T. Pauly,³⁴ C. Paus,²⁶ D. Pellett,⁵ A. Penzo,⁴⁷ T.J. Phillips,¹² G. Piacentino,³⁷ J. Piedra,⁸ K.T. Pitts,²⁰ A. Pompos,³⁹ L. Pondrom,⁵¹ G. Pope,³⁸ O. Poukov,¹¹ T. Pratt,³⁴ F. Prokoshin,¹¹ J. Proudfoot,² F. Ptohos,¹⁵ G. Punzi,³⁷ J. Rademacker,³⁴ A. Rakitine,²⁶ F. Ratnikov,⁴³ H. Ray,²⁷ A. Reichold,³⁴ P. Renton,³⁴ M. Rescigno,⁴² F. Rimondi,³ L. Ristori,³⁷ W.J. Robertson,¹² T. Rodrigo,⁸ S. Rolli,⁴⁹ L. Rosenson,²⁶ R. Roser,¹³ R. Rossin,³⁵ C. Rott,³⁹ A. Roy,³⁹ A. Ruiz,⁸ D. Ryan,⁴⁹ A. Safonov,⁵ R. St. Denis,¹⁷ W.K. Sakumoto,⁴⁰ D. Saltzberg,⁶ C. Sanchez,³² A. Sansoni,¹⁵ L. Santi,⁴⁷ S. Sarkar,⁴² P. Savard,⁴⁶ A. Savoy-Navarro,¹³ P. Schlabach,¹³ E.E. Schmidt,¹³ M.P. Schmidt,⁵² M. Schmitt,³¹ L. Scodellaro,³⁵ A. Scribano,³⁷ A. Sedov,³⁹ S. Seidel,³⁰ Y. Seiya,⁴⁸ A. Semenov,¹¹ F. Semeria,³ M.D. Shapiro,²⁴ P.F. Shepard,³⁸ T. Shibayama,⁴⁸ M. Shimojima,⁴⁸ M. Shochet,¹⁰ A. Sidoti,³⁵ A. Sill,⁴⁵ P. Sinervo,⁴⁶ A.J. Slaughter,⁵² K. Sliwa,⁴⁹ F.D. Snider,¹³ R. Snihur,²⁵ M. Spezziga,⁴⁵ L. Spiegel,¹³ F. Spinella,³⁷ M. Spiropulu,⁷ A. Stefanini,³⁷ J. Strologas,³⁰ D. Stuart,⁷ A. Sukhanov,¹⁴ K. Sumorok,²⁶ T. Suzuki,⁴⁸ R. Takashima,¹⁹ K. Takikawa,⁴⁸ M. Tanaka,² M. Tecchio,²⁷ P.K. Teng,¹ K. Terashi,⁴¹ R.J. Tesarek,¹³ S. Tether,²⁶ J. Thom,¹³ A.S. Thompson,¹⁷ E. Thomson,³² P. Tipton,⁴⁰ S. Tkaczyk,¹³ D. Toback,⁴⁴ K. Tollefson,²⁸ D. Tonelli,³⁷ M. Tönnesmann,²⁸ H. Toyoda,³³ W. Trischuk,⁴⁶ J. Tseng,²⁶ D. Tsybychev,¹⁴ N. Turini,³⁷ F. Ukegawa,⁴⁸ T. Unverhau,¹⁷ T. Vaiculis,⁴⁰ A. Varganov,²⁷ E. Vataga,³⁷ S. Vejčik III,¹³ G. Velev,¹³ G. Veramendi,²⁴ R. Vidal,¹³ I. Vila,⁸ R. Vilar,⁸ I. Volobouev,²⁴ M. von der Mey,⁶ R.G. Wagner,² R.L. Wagner,¹³ W. Wagner,²² Z. Wan,⁴³ C. Wang,¹² M.J. Wang,¹ S.M. Wang,¹⁴ B. Ward,¹⁷ S. Waschke,¹⁷ D. Waters,²⁵ T. Watts,⁴³ M. Weber,²⁴ W.C. Wester III,¹³ B. Whitehouse,⁴⁹ A.B. Wicklund,² E. Wicklund,¹³ H.H. Williams,³⁶ P. Wilson,¹³ B.L. Winer,³² S. Wolbers,¹³ M. Wolter,⁴⁹ S. Worm,⁴³ X. Wu,¹⁶ F. Würthwein,²⁶ U.K. Yang,¹⁰ W. Yao,²⁴ G.P. Yeh,¹³ K. Yi,²¹

(CDF Collaboration)

- ¹ *Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*
- ² *Argonne National Laboratory, Argonne, Illinois 60439*
- ³ *Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*
- ⁴ *Brandeis University, Waltham, Massachusetts 02254*
- ⁵ *University of California at Davis, Davis, California 95616*
- ⁶ *University of California at Los Angeles, Los Angeles, California 90024*
- ⁷ *University of California at Santa Barbara, Santa Barbara, California 93106*
- ⁸ *Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*
- ⁹ *Carnegie Mellon University, Pittsburgh, Pennsylvania 15213*
- ¹⁰ *Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*
- ¹¹ *Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*
- ¹² *Duke University, Durham, North Carolina 27708*
- ¹³ *Fermi National Accelerator Laboratory, Batavia, Illinois 60510*
- ¹⁴ *University of Florida, Gainesville, Florida 32611*
- ¹⁵ *Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*
- ¹⁶ *University of Geneva, CH-1211 Geneva 4, Switzerland*
- ¹⁷ *Glasgow University, Glasgow G12 8QQ, United Kingdom*
- ¹⁸ *Harvard University, Cambridge, Massachusetts 02138*
- ¹⁹ *Hiroshima University, Higashi-Hiroshima 724, Japan*
- ²⁰ *University of Illinois, Urbana, Illinois 61801*
- ²¹ *The Johns Hopkins University, Baltimore, Maryland 21218*
- ²² *Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- ²³ *Center for High Energy Physics: Kyungpook National University, Taegu 702-701; Seoul National University, Seoul 151-742; and SungKyunKwan University, Suwon 440-746; Korea*
- ²⁴ *Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720*
- ²⁵ *University College London, London WC1E 6BT, United Kingdom*
- ²⁶ *Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*
- ²⁷ *University of Michigan, Ann Arbor, Michigan 48109*
- ²⁸ *Michigan State University, East Lansing, Michigan 48824*
- ²⁹ *Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- ³⁰ *University of New Mexico, Albuquerque, New Mexico 87131*
- ³¹ *Northwestern University, Evanston, Illinois 60208*
- ³² *The Ohio State University, Columbus, Ohio 43210*
- ³³ *Osaka City University, Osaka 588, Japan*
- ³⁴ *University of Oxford, Oxford OX1 3RH, United Kingdom*
- ³⁵ *Universita di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy*
- ³⁶ *University of Pennsylvania, Philadelphia, Pennsylvania 19104*
- ³⁷ *Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy*
- ³⁸ *University of Pittsburgh, Pittsburgh, Pennsylvania 15260*
- ³⁹ *Purdue University, West Lafayette, Indiana 47907*
- ⁴⁰ *University of Rochester, Rochester, New York 14627*
- ⁴¹ *Rockefeller University, New York, New York 10021*
- ⁴² *Instituto Nazionale de Fisica Nucleare, Sezione di Roma, University di Roma I, "La Sapienza," I-00185 Roma, Italy*
- ⁴³ *Rutgers University, Piscataway, New Jersey 08855*
- ⁴⁴ *Texas A&M University, College Station, Texas 77843*
- ⁴⁵ *Texas Tech University, Lubbock, Texas 79409*
- ⁴⁶ *Institute of Particle Physics, University of Toronto, Toronto M5S 1A7, Canada*
- ⁴⁷ *Istituto Nazionale di Fisica Nucleare, University of Trieste/ Udine, Italy*
- ⁴⁸ *University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁴⁹ *Tufts University, Medford, Massachusetts 02155*
- ⁵⁰ *Waseda University, Tokyo 169, Japan*
- ⁵¹ *University of Wisconsin, Madison, Wisconsin 53706*
- ⁵² *Yale University, New Haven, Connecticut 06520*

(Dated: March 30, 2005)

We present the first measurement of the A_2 and A_3 angular coefficients of the W boson produced in proton-antiproton collisions. We study $W \rightarrow e\nu_e$ and $W \rightarrow \mu\nu_\mu$ candidate events produced in association with at least one jet at CDF, during Run Ia and Run Ib of the Tevatron at $\sqrt{s}=1.8$ TeV. The corresponding integrated luminosity was 110 pb^{-1} . The jet balances the transverse momentum of the W and introduces QCD effects in W boson production. The extraction of the angular coefficients is achieved through the direct measurement of the azimuthal angle of the charged lepton in the Collins-Soper rest-frame of the W boson. The angular coefficients are measured as a function of the transverse momentum of the W boson. The electron, muon, and combined results are in good agreement with the Standard Model prediction, up to order α_s^2 in QCD.

PACS numbers: 11.30.Er, 11.80.Cr, 12.15.-y, 12.38.-t, 12.38.Qk, 13.38.Be, 13.87.-a, 13.88.+e, 14.70.Fm

I. INTRODUCTION

Measurements of the W boson differential cross section, as a function of energy and direction, provide information about the nature of both the underlying electroweak interaction, and the effects of chromodynamics (QCD). This differential cross section can be expressed as a function of the helicity cross sections of the W , allowing us to study the W polarization and associated asymmetries. Because of the difficulties in fully reconstructing a W boson in three-dimensions at a hadron collider, the complete angular distribution of the W has not been determined yet. In this paper we present the first measurement of two of the four significant leading angular coefficients of the W boson produced at a hadron collider.

The total differential cross section for W boson production in a hadron collider is given by

$$\begin{aligned} \frac{d\sigma}{d(p_T^W)^2 dy d\cos\theta d\phi} &= \frac{3}{16\pi} \frac{d\sigma^u}{d(p_T^W)^2 dy} [(1 + \cos^2\theta) \\ &+ \frac{1}{2}A_0(1 - 3\cos^2\theta) + A_1 \sin 2\theta \cos\phi \\ &+ \frac{1}{2}A_2 \sin^2\theta \cos 2\phi + A_3 \sin\theta \cos\phi \\ &+ A_4 \cos\theta + A_5 \sin^2\theta \sin 2\phi \\ &+ A_6 \sin 2\theta \sin\phi + A_7 \sin\theta \sin\phi] \quad (1) \end{aligned}$$

where p_T^W and y are the transverse momentum and the rapidity of the W in the laboratory frame, and θ and ϕ are the polar and azimuthal angles of the charged lepton from W boson decay in the Collins-Soper (CS) frame [1]. The factors $A_i(p_T^W, y)$ are the angular coefficients of the W boson, which are ratios of the helicity cross sections of the W and its total unpolarized cross section $d\sigma^u/d(p_T^W)^2 dy$. The CS frame [2] is the rest-frame of the W with a z -axis that bisects the angle between the proton direction and the direction opposite that of the antiproton (Figure 1), and it is used because in this frame we can in principle exactly reconstruct the azimuthal angle ϕ and the polar quantity $|\cos\theta|$. Our ignorance of the W boson longitudinal momentum, which is due to our inability to measure the longitudinal momentum of the neutrino, only introduces a two-fold ambiguity on the sign of $\cos\theta$. It is common to integrate Equation (1) over

y and study the variation of the angular coefficients as a function of p_T^W .

To study the angular distribution of the W we must choose a particular charge for the boson. In this paper we consider the W^- bosons; the W^+ bosons in our samples are CP transformed to be treated as W^- bosons. The angular coefficients for the W^+ are obtained by CP transforming Equation (1) [3].

If the W is produced with no transverse momentum, it is polarized along the beam axis, due to the V-A nature of the weak interactions and helicity conservation. In that case A_4 is the only non-zero coefficient. If only valence quarks contributed to W production, A_4 would equal 2, and the angular distribution given by Equation (1) would be $\sim (1 + \cos\theta)^2$, a result that was first verified by the UA1 experiment [4].

If the W is produced with non-negligible transverse momentum, balanced by the associated production of jets, the rest of the angular coefficients are present, and the cross section depends on the azimuthal angle ϕ as well. The last three angular coefficients A_5 , A_6 , and A_7 are non-zero only if gluon loops are present in the production of the W boson. Hence, in order to study all the angular coefficients and associated helicity cross sections of the W boson in a hadron collider, we must consider the production of the W with QCD effects up to order α_s^2 .

The importance of the determination of the W angular coefficients is discussed in [5], and summarized here. It allows us to measure for the first time the full differential cross section of the W and study its polarization, since the angular coefficients are directly related to the helicity cross sections. It also helps us verify the QCD effects in the production of the W up to order α_s^2 . For example, according to the Standard Model (SM), A_2 is not equal to A_0 only if the effects of gluon loops are taken into account. In addition, A_3 is only affected by the gluon-quark interaction and its measurement can be used to constrain the gluon parton distribution functions. Moreover, the next-to-leading order angular coefficients A_5 , A_6 , and A_7 are P -odd and T -odd and may play an important role in direct CP violation effects in W production and decay [6]. Finally, quantitative understanding of the W angular distribution could be used to test new theoretical models and to facilitate new discoveries.




FIG. 1: Transforming from the laboratory frame to the Collins-Soper frame. We first boost to the W rest-frame, then rotate the $x - z$ plane so that it coincides with the $p - \bar{p}$ plane. Finally we rotate the frame around the y -axis so that the z -axis bisects the angle between \vec{p} and $-\vec{p}$. The positive y axis is selected to have the same direction as $\vec{p}_{CS} \times \vec{l}_{CS}$.

In this paper we present the first measurement of the A_2 and A_3 angular coefficients of the W boson. These coefficients fully describe the azimuthal differential cross section of the W boson, and they are two of the four significant coefficients that describe the total differential cross section of the W , given that A_1 and the next-to-leading order angular coefficients have considerably lower values [5, 7]. This measurement is accomplished using the azimuthal angle of the charged lepton in the CS W rest-frame [8], and is presented as a function of the transverse momentum of the W boson. The CS polar angle analysis is more sensitive to the A_0 and A_4 angular coefficients (see [9, 10] for a measurement of A_0). Because Equation (1) arises solely from quantum field theory, without input from any specific theoretical model of W boson production, our experimental results are thus model-independent.

II. THE CDF DETECTOR AND EVENT SELECTION

A. The CDF detector

The CDF detector is described in detail in [11]. It is a general purpose detector of charged leptons, hadrons, jets, and photons, produced from proton-antiproton collisions at the Tevatron accelerator at Fermilab. The W and Z bosons are detected through their decay leptons, while the transverse momentum of the neutrinos is estimated from the missing transverse energy of the events (\cancel{E}_T).

The z -axis of the detector coincides with the direction of the proton beam and defines the polar angle θ_{lab} in the laboratory frame. The y -axis points vertically upward and the x -axis is in the horizontal plane, so as to form a right-handed coordinate system. The pseudorapidity, $\eta_{lab} = -\ln[\tan(\theta_{lab}/2)]$, and the azimuthal angle ϕ_{lab} are used to specify detector physical areas.

The tracking system of CDF consists of the silicon vertex detector (SVX), the vertex time projection chamber (VTX) and the central tracking chamber (CTC), all immersed in a 1.4 T magnetic field produced by a superconducting solenoid of length 4.8 m and radius 1.5 m. The SVX, a four layer silicon micro-strip vertex detector, is located immediately outside the beampipe. It is used to find secondary vertices and provides the impact parameter of tracks in the transverse $r - \phi_{lab}$ plane. The VTX, located outside the SVX, is a vertex time projection chamber that provides $r - z$ tracking information up to a radius of 22 cm and pseudorapidity $|\eta_{lab}| \leq 3.5$. It measures the z -position of the primary vertex. Finally, surrounding the SVX and the VTX is the CTC, a 3.2 m long cylindrical drift chamber containing 84 layers of sense wires arranged in five superlayers of axial wires and four superlayers of stereo wires. The axial superlayers have 12 radially separated layers of sense wires, parallel to the z -axis, that measure the $r - \phi_{lab}$ position of the tracks. The stereo superlayers have six layers of sense wires with alternate $\sim \pm 3^\circ$ stereo angles with respect to the beamline, and measure a combination of $r - \phi_{lab}$ and z information. The stereo and axial data are combined to reconstruct the 3-dimensional track. The CTC covers the pseudorapidity interval $|\eta_{lab}| < 1.0$ and transverse momentum $p_T \geq 0.4$ GeV [12]. The combined momentum resolution of the tracking system is $\delta p_T/p_T = \sqrt{(0.0009 p_T)^2 + (0.0066)^2}$, where p_T is the transverse momentum in GeV.

The solenoid is surrounded by sampling calorimeters used to measure the electromagnetic and hadronic energy of electrons, photons, and jets. The calorimeters cover the pseudorapidity range $|\eta_{lab}| \leq 4.2$ and the azimuthal angle range $0 \leq \phi_{lab} \leq 2\pi$. They are segmented in $\eta_{lab} - \phi_{lab}$ towers pointing to the nominal interaction point at the center of the detector. The tower granularity is $(\Delta\eta_{lab} \times \Delta\phi_{lab}) = (0.1 \times 15^\circ)$ in the central region ($0 \leq |\eta_{lab}| \leq 1.1$) and $(0.1 \times 5^\circ)$ in the plug ($1.1 < |\eta_{lab}| \leq 2.4$) and forward ($2.4 < |\eta_{lab}| \leq 4.2$) regions. Each region has an electromagnetic calorimeter (CEM in the central region, PEM in the plug region, and FEM in the forward region) followed by a hadron calorimeter at larger radius from the beam (CHA, PHA, and FHA respectively). The central calorimeters are segmented in 24 wedges per each half of the detector ($-1.1 \leq \eta_{lab} \leq 0$ and $0 \leq \eta_{lab} \leq 1.1$). The CEM is an 18 radiation length lead-scintillator stack with a position resolution of 2 mm and an energy resolution of $\delta E_T/E_T = \sqrt{(13.5\%/\sqrt{E_T})^2 + (2\%)^2}$, where E_T is the transverse energy in GeV. Located six radiation lengths deep inside the CEM calorimeter (184 cm from

the beamline), proportional wire chambers (CES) with additional cathode strip read-out provide shower position measurements in the z and $r - \phi_{\text{lab}}$ directions. The central hadron calorimeter (CHA) is an iron-scintillator stack which is 4.5 interaction lengths thick and provides energy measurement with a resolution of $\delta E_T/E_T = \sqrt{(50\%/\sqrt{E_T})^2 + (3\%)^2}$, where E_T is the transverse energy in GeV.

The central muon system consists of three components and is capable of detecting muons with transverse momentum $p_T \geq 1.4$ GeV and pseudorapidity $|\eta_{\text{lab}}| < 1.0$. The Central Muon Chambers (CMU) cover the region $|\eta_{\text{lab}}| < 0.6$ and consist of four layers of planar drift chambers outside the hadron calorimeter, allowing the reconstruction of the muons which typically pass the five absorption lengths of material. Outside the CMU there are three additional absorption lengths of material (0.6 m of steel) followed by four layers of drift chambers, the Central Muon Upgrade (CMP). The CMP chambers cover the same pseudorapidity region as the CMU, and they were introduced to limit the background caused from punch-through pions. Finally, the Central Muon Extension chambers (CMX) cover the region $0.6 \leq \eta_{\text{lab}} \leq 1.0$. These drift chambers are sandwiched between scintillators (CSX). Depending on the incident angle, particles have to penetrate six to nine absorption lengths of material to be detected in the CMX. The particle candidate stub provided by the muon system is matched with a track from the CTC in order to successfully reconstruct a muon.

B. The CDF triggers

CDF has a three-level trigger system designed to select events that can contain electrons, muons, jets, and \cancel{E}_T . The first two levels are implemented in hardware, while the third is a software trigger which uses a version of the offline reconstruction software optimized for speed and implemented by a CPU farm.

At level-1, electrons were selected by the presence of an electromagnetic trigger tower with energy above 6 GeV (Run Ia) or 8 GeV (Run Ib), where one trigger tower consisted of two adjacent physical towers (in pseudorapidity). Muons were selected by the presence of a track stub in the CMU or CMX, where there was also signal in the CMP.

At level-2, electrons satisfied one of several triggers. In Run Ia, the event passed the trigger if the energy cluster in the CEM was at least 9 GeV with a seed tower of at least 7 GeV, and a matching track with $p_T > 9.2$ GeV was found by the Central Fast Tracker (CFT), the fast hardware processor that matched CTC tracks in the $r - \phi_{\text{lab}}$ plane with signals in the calorimeters and muon chambers. It also passed the trigger if there was an isolated cluster in the CEM calorimeter of at least 16 GeV. The most common Run Ib level-2 electron trigger requires the existence of a cluster in the CEM with at least 16

GeV and the existence of a matching track in the CFT with $p_T > 12$ GeV. The muon trigger at level-2 required a track of at least 9 GeV (Run Ia) or 12 GeV (Run Ib) that matched a CMX stub (CMX triggers), both CMU and CMP stubs (CMUP triggers), or a CMU stub but no CMP stub (CMNP triggers).

At level-3, reconstruction programs performed 3-dimensional track reconstruction. In the Run Ia level-3 electron trigger, most of the accepted events passed the requirement that the CEM cluster had $E_T > 18$ GeV, and was associated with a track of $p_T > 13$ GeV. The transverse energy of the cluster is defined as $E_T = E \sin \theta$, where E is the total energy deposited in the CEM, and θ is the polar angle measured from the event vertex to the centroid of the cluster. Cuts were applied on the shape of the electron shower profile and the energy deposition patterns. In the Run Ib level-3 electron trigger, CEM $E_T > 18$ GeV and CFT $p_T > 13$ GeV requirements were applied. The muon trigger at level-3 required that the CFT transverse momentum was greater than 18 GeV, the energy deposited in the hadron calorimeter was less than 6 GeV, the energy deposited in the electromagnetic calorimeter was less than 2 GeV, and the extrapolated CTC track was no more than 2 centimeters away from the muon stub in the CMU chambers and 5 centimeters in the CMP or CMX chambers in the x direction. Events that pass the level-3 trigger were recorded to tape for offline analysis.

C. The datasets

The events passing the three levels of our trigger system constitute the inclusive high- p_T electron and muon data samples. We apply kinematic and lepton identification cuts, described in Sections II C 1 and II C 2 to obtain the inclusive W electron and muon datasets respectively. Using these datasets we arrive at the W +jet datasets by applying the jet selection cuts described in Section II C 3.

1. Inclusive W Electron Selection

After passing the three levels of trigger requirements, the following event selection cuts are applied to the inclusive electron data sample:

- The event must belong to a good run.
 - $E_T^e \geq 20$ GeV,
- where E_T^e is the transverse energy of the CEM cluster, corrected for differences in response, non-linearities, and time-dependent changes.
- $|\eta_{\text{lab}}^e| \leq 1$,
- where η_{lab}^e is the pseudorapidity of the electron.
- The electron must fall in a fiducial part of the CEM calorimeter.
 - $\text{ISO}(0.4) \equiv E_{\Delta R=0.4}^{\text{Excess}}/E_T^{\text{cluster}} < 0.1$,
- where $E_{\Delta R=0.4}^{\text{Excess}}$ is the excess transverse energy in a cone

of size $\Delta R = \sqrt{(\Delta\phi_{\text{lab}})^2 + (\Delta\eta_{\text{lab}})^2}$ centered on the direction of the electromagnetic cluster, and E_T^{cluster} is the transverse energy of that cluster.

- $E^{\text{HAD}}/E^{\text{EM}} < 0.055 + 0.00045E^e$,

where E^{HAD} is the energy deposited in the hadron calorimeter, and E^{EM} is the energy deposited in the electromagnetic calorimeter.

- $L_{\text{SHR}} \equiv 0.14 \sum_i \frac{E_i^{\text{meas}} - E_i^{\text{exp}}}{\sqrt{(0.14)^2 E_i^{\text{meas}} + (\Delta E_i^{\text{exp}})^2}} < 0.2$,

where L_{SHR} is the *lateral shower profile*, E_i^{meas} is the energy measured in the i^{th} -tower adjacent to the seed tower, E_i^{exp} is the expectation for the energy in that tower, ΔE_i^{exp} is the uncertainty on the expected energy, and $0.14\sqrt{E^{\text{meas}}}$ is the uncertainty in the measurement of the cluster energy.

- $\chi_{\text{CES}}^2 < 10$.

We measure the shower profile along the z direction using the CES strips and the shower profile along the x direction using the CES wires. By comparing the measured x -shape and z -shape to the ones determined from test-beam studies we extract the chi-squared quantities for the two directions. The chi-squared we use is the average of the two.

- $0.5 \leq E^e/p^e \leq 2.0$,

where E^e is the corrected energy of the electron, and p^e is the beam-constrained momentum of the electron, i.e., the momentum determined when the fit trajectory of the CTC hits is constrained to pass through the beam line.

- $|\Delta X| < 1.5$ cm and $|\Delta Z| < 3.0$ cm,

where ΔX and ΔZ are the difference in the x and z directions respectively, between the extrapolated CTC track and the CES position of the shower.

- $|Z_{\text{VTX}}| \leq 60$ cm,

where Z_{VTX} is the z position of the primary vertex.

- Photon conversions are removed.

We next apply the following cuts:

- $\cancel{E}_T > 20$ GeV,

where \cancel{E}_T is the missing transverse energy in the event, calculated from the energy imbalance in the calorimeters, with a correction for the unclustered energy – calorimeter energy not taken into account by the jet clustering algorithm – and possible presence of muons.

- $M_T^W > 40$ GeV,

where M_T^W is the W transverse mass. This cut removes the background from W bosons decaying into tau leptons which subsequently decay into electrons.

- The event must not be consistent with a Z decaying into two observed leptons, or a Z in which one of the decay tracks has not been identified.

The 73363 events passing these cuts constitute our inclusive W electron data sample (Run Ia: 13290 events and Run Ib: 60073 events), corresponding to an integrated luminosity of 110 pb^{-1} (Run Ia: $19.65 \pm 0.71 \text{ pb}^{-1}$ and Run Ib: $90.35 \pm 3.70 \text{ pb}^{-1}$).

2. Inclusive W Muon Selection

After passing the three levels of trigger requirements, the following event selection cuts are applied to the inclusive muon data sample:

- The event must belong to a good run.
- $p_T^\mu \geq 20$ GeV,

where p_T^μ is the beam-constrained transverse momentum of the muon (determined by a fit to the CTC hits, constrained by the beam line).

- The muon must be fiducial and central (pseudorapidity $|\eta_{\text{lab}}^\mu| \leq 1$).

- $\text{ISO}(0.4) \equiv E_{\Delta R=0.4}^{\text{Excess}}/p_T^\mu < 0.1$, where $E_{\Delta R=0.4}^{\text{Excess}}$ is the excess transverse energy in a cone of size $\Delta R = \sqrt{(\Delta\phi_{\text{lab}})^2 + (\Delta\eta_{\text{lab}})^2}$ centered on the direction of the muon.

- $E_{\text{HAD}} \leq 6$ GeV,

where E_{HAD} is the energy deposited in the hadron calorimeter tower traversed by the muon.

- $E_{\text{EM}} \leq 2$ GeV,

where E_{EM} is the energy deposited in the electromagnetic calorimeter tower traversed by the muon.

- $|\Delta X_{\text{CMU}}| < 2$ cm, $|\Delta X_{\text{CMP}}| < 5$ cm, $|\Delta X_{\text{CMX}}| < 5$ cm,

where ΔX_{CMU} , ΔX_{CMP} and ΔX_{CMX} are the differences between the x position of the stub in the muon chambers and the extrapolation of the CTC track to these muon chambers.

- $|Z_{\text{VTX}}| \leq 60$ cm.

- The event must pass the cosmic ray filter.

- The impact parameter must be $|d_0| \leq 0.2$ cm.

- $|Z_0 - Z_{\text{VTX}}| \leq 5$ cm,

where Z_0 is the z -position of the muon track. This cut, combined with the previous two, significantly reduces the cosmic muon background.

We next apply the following cuts, as in the electron case:

- $\cancel{E}_T > 20$ GeV.

- $M_T^W > 40$ GeV.

- The event must not be consistent with a Z decaying into two observed leptons, or a Z in which one of the decay tracks has not been identified.

The 38601 events passing these cuts constitute our inclusive W muon data sample [Run Ia (CMUP): 4441 events, Run Ia (CMNP): 955 events, Run Ib (CMUP): 20527 events, Run Ib (CMNP): 3273 events, and Run Ib (CMX): 9405 events], corresponding to an integrated luminosity of 107 pb^{-1} [Run Ia (CMUP): $18.33 \pm 0.66 \text{ pb}^{-1}$, Run Ia (CMNP): $19.22 \pm 0.69 \text{ pb}^{-1}$, Run Ib (CMUP): $88.35 \pm 3.62 \text{ pb}^{-1}$, Run Ib (CMNP): $89.20 \pm 3.66 \text{ pb}^{-1}$ and Run Ib (CMX): $88.98 \pm 3.65 \text{ pb}^{-1}$].

3. Inclusive W +jet Event Selection

Our final analysis dataset consists of those W events which include at least one jet with $E_T^{\text{jet}} > 15$ GeV, $|\eta_{\text{lab}}^{\text{jet}}| < 2.4$, and $\Delta R_{i-j}^{\text{lab}} > 0.7$, where $\Delta R_{i-j}^{\text{lab}} \equiv$

TABLE I: The electron (N_e) and muon (N_μ) CDF data event yields for inclusive W +jet production. The muon event yields are lower relative to the electron event yields, due to lower muon efficiencies and acceptances.

| Data event yields for inclusive W +jet production | | |
|---|-------|---------|
| p_T^W (GeV) | N_e | N_μ |
| 15–25 | 5166 | 2821 |
| 25–35 | 3601 | 1869 |
| 35–65 | 3285 | 1880 |
| 65–105 | 624 | 371 |

$\sqrt{(\Delta\eta_{l-j}^{\text{lab}})^2 + (\Delta\phi_{l-j}^{\text{lab}})^2}$, and $\Delta\eta_{l-j}^{\text{lab}}$ and $\Delta\phi_{l-j}^{\text{lab}}$ are the differences in pseudorapidity and polar angle between the charged lepton and the jet in the laboratory frame. The results of the analysis pertain to the W^- boson. All W^+ bosons in the sample are CP transformed to be treated as W^- bosons [13].

These requirements leave 12676 electron W +jet events and 6941 muon W +jet events, with $15 < p_T^W < 105$ GeV, where p_T^W is the transverse momentum of the W boson, defined as the vector sum of the \cancel{E}_T and charged lepton transverse momentum. The data event yields for the four p_T^W bins ($15 < p_T^W < 25$ GeV, $25 < p_T^W < 35$ GeV, $35 < p_T^W < 65$ GeV, and $65 < p_T^W < 105$ GeV) are presented in Table I.

The actual number of events is not of critical importance for us, because we are interested in the shape of the distributions and not the absolute event yields. We thus analyze the distributions normalized to unity. We will come back to the actual event yields after the inclusion of the background (Section V) and systematic uncertainties (Section IX).

III. THE MONTE CARLO SIMULATION

A. The DYRAD Monte Carlo event generator

DYRAD [14] is the next-to-leading order W +jet event generator used to establish the SM prediction. We include the “1-loop” processes, since these affect the next-to-leading order angular coefficients and more completely simulate the events we study. This generator is of order α_s^2 in QCD, generating up to two jets passing the minimal requirement of $E_T^{\text{jet}} > 10$ GeV if the Feynman diagram does not contain any gluon loops, and generates up to one jet with the same requirement if a gluon loop is present in the Feynman diagram. As a result, DYRAD does not appropriately model events with more than two jets. These extra jets in the data occupy low and high values of the azimuthal angle ϕ in our CS frame. We are careful not to bias our measurement due to this effect (see Section VIII). The jet transverse energy cut of 10 GeV is required because the theoretical calculations are unreliable for small jet transverse energies due to infrared and collinear divergencies. A jet-jet angular

separation cut of greater than 0.7 in $\eta_{\text{lab}}-\phi_{\text{lab}}$ space is imposed, which is important for the definition of a jet. No additional kinematic cuts for the jet, charged lepton, and neutrino are required in order to obtain a reliable theoretical prediction of the angular distribution of the W . The cross section for inclusive W +jet production calculated up to order α_s^2 is 722.51 ± 3.89 pb for W^- and W^+ bosons combined. This simulation uses $Q^2 = (M_W^{\text{pole}})^2$, where $M_W^{\text{pole}} = 80.3$ GeV is the pole mass of the W , CTEQ4M($\Lambda = 0.3$ GeV) parton distribution functions [15], and 0.7-cone jets in $\eta_{\text{lab}}-\phi_{\text{lab}}$ space.

In order to obtain smooth SM kinematic distributions up to $p_T^W = 100$ GeV, and especially smooth $\cos\theta$ vs. ϕ distributions for different p_T^W regions, we generated a large sample of DYRAD events (~ 250 M). This Monte Carlo event sample size was required since events with negative weights, corresponding to the gluon loop matrix elements, produce significant fluctuations in the kinematic distributions with limited statistics. The DYRAD simulation allows us to establish the SM prediction for the ϕ distribution of the charged lepton and the predictions for the angular coefficients and helicity cross sections of the W up to order α_s^2 [7]. The expected ϕ distributions for four p_T^W bins are shown in Figure 2. For zero p_T^W we expect a flat distribution, whereas the QCD effects at higher p_T^W result in two minima. In order to simulate the detector response, we pass the generator events through the fast Monte Carlo detector simulator, described in the next section.

B. The fast Monte Carlo detector simulation

The fast Monte Carlo (FMC) CDF detector simulation includes the detailed geometry of the detector, geometrical and kinematic acceptances of all subdetectors, detector resolution effects parameterized using gaussians obtained explicitly from data, detailed magnetic field map, and multiple Coulomb scattering effects. The integrated luminosities, lepton identification and trigger efficiencies, and all experimental cuts imposed on the W , leptons, \cancel{E}_T , and jets are incorporated. The effect of the underlying event, caused by interacting spectator quarks, is also included. The FMC program receives the particle four-momenta for each generated DYRAD event along with the next-to-leading order cross section prediction from DYRAD (which includes gluon loop effects) and produces kinematic distributions smeared by detector resolution, and sculpted by geometrical and kinematic acceptances and efficiencies. The FMC also reports event yield predictions. The FMC successfully reproduces the kinematic features of inclusive W and Z boson production, as well as the features of vector boson production in association with a jet [8, 16].

For the W +jet data, we additionally require at least one “good” jet ($E_T^{\text{jet}} > 15$ GeV and $|\eta_{\text{lab}}^{\text{jet}}| < 2.4$) that also passes the $\Delta R_{l-j}^{\text{lab}} > 0.7$ cut, where $\Delta R_{l-j}^{\text{lab}}$ is the opening angle in $\eta_{\text{lab}}-\phi_{\text{lab}}$ space between the lepton and the




FIG. 2: The theoretical charged lepton ϕ distribution in the Collins-Soper W rest-frame for the four p_T^W regions, as generated by DYRAD. The distributions are normalized to unity.

TABLE II: The electron (N_e) and muon (N_μ) FMC event yields for inclusive W +jet production up to order α_s^2 . Contributions from backgrounds are not included. The uncertainties in the event yield predictions are dominated by the uncertainties associated with integrated luminosities, lepton identification and trigger efficiencies and the DYRAD prediction of the W +jet production cross section. Systematic uncertainties associated with PDF choice and Q^2 scale variation are not included.

| FMC event yields for inclusive W +jet production | | |
|--|----------------|----------------|
| p_T^W (GeV) | N_e | N_μ |
| 15–25 | 3867 ± 137 | 2027 ± 102 |
| 25–35 | 2632 ± 93 | 1384 ± 66 |
| 35–65 | 2474 ± 87 | 1314 ± 67 |
| 65–105 | 518 ± 18 | 279 ± 14 |

leading “good” jet. The FMC event yields for inclusive W +jet production up to order α_s^2 are presented in Table II. The Parton Distribution Functions (PDF) systematics and the renormalization and factorization scale (Q^2) systematics will be included in Section VI.

The FMC detector simulation, along with DYRAD, shows how the acceptances and efficiencies of the detector and the analysis cuts affect the ϕ distributions that are experimentally observed. Figure 3 shows the expected measurement of the ϕ distributions for the electron dataset (the muon distributions are almost identical) for the four p_T^W bins. The effects of the acceptances and efficiencies are significant; instead of two minima we




FIG. 3: The expected CDF electron ϕ distribution in the Collins-Soper W rest-frame for the four p_T^W regions, after experimental cuts and detector smearing, as generated by the FMC. The distributions are normalized to unity. The muon distributions are almost identical.

observe two maxima. The main reason for this is the charged lepton and neutrino p_T cuts, which limit the allowed $(\cos\theta, \phi)$ phase space considerably. The FMC plots are normalized to the FMC signal event yields, and all experimental cuts have been applied.

IV. ACCEPTANCES AND EFFICIENCIES

The lepton identification and trigger efficiencies are measured by using the leptons from CDF Run Ia and Ib Z data and by studying random cone distributions of leptonic W and Z decay Run Ia and Ib data samples. The kinematic and geometrical acceptances are calculated using the DYRAD event generator, which produces the SM prediction, and the FMC detector simulation, which produces the CDF experimental expectation.

We are especially interested in the product of overall acceptance times efficiency (ae) as a function of $(\cos\theta, \phi)$ associated with each of the four p_T^W bins. We create 2-dimensional histograms of $\cos\theta$ vs. ϕ for each of the four p_T^W bins, using the DYRAD simulation. This procedure is repeated after the events pass the FMC simulation, where the appropriate mixture of Run Ia and Run Ib W leptons is used, based on FMC event yield predictions for all subdetectors.

The resulting plots are shown in Figures 4, 5, and 6, for DYRAD, FMC-electrons, and FMC-muons respec-




FIG. 4: The $\cos\theta$ vs. ϕ phase space for the four p_T^W bins, for the DYRAD generator simulation (arbitrary units).




FIG. 6: The $\cos\theta$ vs. ϕ phase space for the four p_T^W bins, for the muon FMC signal simulation (arbitrary units).




FIG. 5: The $\cos\theta$ vs. ϕ phase space for the four p_T^W bins, for the electron FMC signal simulation (arbitrary units).




FIG. 7: Acceptance times efficiency for the electrons as a function of $\cos\theta$ and ϕ in the Collins-Soper frame.

tively. We subsequently divide the FMC 2-dimensional histograms by the corresponding DYRAD ones, producing the 2-dimensional differential acceptance times efficiency $ae(\cos\theta, \phi)$ of Figures 7 and 8, for electrons and

muons respectively. The overall acceptance times efficiency is higher for the electrons. These $ae(\cos\theta, \phi)$ values are used for the $\cos\theta$ -integration of the cross section, as described in Section VII.




FIG. 8: Acceptance times efficiency for the muons as a function of $\cos\theta$ and ϕ in the Collins-Soper frame.

V. BACKGROUND ESTIMATION

The main sources of background in the $(W \rightarrow e\nu_e)+\text{jet}$ and $(W \rightarrow \mu\nu_\mu)+\text{jet}$ processes are $Z+\text{jets}$ events where the Z is misidentified as a W (“one-legged” Z), $(W \rightarrow \tau\nu_\tau)+\text{jet}$ events, and QCD background resulting from jet misidentification. The $W + \gamma$ background is only a few events with run and event numbers the same as the $W + \gamma$ events of [16]; it is treated as a systematic uncertainty, which also gives an indication of the radiative effects in the measurement. This uncertainty is very small (see Section IX).

A small background contribution arises from $t\bar{t}$ production, where one of the produced W bosons decays leptonically and the other W boson decays hadronically to jets. This background is estimated to be 30 ± 7 events for the electron sample [17] and 16 ± 3 events for the muon sample, a 0.3% effect. An equally small background is the $Z \rightarrow \tau^+\tau^-$ production, where one of the tau leptons decays hadronically and the other one leptonically. This background is estimated to be 47 ± 1 events in the electron sample [17], and 25 ± 1 events in the muon sample, a 0.5% effect. To demonstrate the insignificance of the $t\bar{t}$ and $Z \rightarrow \tau^+\tau^-$ backgrounds, we perform our analysis including the charged lepton ϕ distribution for these background events, in several possible shapes, for the four p_T^W bins. The resulting change in the extracted values of the angular coefficients is negligible compared to our systematic and statistical uncertainties. Thus, we ignore the backgrounds associated with $t\bar{t}$ production and

$Z \rightarrow \tau^+\tau^-$ decays.

Finally, the cosmic ray background in the muon $W+\text{jet}$ datasets is estimated to be significantly less than 0.1%, and is therefore neglected.

A. One-legged Z background

To study this background we generate a DYRAD sample of $Z+\text{jet}$ events and pass it through the FMC Monte Carlo simulation and the subsequent analysis program. This predicts how many Z bosons are misidentified as W bosons. In these cases, the Z bosons satisfy all kinematic and lepton identification cuts for W bosons, but one of their decay leptons, or *legs*, is undetected. The DYRAD cross section for $Z+\text{jet}$ up to order α_s^2 is 68.21 ± 0.37 pb. For this DYRAD simulation we used $Q^2 = (M_Z^{\text{pole}})^2 = 91.2$ GeV, the CTEQ4M($\Lambda = 0.3$ GeV) parton distribution functions, 0.7-cone jets, jet-jet angular separation greater than 0.7 in $\eta_{\text{lab}}-\phi_{\text{lab}}$ space, and $E_T^j > 10$ GeV. At the FMC level, we impose our usual W boson event selection cuts and additionally require at least one “good” jet ($E_T^{\text{jet}} > 15$ GeV and $|\eta_{\text{lab}}^{\text{jet}}| < 2.4$) that also passes the $\Delta R_{l-j}^{\text{lab}} > 0.7$ cut. These results are summarized in Table III. Overall we expect 123 ± 5 electron one-legged- $Z+\text{jet}$ events and 337 ± 18 muon one-legged- $Z+\text{jet}$ events passing the $W+\text{jet}$ cuts, without applying any cut on the W transverse momentum. Comparing these numbers to the FMC event yields for $W+\text{jet}$, the one-legged- $Z+\text{jet}$ background is $(1.14 \pm 0.06)\%$ for the electron $W+\text{jet}$ and $(5.90 \pm 0.43)\%$ for the muon $W+\text{jet}$ sample. This background is higher for the muon sample, because of the limited coverage of the muon chambers, which is responsible for higher yields of one-legged muon Z bosons.

To examine how this background affects the $W+\text{jet}$ lepton ϕ distribution, we plot the ϕ distribution for the leptons from these processes for the four p_T^W bins (Figures 9 and 10). We see that the same pattern of two maxima at $\frac{\pi}{2}$ and $\frac{3\pi}{2}$ is present. The background plots are normalized to the expected event yields from the FMC, multiplied by a factor of five (to make them visible), and superimposed on the signal FMC distributions, normalized to the signal FMC event yields. We include the one-legged Z FMC ϕ distribution in the complete theoretical prediction of the ϕ distributions, in order to correctly extract the angular coefficients.

B. $(W \rightarrow \tau\nu)+\text{jet}$ background

If the W boson decays to a τ that subsequently decays leptonically, the three final neutrinos contribute to the \cancel{E}_T , which is incorrectly associated with a single neutrino. The signal of one charged lepton along with the \cancel{E}_T mimics that of a W directly decaying to the charged lepton. Most of the tau background is removed when we utilize the fact that the charged lepton and \cancel{E}_T coming




FIG. 9: Electron ϕ distributions for the four p_T^W bins for electron W +jet FMC events (solid histogram) and for Z +jet FMC background (multiplied by 5), where one of the electrons from the Z decay is undetected and the other one passes the detection and analysis requirements (dashed histogram). The histograms are normalized to the electron FMC signal event yields.

TABLE III: Monte Carlo background estimation of the number of electron and muon one-legged Z +jet events. The background fractions are calculated with respect to the FMC W +jet event yields.

| One-legged Z +jet background | | | | |
|--------------------------------|------------|--------------------|-------------|--------------------|
| p_T^W (GeV) | N_e | Fraction | N_μ | Fraction |
| 15–25 | 47 ± 2 | $1.22 \pm 0.07 \%$ | 127 ± 7 | $6.26 \pm 0.47 \%$ |
| 25–35 | 30 ± 1 | $1.14 \pm 0.05 \%$ | 82 ± 4 | $5.92 \pm 0.40 \%$ |
| 35–65 | 25 ± 1 | $1.01 \pm 0.05 \%$ | 72 ± 4 | $5.48 \pm 0.41 \%$ |
| 65–105 | 5 ± 0 | $0.96 \pm 0.03 \%$ | 12 ± 1 | $4.30 \pm 0.42 \%$ |

from the τ decay are soft. As a result, the W transverse mass in the τ events is significantly smaller than that in the electron or muon events. By applying the p_T cuts for the leptons and the W transverse mass cut, we remove 92% of the tau W +jet events at the DYRAD generator level.

To study the remaining tau background we start with a tau W +jet DYRAD sample ($Q^2 = (M_W^{\text{pole}})^2$, CTEQ4M($\Lambda = 0.3$ GeV) parton distribution function and 0.7-cone jets in $\eta_{\text{lab}}-\phi_{\text{lab}}$ space), and we let the tau decay to an electron or a muon. We then vector-sum the three neutrinos resulting from the W and tau decays to form a single \cancel{E}_T . Subsequently, we pass the events through the FMC detector simulator to see how many




FIG. 10: Muon ϕ distributions for the four p_T^W bins for muon W +jet FMC events (solid histogram) and for Z +jet FMC background (multiplied by 5), where one of the muons from the Z decay is undetected and the other one passes the detection and analysis requirements (dashed histogram). The histograms are normalized to the muon FMC signal event yields.

events pass the W +jet cuts after they are weighted by the detector acceptances and efficiencies. The branching ratios for the tau decays we use are 17.83 % for electrons and 17.37 % for muons [18]. At the FMC level, we require at least one “good” jet ($E_T^{\text{jet}} > 15$ GeV and $|\eta_{\text{lab}}^{\text{jet}}| < 2.4$) that also passes the $\Delta R_{l-j}^{\text{lab}} > 0.7$ cut. The tau background results are presented in Table IV. Overall we expect 247 ± 9 tau electrons and 130 ± 7 tau muons to infiltrate the W +jet samples, without applying any cut on the W transverse momentum. Comparing these numbers to the FMC event yields for the electron and muon W +jet samples, the tau background is $(2.28 \pm 0.12)\%$ for the electron W +jet sample, and $(2.28 \pm 0.17)\%$ for the muon W +jet sample.

To see how this background affects the W +jet lepton ϕ distribution, we plot the ϕ distribution for the leptons resulting from leptonic tau decays in W +jets events for the four p_T^W bins (Figures 11 and 12). We see that the pattern of two maxima at $\frac{\pi}{2}$ and $\frac{3\pi}{2}$ is again present. The background plots are normalized to the expected event yields from the FMC, multiplied by a factor of five, and superimposed on the signal FMC distributions, normalized to the signal FMC event yields. We include the τ -background FMC ϕ distribution in the complete theoretical prediction of the ϕ distributions, in order to correctly extract the angular coefficients.




FIG. 11: Electron ϕ distributions for the four p_T^W bins for electron W +jet FMC events (solid histogram) and for tau W +jet FMC background (multiplied by 5), where the tau decays to an electron (dashed histogram). The histograms are normalized to the FMC signal event yields.

TABLE IV: Monte Carlo background estimation of the number of electron and muon W +jet events, where the W decays to a tau and the electron or muon is the decay product of the tau. The fractions of the backgrounds are calculated with respect to the FMC W +jet event yields.

| $(W \rightarrow \tau \bar{\nu}_\tau \rightarrow \bar{\nu}_\tau \nu_\tau \bar{\nu}_{e/\mu} e/\mu) + \text{jet background}$ | | | | |
|---|------------|--------------------|------------|--------------------|
| p_T^W (GeV) | N_e | Fraction | N_μ | Fraction |
| 15–25 | 86 ± 3 | $2.22 \pm 0.11 \%$ | 45 ± 2 | $2.22 \pm 0.15 \%$ |
| 25–35 | 57 ± 2 | $2.16 \pm 0.10 \%$ | 30 ± 2 | $2.17 \pm 0.18 \%$ |
| 35–65 | 56 ± 2 | $2.26 \pm 0.11 \%$ | 30 ± 2 | $2.28 \pm 0.19 \%$ |
| 65–105 | 15 ± 1 | $2.89 \pm 0.22 \%$ | 8 ± 0 | $2.87 \pm 0.14 \%$ |

C. QCD background

The QCD background in the case of inclusive W production and decay consists predominantly of dijet events, where one of the jets is misidentified as a lepton and the other one is not detected, resulting in the creation of \cancel{E}_T . In the W +jet case, the QCD background is multi-jet events, where one of the jets is detected, one is lost or mismeasured (resulting in \cancel{E}_T) and one is misidentified as a charged lepton to erroneously reconstruct a W . The number and distribution of QCD background events in the four p_T^W bins are determined from the Run Ia and Run Ib CDF data.

To measure the expected number of QCD background events in our data samples we look at leptons with isola-




FIG. 12: Muon ϕ distributions for the four p_T^W bins for muon W +jet FMC events (solid histogram) and for tau W +jet FMC background (multiplied by 5), where the tau decays to a muon (dashed histogram). The histograms are normalized to the FMC signal event yields.

tion (ISO), defined in Section II, greater than 0.2. Our signal is in the ISO < 0.1 region and most of the events with lepton ISO > 0.2, but not all of them, are QCD background events. The upper histogram of Figure 13 shows the isolation distribution of the electrons from W +jet events, for the first p_T^W bin. When plotted on a semi-log scale, the ISO < 0.1 and the ISO > 0.2 regions can be approximated with two straight lines. The technique we use extrapolates the ISO > 0.2 line into the ISO < 0.1 signal region to calculate its integral and obtain the number of events in the signal region, using the assumption that the QCD background shape is not altered in that region. This method would give us the true number of QCD background events, if the ISO > 0.2 region was filled exclusively with QCD events. In reality, only a fraction of these events are true QCD background, the rest being W +jet events. Since we expect to have some W +jet events in the region of lepton isolation from 0.1 to 0.2, we fit the area above 0.2 with a straight line (in the semi-log histogram), which describes the QCD background. We also fit five continuous regions of lepton isolation, around the central region of ISO=0.20 to ISO=0.65 (namely 0.15-0.65, 0.25-0.65, 0.15-0.60, 0.20-0.65, and 0.25-0.70) to obtain a systematic uncertainty for this procedure.

Since not all of the extrapolated region is QCD background, we obtain a measurement of the percentage of the true QCD background in the electron W +jet sample

TABLE V: The linear least-squares fit parameterization of the $\Delta\phi_{l-j}^{\text{lab}}$ distribution for $\text{ISO} > 0.2$ electron W +jet events (second column) allows us to estimate the number of $\text{ISO} > 0.2$ W +jet events in the $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ region. The integral of this line, divided by the bin width ($\pi/30$) of the $\Delta\phi_{l-j}^{\text{lab}}$ histogram, is the number of W +jet events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ shown in the third column. These events are subtracted from the total number of (W +jet + QCD) background events in the $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ region (fourth column). The result is divided by the total number of events in the $\text{ISO} > 0.2$ region, to obtain an estimate of the fraction of true QCD background events (fifth column).

| p_T^W (GeV) | Fit parameterization of electron W +jet events with $\text{ISO} > 0.2$ | Electron W +jet events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ | (W +jet)+QCD events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ | Fraction of true QCD events |
|---------------|--|--|--|-----------------------------|
| 15–25 | $2.21 \times \Delta\phi_{l-j}^{\text{lab}} - 0.05$ | 37.9 | 257 | $0.66 = (257 - 37.9) / 332$ |
| 25–35 | $0.70 \times \Delta\phi_{l-j}^{\text{lab}} + 0.78$ | 16.9 | 98 | $0.58 = (98 - 16.9) / 141$ |
| 35–65 | $1.04 \times \Delta\phi_{l-j}^{\text{lab}} + 0.45$ | 20.6 | 49 | $0.31 = (49 - 20.6) / 91$ |
| 65–105 | $0.17 \times \Delta\phi_{l-j}^{\text{lab}} + 0.73$ | 7.5 | 10 | $0.13 = (10 - 7.5) / 20$ |

TABLE VI: The number of electron W +jet events, the number of QCD background events before correction and their percentage in the signal region, and the fraction of true QCD background events and their percentage in the signal region, for the four p_T^W bins (see text for details).

| p_T^W (GeV) | Number of electron W +jet | QCD events before correction | Percentage of QCD before correction | Fraction of true QCD events | Percentage of QCD background |
|---------------|-----------------------------|------------------------------|-------------------------------------|-----------------------------|------------------------------|
| 15–25 | 5166 | 423^{+77}_{-42} | $8.18^{+1.11}_{-0.81}\%$ | $0.66^{+0.04}_{-0.04}$ | $5.40^{+0.80}_{-0.63}\%$ |
| 25–35 | 3601 | 353^{+26}_{-148} | $9.80^{+0.74}_{-4.17}\%$ | $0.58^{+0.07}_{-0.08}$ | $5.68^{+0.81}_{-2.50}\%$ |
| 35–65 | 3285 | 54^{+151}_{-25} | $1.64^{+4.60}_{-0.75}\%$ | $0.31^{+0.13}_{-0.13}$ | $0.51^{+1.44}_{-0.32}\%$ |
| 65–105 | 624 | 14^{+8}_{-3} | $2.24^{+1.24}_{-0.27}\%$ | $0.13^{+0.87}_{-0.13}$ | $0.29^{+1.96}_{-0.29}\%$ |




FIG. 13: Isolation of the electron in W +jet events (upper histogram) and the difference in the azimuthal angles of the electron and the jet for high isolation events (lower histogram). These histograms are used to estimate the QCD background in the electron W +jet data, for the $15 \leq p_T^W \leq 25$ GeV bin (see text). Corresponding histograms are used for the estimation of the QCD background in the three other electron W +jet p_T^W bins.

above electron isolation of 0.1, by making a histogram of $\Delta\phi_{l-j}^{\text{lab}}$ for the events with $\text{ISO} > 0.2$, where $\Delta\phi_{l-j}^{\text{lab}}$ is the difference in the ϕ angle between the electron and the highest- E_T jet, with no other requirements for that jet. We expect the $\Delta\phi_{l-j}^{\text{lab}}$ distribution to be almost flat for the W +jet events, because no correlation exists between the jet and the lepton ϕ directions. In reality, this distribution decreases at low $\Delta\phi_{l-j}^{\text{lab}}$, due to the application of the lepton isolation cut in our data. For QCD background, we expect the $\Delta\phi_{l-j}^{\text{lab}}$ between the highest E_T jet and the jet resembling the lepton to peak at π . The lower histogram of Figure 13 shows the $\Delta\phi_{l-j}^{\text{lab}}$ for the events with lepton isolation greater than 0.2 for electron W +jet events and for the first p_T^W bin. We fit the region $\Delta\phi_{l-j}^{\text{lab}} \leq 2.5$ (W +jet contribution) with a straight line. The region of the histogram $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ above that line corresponds to true QCD background. By dividing this part of the histogram by the total number of events with $\text{ISO} > 0.2$, we determine the true fraction of QCD background in the $\text{ISO} > 0.2$ region. We expect the same fraction to be valid in the signal region ($\text{ISO} < 0.1$). Therefore, the number of true QCD background events is obtained by multiplying the number of $\text{ISO} < 0.1$ events (as obtained by extrapolating the $\text{ISO} > 0.2$ line into the signal region of the lepton isolation plot) by the QCD background fraction obtained from the $\Delta\phi_{l-j}^{\text{lab}}$ plot. The procedure is repeated for the four p_T^W bins. Table V shows the extracted fraction of QCD background in the $\text{ISO} > 0.2$ region for the four p_T^W bins. The electron W +jet QCD background results are presented in Table VI.

In the study of the QCD background in the muon sam-




FIG. 14: Isolation of the muon in W +jet events (upper histogram) and the difference in the azimuthal angles of the muon and the jet for high isolation events (lower histogram). These histograms are used to estimate the QCD background in the muon W +jet data, for the $15 \leq p_T^W \leq 25$ GeV bin (see text). Corresponding histograms are used for the estimation of the QCD background in the three other muon W +jet p_T^W bins. At low $\Delta\phi_{l-j}^{\text{lab}}$ the distribution increases, due to bremsstrahlung processes associated with residual muon Z +jet background in the muon W +jet sample. These events disappear if we apply the `zmuo_veto` cut. (see text).

ple we face a new problem. We originally apply a cut to muon W +jet data (`zmuo_veto`) in order to remove events that are consistent with the production of a Z boson, where one of the muons is non-isolated because it fails one (and only one) of the following cuts:

- The muon isolation cut $\text{ISO} < 0.1$
- The electromagnetic calorimeter cut $E^{\text{EM}} < 2$ GeV
- The hadron calorimeter cut $E^{\text{HAD}} < 6$ GeV

These dimuon events are true Z bosons that look like W bosons because one muon does not pass one of the above cuts due to inner bremsstrahlung or bremsstrahlung in the electromagnetic or hadronic calorimeters. The `zmuo_veto` cut mainly affects the tail of the muon isolation distribution ($\text{ISO} > 0.2$) and causes us to underestimate the QCD background, since we use that tail to estimate it. Therefore, for the muon W +jet samples, for the purposes of determination of QCD background, we neglect this cut, in order to remove this bias at high muon isolation ($\text{ISO} > 0.2$) and make the transition from the

low to high isolation smooth. Some of the muon Z background is thus counted as QCD background; however we do not expect it to radically affect our QCD background estimation. In the isolation method we fit the background starting from $\text{ISO}=0.17$ to $\text{ISO}=0.40$, to increase the statistical significance of our estimation. We also fit five continuous regions of lepton isolation, around the central region of $\text{ISO}=0.17$ to $\text{ISO}=0.40$ (namely 0.16-0.40, 0.18-0.40, 0.16-0.35, 0.17-0.40, and 0.18-0.45) to obtain a systematic uncertainty for this procedure. The upper histogram of Figure 14 shows the isolation distribution and fits for the muon W +jet events and for the first p_T^W bin.

We obtain a measurement of the percentage of the true QCD background in the muon W +jet sample above muon isolation of 0.2, by making a histogram of $\Delta\phi_{l-j}^{\text{lab}}$ for the events with $\text{ISO} > 0.2$, where $\Delta\phi_{l-j}^{\text{lab}}$ is the difference in the ϕ angle between the muon and the highest- E_T jet, with no other requirements for that jet. The lower histogram of Figure 14 shows $\Delta\phi_{l-j}^{\text{lab}}$ for the events with isolation greater than 0.2 for muon W +jet events, for the first p_T^W bin. The peak in the $\Delta\phi_{l-j}^{\text{lab}} = 0$ region is due to the muon bremsstrahlung processes that are not suppressed after we relax the `zmuo_veto` cut. We ignore these events when we fit to the straight line describing the W +jet events with high isolation muons. Table VII shows the extracted fraction of QCD background in the $\text{ISO} > 0.2$ region for the four p_T^W bins. The muon W +jet QCD background results are presented in Table VIII. For the highest muon p_T^W bin the predicted number of true W +jet events is greater than the total number of events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$, which results in a fraction of true QCD background events above $\text{ISO} > 0.2$ equal to zero.

After we calculate the percentage of the QCD background in the signal region, we multiply it by the CDF W +jet event yields to obtain the absolute prediction of the number of QCD background events in each of the four p_T^W bins, for both and electron and muon W +jet data. The results are presented in Table IX.

To complete the study of the QCD background we need to estimate its shape to properly include this background in the Standard Model prediction of the lepton ϕ distribution in the CS frame, for each of the four p_T^W bins. We plot ϕ for the events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ for the electrons and muon datasets, as shown in Figures 15 and 16, respectively. We fit the distributions to the sum of two Gaussians and two straight lines. For the last p_T^W bin of the electrons and the last two p_T^W bins of the muons, there are not enough statistics for the fit, so we use the total distributions (for $15 \leq p_T^W \leq 105$ GeV) normalized to the number of events for those high p_T^W bins. We do not expect the shape of the QCD background to be significantly altered with increasing p_T^W . We assume that these distributions are the same as the ones in the signal region ($\text{ISO} < 0.1$) after they are properly normalized. We use these distributions to add the QCD background to the Standard Model prediction, after they are normal-

TABLE VII: The linear least-squares fit parameterization of the $\Delta\phi_{l-j}^{\text{lab}}$ distribution for $\text{ISO} > 0.2$ muon $W+\text{jet}$ events (second column) allows us to estimate the number of $\text{ISO} > 0.2$ $W+\text{jet}$ events in the $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ region. The integral of this line, divided by the bin width ($\pi/30$) of the $\Delta\phi_{l-j}^{\text{lab}}$ histogram, is the number of $W+\text{jet}$ events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ shown in the third column. These events are subtracted from the total number of ($W+\text{jet} + \text{QCD}$) background events in the $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ region (fourth column). The result is divided by the total number of events in the $\text{ISO} > 0.2$ region, to obtain an estimate of the fraction of true QCD background events (fifth column).

| p_T^W (GeV) | Fit parameterization of muon $W+\text{jet}$ events with $\text{ISO} > 0.2$ | Muon $W+\text{jet}$ events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ | ($W+\text{jet}$)+QCD events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ | Fraction of true QCD events |
|---------------|--|--|---|-----------------------------|
| 15–25 | $0.29 \times \Delta\phi_{l-j}^{\text{lab}} + 1.31$ | 13 | 164 | $0.52 = (164-13)/288$ |
| 25–35 | $0.46 \times \Delta\phi_{l-j}^{\text{lab}} + 0.79$ | 12.8 | 69 | $0.40 = (69-12.8)/140$ |
| 35–65 | $0.25 \times \Delta\phi_{l-j}^{\text{lab}} + 1.02$ | 10.5 | 19 | $0.14 = (19-10.5)/61$ |
| 65–105 | $0 \times \Delta\phi_{l-j}^{\text{lab}} + 1$ | 6.1 | 1 | $0 = 0/6$ |

TABLE VIII: The number of muon $W+\text{jet}$ events without the application of the `zmuo_veto` cut, the number of QCD background events before correction and their percentage in the signal region, and the fraction of true QCD background events and their percentage in the signal region, for the four p_T^W bins (see text for details).

| p_T^W (GeV) | Number of muon $W+\text{jet}$ | QCD events before correction | Percentage of QCD before correction | Fraction of true QCD events | Percentage of QCD background |
|---------------|-------------------------------|------------------------------|-------------------------------------|-----------------------------|------------------------------|
| 15–25 | 2779 | 280^{+102}_{-25} | $10.07^{+3.68\%}_{-0.90\%}$ | $0.52^{+0.05}_{-0.04}$ | $5.24^{+1.98\%}_{-0.62\%}$ |
| 25–35 | 1943 | 103^{+129}_{-4} | $5.30^{+6.64\%}_{-0.21\%}$ | $0.40^{+0.08}_{-0.08}$ | $2.12^{+2.63\%}_{-0.43\%}$ |
| 35–65 | 2002 | 139^{+50}_{-21} | $6.94^{+2.50\%}_{-1.05\%}$ | $0.14^{+0.22}_{-0.14}$ | $0.97^{+1.37\%}_{-0.97\%}$ |
| 65–105 | 389 | 11^{+0}_{-0} | $2.70^{+0.00\%}_{-0.00\%}$ | 0^{+1}_{-0} | $0^{+2.7\%}_{-0\%}$ |




FIG. 15: The Collins-Soper ϕ distribution of electrons from $W+\text{jet}$ events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ for each of the four p_T^W bins. These events are predominantly QCD background events. We fit the distribution of the first three p_T^W bins with two Gaussians on top of two straight lines. For the highest p_T^W bin we use the distribution of the total QCD background, normalized to the number of the QCD events in this bin. (see text for details).




FIG. 16: The Collins-Soper ϕ distribution of muons from $W+\text{jet}$ events with $\text{ISO} > 0.2$ and $\Delta\phi_{l-j}^{\text{lab}} > 2.5$ for each of the four p_T^W bins. These events are predominantly QCD background events. We fit the distribution of the first two p_T^W bins with two Gaussians on top of two straight lines. For the two highest p_T^W bins we use the distribution of the total background, normalized to the number of the QCD events in those bins.

TABLE IX: QCD background estimation for the electron and muon W +jet events. The fractions of the backgrounds are calculated with respect to the CDF Data W +jet events.

| QCD background | | | | | |
|----------------|-------------------|--------------------------|-------------------|--------------------------|--|
| p_T^W (GeV) | N_e | Fraction | N_μ | Fraction | |
| 15–25 | 279^{+41}_{-33} | $5.40^{+0.80}_{-0.63}$ % | 148^{+56}_{-17} | $5.24^{+1.98}_{-0.62}$ % | |
| 25–35 | 205^{+29}_{-90} | $5.68^{+0.81}_{-2.50}$ % | 40^{+50}_{-8} | $2.12^{+2.69}_{-0.43}$ % | |
| 35–65 | 17^{+47}_{-11} | $0.51^{+1.44}_{-0.32}$ % | 18^{+30}_{-18} | $0.97^{+1.57}_{-0.97}$ % | |
| 65–105 | 2^{+12}_{-2} | $0.29^{+1.36}_{-0.29}$ % | 0^{+10}_{-0} | $0^{+2.7}_{-0}$ % | |

TABLE X: Summary of electron W +jet backgrounds. The background fractions are calculated with respect to the FMC signal event yields for the electroweak backgrounds and with respect to the data for the QCD background.

| Electron W +jet Backgrounds | | | | |
|-------------------------------|----------------------------|----------------------------|---------------------------|-------------------------|
| Background | $p_T^W=15-25$ GeV | $p_T^W=25-35$ GeV | $p_T^W=35-65$ GeV | $p_T^W=65-105$ GeV |
| $W \rightarrow \tau\nu_\tau$ | 86 ± 3 (2.22 %) | 57 ± 2 (2.16 %) | 56 ± 2 (2.26 %) | 15 ± 1 (2.89%) |
| $Z \rightarrow e^+e^-$ | 47 ± 2 (1.22 %) | 30 ± 1 (1.14 %) | 25 ± 1 (1.01 %) | 5 ± 0 (0.96 %) |
| QCD | 279^{+41}_{-33} (5.40 %) | 205^{+29}_{-90} (5.68 %) | 17^{+47}_{-11} (0.51 %) | 2^{+12}_{-2} (0.29 %) |

TABLE XI: Summary of muon W +jet backgrounds. The background fractions are calculated with respect to the FMC signal event yields for the electroweak backgrounds and with respect to the data for the QCD background.

| Muon W +jet Backgrounds | | | | |
|------------------------------|----------------------------|--------------------------|---------------------------|----------------------|
| Background | $p_T^W=15-25$ GeV | $p_T^W=25-35$ GeV | $p_T^W=35-65$ GeV | $p_T^W=65-105$ GeV |
| $W \rightarrow \tau\nu_\tau$ | 45 ± 2 (2.22 %) | 30 ± 2 (2.17 %) | 30 ± 2 (2.28 %) | 8 ± 0 (2.87%) |
| $Z \rightarrow \mu^+\mu^-$ | 127 ± 7 (6.26 %) | 82 ± 4 (5.92 %) | 72 ± 4 (5.48 %) | 12 ± 1 (4.30 %) |
| QCD | 148^{+56}_{-17} (5.24 %) | 40^{+50}_{-8} (2.12 %) | 18^{+30}_{-18} (0.97 %) | 0^{+10}_{-0} (0 %) |

TABLE XII: The expected total event yields for inclusive W +jet production. The signal and electroweak backgrounds are calculated up to order α_s^2 . The PDF and Q^2 systematics have also been included (second set of uncertainties).

| p_T^W (GeV) | Expected signal+background event yields for inclusive W +jet production | | | | | |
|---------------|---|--------------------|-------------------------------|------------------|----------------------|-------------------------------|
| | Electrons | | | Muons | | |
| | N_e (Signal) | N_e (Background) | N_e (Total prediction) | N_μ (Signal) | N_μ (Background) | N_μ (Total prediction) |
| 15–25 | 3867 ± 137 | 412^{+41}_{-33} | $4279^{+148}_{-146}+880/-660$ | 2027 ± 102 | 320^{+57}_{-19} | $2347^{+124}_{-112}+484/-330$ |
| 25–35 | 2632 ± 93 | 292^{+29}_{-90} | $2924^{+100}_{-132}+598/-408$ | 1384 ± 66 | 152^{+50}_{-10} | $1536^{+88}_{-72}+329/-224$ |
| 35–65 | 2474 ± 87 | 98^{+47}_{-11} | $2572^{+102}_{-91}+562/-383$ | 1314 ± 67 | 120^{+31}_{-19} | $1434^{+79}_{-75}+312/-213$ |
| 65–105 | 518 ± 18 | 22^{+12}_{-2} | $540^{+22}_{-19}+118/-81$ | 279 ± 14 | 20^{+10}_{-1} | $299^{+18}_{-15}+66/-45$ |

ized to the expected number of QCD background events, given by Table IX.

cluded (see Section IX).

D. Summary of backgrounds and Standard Model event yields prediction.

Backgrounds for electron and muon W +jet events for each of the four p_T^W bins are summarized in Tables X and XI respectively. We obtain the total W +jet event yield prediction by adding these backgrounds to the FMC W +jet signal prediction of Table II. To obtain the final uncertainties, we add linearly the uncertainties associated with the W +jet signal and electroweak background and add the result to the QCD background uncertainty in quadrature. The total W +jet event yields after the inclusion of the backgrounds are presented in Table XII. The PDF and Q^2 systematic uncertainties are also in-

VI. COMPARISON BETWEEN EXPECTED AND OBSERVED W DISTRIBUTIONS

We study the expected (FMC) W kinematical distributions after the inclusion of backgrounds and compare them to the experimental distributions. Figures 17 and 19 show the W transverse momentum for electrons and muons respectively. The observed and simulated distributions have been normalized to unity. We observe good agreement between the observed and simulated p_T^W distributions. Figure 18 shows the W transverse mass distribution for the electron W +jet dataset and for the DYRAD events passed through the FMC detector simulation. Figure 20 shows the same distributions for the four p_T^W bins. Figures 21 and 22 show the same distributions for the muon W +jet datasets. The observed and simulated dis-




FIG. 17: The transverse momentum of the W for the electron W +jet data sample (points) along with the FMC signal simulation including backgrounds (solid histogram). The backgrounds are multiplied by 5, to be visible. The data and expected signal+background distributions are normalized to unity.




FIG. 19: The transverse momentum of the W for the muon W +jet data sample (points) along with the FMC signal simulation including backgrounds (solid histogram). The backgrounds are multiplied by 5, to be visible. The data and expected signal+background distributions are normalized to unity.




FIG. 18: The transverse mass of the W for the electron W +jet data sample (points) along with the FMC signal simulation including backgrounds (solid histogram). The backgrounds are multiplied by 5, to be visible. The data and expected signal+background distributions are normalized to unity.




FIG. 20: The transverse mass of the W for the electron W +jet data sample (points) along with the FMC signal simulation including backgrounds (histogram), for the four p_T^W bins. The data and expected signal+background distributions are normalized to unity.




FIG. 21: The transverse mass of the W for the muon W +jet data sample (points) along with the FMC signal simulation including backgrounds (solid histogram). The backgrounds are multiplied by 5, to be visible. The data and expected signal+background distributions are normalized to unity.

tributions are again normalized to unity. In all of the above plots, the FMC distributions are produced with properly weighted signal and background contributions, for electron and muon detector regions.

VII. DIRECT MEASUREMENT OF THE AZIMUTHAL ANGLE OF THE CHARGED LEPTONS FROM W DECAYS IN THE COLLINS-SOPER FRAME

For each W event we boost to the W rest-frame to calculate the azimuthal angle of the charged lepton. The longitudinal momentum of the W (p_z^W) is not known, because the longitudinal momentum of the neutrino is not measurable, so we use the mass of the W to constrain it. For a particular event, the longitudinal momentum of the neutrino is constrained by the mass of the W , according to the equation:

$$p_z^\nu = \frac{1}{(2p_T^l)^2} \left(A p_z^l \pm E^l \sqrt{A^2 - 4(p_T^l)^2 (p_T^\nu)^2} \right), \quad (2)$$

where

$$A = M_W^2 + (p_T^W)^2 - (p_T^l)^2 - (p_T^\nu)^2, \quad (3)$$

E^l is the energy of the charged lepton, p_T^l is its transverse momentum, p_z^l is its longitudinal momentum, p_T^ν is the neutrino transverse momentum, and p_T^W is the transverse




FIG. 22: The transverse mass of the W for the muon W +jet data sample (points) along with the FMC signal simulation including backgrounds (histogram), for the four p_T^W bins. The data and expected signal+background distributions are normalized to unity.

momentum of the W . This equation is unique for every event, since the kinematics of the lepton and neutrino, as well as the mass of the W , contribute to the shape of the curve $p_z^\nu = f(M_W)$. If the mass of the W was known on an event by event basis, there would be a two-fold ambiguity in the value of p_z^ν of the neutrino in the laboratory frame. Because the W boson has a finite width given by a PDF-convoluted Breit-Wigner distribution, $BW(M)$, we actually have two distributions of possible values of p_z^ν , $BW(M(p_z^\nu))$, where $M(p_z^\nu)$ is the mass of the W as a function of the neutrino longitudinal momentum for the particular kinematics of the event.

The choice of one of the two neutrino longitudinal momentum solutions does not affect the ϕ analysis, since both solutions result in the same charged lepton ϕ in the CS frame. For this analysis, only the choice of the W mass is of interest. The choice is made based on the 2-dimensional M^W vs. M_T^W histograms constructed with DYRAD events. For a specific M_T^W we use a probability distribution of W masses and randomly select one for each event, based on that distribution. This method was devised to better reconstruct the $|\cos(\theta)|$ distribution [8], since the polar angle is very sensitive to the selection of the W mass. In our analysis, the azimuthal angle is not affected by the choice of mass, so the answer is almost the same even if we choose a mass based on the Breit-Wigner distribution and the requirement that the mass is greater than the measured transverse mass.

After obtaining a ϕ for every event, we proceed to an-

alyze our sample. Theoretically, the W differential cross section, integrated over $\cos\theta$ and y is given by:

$$\frac{d\sigma}{d(p_T^W)^2 d\phi} = C(1 + \beta_1 \cos\phi + \beta_2 \cos 2\phi + \beta_3 \sin\phi + \beta_4 \sin 2\phi), \quad (4)$$

where

$$C = \frac{1}{2\pi} \frac{d\sigma}{d(p_T^W)^2}, \quad \beta_1 = \frac{3\pi}{16} A_3(p_T^W), \quad \beta_2 = \frac{A_2(p_T^W)}{4}, \\ \beta_3 = \frac{3\pi}{16} A_7(p_T^W), \quad \beta_4 = \frac{A_5(p_T^W)}{2}. \quad (5)$$

The theoretical ϕ distributions for the charged lepton from W boson decay in W +jet production are shown in Figure 2.

From Equations (4) and (5), the reader might conclude that only the A_2 , A_3 , A_5 , and A_7 coefficients are measurable with the ϕ analysis, since the other angular coefficients are integrated out. However, in the actual W +jet data samples, what we measure is the number of events:

$$N(p_T^W, \phi) = \int \frac{d\sigma}{d(p_T^W)^2 d\phi d\cos\theta} ae(p_T^W, \cos\theta, \phi) d\cos\theta \int \mathcal{L} dt + N_{bg}(p_T^W, \phi), \quad (6)$$

where \mathcal{L} is the instantaneous luminosity and $ae(p_T^W, \cos\theta, \phi)$ is the overall acceptance times efficiency, determined in Section IV, for a particular W transverse momentum and region in the $(\cos\theta, \phi)$ phase space. The quantity $N_{bg}(p_T^W, \phi)$ is the background for the given ϕ bin and p_T^W , estimated in Section V. Combining Equations (6) and (1), the measured distribution is

$$N(p_T^W, \phi) = C'(f_{-1}(p_T^W, \phi) + \sum_{i=0}^7 A_i(p_T^W) f_i(p_T^W, \phi)) + N_{bg}(p_T^W, \phi), \quad (7)$$

where $C' = C \int \mathcal{L} dt$. The f_i are fitting functions, which are integrals of the product of the explicit functions $g_i(\cos\theta, \phi)$ and $ae(\cos\theta, \phi)$:

$$f_i(p_T^W, \phi) = \int_0^\pi g_i(\theta, \phi) ae(p_T^W, \cos\theta, \phi) d\cos\theta, \quad (8) \\ i = -1, \dots, 7$$

where

$$\begin{aligned} g_{-1}(\theta, \phi) &= 1 + \cos^2\theta \\ g_0(\theta, \phi) &= \frac{1}{2}(1 - 3\cos^2\theta) \\ g_1(\theta, \phi) &= \sin 2\theta \cos\phi \\ g_2(\theta, \phi) &= \frac{1}{2} \sin^2\theta \cos 2\phi \\ g_3(\theta, \phi) &= \sin\theta \cos\phi \\ g_4(\theta, \phi) &= \cos\theta \\ g_5(\theta, \phi) &= \sin^2\theta \sin 2\phi \\ g_6(\theta, \phi) &= \sin 2\theta \sin\phi \\ g_7(\theta, \phi) &= \sin\theta \sin\phi \end{aligned} \quad (9)$$

Because we multiply the $g_i(\theta, \phi)$ functions by $ae(p_T^W, \cos\theta, \phi)$ before integrating over $\cos\theta$, no f_i is exactly zero and all of the angular coefficients A_i are in principle measurable. We have verified that the FMC-simulated ϕ distributions, fitted with a linear combination of f_i , result in angular coefficient values consistent with the SM predictions [7]. This result supports the self-consistency of the method.

We use Simpson integration for the calculation of the f_i fitting functions given by Equation (8). The explicit functions $g_i(\theta, \phi)$ are integrated over $\cos\theta$, after they are weighted with the value of $ae(p_T^W, \cos\theta, \phi)$ extracted from the 2-dimensional histograms of Figures 7 and 8.

Although the use of Equation (7) allows us in principle to measure all of the angular coefficients, in reality, the current statistics do not allow us to make a significant measurement of angular coefficients other than A_2 and A_3 . This is due to the fact that the fitting functions $f_{i \neq 2,3}$ are small, and the ϕ distributions are insensitive to large variations of the corresponding angular coefficients. Figure 23 shows how the expected electron ϕ distributions are modified as the angular coefficients A_i are varied, one coefficient at a time (the muon ϕ distributions are almost identical). Using Equation (7), we vary A_0 , A_2 , and A_3 from 0 to 1 with a step size of 0.1, and A_4 from 0 to 2 with a step size of 0.2. We find that only A_2 and A_3 strongly affect the azimuthal distributions, thus only these two angular coefficients are measurable with the our current ϕ analysis. Large variations of A_0 and A_4 result in small changes in the ϕ distributions, hence the uncertainties associated with the measurement of A_0 and A_4 are large; these two coefficients cannot be measured in a statistically significant manner with the ϕ analysis. The same is true for A_1 , A_5 , A_6 , and A_7 , all of which are consistent with zero for our current experimental precision.

Figure 24 shows the observed CS electron ϕ distributions for CDF electron W +jet data for the four p_T^W bins. Figure 25 shows the corresponding ϕ distributions of the CDF muon W +jet data. The solid lines are the SM theoretical predictions including backgrounds, whereas the points correspond to CDF W +jet data (the error bars




FIG. 23: The predicted electron ϕ distributions in the Collins-Soper W rest-frame for the first p_T^W bin, varying only one angular coefficient at a time and keeping the other angular coefficients at their Standard Model values. Only A_2 and A_3 significantly affect the shape of these distributions. The same is true for the other p_T^W bins and also for the muon ϕ distributions.

are statistical only). The theoretical prediction for the ϕ distributions is constructed using Equations (7) and (8). The free parameters are the angular coefficients A_i . The background ϕ shapes are given by Figures 9, 11, and 15, for electrons and Figures 10, 12, and 16 for muons, normalized to the event yields of Tables X and XI respectively. The expected signal is normalized to the FMC signal event yields of Table XII, and subsequently the backgrounds are added to construct $N(p_T^W, \phi)$, according to Equation (7). The total theoretically predicted distributions along with the experimental ones, are finally normalized to unity. The experimental results are in good agreement with the Standard Model prediction, which includes the effects of W polarization and QCD contributions up to order α_s^2 .

VIII. MEASUREMENT OF THE ANGULAR COEFFICIENTS

The values of the angular coefficients A_2 and A_3 are extracted using the least-squares fitting method and the data associated with Figures 24 and 25. The least-squares fit is performed over the negative x -axis of the CS frame ($\pi/2 < \phi < 3\pi/2$) for the following two reasons.

Firstly, if a single jet perfectly balances the W boson,




FIG. 24: The ϕ distributions for the electron CDF data (points) the SM Monte Carlo (solid lines) and the result of the fit (dashed lines) for the four bins of p_T^W . The errors are only statistical. The fit is performed from $\pi/2$ to $3\pi/2$ and resulted in χ^2/dof equal to 2.32, 1.80, 2.18, and 1.11 for the four bins respectively (11 degrees of freedom). All distributions are normalized to unity.

its momentum will be placed on the positive x -axis in the CS frame. In reality, the leading jet will be in the $x > 0$ region of the $z-x$ plane, in proximity to the x -axis, as seen in Figure 26 for the electron W +jet data. The leading jet's ϕ in the CS frame will almost always be less than $\pi/2$ or greater than $3\pi/2$, as shown in Figure 26. A kinematic correlation exists between the angular separation ΔR between the jet and the lepton in the $\phi_{\text{lab}} - \eta_{\text{lab}}$ space and the CS ϕ of the lepton, as shown in Figure 26. The situation is similar for the possible subleading jets in the W +jet events (Figure 27). W +jet events with more than two jets are not modeled in DYRAD simulation; their presence in the data creates extra biases in the low and high regions of the lepton ϕ distributions. Because of the lepton-jet angular separation and lepton isolation requirements in our W +jet datasets we obtain a bias-free measurement of the angular coefficients A_2 and A_3 if we exclude the positive- x half-plane region of the CS frame.

Secondly, the term $A_3 f_3(\phi)$ in Equation 7 is the smallest measurable term with our data. Therefore, a more significant measurement of the angular coefficient A_3 is obtained in the CS ϕ region where the rest of the terms (and mainly the predominant $A_4 f_4(\phi)$ term), contribute less. The ratio $A_3 f_3(\phi) / A_4 f_4(\phi)$ is significantly larger in the $\pi/2 < \phi < 3\pi/2$ region, and thus a more sensitive measurement of A_3 is obtained in this region. We




FIG. 25: The ϕ distributions for the muon CDF data (points) the SM Monte Carlo (solid lines) and the result of the fit (dashed lines) for the four bins of p_T^W . The errors are only statistical. The fit is performed from $\pi/2$ to $3\pi/2$ and resulted in χ^2/dof equal to 0.41, 1.39, 1.33, and 1.71 for the four bins respectively (11 degrees of freedom). All distributions are normalized to unity.

normalize the theory to data from $\pi/2$ to $3\pi/2$ before we start the fitting procedure, which is carried out in the $x < 0$ region of the $z - x$ plane.

We use the MINUIT χ^2 minimization program [19] to fit the electron and muon ϕ distributions to the fitting functions f_i . Since these functions are not linearly independent, we cannot fit with all parameters free. For this reason we keep the angular coefficients A_0 and A_4 fixed at their SM values and allow A_2 and A_3 to vary. After we extract values for A_2 and A_3 , we fix these coefficients at these values, and we repeat the fit procedure varying only the A_0 and A_4 angular coefficients. The angular coefficients A_1 , A_5 , A_6 , and A_7 are always kept fixed at their SM values, since the theoretical prediction for these coefficients is very close to zero and the variation for the first 100 GeV of p_T^W is small in comparison to the experimental precision. We expect large statistical uncertainties for the extracted values of A_0 and A_4 , since they do not significantly affect the ϕ distribution. Large variations in their value only slightly alter the leptons' ϕ angular distribution.

The results of the MINUIT fits are shown as dashed histograms in Figure 24 for the electron W +jet data and Figure 25 for the muon W +jet data. Our measurements of the angular coefficients for the electron and muon W +jet data are presented in Figures 28 and 29




FIG. 26: The p_z vs. p_x , p_y vs. p_x , and ϕ of the leading jet in the CS frame and the ΔR between the jet and the lepton in the laboratory frame vs. lepton ϕ in the CS frame, for electron data.




FIG. 27: The p_z vs. p_x , p_y vs. p_x , and ϕ of the extra jets in the CS frame and the ΔR between the jets and the lepton in the laboratory frame vs. lepton ϕ in the CS frame, for electron data.




FIG. 28: The measurement of the angular coefficients for the W +jet electron data (points) and the SM prediction up to order α_s^2 (line). The errors are only statistical.




FIG. 29: The measurement of the angular coefficients for the W +jet muon data (points) and the SM prediction up to order α_s^2 (line). The errors are only statistical.

respectively. The bin centers are determined using the average value of p_T^W for the range of the four p_T^W bins. The measured angular coefficients associated with the electron and muon W +jet data agree with the SM prediction and with each other. We emphasize that the SM prediction is only up to order α_s^2 in QCD. The statistical uncertainties for A_0 and A_4 are very large, as expected, making the measurement of these coefficients unrealistic using the azimuthal angle analysis.

Assuming weak-interaction lepton-universality, we combine the measurements of the angular coefficients obtained from the electron and muon W +jet datasets, treating them as the results of two separate experiments. If A_e and A_μ are the electron and muon measurements with statistical uncertainties σ_e and σ_μ respectively, then the combined measurement is $A_{\text{comb}} = (A_e/\sigma_e^2 + A_\mu/\sigma_\mu^2)/(1/\sigma_e^2 + 1/\sigma_\mu^2)$, with statistical uncertainty $\sigma_{\text{comb}} = (1/\sigma_e^2 + 1/\sigma_\mu^2)^{-1/2}$. The result of this statistical combination, along with the SM prediction, is presented in Figure 30.

IX. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties associated with the measurement of the angular coefficients A_i are related to the jet definition and energy scale, the selection of the W mass on an event-by-event basis, the background estimation, possible presence of $W + \gamma$ events in our datasets,




FIG. 30: The measurement of the angular coefficients for the combination of electrons and muons (points) and the SM prediction up to order α_s^2 (line). The errors are only statistical.

the assumed values of A_0 and A_4 , the choice of parton distribution functions, and the renormalization and factorization scale Q^2 of the event. The jet systematic uncertainties, the variation of the A_0 and A_4 values, and Q^2 scale uncertainty are the dominant sources of systematics.

A. Jet systematic uncertainties

The number of data events passing the jet cuts is affected by the systematic uncertainties associated with the jet E_T scale and the rapidity requirement. The same systematic uncertainty has an effect on the measurement of the angular coefficients.

The uncertainty on jet E_T scale depends on the calorimeter stability, relative energy scale corrections, extra interactions, and underlying event corrections. The total uncertainty is a quadratic sum of these effects. The systematic uncertainty in the jet energy scale affects the reconstruction of the \cancel{E}_T and the W boson. For every FMC W +jet event, we shift the energy of the jet by $\sigma_+ = 85\%\sqrt{E^{\text{jet}}}$, where E^{jet} is the energy of the jet in GeV, without changing its direction. We then correct the \cancel{E}_T value and recalculate all the kinematic variables associated with the W boson, jet, and \cancel{E}_T . We subsequently extract the new acceptance times efficiency $ae(\cos\theta, \phi)$ and analyze the data. We repeat this procedure for the energy shifted by $\sigma_- = -85\%\sqrt{E^{\text{jet}}}$ and calculate the systematic effect of the jet energy scale on the measurement of the angular coefficients, presented in Table XIV for the electron, muon, and the combination of the two results. To obtain the combined results, we combine the electron and muon measurements for each p_T^W bin and for each choice of E_T^{jet} energy shift, using the statistical uncertainties of the central measurements. The difference between the shifted combined values and the central combined value determines the systematic uncertainty on the combined measurement. The same method is used for all the systematic uncertainty estimates.

We vary the jet E_T cut by $(\delta E_T) = \pm 850$ MeV in both data and MC and repeat the analysis each time, to determine its effect on the measurement of the angular coefficients and on the FMC-prediction of signal event yields. Table XIII shows the systematic uncertainty in the measurement of the W +jet event yields associated with the jet E_T cut variation, for the four p_T^W bins. Overall, there is a $+6.4\%/ -5.8\%$ effect in the electron event yields and a $+6.0\%/ -5.7\%$ effect in the muon event yields due to the jet E_T cut.

The uncertainty on the rapidity η_{lab} of the jet is $\delta\eta_{\text{lab}} = \pm 0.2$. We vary the jet η_{lab} cut from 2.2 to 2.6 to obtain the variation in the data event yields presented in Table XIII, for the four p_T^W bins. Overall, there is a $+2.2\%/ -2.4\%$ effect in the electron event yields and a $+0.7\%/ -2.2\%$ effect in the muon event yields due to the jet $\eta_{\text{lab}}^{\text{jet}}$ cut.

In order to obtain an estimate of the systematic uncertainty in the measurement of the angular coefficients associated with the jet E_T and η_{lab} cuts, we run the analysis for 11 values of the E_T^{jet} cut, from 14.15 GeV to 15.85 GeV, and for five values of $|\eta_{\text{lab}}^{\text{jet}}|$ cut, from 2.2 to 2.6. We record the variations in the measurement of the angular coefficients for electrons, muons, and the combination of the two results. The results for the four p_T^W bins are presented in Table XIV.

B. Systematic uncertainty due to W mass selection

As previously discussed, in order to boost to the W rest-frame, a mass value is selected for the W boson. We have four different methods for selecting this mass on an event-by-event basis. We investigate how each mass selection method affects our angular coefficients measurement. The first method selects a Breit-Wigner mass, which is greater than the measured transverse mass of the W boson. The second method selects the greater of the pole mass and the transverse mass. In the third method we select the pole mass or, in case it is less than the transverse mass, we select a Breit-Wigner mass, which is greater than the transverse mass. Finally the fourth method (default) selects a mass based on the distribution that results from the slice of the theoretical (DYRAD) M^W vs. M_T^W 2-dimensional histogram (for W +jet events) at the measured transverse mass of the W boson. This last method is preferred because it removes some biases in the measurement of the polar angle θ . In the ϕ analysis, the systematic uncertainty on the azimuthal angle ϕ due to the selection of the mass of the W is minimal. We run the analysis for the four mass selection methods and record the variations in the measurement of the angular coefficients for electrons, muons, and the combination of the two results. All methods give almost identical measurements of ϕ . The systematic uncertainties for the four p_T^W bins are presented in Table XIV.

C. Backgrounds estimate systematic uncertainty

There is an uncertainty in the estimation of the backgrounds, given by the uncertainties in Tables X and XI. We vary our prediction from the highest value to the lowest possible value for every background as well as the FMC signal event yields. These uncertainties do not include the PDF and Q^2 systematics. For each variation, we rerun the analysis programs for the electron and muon case, and we also combine the results. The systematic uncertainties are presented for the four p_T^W bins in Table XIV.

TABLE XIII: Systematic uncertainties on W +jet event yields due to the E_T^{jet} and $\eta_{\text{lab}}^{\text{jet}}$ cuts, and the total systematic uncertainty due to these sources, for the four p_T^W bins.

| W+jet event yield systematic uncertainties due to the E_T^{jet} and $\eta_{\text{lab}}^{\text{jet}}$ cuts | | | | |
|--|----------------|--------------------------------------|--|------------------------------|
| p_T^W (GeV) | Charged Lepton | δN due to E_T^{jet} | δN due to $\eta_{\text{lab}}^{\text{jet}}$ | Total systematic uncertainty |
| 15–25 | electron | +593 | +156 | +613 |
| | muon | -515 +308 -290 | -154 +34 -81 | -538 +310 -301 |
| 25–35 | electron | +179 | +73 | +195 |
| | muon | -175 +95 -79 | -104 +12 -44 | -204 +96 -90 |
| 35–65 | electron | +38 | +50 | +63 |
| | muon | -50 +12 -29 | -43 +3 -29 | -66 +12 -41 |
| 65–105 | electron | +0 | +1 | +1 |
| | muon | -1 +0 -0 | -2 +0 -0 | -2 +0 -0 |

D. $W + \gamma$ systematic uncertainty

The W +jet angular distribution can be affected by $W + \gamma$ production, for a hard γ well-separated from the charged lepton from the W decay. Some of the events in our datasets are consistent with $W + \gamma$ production, according to [16]. We remove those events and remeasure A_2 and A_3 . The variation from the original measurement is treated as a systematic uncertainty. The systematic uncertainties for the four p_T^W bins are presented in Table XIV.

E. A_0 and A_4 variation systematic uncertainty

In our analysis we keep A_0 and A_2 fixed at their SM values. To check how this affects our measurement, we set A_0 and A_2 at minimum and maximum values in all possible combinations and repeat the analysis four times ($A_0(\text{min})=0$, $A_0(\text{max})=1$, $A_4(\text{min})=0$ and $A_4(\text{max})=2$). The systematic uncertainties for the four p_T^W bins are presented in Table XIV.

F. PDF systematic uncertainty

To study the uncertainty associated with the parton distribution functions, we use the MRSA' [$\alpha_s(M_Z) = 0.105$ and $\Lambda = 0.150$] PDF [20] and repeat the analysis. The systematic uncertainties for the four p_T^W bins are presented in Table XIV. When we use all PDFs of the MRSA and CTEQ families, we end up with a systematic uncertainty of $\pm 11\%$ on the DYRAD cross section, which affects both the central FMC signal event yields and the electroweak backgrounds. These variations are used for the estimation of the total FMC event yields systematic uncertainty due to choice of PDF.

G. Q^2 Systematic uncertainty

Finally we change the renormalization and factorization scale Q^2 so that it is equal to the square of the

transverse momentum of the W , instead of the default square of the pole mass of the W boson. The systematic uncertainties for the four p_T^W bins are presented in Table XIV. If we try all Q^2 choices provided by DYRAD (total invariant mass squared, dynamic mass squared, total energy of the W squared, and transverse energy of the leading jet squared, in addition to the two mentioned above), we end up with a systematic uncertainty of $+19\% / -10\%$ on the DYRAD cross section, which affects both the central FMC signal event yields and the electroweak backgrounds. These variations are used for the estimation of the total FMC event yields systematic uncertainty due to Q^2 scale variation.

H. Overall analysis systematic uncertainties

Table XV summarizes the total systematic uncertainties for the A_2 and A_3 measurement, for the four p_T^W bins and for the electron, muon, and combined results. To populate this table, we combine the systematics described above and presented in Table XIV.

I. Overall systematic uncertainties in data and Monte Carlo event yields

Combining the data event yield systematics due to E_T^{jet} and $\eta_{\text{lab}}^{\text{jet}}$ cut variations in quadrature, we get the final data event yields presented in Table XVI. Comparing with the FMC event yields of Table XII, we see that there is a reasonable agreement with the SM prediction. In Table XII we have also included the PDF and Q^2 FMC systematic uncertainties described earlier, combined in quadrature to give a systematic uncertainty of $+22\% / -15\%$ on the FMC signal event yields and electroweak background. We do not expect perfect agreement since the DYRAD generator produces up to two jets with $E_T^{\text{jet}} > 10$ GeV (order α_s^2), while in the data we have many events with more than two jets with $E_T^{\text{jet}} > 10$ GeV. If we impose a cut on the number of jets in the CDF data, by not accepting more than two jets in an event, and applying strict cuts on at least one jet, the disagree-

TABLE XIV: Systematic uncertainties in the measurement of A_2 and A_3 along with their sources, for electron and muon W +jet events and the combination of the electron and muon results.

| Source of systematic uncertainty | p_T^W (GeV) | Electrons | | Muons | | Combination | |
|----------------------------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | δA_2 | δA_3 | δA_2 | δA_3 | δA_2 | δA_3 |
| Jet E_T cut | 15–25 | +0.0275 | +0.0000 | +0.0146 | +0.0036 | +0.0230 | +0.0000 |
| | | -0.0533 | -0.0064 | -0.0502 | -0.0032 | -0.0523 | -0.0043 |
| | 25–35 | +0.0395 | +0.0047 | +0.0676 | +0.0073 | +0.0490 | +0.0050 |
| | | -0.0109 | -0.0039 | -0.0559 | -0.0017 | -0.0261 | -0.0027 |
| | 35–65 | +0.0433 | +0.0030 | +0.0177 | +0.0042 | +0.0357 | +0.0031 |
| | -0.0000 | -0.0033 | -0.0131 | -0.0011 | -0.0000 | -0.0022 | |
| | 65–105 | +0.0000 | +0.0167 | +0.1519 | +0.0055 | +0.0556 | +0.0024 |
| | | -0.0617 | -0.0153 | -0.0000 | -0.0808 | -0.0299 | -0.0374 |
| Jet η_{lab} cut | 15–25 | +0.0043 | +0.0000 | +0.0098 | +0.0044 | +0.0002 | +0.0000 |
| | | -0.0324 | -0.0036 | -0.0374 | -0.0000 | -0.0175 | -0.0018 |
| | 25–35 | +0.0007 | +0.0012 | +0.0299 | +0.0000 | +0.0106 | +0.0000 |
| | | -0.0346 | -0.0022 | -0.0221 | -0.0085 | -0.0304 | -0.0032 |
| | 35–65 | +0.0093 | +0.0023 | +0.0070 | +0.0038 | +0.0070 | +0.0013 |
| | -0.0049 | -0.0005 | -0.0075 | -0.0028 | -0.0032 | -0.0000 | |
| | 65–105 | +0.0000 | +0.0022 | +0.1021 | +0.0000 | +0.0112 | +0.0000 |
| | | -0.0617 | -0.0052 | -0.0000 | -0.0557 | -0.0258 | -0.0203 |
| Jet energy scale | 15–25 | +0.0594 | +0.0042 | +0.0499 | +0.0075 | +0.0560 | +0.0052 |
| | | -0.0588 | -0.0014 | -0.0694 | -0.0044 | -0.0626 | -0.0023 |
| | 25–35 | +0.0851 | +0.0096 | +0.0860 | +0.0261 | +0.0854 | +0.0147 |
| | | -0.1305 | -0.0020 | -0.0994 | -0.0141 | -0.1065 | -0.0057 |
| | 35–65 | +0.0017 | +0.0204 | +0.0142 | +0.0183 | +0.0053 | +0.0197 |
| | -0.0514 | -0.0069 | -0.0850 | -0.0102 | -0.0614 | -0.0080 | |
| | 65–105 | +0.2468 | +0.0836 | +0.2989 | +0.0382 | +0.2660 | +0.0662 |
| | | -0.3821 | -0.0489 | -0.3186 | -0.0176 | -0.3587 | -0.0369 |
| M_W selection | 15–25 | +0.0030 | +0.0000 | +0.0000 | +0.0001 | +0.0019 | +0.0000 |
| | | -0.0000 | -0.0011 | -0.0016 | -0.0000 | -0.0000 | -0.0007 |
| | 25–35 | +0.0018 | +0.0010 | +0.0034 | +0.0007 | +0.0024 | +0.0007 |
| | | -0.0000 | -0.0000 | -0.0000 | -0.0000 | -0.0000 | -0.0000 |
| | 35–65 | +0.0068 | +0.0034 | +0.0000 | +0.0038 | +0.0012 | +0.0035 |
| | -0.0046 | -0.0002 | -0.0120 | -0.0000 | -0.0040 | -0.0000 | |
| | 65–105 | +0.0000 | +0.0032 | +0.0560 | +0.0000 | +0.0000 | +0.0013 |
| | | -0.0561 | -0.0000 | -0.0000 | -0.0598 | -0.0149 | -0.0210 |
| τ background | 15–25 | +0.0009 | +0.0001 | +0.0014 | +0.0001 | +0.0011 | +0.0001 |
| | | -0.0009 | -0.0000 | -0.0013 | -0.0001 | -0.0011 | -0.0001 |
| | 25–35 | +0.0008 | +0.0000 | +0.0023 | +0.0002 | +0.0013 | +0.0001 |
| | | -0.0010 | -0.0001 | -0.0018 | -0.0002 | -0.0013 | -0.0002 |
| | 35–65 | +0.0008 | +0.0001 | +0.0018 | +0.0002 | +0.0011 | +0.0001 |
| | -0.0012 | -0.0002 | -0.0018 | -0.0002 | -0.0014 | -0.0002 | |
| | 65–105 | +0.0024 | +0.0003 | +0.0000 | +0.0000 | +0.0015 | +0.0002 |
| | | -0.0025 | -0.0005 | -0.0000 | -0.0000 | -0.0016 | -0.0003 |
| Z background | 15–25 | +0.0000 | +0.0001 | +0.0000 | +0.0001 | +0.0000 | +0.0001 |
| | | -0.0009 | -0.0000 | -0.0017 | -0.0000 | -0.0012 | -0.0000 |
| | 25–35 | +0.0000 | +0.0001 | +0.0000 | +0.0002 | +0.0000 | +0.0001 |
| | | -0.0010 | -0.0000 | -0.0037 | -0.0000 | -0.0019 | -0.0000 |
| | 35–65 | +0.0000 | +0.0001 | +0.0000 | +0.0002 | +0.0000 | +0.0001 |
| | -0.0012 | -0.0001 | -0.0029 | -0.0000 | -0.0017 | -0.0000 | |
| | 65–105 | +0.0000 | +0.0003 | +0.0001 | +0.0006 | +0.0000 | +0.0004 |
| | | -0.0025 | -0.0000 | -0.0000 | -0.0007 | -0.0016 | -0.0001 |
| QCD background | 15–25 | +0.0140 | +0.0009 | +0.0329 | +0.0010 | +0.0206 | +0.0009 |
| | | -0.0137 | -0.0009 | -0.0125 | -0.0030 | -0.0126 | -0.0015 |
| | 25–35 | +0.0092 | +0.0013 | +0.0240 | +0.0009 | +0.0143 | +0.0012 |
| | | -0.0326 | -0.0003 | -0.0076 | -0.0039 | -0.0242 | -0.0014 |
| | 35–65 | +0.0018 | +0.0010 | +0.0062 | +0.0013 | +0.0000 | +0.0004 |
| | -0.0149 | -0.0001 | -0.0087 | -0.0015 | -0.0086 | -0.0000 | |
| | 65–105 | +0.0134 | +0.0002 | +0.0243 | +0.0039 | +0.0174 | +0.0006 |
| | | -0.0050 | -0.0014 | -0.0000 | -0.0007 | -0.0031 | -0.0001 |
| FMC signal event yield | 15–25 | +0.0031 | +0.0000 | +0.0000 | +0.0012 | +0.0019 | +0.0000 |
| | | -0.0065 | -0.0023 | -0.0004 | -0.0032 | -0.0043 | -0.0016 |
| | 25–35 | +0.0190 | +0.0003 | +0.0355 | +0.0020 | +0.0246 | +0.0008 |
| | | -0.0000 | -0.0016 | -0.0000 | -0.0017 | -0.0000 | -0.0016 |
| | 35–65 | +0.0173 | +0.0023 | +0.0030 | +0.0039 | +0.0115 | +0.0028 |
| | -0.0000 | -0.0033 | -0.0022 | -0.0000 | -0.0000 | -0.0022 | |
| | 65–105 | +0.0000 | +0.0009 | +0.0830 | +0.0000 | +0.0110 | +0.0000 |
| | | -0.0540 | -0.0052 | -0.0000 | -0.0557 | -0.0267 | -0.0246 |
| $W+\gamma$ | 15–25 | +0.0000 | +0.0000 | +0.0022 | +0.0021 | +0.0000 | +0.0000 |
| | | -0.0060 | -0.0031 | -0.0000 | -0.0000 | -0.0031 | -0.0015 |
| | 25–35 | +0.0000 | +0.0019 | +0.0050 | +0.0000 | +0.0000 | +0.0001 |
| | | -0.0064 | -0.0000 | -0.0000 | -0.0039 | -0.0025 | -0.0000 |
| | 35–65 | +0.0103 | +0.0023 | +0.0026 | +0.0015 | +0.0080 | +0.0020 |
| | -0.0000 | -0.0000 | -0.0000 | -0.0000 | -0.0000 | -0.0000 | |
| | 65–105 | +0.0000 | +0.0024 | +0.0000 | +0.0000 | +0.0000 | +0.0000 |
| | | -0.0704 | -0.0000 | -0.0060 | -0.0166 | -0.0467 | -0.0049 |
| A_0 and A_4 variation | 15–25 | +0.0296 | +0.0074 | +0.0323 | +0.0080 | +0.0305 | +0.0076 |
| | | -0.2024 | -0.0015 | -0.1965 | -0.0016 | -0.2003 | -0.0015 |
| | 25–35 | +0.0691 | +0.0155 | +0.0553 | +0.0240 | +0.0644 | +0.0181 |
| | | -0.2326 | -0.0051 | -0.1745 | -0.0081 | -0.2129 | -0.0060 |
| | 35–65 | +0.1053 | +0.0218 | +0.0917 | +0.0170 | +0.1013 | +0.0202 |
| | -0.1610 | -0.0158 | -0.1390 | -0.0122 | -0.1545 | -0.0146 | |
| | 65–105 | +0.1189 | +0.0150 | +0.0963 | +0.0167 | +0.1106 | +0.0157 |
| | | -0.0708 | -0.0277 | -0.0553 | -0.0312 | -0.0651 | -0.0290 |
| PDF variation | 15–25 | +0.0000 | +0.0013 | +0.0000 | +0.0017 | +0.0000 | +0.0014 |
| | | -0.0159 | -0.0000 | -0.0419 | -0.0000 | -0.0251 | -0.0000 |
| | 25–35 | +0.0152 | +0.0000 | +0.0000 | +0.0000 | +0.0004 | +0.0000 |
| | | -0.0000 | -0.0034 | -0.0018 | -0.0003 | -0.0000 | -0.0024 |
| | 35–65 | +0.0000 | +0.0011 | +0.0759 | +0.0000 | +0.0178 | +0.0006 |
| | -0.0068 | -0.0000 | -0.0000 | -0.0003 | -0.0000 | -0.0000 | |
| | 65–105 | +0.0022 | +0.0134 | +0.4001 | +0.0000 | +0.1486 | +0.0000 |
| | | -0.0000 | -0.0000 | -0.0000 | -0.0931 | -0.0000 | -0.0275 |
| Q^2 variation | 15–25 | +0.1043 | +0.0028 | +0.1200 | +0.0000 | +0.1098 | +0.0000 |
| | | -0.0000 | -0.0000 | -0.0000 | -0.0121 | -0.0000 | -0.0018 |
| | 25–35 | +0.1319 | +0.0000 | +0.0668 | +0.0000 | +0.1099 | +0.0000 |
| | | -0.0000 | -0.0104 | -0.0000 | -0.0076 | -0.0000 | -0.0096 |
| | 35–65 | +0.1347 | +0.0000 | +0.0386 | +0.0000 | +0.1062 | +0.0000 |
| | -0.0000 | -0.0267 | -0.0000 | -0.0180 | -0.0000 | -0.0239 | |
| | 65–105 | +0.0000 | +0.0101 | +0.0000 | +0.0162 | +0.0000 | +0.0124 |
| | | -0.0959 | -0.0000 | -0.2251 | -0.0000 | -0.1434 | -0.0000 |

TABLE XV: Total systematic uncertainties in the measurement of A_2 and A_3 , for electron and muon W +jet events and the combination of the electron and muon results. The systematic uncertainties of Table XIV are combined in quadrature.

| Total systematic uncertainties | | | | | | |
|--------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| p_T^W (GeV) | Electrons | | Muons | | Combination | |
| | δA_2 | δA_3 | δA_2 | δA_3 | δA_2 | δA_3 |
| 15–25 | +0.1275 | +0.0092 | +0.1390 | +0.0128 | +0.1307 | +0.0094 |
| 25–35 | -0.2209 | -0.0087 | -0.2220 | -0.0140 | -0.2189 | -0.0064 |
| 35–65 | +0.1773 | +0.0191 | +0.1492 | +0.0362 | +0.1641 | +0.0239 |
| 65–105 | -0.2622 | -0.0131 | -0.2098 | -0.0208 | -0.2428 | -0.0137 |
| | +0.1773 | +0.0305 | +0.1276 | +0.0263 | +0.1530 | +0.0289 |
| | -0.1699 | -0.0321 | -0.1643 | -0.0243 | -0.1665 | -0.0293 |
| | +0.2743 | +0.0883 | +0.5503 | +0.0453 | +0.3297 | +0.0692 |
| | -0.4229 | -0.0587 | -0.3941 | -0.1629 | -0.3977 | -0.0764 |

TABLE XVI: The electron and muon CDF data event yields for inclusive W +jet production, with statistical and systematic uncertainties. The systematic uncertainties are due to E_T^{jet} and $\eta_{\text{lab}}^{\text{jet}}$ cuts.

| Data event yields for inclusive W +jet production | | | |
|---|---------------|---------------|------------------|
| p_T^W (GeV) | N_e | N_μ | |
| 15–25 | 5166 ± 72 | 2821 ± 53 | $^{+310}_{-301}$ |
| 25–35 | 3601 ± 60 | 1869 ± 43 | $^{+96}_{-90}$ |
| 35–65 | 3285 ± 57 | 1880 ± 43 | $^{+12}_{-41}$ |
| 65–105 | 624 ± 25 | 371 ± 19 | $^{+0}_{-0}$ |

ment is reduced by more than 50%. Nevertheless, we prefer not to constrain the dataset in such a manner. Note that the event yield measurements do not affect the angular coefficient measurements, since we are interested only in the *shapes* of the distributions, in the latter case.

X. FINAL RESULTS

Combining the statistical and systematic uncertainties associated with the A_2 and A_3 measurement, we obtain our final results, presented in Tables XVII and XVIII. Figure 31 shows the measurement of A_2 and A_3 for the electron W +jet data and Figure 32 shows the measurement of A_2 and A_3 for the muon W +jet data. The combination of the electron and muon measurements of the two angular coefficients is presented in Figure 33. The Standard Model predictions for these angular coefficients, up to order α_s^2 , are also presented.

XI. CONCLUSIONS

We have made the first measurement of the A_2 and A_3 angular coefficients of W boson production and decay, using the CDF Run Ia and Run Ib electron and muon W +jet data. Our datasets include at least one jet, satisfying the energy and pseudorapidity requirements. Due to finite statistical analyzing power of our W +jet datasets and the characteristics of the W decay, only the measurement of A_2 and A_3 angular coefficients is statistically significant, with the analysis of the azimuthal

TABLE XVII: The measurement of the A_2 coefficient along with the statistical and systematic uncertainties for electron and muon W +jet events and the combination of the electron and muon results. The SM values up to order α_s^2 are also included.

| Measurement of A_2 angular coefficient | | |
|--|---------------|-----------------|
| | p_T^W (GeV) | A_2 |
| Electrons | 15–25 | 0.02 ± 0.14 |
| | 25–35 | 0.14 ± 0.15 |
| | 35–65 | 0.45 ± 0.13 |
| | 65–105 | 1.24 ± 0.29 |
| Muons | 15–25 | 0.14 ± 0.19 |
| | 25–35 | 0.55 ± 0.21 |
| | 35–65 | 0.55 ± 0.20 |
| | 65–105 | 0.98 ± 0.38 |
| Combination | 15–25 | 0.06 ± 0.11 |
| | 25–35 | 0.28 ± 0.12 |
| | 35–65 | 0.48 ± 0.11 |
| | 65–105 | 1.15 ± 0.23 |
| Standard Model | 15–25 | 0.09 |
| | 25–35 | 0.19 |
| | 35–65 | 0.35 |
| | 65–105 | 0.60 |

TABLE XVIII: The measurement of the A_3 coefficient along with the statistical and systematic uncertainties for electron and muon W +jet events and the combination of the electron and muon results. The SM values up to order α_s^2 are also included.

| Measurement of A_3 angular coefficient | | |
|--|---------------|-----------------|
| | p_T^W (GeV) | A_3 |
| Electrons | 15–25 | 0.03 ± 0.06 |
| | 25–35 | 0.07 ± 0.06 |
| | 35–65 | 0.13 ± 0.07 |
| | 65–105 | 0.21 ± 0.15 |
| Muons | 15–25 | 0.03 ± 0.09 |
| | 25–35 | 0.09 ± 0.09 |
| | 35–65 | 0.13 ± 0.10 |
| | 65–105 | 0.33 ± 0.19 |
| Combination | 15–25 | 0.03 ± 0.05 |
| | 25–35 | 0.08 ± 0.05 |
| | 35–65 | 0.13 ± 0.06 |
| | 65–105 | 0.26 ± 0.12 |
| Standard Model | 15–25 | 0.02 |
| | 25–35 | 0.04 |
| | 35–65 | 0.08 |
| | 65–105 | 0.16 |

angle of the charged lepton in the W rest-frame. The A_0 and A_4 coefficients are preferably measurable with a polar angle analysis, while A_1 and the next-to-leading order coefficients – A_5 , A_6 , and A_7 – are not measurable, with any meaningful statistical significance, with Run I W +jet data.

At leading order, the A_2 and A_3 angular coefficients fully describe the azimuthal W angular distribution in




FIG. 31: Measured A_2 and A_3 using electron W +jet events. The total (outer) and statistical (inner) uncertainties are shown along with the Standard Model 1-loop prediction up to order α_s^2 (dashed line).




FIG. 32: Measured A_2 and A_3 using muon W +jet events. The total (outer) and statistical (inner) uncertainties are shown along with the Standard Model 1-loop prediction up to order α_s^2 (dashed line).

the Collins-Soper W rest-frame. These angular coefficients are also part of the total W differential cross section, and can be expressed as ratios of the corresponding helicity cross sections of the W to its total unpolarized cross section. This measurement tests the Standard Model prediction for W polarization, and the associated QCD corrections present in the production of W bosons at high transverse momenta. We observe good agreement with the Standard Model prediction up to order α_s^2 in QCD.

Acknowledgments

We thank Walter Giele for DYRAD support and useful discussions. We also thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium fuer Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, UK;




FIG. 33: Measured A_2 and A_3 using the combination of electron and muon measurements. The total (outer) and statistical (inner) uncertainties are shown along with the Standard Model 1-loop prediction up to order α_s^2 (dashed line).

the Russian Foundation for Basic Research; the Comision Interministerial de Ciencia y Tecnologia, Spain; and in part by the European Community's Human Poten-

tial Programme under contract HPRN-CT-2002-00292, Probe for New Physics.

-
- [1] E. Mirkes, Nucl. Phys. B **387**, 3 (1992).
- [2] J. C. Collins and D. E. Soper, Phys. Rev. D **16**, 2219 (1977).
- [3] If we CP transform the W^- into W^+ keeping the momenta of the colliding hadrons fixed (in order to have a fixed CS frame), Equation (1) describes the W^+ if the A_1 , A_4 and A_6 coefficients are multiplied by (-1) ; these coefficients multiply P -odd functions of θ and ϕ , since the Parity transformation takes $\phi \rightarrow \phi$ and $\theta \rightarrow \pi - \theta$ in the CS frame.
- [4] C. Albajar et al. (UA1 Collaboration), Z. Phys. C - Particles and Fields **44**, 15 (1989).
- [5] E. Mirkes and J. Ohnemus, Phys. Rev. D **50**, 5692 (1994).
- [6] K. Hagiwara, K. Hikasa, and N. Kai, Phys. Rev. Lett. **52**, 1076 (1984).
- [7] J. Strologas and S. Errede, submitted to Phys. Rev. D (2005), arXiv:hep-ph/0503291.
- [8] J. Strologas, Ph.D. thesis, University of Illinois at Urbana-Champaign (2002).
- [9] B. Abbott et al. (D0 Collaboration), Phys. Rev. D **63**, 072001 (2001).
- [10] D. Acosta et al. (CDF Collaboration), Phys. Rev. D **70**, 032004 (2004).
- [11] F. Abe et al. (CDF Collaboration), Nucl. Instrum. Methods Phys. Res. A **271**, 387 (1988).
- [12] In this paper, we use natural units, according to which $c = \hbar = 1$. Therefore, the energy, momentum, and mass are measured in eV.
- [13] This analysis has been performed on W^+ and W^- separately with consistent results.
- [14] W. T. Giele, E. W. N. Glover, and D. A. Kosower, Nucl. Phys. B **403**, 633 (1993).
- [15] H. L. Lai et al., Phys. Rev. D **55**, 1280 (1997).
- [16] L. Christofek, Ph.D. thesis, University of Illinois at Urbana-Champaign (2001).
- [17] J. Dittmann, Ph.D. thesis, Duke University (1998).
- [18] K. Hagiwara et al. (Particle Data Group), Phys. Rev. D **66**, 010001 (2002).
- [19] Minimization routine from CERN library.
- [20] A. D. Martin, W. J. Stirling, and R. G. Roberts, Phys. Lett. B **354**, 155 (1995).