

# Urban Ecosystems

## Effects of garden management practices, by different types of gardeners, on human wellbeing and ecological and soil sustainability in Swiss cities --Manuscript Draft--

<b>Manuscript Number:</b>	UECO-D-17-00192R1	
<b>Full Title:</b>	Effects of garden management practices, by different types of gardeners, on human wellbeing and ecological and soil sustainability in Swiss cities	
<b>Article Type:</b>	Manuscript	
<b>Keywords:</b>	Gardening; Wellbeing; Species richness; Soil quality; Management practices; Gardener typology.	
<b>Corresponding Author:</b>	Robert Home Forschungsinstitut für biologischen Landbau Frick, SWITZERLAND	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	Forschungsinstitut für biologischen Landbau	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Robert Home	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Robert Home	
	Olivia Lewis	
	Nicole Bauer	
	Andreas Fliessbach	
	David Frey	
	Stéphanie Lichtsteiner	
	Marco Moretti	
	Simon Tresch	
	Christopher Young	
	Andrea Zanetta	
	Matthias Stolze	
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (CH) (CRSII1_154416)	Dr Matthias Stolze
<b>Abstract:</b>	<p>Gardens have effects on the local ecology as well as on the wellbeing of the gardener, but few studies have attempted to study gardens using both ecological and social outcome variables. The aim of this exploratory study is to address this research gap by identifying the characteristics of gardens and the management practices of gardeners that enhance the outcomes of gardening, which we separate into three dimensions: human wellbeing, biodiversity, and soil quality. Data were collected from 18 gardens in Zurich, Switzerland and a typology of gardeners was identified, which included 'conservationist', 'functional', 'minimum effort', 'child-friendly', and 'aesthetic' gardeners. The conservationist gardeners were found to have, on average, the highest species richness in their gardens, while the minimum effort gardeners had the lowest, which suggests that some degree of management can enhance species richness. The</p>	

conservationist and minimum effort gardeners had, on average, the highest values for stable aggregates, while the minimum effort gardeners had the highest phosphorous content in their soil. The wellbeing of the minimum effort gardeners was lower than the other groups, which suggests it is the act of gardening, rather than merely having a garden, which leads to wellbeing outcomes. The results suggest that ecologically friendly gardening is compatible with desired social outcomes and furthermore that the beneficial effects of gardens are indeed related to the practices implemented by the gardeners, which are influenced by their attitudes towards gardening and the role of gardens in their lives.

[Click here to view linked References](#)

1 **Authors:** Robert Home\*<sup>1</sup>, Olivia Lewis<sup>1</sup>, Nicole Bauer<sup>2</sup>, Andreas Fliessbach<sup>1</sup>, David Frey<sup>2</sup>,  
2 Stéphanie Lichtsteiner<sup>1</sup>, Marco Moretti<sup>2</sup>, Simon Tresch,<sup>1,2</sup> Christopher Young<sup>2</sup>, Andrea  
3 Zanetta<sup>2</sup>, Matthias Stolze<sup>1</sup>

4  
5 **Title:** Effects of garden management practices, by different types of gardeners, on  
6 human wellbeing and ecological and soil sustainability in Swiss cities

7  
8 **Affiliations and addresses:**

9  
10 <sup>1</sup> Research Institute of Organic Agriculture (FiBL), Ackerstrasse 113, 5070 Frick,  
11 Switzerland

12  
13 <sup>2</sup> Federal Research Institute for Forest Snow and Landscape Research (WSL),  
14 Zurcherstrasse 111, 8902 Birmensdorf, Switzerland

15  
16  
17  
18  
19 **\*Corresponding author:**

20  
21 Email address: robert.home@fiBL.org

22 Telephone number: +41 62 865 7215

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

As the European urban population has increased, and urban planners have been reluctant to allow urban sprawl, cities and towns have increased in density. This high-density urban living means that urban green spaces are increasingly becoming important nodes of contact with nature as well as providing nearby outdoor recreation areas (Frick et al., 2007). The associated increase in demand for housing and infrastructure puts pressure on these spaces, which are an attractive target for development. Consequently, a growing number of European urban residents are becoming increasingly disconnected from natural environments (Kiesling and Manning, 2010).

Despite the pressure from urbanization and densification of cities, urban green spaces have persisted and contribute to the socio-ecological environment of a city through direct and indirect social and environmental impacts (Drescher et al., 2006). Indeed, for many people, urban green spaces, and especially gardens, provide their only, or at least their primary, regular contact with the natural environment (Freeman et al., 2012).

Privately managed gardens are a particular type of urban green space in which the gardener can create social and environmental outcomes from their activities. Such gardens collectively account for a considerable proportion of urban green spaces in most European countries (Van Heezik et al., 2013) including Switzerland (Lindemann-Matthies and Marty, 2013).

Home et al. (2012) observed that there have been many studies of the social benefits of urban green spaces, including gardens, and the almost universal opinion is that urban green spaces contribute to the perceived quality of urban landscapes and to the quality of life of urban residents. The sustainable management of urban green spaces and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45 garden areas therefore can be regarded as crucial for achieving a resilient urban future  
46 (Goddard et al., 2013). The agreement about the social benefits of urban nature has led  
47 to recommendations that there should simply be more green spaces, with Sullivan et al.  
48 (2004) suggesting that benefits for city residents would be maximised, and quality of life  
49 enhanced, if there were nature at every doorstep.

50 In addition to social benefits, urban green spaces have also been shown to be of  
51 ecological value. They support high biodiversity (Sattler et al., 2010), enhance ecosystem  
52 functions, such as pollination (Hall et al., 2017), and provide a wide range of  
53 environmental benefits (Ziter, 2016). Gardeners can influence the diversity and  
54 abundance of plant and animal species by modifying the amount of habitats and  
55 resources for wildlife in the gardens (Gaston et al. 2005) through their choices of how to  
56 build and manage their gardens (Van Heezik et al., 2013). Provision of different habitat  
57 patches and structural elements, such as hedges, woodpiles, or ponds, increases the  
58 habitat variety and the attractiveness of garden areas (Davies et al., 2009). Smith et al.  
59 (2006) found that human influence, in terms of management practices and the  
60 installation of structural elements in the gardens, has a significant impact on species  
61 abundance and richness in the gardens.

62 The importance of gardens, and of garden management, as contributors to the social and  
63 ecological quality of urban environments suggests the value of gaining some  
64 understanding of the interactions between social and ecological characteristics of  
65 gardens. However, most studies have remained within disciplines, and few have  
66 included both ecological and social variables (Van Heezik et al., 2013). The aim of this  
67 study is to address this research gap by identifying the characteristics of gardens and  
68 the management practices of gardeners that enhance the outcomes of gardening, which  
69 we separate into three dimensions: human wellbeing, biodiversity, and soil quality. Once

70 these characteristics and management practices have been identified, strategies can be  
71 suggested to encourage garden management that enhances the quality of gardens.  
72 Furthermore, strategies are likely to be more effective if they are tailored to the agent  
73 who is intended to implement them, so an additional aim is to identify whether  
74 particular types of garden managers exist.

## 75 **2. Typologies of gardeners**

76 There has been little multidimensional empirical research into urban gardens, so we  
77 chose a case study approach, which is a descriptive, exploratory or explanatory analysis  
78 of, in this instance, a selected set of gardens, that explores causation to find underlying  
79 principles (Yin, 2009). In this study, we focus on two types of privately managed  
80 gardens: Domestic gardens, which are usually next to people's houses; and allotment  
81 gardens, which are usually separate from houses, are organised into plots that are  
82 divided from a larger piece of land, and were primarily intended to provide a garden for  
83 those who would otherwise have no access to one. Van Heezik et al. (2014) pointed out  
84 that most studies of urban vegetation and ecosystem function have focussed only on  
85 vegetation in front gardens or on trees: mainly because of ease of visibility from the  
86 street. They further point out the value of examining whole gardens, with a focus on  
87 woody vegetation because of its contribution to ecosystem services and to habitats (Van  
88 Heezik et al., 2014).

89 Within urban gardens, there is a wide range of different gardening management  
90 approaches: Thompson (2007) distinguishes between conventional gardening  
91 management approaches and near-natural gardening and considers them as two ends of  
92 a gradient with a large variety of possibilities between the extremes. Lindemann-  
93 Matthies and Marty (2013) state that the majority of Swiss gardeners apply

94 management practices that are towards the conventional end of the spectrum. These  
95 management approaches are presumably affected by the motivations of the gardener, as  
96 illustrated by Zagorski et al. (2004), who grouped gardeners into four attitudinal groups  
97 of: 'functional' gardeners, who commit significant time and work to creating a traditional  
98 gardenesque type garden, which is characterised by high degrees of management, large  
99 proportions of exotic plants, and geometrically shaped garden beds (Turner, 1986);  
100 'conservationist' gardeners who aim for near-natural gardens; 'romantic' gardeners who  
101 value privacy and tend to create woody gardens; and 'minimum effort' gardeners who  
102 like gardens to create themselves, and preferred to minimize the activity of gardening.  
103 Kettle (2014) created a typology of five gardener types in Irish allotment gardens based  
104 on motivations: the 'Practical Gardener', the 'Idealist/Eco-Warrior', the 'Socio-Organic  
105 Gardener', the 'Gucci Gardener', and the 'Non-Gardening Gardener', and identified  
106 different gardening practices and structural differences in the gardens between types.  
107 The implication of these studies is that the motivations of the gardener influence which  
108 practices are chosen, which subsequently influences the characteristics of the garden  
109 and the social and environmental outcomes of the practices. There is, however,  
110 insufficient evidence in the literature as to whether these classifications are applicable  
111 in the Swiss context, so primary research is needed.

### 3. Methodology

113 The study was undertaken in two distinct phases. The first phase was to use Q-  
114 methodology to identify latent groups within the sample and thus identify a typology of  
115 gardeners based on their attitudes towards gardens and gardening. The second phase  
116 was to examine the gardening outcomes from gardeners in the different latent groups. In  
117 this study, we follow the example given by Van Heezik et al. (2014), and include the

118 whole garden, but expand the evaluation to include assessment of outcomes for  
1  
2 119 biodiversity and soil. Furthermore, we follow the suggestion of Goddard et al. (2013) to  
3  
4  
5 120 include the perspectives of the gardeners. Specifically, the evaluated outcomes included:  
6  
7 121 garden practices, including which structural elements and habitat types had been  
8  
9  
10 122 installed; the wellbeing of the gardener in relation to the gardening experience; plant  
11  
12 123 and arthropod species richness; and soil parameters. However, inclusion of several  
13  
14  
15 124 dimensions of outcomes means that the study sites were subject to intensive  
16  
17 125 examination, which is also resource intensive. The available resources allowed a  
18  
19  
20 126 relatively small sample size, so this study should be considered to be an exploratory case  
21  
22 127 study.

### 25 128 **3.1 Sample**

27  
28 129 The sample consisted of 18 gardens (9 allotments and 9 domestic gardens) in the city of  
29  
30  
31 130 Zurich, Switzerland. Zurich is a medium sized, central European city with 410 000  
32  
33 131 residents and is the centre of the largest metropolitan area of Switzerland, with 1.3  
34  
35  
36 132 million residents. Domestic gardens, allotment gardens, and parks cover around 1500  
37  
38 133 ha, which is approximately 15% of the city's administrative area (Gruen Stadt Zürich,  
39  
40  
41 134 2010).

42  
43  
44 135 The selected gardens were all approximately 200m<sup>2</sup> in size, with variance in garden size  
45  
46  
47 136 kept as low as possible to minimize a size effect. They were selected based on two main  
48  
49 137 factors that were expected to affect soil characteristics and biodiversity as well as  
50  
51  
52 138 ecosystem services and underlying processes: the habitat heterogeneity (i.e. number of  
53  
54 139 habitat patches and structures) and management intensity (i.e. degree of disturbance)  
55  
56  
57 140 within the gardens; and the surrounding landscape composition and configuration (i.e.  
58  
59 141 gradient of urbanization).



## 142 **3.2 Indicators of Outcomes**

1  
2  
3 143 To evaluate the outcomes of gardening practices, and thereby to address the aim of this  
4  
5  
6 144 study, indicators are needed. For the purposes of this study, we elected to evaluate the  
7  
8 145 outcomes in three dimensions: social wellbeing, biodiversity, and soil quality.

10  
11  
12 146 Indicators of wellbeing were based on the results of Irvine et al. (2013), who identified a  
13  
14 147 range of constructs, which they classified into seven domains, with which park users  
15  
16  
17 148 conceptualize green spaces as a resource for health and wellbeing. From this taxonomy,  
18  
19 149 we created a 12-item scale, with items indicating the concepts (shown in table 1).

### 23 150 **Table 1: About here**

24  
25  
26 151 Indicators of biodiversity were based on species richness, which is often positively  
27  
28  
29 152 correlated with ecosystem functioning (Cardinale et al., 2012) and has been shown to be  
30  
31 153 a useful biodiversity indicator in the cases of vascular plants (Van Heezik et al., 2014);  
32  
33  
34 154 ground dwelling arthropods (Braaker et al., 2014); and flying insects (Sattler et al.,  
35  
36 155 2011). We are aware that use of three indicators gives only an approximation of overall  
37  
38  
39 156 complexity, of which there is certainly no straightforward way of measuring, but argue  
40  
41 157 that this is acceptable in this exploratory study.

42  
43  
44  
45 158 Indicators of soil quality should be able to readily show changes in soil conditions  
46  
47 159 (Brejda et al., 2000), so we selected indicators of physical, chemical and biological  
48  
49  
50 160 properties, which are commonly used for holistic soil quality assessments (Karlen et al.,  
51  
52 161 2008). Indicators that have proven useful in previous study are aggregate stability  
53  
54  
55 162 (Zornoza et al., 2015); phosphorous concentration (Haynes and Tregurtha, 1999); and  
56  
57 163 microbial biomass carbon (Muscolo et al., 2015).

## 61 164 **3.3 Data collection and analysis**

165 The Q-methodology was applied by means of face to face interviews with the  
1  
2 166 owner/manager of each of the sample gardens. Each participant was presented with a  
3  
4  
5 167 set of 33 or 30 statements (for allotment gardeners and domestic gardeners,  
6  
7 168 respectively), and asked to sort them into a matrix with empty cells in roughly the shape  
8  
9  
10 169 of a normal distribution, according to their relative level of agreement with each  
11  
12 170 statement. The Q data was organized into a correlation matrix, which was then used to  
13  
14  
15 171 perform a factor analysis. Q sorts that are highly correlated with one another may be  
16  
17 172 considered to have a family resemblance (Brown 1996), which allows an identification  
18  
19  
20 173 of the latent “types” of gardener. The analysis was conducted using the program  
21  
22 174 PQMethod, with varimax rotation.

25 175 Data on gardener wellbeing were gathered from garden owners by using a  
26  
27  
28 176 questionnaire, which was mailed to the participating gardeners in May 2017. The  
29  
30  
31 177 questionnaire included a 12-item scale with one item for each of the indicators identified  
32  
33 178 by Irvine et al. (2013). Participants responded to each question on a scale of 1-5, with 1  
34  
35  
36 179 indicating “fully disagree” and 5 indicating “fully agree”. An overall wellbeing index was  
37  
38 180 also calculated for each gardener type: calculated as the sum of the unweighted averages  
39  
40  
41 181 of each of the above 12 indicators, and reduced to a scale of 1-5, with 5 indicating high  
42  
43 182 wellbeing. This questionnaire also included questions about management practices and  
44  
45  
46 183 which landscape elements had been installed in the garden. Responses were received  
47  
48 184 from 14 of the 18 gardeners, with the remainder indicating that they were incapable of  
49  
50  
51 185 completing the questionnaire due to age or illness.

54 186 Ground-dwelling arthropods were sampled using two triplets of 7-cm diameter pitfall  
55  
56  
57 187 traps placed in two of the most common garden habitat types, such as lawn, flower beds,  
58  
59 188 vegetable beds and berry cultivations within each garden (Duelli et al. 1999). Flying  
60  
61 189 insects were sampled using coloured pan traps, i.e. three 1-litre bowls (UV-bright blue,  
62  
63  
64  
65

190 white, and yellow) fixed on a 1.5m tall pole. Both pitfall and pan traps were filled with a  
191 0.2% Rocima solution (a bactericide and fungicide from Acima, Buchs, Switzerland) and  
192 emptied on a weekly basis between May 18, 2015 and August 19, 2015, which covers the  
193 main activity season of most taxa (Obrist and Duelli 2010). In total, 20 taxonomic groups  
194 of arthropods representative of the main trophic levels and locomotion modes were  
195 sorted in the lab following standard procedures (Duelli et al. 1999), and identified to the  
196 species level. Plant species richness was assessed by two complementary methods: Two  
197 vegetation relevés of 10m<sup>2</sup> (in July) centred around pitfall trap sites; and a total garden  
198 flora inventory repeated three to four times in 2015 (early spring, spring, summer and  
199 early autumn).

200 Soils were sampled in March 2015 in the 18 selected gardens. Within the gardens,  
201 samples were taken near annual herbaceous plants (vegetables and flowers), which we  
202 regard as 'high disturbance'; and near perennial plants (berry cultures and lawn and  
203 perennial flowers), which we regard as 'low disturbance'. These paired samples were  
204 taken as a bulk sample of six soil cores from 0-20 cm. Soils were sieved and dried or  
205 adjusted to soil moisture corresponding to 40-50% of the maximum water holding  
206 capacity. Bulk density was determined in undisturbed ring samples after drying at  
207 105°C. Soil organic carbon, pH, and phosphorous content were analysed in dried soils.  
208 Soil microbial biomass was determined in moist samples. All analyses were done  
209 according to Swiss reference methods (Agroscope, 1996).

## 4. Results

### 4.1 Gardener types

The Q-methodology enabled the identification of five factors, with a total explained  
variance of 77%, and with all sorts accounted for by these factors. The statements that

214 were used in the Q-sort, along with scores indicating where the statement is placed on a  
1  
2 215 representative Q-sort for each of the 5 gardener types are shown in Table 2.  
3  
4

5  
6 216 **Table 2: About here**  
7

8  
9 217 The placement of each statement is shown by a number, ranging from +3 “I agree the  
10  
11 218 most” to -3 “I agree the least”, which indicates in which column of the representative  
12  
13 219 matrix the statement falls. Each gardener type corresponds to a representative Q sort  
14  
15 220 (i.e. statements placed into matrix in a way that represents the views of the type) and  
16  
17 221 has been named. We adopted the same nomenclature for three of the groups identified  
18  
19 222 in this study: ‘conservationist’, ‘functional’, and ‘minimum effort’ gardeners, as those  
20  
21 223 given by Zagorksi et al. (2004) to three of their four ‘attitude groups’ because these  
22  
23 224 groups closely mirrored each other in the two studies. We identified two further groups,  
24  
25 225 which we call ‘aesthetic’ gardeners and ‘child-friendly’ gardeners. These names will be  
26  
27 226 used throughout the presentation of the results.  
28  
29  
30

31  
32  
33  
34 227 Although standard Q-methodology encourages creating factors with two or more  
35  
36 228 exemplars (a sort loading significantly on the factor), we chose to accept a factor with  
37  
38 229 one exemplar because we suspect that child-friendly gardening is a shared orientation,  
39  
40 230 but that our sample size was too small to capture more than one such gardener (see  
41  
42 231 Watts and Stenner, 2005, on accepting one-factor exemplars). Descriptions of each type  
43  
44 232 are as follows.  
45  
46  
47  
48  
49

50  
51 233 ***Conservationist gardeners***  
52

53  
54 234 Of the 9 gardeners in this group, 7 are allotment gardeners and 2 are domestic  
55  
56 235 gardeners. These gardeners are motivated by their belief that everything in nature is  
57  
58 236 interconnected and interdependent. They get joy from having biodiversity in the garden  
59  
60 237 and want to contribute to the preservation and promotion of biodiversity. They are also  
61  
62  
63  
64  
65

238 motivated by producing healthy food in the garden. They do not believe that pesticides  
1  
2 239 and artificial fertilizers are necessary. Their practices are not motivated by the  
3  
4  
5 240 *expectations* of their neighbours nor because other gardeners do so. They are less  
6  
7 241 concerned than others about having a lawn or having a neat garden.  
8  
9

10  
11 242 ***Functional gardeners***  
12  
13

14 243 Both of the gardeners in the second group are domestic gardeners. They choose their  
15  
16 244 practices because they've had good experiences with them so far. They are convinced it  
17  
18  
19 245 is the right way to garden. They decide what to do in the garden because they like to  
20  
21 246 learn about gardening, are happy to see biodiversity in the garden and find the garden  
22  
23  
24 247 beautiful to look at. They do not garden in a certain way because they learned it from  
25  
26 248 their family or because they are trying to mimic other gardeners. They are not motivated  
27  
28  
29 249 to garden to produce healthy food in the garden or to harvest a lot of food. They are not  
30  
31 250 motivated by worry about the consequences of artificial pesticides and fertilizers and  
32  
33  
34 251 believe more than other groups that the garden would not work if they stopped using  
35  
36 252 these products.  
37  
38  
39

40 253 ***Minimum effort gardeners***  
41  
42

43 254 Of the 2 gardeners in the third group, both are domestic gardeners. These gardeners are  
44  
45  
46 255 motivated to garden in a way in which they do not have to put in a lot of physical  
47  
48 256 exertion or pay big investment costs. They choose to keep gardening in the same way  
49  
50  
51 257 that they have always done. They like to have a neat garden and be a model for children  
52  
53 258 through their gardening practices but they are not motivated to garden in a way that  
54  
55  
56 259 actively allows children to play in the garden without danger. They are more motivated  
57  
58 260 than other groups to garden in a way that saves time. They do not choose gardening  
59  
60  
61 261 practices to meet their neighbours' expectations, because other gardeners  
62  
63  
64  
65

262 recommended it to them or to get social recognition. They do not think the garden

1

2 263 would not work if they stopped using artificial pesticides and fertilizers.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

264 ***Child-friendly gardeners***

265 The gardener in the fourth group is a domestic gardener. (S)he is motivated to create a

266 neat garden and to have a lawn. (S)he gardens in a certain way because (s)he is

267 convinced that that's the right way. (S)he does not garden to learn about new gardening

268 practices and implement the knowledge or to use practices recommended by other

269 gardeners, mimic what other gardeners do, or replicate practices that were passed down

270 in the family. (S)he gardens in his/her way because (s)he believes that it would not work

271 without pesticides and artificial fertilizers. (S)he is more concerned than other groups

272 with the ability for children to play in the garden without danger and is neither

273 motivated by saving time in the garden nor enjoying the challenge provided by the way

274 of gardening.

275 ***Aesthetic gardeners***

276 Of the 4 gardeners in the fifth group, 3 are allotment gardeners and 1 is a domestic

277 gardener. The aesthetic gardeners garden in a certain way because they want to

278 contribute to preserving and promoting biodiversity, which they get joy from having in

279 the garden. It is important to them to garden in a way that keeps the garden looking neat

280 and beautiful. They use their practices because they have an inner conviction that it is

281 the right way to garden. They are not motivated by having a lawn, by meeting the

282 expectations of their neighbours, nor to harvest a lot of food. More than other groups,

283 their practices are influenced by what they learned from their family and what other

284 gardeners do.

285 **4.2 Gardener Wellbeing**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

286 The results of the responses to the wellbeing indicators are presented, by gardener type,  
287 in figure 1.

288 **Figure 1: about here**

289 **4.3 Structural elements in gardens**

290 The number of gardens in which selected structural elements had been installed is  
291 shown in table 3.

292 **Table 3: about here**

293 **4.4 Biodiversity in gardens**

294 The results of the appraisal of the biodiversity indicators are shown in Table 4.

295 **Table 4: about here**

296 **4.5 Soil quality in gardens**

297 The results of the appraisal of the soil quality indicators are shown in Table 5.

298 **Table 5: about here**

299 **5. Discussion**

300 Zagorski et al.'s (2004, p.212) 'conservationist' gardeners were "distinguished from  
301 others by their devotion to habitat preservation, trees, native plants", so this name  
302 appears appropriate. The 'conservationist' gardeners group is similar to the  
303 'idealist/eco-warrior' gardener type identified by Kettle (2014, p. 43), whose  
304 motivations "are part of wider concerns for the environment and ecological  
305 sustainability". The 'functional gardeners' group appears to be analogous to Zagorski et  
306 al.'s (2004, p.211) 'functional gardeners' who "are best discriminated from gardeners in  
307 other groups by their attachment to functionality (or practicality) and their pleasure in  
308 working with the garden". Zagorski et al.'s (2004, p.211) 'minimum effort' gardeners

309 similarly liked gardens to create themselves and preferred to minimize the act of  
1  
2 310 gardening. We similarly identified a group of gardeners who are motivated to minimize  
3  
4  
5 311 physical exertion, investment costs, and time, and keep gardening in the same way they  
6  
7 312 have always done. These are also similar to the ‘non-gardening’ gardeners identified by  
8  
9  
10 313 Kettle (2014).  
11  
12  
13 314 ‘Child-friendly’ gardeners, were not detected in the study of Zagorski et al. (2004)  
14  
15  
16 315 However, their distinguishing motivation (ability for children to play in the garden  
17  
18 316 without danger) is similar to those of the ‘non-gardening’ gardeners observed by Kettle  
19  
20  
21 317 (2014, p.56), for whom the gardens “play an important social role for them, and in  
22  
23 318 particular, their children [...they] place a high value on the social and pedagogic value of  
24  
25  
26 319 the allotment landscape”. In contrast, Kettle (2014) observed the gardens appeared  
27  
28 320 ‘neglected or abandoned’, whereas the ‘child-friendly’ gardener in our study placed high  
29  
30  
31 321 importance on keeping the garden neat and having a lawn. Further research, with a  
32  
33 322 larger sample size, may support the idea of this group, potentially by investigating child-  
34  
35  
36 323 friendly features like sandpits as seen in the Irish ‘non-gardening’ gardens (Kettle,  
37  
38 324 2014). The ‘child-friendly’ gardener valued children being able to play safely in the  
39  
40  
41 325 garden, and therefore it is unsurprising that they had a lawn and flower beds but no  
42  
43 326 other features, such as a water feature, free hedge, or dry stone wall, which could be  
44  
45  
46 327 potentially hazardous to playing children. The similarity of the ‘child friendly’ gardeners  
47  
48 328 to the ‘non-gardening’ gardeners identified by Kettle (2014) suggest that the ‘child-  
49  
50  
51 329 friendly’ gardeners might be a subset of ‘non-gardening’ gardeners, but differentiated by  
52  
53 330 their focus on children in the garden.  
54  
55  
56 331 The ‘aesthetic’ gardeners have similarities to both Zagorski et al.’s (2004, p.211)  
57  
58  
59 332 ‘romantic’ gardeners who were “distinguished by their attachment to romance,  
60  
61  
62 333 sentimentality, privacy and space”, by valuing aesthetics, and liking fauna/biodiversity  
63  
64  
65



334 in the garden; and to the 'Gucci gardeners' described by Kettle (2014), through focus on  
1  
2 335 certain design principles. The two groups contrast slightly: 'romantic' gardeners desire  
3  
4  
5 336 privacy and space, whereas 'Gucci' gardeners desire interaction and belonging, but also  
6  
7 337 want to have a green space of their own. However, Kettle (2014, p. 53) describes the  
8  
9  
10 338 Gucci Gardeners as 'in all probability [...] a passing trend', but neither our results nor the  
11  
12 339 results of Zagorski et al. (2004) indicate this, so the nomenclature of 'Gucci Gardeners' is  
13  
14  
15 340 not used; we also do not know if the 'aesthetic' gardeners in this study had a particularly  
16  
17 341 romantic aesthetic, hence the more generalized name of 'aesthetic' gardeners.

18  
19  
20  
21 342 The 'conservationist' gardeners were the only group to include water features in their  
22  
23 343 gardens, although water features are commonly included in gardenesque type gardens  
24  
25  
26 344 (Turner, 1986) favoured by the functional gardeners. The 'conservationist' gardeners  
27  
28 345 also commonly featured a wild area in their gardens, as did one of the 'minimum effort'  
29  
30  
31 346 gardeners. Interestingly, three of the four 'aesthetic' type gardeners also included a wild  
32  
33 347 area, which suggests that there is an aesthetic appeal to wildness, which is in agreement  
34  
35  
36 348 with the results of Home et al. (2010) who found that cultivated wildness was  
37  
38 349 considered attractive by Swiss residents. Neither 'functional' gardeners nor the 'child  
39  
40  
41 350 friendly' gardener cultivated vegetables in their garden, while one of the 'minimum  
42  
43 351 effort' gardeners did cultivate vegetables, which was not expected. However, the finding  
44  
45  
46 352 that 'child friendly' gardeners did not cultivate vegetables further supports the  
47  
48 353 hypothesis that 'child friendly' gardeners are a subset of Kettle's (2014) 'non-gardening'  
49  
50  
51 354 gardeners.

52  
53  
54 355 Generally, respondents indicated high levels of wellbeing across all indicators. However,  
55  
56  
57 356 the overall wellbeing index for the 'minimum effort' gardeners (2.33) is much lower  
58  
59 357 than that for the other gardener types (all >4.20). Although it cannot be statistically  
60  
61 358 tested with a sample size of 14, the results suggest that just having a garden does not in  
62  
63  
64  
65

359 itself enhance well-being because, if that were the case, it could be expected that all  
1  
2 360 gardener types would have the same wellbeing outcomes. The results rather suggest  
3  
4  
5 361 that it is the actual act of gardening: in which the 'minimum effort' gardeners are less  
6  
7 362 involved in than the others, is important for increasing wellbeing. Conversely, the result  
8  
9  
10 363 might also be due to self-selection: If places other than the garden are better for  
11  
12 364 enhancing the wellbeing of a 'minimum-effort' gardener, they will only invest minimum  
13  
14  
15 365 time and effort in the garden.

16  
17  
18 366 Biodiversity indicator results according to gardener type showed a pattern, indicating  
19  
20  
21 367 possible correlation between gardener type and biodiversity, which is likely mediated  
22  
23 368 by management practices and variety in habitat types and structural elements:  
24  
25  
26 369 especially for plants and ground-dwelling species. The 'conservationist' gardeners had,  
27  
28 370 on average, a higher variety of plants, ground dwelling arthropods and flying insects  
29  
30  
31 371 than the other groups. These gardeners also reported being highly motivated to promote  
32  
33 372 and preserve biodiversity, and this may be reflected in the outcomes of their garden.

34  
35  
36  
37 373 Interestingly, the 'minimum effort' gardeners, who do not focus on cultivation, (Kettle,  
38  
39 374 2014) had lower biodiversity than the 'conservationist' gardeners for all 3 indicators,  
40  
41  
42 375 and the lowest flying insect diversity of any group, which suggests that management can  
43  
44 376 enhance biodiversity. These results fit with the Intermediate Disturbance Hypothesis  
45  
46  
47 377 (Catford et al., 2012), which suggests that both extremes of a management gradient (no-  
48  
49 378 management and extreme intense management) lead to a reduction of available niches  
50  
51  
52 379 and thus to a lower number of coexisting species.

53  
54  
55 380 Gardens managed by the two 'functional' gardeners were remarkably similar in species  
56  
57  
58 381 numbers of vascular plants and ground dwelling arthropods (mean=42.5), with the  
59  
60 382 latter similar in number to that found in the 'minimum effort' gardeners' gardens  
61  
62  
63  
64  
65

383 (mean=47). The Q-methodology revealed that these gardeners were not motivated by  
1  
2 384 worry about the effects of pesticides and herbicides, although the minimum number of  
3  
4  
5 385 ground dwelling arthropods (28) in a 'conservationist' gardener's garden was found to  
6  
7 386 be lower than the minimum found in either the 'minimum effort' (40) or 'functional'  
8  
9  
10 387 gardeners' (41) gardens.

11  
12  
13 388 The results show that 'conservationist' and 'minimum effort' gardeners had, on average,  
14  
15  
16 389 the highest values for stable aggregates, for which high values have several benefits for  
17  
18 390 the soil functioning as detailed by Karlen et al. (2008). The high values for stable  
19  
20  
21 391 aggregates in gardens managed by 'minimum effort' gardeners can be explained because  
22  
23 392 these gardeners either never, or no more than once a year, dig up their vegetable or  
24  
25  
26 393 flower beds or break up the soil there with a fork. However, all garden types had  
27  
28 394 average stable aggregate values well above 65%, the common value of top soil for no  
29  
30  
31 395 tillage systems stated by Beare et al. (1994), which indicates that soil structure is  
32  
33 396 enhanced compared to agricultural sites: even within the groups who dig up or break up  
34  
35  
36 397 the soil more often.

37  
38  
39 398 The phosphorous levels were highest for 'minimum effort' gardeners. These levels  
40  
41  
42 399 suggest the addition of mineral fertilizer: especially in the minimum effort and aesthetic  
43  
44 400 gardener groups. Soluble phosphorus contents were significantly higher in the disturbed  
45  
46  
47 401 areas, which suggest that they had received more fertilizers than the undisturbed areas  
48  
49 402 such as lawn and berry cultures.

50  
51  
52  
53 403 For all groups, the mean values of microbial biomass were between 600-900 mg/kg,  
54  
55 404 which is approximately equivalent to levels found in organic no-till systems (Karlen et  
56  
57  
58 405 al. 1994) and higher than in conventional Swiss agricultural soils (Krauss et al., 2017).  
59  
60 406 The values were higher in the undisturbed garden areas, which may be explained by the

407 higher rooting density and the perennial cover that permanently provides food  
1  
2 408 resources, such as rhizoexudates, to soil microorganisms and act as hot spots for  
3  
4  
5 409 microbial activity in soils.  
6  
7

8 410 One of the main characteristics of urban soils is their relatively high variability and the  
9  
10  
11 411 patchiness of soil quality due to the high number of cultivated plant species and changes  
12  
13 412 in the soil types due to construction or land use history (Edmondson et al., 2014).  
14

15  
16 413 However, in the case of urban garden soils, gardening activities have the potential to  
17  
18 414 improve soil quality significantly (Tresch et al., 2018). This may also be reflected in our  
19  
20  
21 415 data set in the relatively high values of soil stable aggregates and microbial biomass.  
22  
23

## 24 416 **6. Conclusions**

25  
26  
27  
28 417 The case study gardens were quite intensively examined, with high demands on the  
29  
30  
31 418 participating gardeners due to multiple visits by ecologists and soil scientists along with  
32  
33 419 lengthy interviews by sociologists, in what is, by definition, the private sphere of the  
34  
35  
36 420 garden owners. Although this study found several apparent links between attitudes  
37  
38 421 towards gardening, management practices, and social, ecological and soil outcomes, the  
39  
40  
41 422 available resources did not allow a larger sample, and the reasonably small sample size  
42  
43 423 makes generalizations difficult. It was also beyond the scope of this exploratory study to  
44  
45  
46 424 increase the number of interviews and consider other factors that potentially influence  
47  
48 425 our indicators, such as demographics, the landscape composition (i.e. relative proportions  
49  
50  
51 426 of land-cover types) and configuration (i.e. spatial distribution of land cover types such as  
52  
53 427 buildings, impervious areas, etc.) (Braaker et al. 2014). It will be the challenge for future  
54  
55  
56 428 research to consider these factors in encompassing studies with larger samples.  
57

58  
59 429 Despite these limitations, we were able to observe some patterns in the collected data.  
60

61 430 The degree of wellbeing experienced by the responding gardeners did not appear to be  
62  
63  
64  
65

431 related to gardener type with the exception of the minimum-effort gardener who  
1  
2 432 reported an overall lower score on the wellbeing scale. This finding makes intuitive  
3  
4  
5 433 sense in light of the degree of control that the gardener has over the design and  
6  
7 434 implementation of their gardens. Gardeners can make their gardens to suit themselves,  
8  
9  
10 435 and it appears that they each achieve some satisfaction with their results. The gardener  
11  
12 436 who let the garden design itself also probably had a lower degree of engagement with  
13  
14  
15 437 the garden. This result suggests that it might be the act of gardening, rather than the  
16  
17 438 ownership of the garden that leads to wellbeing.

18  
19  
20  
21 439 The garden types did however appear to be related to the measures of biodiversity and  
22  
23 440 soil quality. The most likely explanation for this connection is that the attitudes of the  
24  
25  
26 441 gardeners are reflected in the practices along with the garden elements that have been  
27  
28 442 installed in the gardens. Indeed, the ‘conservationist’ gardeners tended to avoid  
29  
30  
31 443 pesticide use more, and to have a larger range of landscape elements, which provide a  
32  
33 444 larger range of habitats, than did the other gardener types. Similarly, the garden types  
34  
35  
36 445 also appear to be related to the measures of soil quality. In this case, the  
37  
38 446 ‘conservationist’ gardeners and the ‘minimum effort’ gardeners appear to enhance soil  
39  
40  
41 447 quality indicators, which appears to primarily reflect practices.

42  
43  
44 448 The study identified some indications that there might indeed be relationships between  
45  
46  
47 449 garden practices and the outcomes on biodiversity, soil, and human well-being. If these  
48  
49 450 links are established quantitatively, there are implications for city management who  
50  
51  
52 451 may have an interest in encouraging practices they consider desirable. We have  
53  
54 452 attempted to measure the outcomes of practices on indicators in three dimensions;  
55  
56  
57 453 however, we suspect interactions between the three dimensions might also occur. For  
58  
59 454 example, the conservationist gardeners’ gardens had the highest mean number of  
60  
61 455 vascular plants and also the highest mean microbial biomass (Cmic), which raises the  
62  
63  
64  
65

456 suspicion that there might be a relationship between them. However, this may also be  
1  
2 457 due to the small sample size, so further study of a larger sample would be needed to  
3  
4  
5 458 confirm this suspicion.  
6  
7

8 459 The findings of this study suggest that the beneficial effects of gardens are indeed  
9  
10  
11 460 related to the practices implemented by the gardeners, which in turn appears to be  
12  
13 461 influenced by the attitudes held by the gardeners. Gardeners design and implement  
14  
15  
16 462 practices in their gardens so that their gardens are how they like them. Gardeners with  
17  
18 463 more positive attitudes towards nature were found to manage gardens with higher plant  
19  
20  
21 464 and animal species richness along with more favourable soil quality measurements. The  
22  
23 465 challenge for ecologists seeking to enhance the ecological quality of urban domestic and  
24  
25  
26 466 allotment gardens will be to convince gardeners that ecologically friendly gardening is  
27  
28 467 compatible with other desired social outcomes.  
29  
30

### 31 **Acknowledgements**

32 468  
33  
34 469 This research was funded by the Sinergia fund of the Swiss National Science Foundation.  
35  
36  
37 470 We are also grateful to the many students who assisted with the data collection and to  
38  
39 471 the 18 gardeners who generously gave up their privacy by opening their gardens to the  
40  
41  
42 472 very many visits by researchers over the course of this project. The study was conducted  
43  
44 473 in compliance with all relevant ethical requirements in Switzerland.  
45  
46  
47 474

## 48 49 **7. List of references**

50  
51  
52  
53 476 Agroscope (1996) Schweizerische Referenzmethoden der Eidgenössischen  
54  
55 477 landwirtschaftlichen Forschungsanstalten. Eidg. Forschungsanstalt für  
56  
57  
58 478 Landwirtschaftlichen Pflanzenbau, FAP,  
59  
60  
61  
62  
63  
64  
65

- 479 Beare M, Hendrix P, Coleman D (1994) Water-stable aggregates and organic matter  
1  
2 480 fractions in conventional-and no-tillage soils Soil Science Society of America  
3  
4  
5 481 Journal 58:777-786  
6
- 7 482 Braaker S, Ghazoul J, Obrist M, Moretti M (2014) Habitat connectivity shapes urban  
8  
9  
10 483 arthropod communities: the key role of green roofs Ecology 95:1010-1021  
11
- 12 484 Brejda JJ, Moorman TB, Smith JL, Karlen DL, Allan DL, Dao TH (2000) Distribution and  
13  
14  
15 485 variability of surface soil properties at a regional scale Soil Science Society of  
16  
17 486 America Journal 64:974-982  
18
- 19  
20 487 Brown SR (1996) Q methodology and qualitative research Qualitative health research  
21  
22 488 6:561-567  
23
- 24  
25 489 Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, Mace GM,  
26  
27 490 Tilman D, Wardle DA, Kinzig AP, Daily GC, Loreau M, Grace JB, Larigauderie A,  
28  
29  
30 491 Srivastava DS, Naeem S (2012) Biodiversity loss and its impact on humanity  
31  
32 492 Nature 486:59-67  
33
- 34  
35 493 Catford JA, Daehler CC, Murphy HT, Sheppard AW, Hardesty BD, Westcott DA, Rejmanek  
36  
37 494 M, Bellingham PJ, Pergl J, Horvitz CC, Hulme PE (2012) The intermediate  
38  
39  
40 495 disturbance hypothesis and plant invasions: Implications for species richness and  
41  
42 496 management Perspectives in Plant Ecology, Evolution and Systematics 14:231-  
43  
44 497 241  
45
- 46  
47 498 Davies ZG, Fuller RA, Loram A, Irvine KN, Sims V, Gaston KJ (2009) A national scale  
48  
49  
50 499 inventory of resource provision for biodiversity within domestic gardens  
51  
52 500 Biological Conservation 142:761-771  
53
- 54  
55 501 Drescher A, Holmer R, Iaquina D (2006) Urban homegardens and allotment gardens for  
56  
57 502 sustainable livelihoods: management strategies and institutional environments.  
58  
59 503 Tropical Homegardens, Springer, The Netherlands, pp 317-338  
60  
61  
62  
63  
64  
65

- 504 Duelli P, Obrist MK, Schmatz DR (1999) Biodiversity evaluation in agricultural  
1  
2 505 landscapes: above-ground insects Agriculture, Ecosystems & Environment 74:33-  
3  
4 506 64  
5  
6  
7 507 Edmondson JL, Davies ZG, Gaston KJ, Leake JR (2014) Urban cultivation in allotments  
8  
9 508 maintains soil qualities adversely affected by conventional agriculture Journal of  
10  
11 509 Applied Ecology 51:880-889  
12  
13  
14 510 Freeman C, Dickinson KJ, Porter S, van Heezik Y (2012) “My garden is an expression of  
15  
16 511 me”: Exploring householders' relationships with their gardens Journal of  
17  
18 512 Environmental Psychology 32:135-143  
19  
20  
21 513 Frick J, Degenhardt B, Buchecker M (2007) Predicting local residents' use of nearby  
22  
23 514 outdoor recreation areas through quality perceptions and recreational  
24  
25 515 expectations For Snow Landsc Res 81:2  
26  
27  
28 516 Gaston KJ, Warren PH, Thompson K, Smith RM (2005) Urban domestic gardens (IV): the  
29  
30 517 extent of the resource and its associated features Biodiversity and conservation  
31  
32 518 14:3327-3349  
33  
34  
35 519 Goddard MA, Dougill AJ, Benton TG (2013) Why garden for wildlife? Social and  
36  
37 520 ecological drivers, motivations and barriers for biodiversity management in  
38  
39 521 residential landscapes Ecological Economics 86:258-273  
40  
41  
42 522 Gruen Stadt Zürich (2010), Biotoptypenkartierung der Stadt Zürich, [https://www.stadt-  
43  
44 523 zuerich.ch/ted/de/index/gsz/planung\\_u\\_bau/inventare\\_und\\_grundlagen/naturschut  
45  
46 524 z-inventar\\_und\\_kartierungen.secure.html](https://www.stadt-zuerich.ch/ted/de/index/gsz/planung_u_bau/inventare_und_grundlagen/naturschutz-inventar_und_kartierungen.secure.html)  
47  
48  
49  
50  
51  
52 525 Hall DM, Camilo GR, Tonietto RK, Ollerton J, Ahrne K, Arduser M, Ascher JS, Baldock  
53  
54 526 KCR, Fowler R, Frankie G, Goulson D, Gunnarsson B, Hanley ME, Jackson JI,  
55  
56 527 Langellotto G, Lowenstein D, Minor ES, Philpott SM, Potts SG, Sirohi MH, Spevak  
57  
58  
59  
60  
61  
62  
63  
64  
65



- 528 EM, Stone GN, Threlfall CG (2017) The city as a refuge for insect pollinators  
1  
2 529 Conservation Biology 31:24-29  
3  
4  
5 530 Haynes R, Tregurtha R (1999) Effects of increasing periods under intensive arable  
6  
7 531 vegetable production on biological, chemical and physical indices of soil quality  
8  
9  
10 532 Biology and Fertility of Soils 28:259-266  
11  
12 533 Home R, Bauer N, Hunziker M (2010) Cultural and biological determinants in the  
13  
14  
15 534 evaluation of urban green spaces Environment and Behavior 42:494-523  
16  
17 535 Home R, Hunziker M, Bauer N (2012) Psychosocial outcomes as motivations for visiting  
18  
19  
20 536 nearby urban green spaces Leisure Sciences 34:350-365  
21  
22 537 Irvine KN, Warber SL, Devine-Wright P, Gaston KJ (2013) Understanding urban green  
23  
24  
25 538 space as a health resource: A qualitative comparison of visit motivation and  
26  
27 539 derived effects among park users in Sheffield, UK International journal of  
28  
29  
30 540 environmental research and public health 10:417-442  
31  
32 541 Karlen D, Wollenhaupt N, Erbach D, Berry E, Swan J, Eash N, Jordahl J (1994) Long-term  
33  
34  
35 542 tillage effects on soil quality Soil and Tillage Research 32:313-327  
36  
37 543 Karlen DL, Andrews SS, Wienhold BJ, Zobeck TM (2008) Soil quality assessment: past,  
38  
39  
40 544 present and future Journal of Integrative Biosciences 6:3-14  
41  
42 545 Kettle P (2014) Motivations for investing in allotment gardening in Dublin: a sociological  
43  
44  
45 546 analysis Irish Journal of Sociology 22:30-63  
46  
47 547 Kiesling FM, Manning CM (2010) How green is your thumb? Environmental gardening  
48  
49  
50 548 identity and ecological gardening practices Journal of Environmental Psychology  
51  
52 549 30:315-327  
53  
54 550 Krauss M, Krause H, Spangler S, Kandeler E, Behrens S, Kappler A, Maeder P, Gattinger A  
55  
56  
57 551 (2017) Tillage system affects fertilizer-induced nitrous oxide emissions Biology  
58  
59  
60 552 and Fertility of Soils 53:49-59  
61  
62  
63  
64  
65

- 553 Lindemann-Matthies P, Marty T (2013) Does ecological gardening increase species  
1  
2 554 richness and aesthetic quality of a garden? *Biological Conservation* 159:37-44  
3  
4  
5 555 Muscolo A, Settineri G, Attinà E (2015) Early warning indicators of changes in soil  
6  
7 556 ecosystem functioning *Ecological Indicators* 48:542-549  
8  
9  
10 557 Obrist M, Duelli P (2010) Rapid biodiversity assessment of arthropods for monitoring  
11  
12 558 average local species richness and related ecosystem services *Biodiversity and*  
13  
14  
15 559 conservation 19:2201-2220  
16  
17 560 Sattler T, Duelli P, Obrist M, Arlettaz R, Moretti M (2010) Response of arthropod species  
18  
19  
20 561 richness and functional groups to urban habitat structure and management  
21  
22 562 *Landscape Ecology* 25:941-954  
23  
24  
25 563 Sattler T, Obrist MK, Duelli P, Moretti M (2011) Urban arthropod communities: Added  
26  
27 564 value or just a blend of surrounding biodiversity? *Landscape and Urban Planning*  
28  
29  
30 565 103:347-361  
31  
32 566 Smith RM, Warren PH, Thompson K, Gaston KJ (2006) Urban domestic gardens (VI):  
33  
34 567 environmental correlates of invertebrate species richness *Biodiversity and*  
35  
36  
37 568 *Conservation* 15:2415-2438  
38  
39  
40 569 Sullivan WC, Kuo FE, Depooter SF (2004) The fruit of urban nature: Vital neighborhood  
41  
42 570 spaces *Environment and Behavior* 36:678-700  
43  
44  
45 571 Thompson P (2007) *The Self-sustaining Garden: The Guide to Matrix Planting*. Timber  
46  
47 572 Press,  
48  
49  
50 573 Tresch S, Moretti M, Le Bayon R-C, Mäder P, Zanetta A, Frey D, Fliessbach A (2018). A  
51  
52 574 Gardener's Influence on Urban Soil Quality. *Frontiers in Environmental Science* 6  
53  
54  
55 575 Turner T (1986) *English garden design: history and style since 1650*. Antique Collectors'  
56  
57 576 Club Ltd.,  
58  
59  
60  
61  
62  
63  
64  
65

577 van Heezik Y, Freeman C, Porter S, Dickinson K (2014) Native and exotic woody  
1  
2 578 vegetation communities in domestic gardens in relation to social and  
3  
4  
5 579 environmental factors *Ecology and Society* 19  
6  
7 580 van Heezik Y, Freeman C, Porter S, Dickinson KJ (2013) Garden size, householder  
8  
9  
10 581 knowledge, and socio-economic status influence plant and bird diversity at the  
11  
12 582 scale of individual gardens *Ecosystems*:1442-1454  
13  
14  
15 583 Watts S, Stenner P (2005) *Doing Q methodology: theory, method and interpretation*  
16  
17 584 *Qualitative research in psychology* 2:67-91  
18  
19  
20 585 Yin RK (2009) *Case Study Research: Design and Methods*. SAGE Publications,  
21  
22 586 Zagorski T, Kirkpatrick J, Stratford E (2004) Gardens and the bush: gardeners' attitudes,  
23  
24  
25 587 garden types and invasives *Geographical Research* 42:207-220  
26  
27 588 Ziter C (2016) The biodiversity–ecosystem service relationship in urban areas: a  
28  
29  
30 589 quantitative review *Oikos* 125:761-768  
31  
32 590 Zornoza R, Acosta J, Bastida F, Domínguez S, Toledo D, Faz A (2015) Identification of  
33  
34  
35 591 sensitive indicators to assess the interrelationship between soil quality,  
36  
37 592 management practices and human health *Soil* 1:173  
38  
39  
40 593  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

*Table 1: Irvine et al.'s (2013) domains, with the corresponding statements and indicator names.*

Domain	Statement	Indicator name
Physical effects	When I work in the garden, I can let go and relax	Relaxed
	My motivation to spend time in garden is to recover	Revitalized
	My motivation to spend time in garden is to get some exercise	Exercise
Affective	My motivation to spend time in garden is to have a useful activity	Useful activity
	When I work in the garden, I can express myself	Express myself
Place attachment	I feel connected to my garden	Connected to garden
	I feel at one with my garden	At one with garden
Spiritual	The garden is a change from my everyday	Change
	I get feelings of satisfaction when I work in the garden	Satisfied
Cognitive	My garden is an escape from unwanted distractions	Escape distractions
	It's easier to concentrate when I work in the garden	Attention restoration
Social	My motivation to spend time in garden is to spend time with friends	Social

Table 2: Scores indicating where the statement is placed on a representative Q-sort for each of the 5 gardener types (from +3 "I agree the most" to -3 "I agree the least")

<i>Statement: I cultivate and manage my garden the way I do</i>		<i>Conservationist</i>	<i>Functional</i>	<i>Minimum effort</i>	<i>Child friendly</i>	<i>Aesthetic</i>
1	because I can harvest more like this	-1	-3	-2	0	-3
2	because it means no large investment costs are necessary	-2	-1	+3	-1	-2
3	because I can get direct environmental benefits like this (e.g. pest control)	+1	-3	-1	0	-1
4	because I've had good experiences with it so far	+1	+3	-1	+1	+1
5	because the garden is aesthetically beautiful to look at	0	+2	0	+1	+2
6	because I have an inner conviction it's the right way	+1	+3	+1	+2	+3
7	because this has been passed down in the family	-1	-3	-1	-2	+1
8	because I want to produce healthy food in the garden	+2	-3	-1	+1	0
9	because I want to contribute to the preservation and promotion of biodiversity	+3	0	+2	+1	+3
10	because I want to conserve natural resources (e.g. water)	0	0	0	-1	0
11	because I worry about the consequences of pesticide use	+1	-2	0	0	-1
12	because I like to have a neat garden	-2	+2	+2	+3	+2
13	because I want to keep the soil fertile	+2	0	-1	0	+1
14	because I am convinced everything in nature is interconnected and interdependent	+3	+1	0	-1	+2
15	because (my) children can play in the garden without danger	0	-2	-3	+3	+1
16	because I like that it has a lawn	-1	+1	+1	+2	-3
17	because my neighbours expect this from me	-3	0	-3	-1	-3
18	because this brings me social recognition by the neighbours	-1	-1	-2	-1	0
19	because I want to be a model for (my) children	0	-1	+2	-3	0
20	because I think that gardeners have an ecological responsibility	+2	+1	1	+2	0
21	because other gardeners do so	-3	-2	-2	-2	+1
22	because this takes little time	-2	-1	+2	-2	-1
23	because I like to learn about gardening and implement this knowledge	+2	+2	0	-3	-1
24	because this is very convenient	0	-1	+1	1	0
25	because it would not work without pesticides and artificial fertilizers	-3	+1	-3	-2	-2
26	because this takes little physical exertion	-2	0	+3	0	-2
27	because I have always done it this way	-1	-1	+3	0	-2
28	because other gardeners recommended this to me	0	0	-2	-3	-1
29	because I get joy from having biodiversity in the garden	+3	+3	+1	+2	+3
30	because I enjoy the challenge that this system provides	+1	1	0	+3	+2

*Table 3: Number of respondents with each structural element in their garden: classified according to gardener type*

	Conservationist (n=9)	Functional (n=2)	Minimum effort (n=2)	Child-friendly (n=1)	Aesthetic (n=4)
Lawn	5	2	1	1	0
Vegetable beds	8	0	1	0	3
Flower beds	8	2	1	1	4
Water feature	5	0	0	0	0
Wild area	7	0	1	0	3
Free hedge	6	0	1	0	1
Dry stone wall	3	1	0	0	0

Table 4: Biodiversity indicators per gardener type.

<b>Conservationist gardeners (n=9)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Biodiversity	Vascular plants	75	192	135	38
	Ground dwelling arthropods	28	79	59.7	17.3
	Flying insects	57	127	98.7	21.2
<b>Functional gardeners (n=2)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Biodiversity	Vascular plants	102	110	106	5.7
	Ground dwelling arthropods	41	44	42.5	2.1
	Flying insects	60	102	81	29.7
<b>Minimum effort gardeners (n=2)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Biodiversity	Vascular plants	68	114	91	32.5
	Ground dwelling arthropods	40	54	47	9.9
	Flying insects	54	74	64	14.1
<b>Child-friendly gardeners (n=1)</b>		<b>Value</b>			
Biodiversity	Vascular plants	71			
	Ground dwelling arthropods	33			
	Flying insects	83			
<b>Aesthetic gardeners (n=4)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Biodiversity	Vascular plants	86	117	99.3	13
	Ground dwelling arthropods	23	56	42.8	14.1
	Flying insects	82	116	95.3	14.7

Table 5: soil quality indicators per gardener type.

<b>Conservationist gardeners (n=9)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Soil quality	Stable aggregate (%)	64	94	84.06	6.02
	Phosphorous (mg/kg)	60.86	460.6	202.8	93.05
	Microbial biomass (Cmic) (mg/kg)	411.7	1343	861.8	207.63
<b>Functional gardeners (n=2)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Soil quality	Stable aggregate (%)	59	93	81.75	8.13
	Phosphorous (mg/kg)	66.51	196.1	119	29.60
	Microbial biomass (Cmic) (mg/kg)	279.9	1026	706.6	338.84
<b>Minimum effort gardeners (n=2)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Soil quality	Stable aggregate (%)	75	92	85.75	6.72
	Phosphorous (mg/kg)	168.3	465.2	305.2	118.84
	Microbial biomass (Cmic) (mg/kg)	468	862.8	684.6	27.18
<b>Child-friendly gardeners (n=1)</b>		<b>Value</b>			
Soil quality	Stable aggregate (%)	80			
	Phosphorous (mg/kg)	32.65			
	Microbial biomass (Cmic) (mg/kg)	846.8			
<b>Aesthetic gardeners (n=4)</b>		<b>Min</b>	<b>Max</b>	<b>Mean</b>	<b>Std.dev</b>
Soil quality	Stable aggregate (%)	47	93	76.62	10.23
	Phosphorous (mg/kg)	27.54	458.4	209.9	162.42
	Microbial biomass (Cmic) (mg/kg)	306.3	853.9	578.5	104.82






Figure 1. Well-being indicators for each gardener type

Comment	Response
<p>Page 3 Line 50 - 57. You state the aim of the study here. However, everything from p.4 Line 17 to p. 9 Line 32 is an extended discussion of methods that you used in your study. This is appropriate, however the 5 pages devoted to this discussion is too much. You should consider how to modify your introduction and methods section so that the introduction has a concise literature review and the methods section covers the approaches you used in your research.</p>	<p>This is a fair point. In the submitted draft, we included background information and justification for selection of each of the indicators, which we agree might have been excessive. The excess text has been removed, and the paper has been re-designed into a more conventional format.</p>
<p>Page 10 Line 17. Please explain what was studied using Q-methodology. One way to do this is to state what you want to determine about gardeners before you describe the analytical method.</p>	<p>We've added the text in (new) line 115: "The study was undertaken in two distinct phases. The first phase was to use Q-methodology to identify latent groups within the sample and thus identify a typology of gardeners based on their attitudes towards gardens and gardening".</p>
<p>In addition, some readers may wonder if Swiss institutions require some type of consent process before conducting research with human subjects. If this is required please state that your project was reviewed and approved.</p>	<p>This is not a requirement in Switzerland. The position of the Swiss Government is that agreement to participate in an interview implies consent to being interviewed. We added the statement "The study was conducted in compliance with all relevant ethical requirements in Switzerland" In the acknowledgements.</p>
<p>The combined results and discussion section is somewhat difficult to follow. The descriptions of the gardener types are very good. However it is difficult to determine what you found for the additional variables measured. Page 18 Line 54 for example, only refers to a table and the detail about what was found about biodiversity is missing. Separating the results and discussion sections would help.</p>	<p>We have separated the results and discussion sections, and we have taken care with internal cross referencing. This restructuring has made the text significantly easier to follow.</p>
<p>The statement numbers are confusing and may explain why the reviewer felt that there were missing citations. Including a data file with the survey text would be helpful, and linking to specific statements is not necessary.</p>	<p>The statement numbers have been removed. We believe it is reasonably clear what is being referred to in the discussion.</p>
<p>The conclusions do not always support what is written in the results section. For example, there are statements about correlations among environmental variables which are not found in the data presented in the tables and figures.</p>	<p>We have taken care to make sure that all conclusions are justified by the results. We do however raise questions about relationships that we suspect may exist, but have added new text to make it clear that the data is insufficient to confirm the suspected relationships.</p>
<p>A major flaw that lends to speculation of accuracy is the lack of proper citations. In multiple instances, citations were missing from statements within the manuscript.</p>	<p>We have paid close attention to ensure that the paper is correctly cited.</p>
<p>Of the 84 references, 33 were not cited within the manuscript itself. This is a serious error and</p>	<p>The superfluous references have been deleted from the reference list.</p>

leads to question the proper citation of statements within the manuscript.	
--	--