


RESEARCH

Open Access



Genetic variants of prospectively demonstrated phenocopies in *BRCA1/2* kindreds

Mev Dominguez-Valentin^{1*} , D. Gareth R. Evans^{2,3}, Sigve Nakken¹, H el ene Tubeuf^{4,5}, Daniel Vodak¹, Per Olaf Ekstr om¹, Anke M. Nissen^{6,7}, Monika Morak^{6,7}, Elke Holinski-Feder^{6,7}, Alexandra Martins⁴, P al M oller^{1,8,9} and Eivind Hovig^{1,10,11}

Abstract

Background: In kindreds carrying *path_BRCA1/2* variants, some women in these families will develop cancer despite testing negative for the family's pathogenic variant. These families may have additional genetic variants, which not only may increase the susceptibility of the families' *path_BRCA1/2*, but also be capable of causing cancer in the absence of the *path_BRCA1/2* variants. We aimed to identify novel genetic variants in prospectively detected breast cancer (BC) or gynecological cancer cases tested negative for their families' pathogenic *BRCA1/2* variant (*path_BRCA1* or *path_BRCA2*).

Methods: Women with BC or gynecological cancer who had tested negative for *path_BRCA1* or *path_BRCA2* variants were included. Forty-four cancer susceptibility genes were screened for genetic variation through a targeted amplicon-based sequencing assay. Protein- and RNA splicing-dedicated in silico analyses were performed for all variants of unknown significance (VUS). Variants predicted as the ones most likely affecting pre-mRNA splicing were experimentally analyzed in a minigene assay.

Results: We identified 48 women who were tested negative for their family's *path_BRCA1* ($n = 13$) or *path_BRCA2* ($n = 35$) variants. Pathogenic variants in the *ATM*, *BRCA2*, *MSH6* and *MUTYH* genes were found in 10% (5/48) of the cases, of whom 15% (2/13) were from *path_BRCA1* and 9% (3/35) from *path_BRCA2* families. Out of the 26 unique VUS, 3 (12%) were predicted to affect RNA splicing (*APC* c.721G > A, *MAP3K1* c.764A > G and *MSH2* c.815C > T). However, by using a minigene assay we here show that *APC* c.721G > A does not cause a splicing defect, similarly to what has been recently reported for the *MAP3K1* c.764A > G. The *MSH2* c.815C > T was previously described as causing partial exon skipping and it was identified in this work together with the *path_BRCA2* c.9382C > T (p.R3128X).

Conclusion: All women in breast or breast/ovarian cancer kindreds would benefit from being offered genetic testing irrespective of which causative genetic variants have been demonstrated in their relatives.

Keywords: *BRCA1*, *BRCA2*, Breast cancer, Gene panel testing, RNA splicing

Background

Breast cancer (BC) is one of the most common human malignancies, accounting for 22% of all cancers in women worldwide [1]. A significant proportion of BC cases can be explained by hereditary predisposition and approximately 30% of this hereditary cancer risk is explained by the currently known high-penetrance

susceptibility genes [2–5]. Notably, carriers of pathogenic *BRCA1* or *BRCA2* variants (*path_BRCA1* or *path_BRCA2*) have an increased risk of developing BC (average lifetime risk of 35–85%) and ovarian cancer (average lifetime risk 11–39%). Further, carriers of pathogenic variants of *ATM*, *CHEK2*, *PALB2*, *NBS1* and *RAD50* have been found to confer two- to five-fold increased risk for developing BC [1, 6]. It is also known that pathogenic variants in *TP53*, *PTEN*, *STK11* and *CDH1*, resulting in Li-Fraumeni syndrome, Cowden syndrome, Peutz–Jeghers syndrome and hereditary diffuse

* Correspondence: mev_dv@yahoo.com

¹Department of Tumor Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway

Full list of author information is available at the end of the article

gastric cancer, respectively, are associated with a high lifetime risk (> 40%) of BC. Moreover, pathogenic variants in *RAD51* paralogs, i.e., *RAD51C*, confer an increased risk of ovarian cancer [7]. The frequency of pathogenic variants in BC-associated genes varies significantly among different populations, as exemplified by the frequently studied founder pathogenic variant c.1100delC in *CHEK2* [6].

The identification of *path_BRCA1* or *path_BRCA2* in an affected BC individual enables access to evidence-based screening for family members, and thus facilitates the implementation of appropriate cancer prevention in these families [1, 5, 6]. However, some women in families with an identified pathogenic variant will develop cancer despite testing negative for the family's pathogenic variant, often denoted as phenocopies [8]. In BC kindreds having a demonstrated *path_BRCA2* variant, the number of phenocopies is reportedly more frequent than expected by chance [8–10]. It has been proposed that these families may have additional genetic variants, which not only may increase the susceptibility of the families' *path_BRCA1/2*, but also be capable of causing cancer in the absence of the *path_BRCA1/2* demonstrated in the families [5–7].

The current practice of genetic counselling for women who do not carry the *path_BRCA1/2* variants of their relatives is challenging since their recognition is crucial for application of proper diagnostic and therapeutic approaches in these families. To discover additional inherited disease-causing variants in *path_BRCA1/2* kindreds, we examined all prospectively detected BC or gynecological cancer cases in these kindreds by next-generation sequencing (NGS) using a panel of 44 cancer susceptibility genes. All detected variants were analyzed by RNA splicing- and protein-dedicated in silico methods. Variants predicted as the most likely to affect splicing were experimentally analyzed by using a cell-based minigene splicing assay.

Methods

Study population

For more than 20 years, we (the Hereditary Cancer Biobank from the Norwegian Radium Hospital, Norway; and the Department of Genomic Medicine from the University of Manchester, United Kingdom) have ascertained BC and breast/ovarian cancer kindreds by family history. The sisters and daughters of cancer patients were initially subjected to follow-up by annual mammography and gynecological examinations as appropriate at that time, and later they were all subjected to genetic testing [11].

Both collaborating outpatient genetic centers identified 48 women with prospectively detected BC or gynecological cancer at follow-up, who were tested negative for their

respective families' *path_BRCA1/2* variants. Clinical data were obtained from pathology reports and clinical files.

Ethical approval for the prospective study was granted from the Norwegian Data Inspectorate and Ethical Review Board (ref 2015/2382). All examined patients had signed an informed consent for their participation in the study.

Targeted sequencing

Genomic DNA was isolated from peripheral blood samples and targeted sequencing was carried out using a TrueSeq amplicon based assay v.1.5 on a MiSeq apparatus, as previously described [12]. The 44-gene panel used in this work includes genes associated with cancer predisposition as described in a prior study [12].

Sequencing data analysis

Paired-end sequence reads were aligned to the human reference genome (build GRCh37) using the BWA-mem algorithm (v.0.7.8-r55) [13]. The initial sequence alignments were converted to BAM format and subsequently sorted and indexed with SAMtools (v.1.1) [13]. Genotyping of single nucleotide variants (SNV) and short indels was performed by GATK's HaplotypeCaller. Filtering of raw genotype calls and assessment of callable regions/loci were done according to GATK's best practice procedures, as described more detail previously [12].

Variants were annotated using ANNOVAR (version November 2015) [14] and were queried against a range of variant databases and protein resources (v29, December 2015), as previously described [12].

Validation by cycling temperature capillary electrophoresis

The pathogenic variants identified in this study were validated by cycling temperature capillary electrophoresis. The method is based on allele separation by cooperative melting equilibrium while cycling the temperature surrounding capillaries [15]. This approach has previously been described and extensively used to detect somatic mutations and single nucleotide polymorphisms (SNPs) [16–19]. The amplicon design was performed by the variant melting profile tool (https://hyperbrowser.uio.no/hb/?tool_id=hb_variant_melting_profiles/) [20]. Primer sequences, PCR reaction conditions and electrophoresis settings are described in Additional file 1.

Genetic variants nomenclature and classification

The nomenclature guidelines of the Human Genome Variation Society (HGVS) were used to describe the detected genetic variants [21]. The recurrence of the identified variants was established by interrogating six databases (in their latest releases as of November 2016): Evidence-based Network for the Interpretation of Germline Mutant

Alleles (ENIGMA), Breast Cancer Information Core Database (BIC), the International Society of Gastrointestinal Hereditary Tumors (InSiGHT) Database, the Leiden Open Variation Database (LOVD), ClinVar, and the Human Gene Mutation Database (HGMD).

Novel variants were considered pathogenic if either one of the following criteria was met: a) introduced a premature stop codon in the protein sequence (nonsense or frameshift); b) occurred at positions +1/+2 or -1/-2 of donor or acceptor splice sites, respectively; and c) represented whole-exon deletions or duplications.

In silico analyses of VUS

Two types of bioinformatics methods were used to predict the impact of selected variants on RNA splicing. First, we used MaxEntScan (MES) and SSF-like (SSFL) to predict variant-induced alterations in 3' and 5' splice site strength, as described by Houdayer et al. [22], except that here both algorithms were interrogated by using the integrated software tool Alamut Batch version 1.5, (Interactive Biosoftware, <http://www.interactive-biosoftware.com>). For prediction of variant-induced impact on exonic splicing regulatory elements (ESR), we resorted to Δ ESRseq- [23], Δ HZeI- [24], and SPANR-based [25] as described by Soukarieh et al. [26]. Score differences (Δ) between variant and wild-type (WT) cases were taken as proxies for assessing the probability of a splicing defect. More precisely, we considered that a variant mapping at a splice site was susceptible of negatively impacting exon inclusion if Δ MES \geq 15% and Δ SSFL \geq 5% [22], whereas an exonic variant located outside the splice sites was considered as a probable inducer of exon skipping if negative Δ scores (below the thresholds described below) were provided by all the 3 ESR-dedicated in silico tools. We chose the following thresholds: <-0.5 for Δ ESRseq-, <-10 for Δ HZeI-, and <-0.2 for SPANR-based scores. In addition, we evaluated the possibility of variant-induced de novo splice sites by taking into consideration local changes in MES and SSFL scores. In this case, we considered that variants located outside the splice sites were susceptible of creating a competing splice site if local MES scores were equal to or greater than those of the corresponding reference splice site for the same exon.

In silico protein impact predictions of VUS were performed with FATHMM (<http://fathmm.biocompute.org.uk>) (v2.3), PolyPhen2-HVAR (v 2.2.2), MutationTaster (data release Nov 2015), MutationAssessor (release 3), SIFT (Jan 2015) and PROVEAN (v1.1 Jan 2015) using dbNSFP v3.4.

Cell-based minigene splicing assays

In order to determine the impact of the *APC* c.721G > A on RNA splicing, we performed functional assays based

on the comparative analysis of the splicing pattern of WT and mutant reporter minigenes [27], as follows. First, the genomic region containing *APC* exon 7 and at least 150 nucleotides of the flanking introns (c.646–169 to c.729 + 247) were amplified by PCR using patient #12470 DNA as template and primers indicated in Additional file 2. Next, the PCR-amplified fragments were inserted into a previously linearized pCAS2 vector [26] to generate the pCAS2-*APC* exon 7 WT and c.721G > A minigenes. All constructs were sequenced to ensure that no unwanted mutations had been introduced into the inserted fragments during PCR or cloning. Then, WT and mutant minigenes were transfected in parallel into HeLa cells grown in 12-well plates (at ~70% confluence) using the FuGENE 6 transfection reagent (Roche Applied Science). Twenty-four hours later, total RNA was extracted using the NucleoSpin RNA II kit (Macherey Nagel) and, the minigene transcripts were analyzed by semi-quantitative RT-PCR using the One-Step RT-PCR kit (QIAGEN), as previously described [26]. The sequences of the RT-PCR primers are shown in Additional file 2. Then, RT-PCR products were separated by electrophoresis on 2.5% agarose gel containing EtBr and visualized by exposure to UV light under saturating conditions using the Gel Doc XR image acquisition system (Bio-Rad), followed by gel-purification and Sanger sequencing for proper identification of the minigenes' transcripts. Finally, splicing events were quantitated by performing equivalent fluorescent RT-PCR reactions followed by capillary electrophoresis on an automated sequencer (Applied Biosystems), and computational analysis by using the GeneMapper v5.0 software (Applied Biosystems).


Results

Family history and clinical characteristics

In total, we identified 48 cases, of whom 18 BC or gynecological cancer patients who did not carry their respective families' *path_BRCA1* or *path_BRCA2* variants ($n = 13$ and $n = 5$, respectively) came from the Hereditary Cancer Biobank from the Norwegian Radium Hospital, while the Department of Genomic Medicine from the University of Manchester identified a total of 30 BC patients, all non-carriers of the family's *path_BRCA2* variants (Fig. 1). The median age at first cancer diagnosis was 53.5 years (range 31–79 years). The incidence was higher for BC (92%), followed by ovarian cancer (4%) and endometrial and cervical cancer (2% each) (Table 1).

Germline findings

In the 48 cases, we identified five (10%) to carry pathogenic variants in *ATM* (c.468G > A, p.Trp156Ter and c.9139C > T, p.Arg3047Ter), *BRCA2* (c.9382C > T, p.Arg3128Ter), *MSH6* (c.2864delC, p.Thr955fs) and



MUTYH (c.1178G > A, p.Gly393Asp). Among these five cases, 2/13 were identified in non-carriers of the family's *path_BRCA1* variant and in 3/35 non-carriers of the family's *path_BRCA2* variant (Fig. 1). Disease type, familial *path_BRCA1/2* and pathogenic variants found in this study are shown in detail in Table 1.

Interestingly, one case with a familial *path_BRCA2* (c.6591_6592delTG) was found to carry another pathogenic variant in the same gene (*BRCA2* c.9382C > T, p.Arg3128Ter), which causes a premature stop in the codon 3128 and is known to be a high risk pathogenic variant (Table 1).

The pathogenic variants in BC-related genes (2 in *ATM* and 1 in *BRCA2*) were found in 3 women with BC or ovarian cancer, while the *MSH6* and the heterozygous *MUTYH* p.Gly393Asp pathogenic variant was found in a woman with endometrial cancer at 57 years and BC diagnosis at 56 years, respectively (Table 1).

Validation of the cancer gene panel output

The presence of the five pathogenic variants detected by targeted NGS was confirmed by cycling temperature capillary electrophoresis, showing 100% correspondence between both methods.

Variants of unknown significance (VUS) and predicted protein alterations

In total, we found 26 unique VUS in 30 out of 48 patients (63%). Common polymorphisms (with an allele frequency $\geq 1\%$ in the general population according to the ExAC database) and benign variants classified according to either ClinVar or the American College of Medical Genetics and Genomics (ACMG) guidelines were excluded from further analyses [41, 58].

The VUS were detected in 17 genes, namely: *AXIN2*, *RAD51B* (in 4 patients each), *MAP3K1* (in 3 patients), *APC*, *ATM*, *MSH2*, *NBN*, *POLE* (in 2 patients each), *BRCA1*, *CDH1*, *CDX2*, *DVL2*, *MRE11A*, *MUTYH*, *NOTCH3*, *PTEN* and *RAD51D* (in 1 patient each) (Table 2). The minor allele frequencies (MAF) of these variants in public databases were very low or no frequency data have been reported (Table 2).

The VUS were furthermore analyzed by using 6 in silico protein prediction tools with different underlying algorithms (Fig. 2). The *MRE11A* c.1139G > A and the *MUTYH* c.881G > A variants were suggested to have a potentially damaging effect on protein level by all six predictions programs. For the variants in the *MSH2*, *NBN*, *POLE* and *BRCA1* genes (*MSH2* c.815C > T, *NBN* c.283G > A, *POLE* c.2459 T > C and *BRCA1* c.1927A > G,

Table 1 Summary of the 48 prospective BC or gynecological cancer patients included in the study

Patient_ID	Institution	Familial path_ BRCA1 or path_BRCA2 variantFamilial path_BRCA1 or path_BRCA2 variant	ICD9 diagnosis (age)	Pathogenic variant identified in the current study
17,161	HCBNRH	<i>BRCA2 c.5217_5223delTTTAAGT</i> (<i>p.Tyr1739Terfs</i>) <i>BRCA2 c.5217_5223delTTTAAGT</i> (<i>p.Tyr1739Terfs</i>)	OC (67)	<i>ATM c.468G > A</i> (<i>p.Trp156Ter</i>)* <i>ATM c.468G > A</i> (<i>p.Trp156Ter</i>)*
6475	HCBNRH	<i>BRCA1 c.1011dupA (p.Val340Glyfs)</i> <i>BRCA1</i> <i>c.1011dupA (p.Val340Glyfs)</i>	BC (52)	<i>ATM c.9139C > T</i> (<i>p.Arg3047Ter</i>) <i>ATM</i> <i>c.9139C > T (p.Arg3047Ter)</i>
13,141	HCBNRH	<i>BRCA1 c.1072delC (p.Leu358Cysfs)</i> <i>BRCA1</i> <i>c.1072delC (p.Leu358Cysfs)</i>	EC (57)	<i>MSH6 c.2864delC</i> (<i>p.Thr955fs</i>)* <i>MSH6</i> <i>c.2864delC (p.Thr955fs)*</i>
1873	HCBNRH	<i>BRCA1 c.1556delA (p.Lys519Argfs)</i> <i>BRCA1</i> <i>c.1556delA (p.Lys519Argfs)</i>	MTHM (56), BC (70)	Not
5378	HCBNRH	<i>BRCA1 c.697_698delGT (p.Val233Asnfs)</i> <i>BRCA1</i> <i>c.697_698delGT</i> (<i>p.Val233Asnfs</i>)	BC (52)	Not
5180	HCBNRH	<i>BRCA1 c.5194-2A > C</i> <i>BRCA1 c.5194-2A > C</i>	BC (39)	Not
22	HCBNRH	<i>BRCA2 c.3847_3848delGT (p.Val1283Lysfs)</i> <i>BRCA2</i> <i>c.3847_3848delGT</i> (<i>p.Val1283Lysfs</i>)	BC (63)	Not
243	HCBNRH	<i>BRCA2 c.3847_3848delGT (p.Val1283Lysfs)</i> <i>BRCA2</i> <i>c.3847_3848delGT</i> (<i>p.Val1283Lysfs</i>)	CVC (41)	Not
5348	HCBNRH	<i>BRCA1 c.1556delA (p.Lys519Argfs)</i> <i>BRCA1</i> <i>c.1556delA (p.Lys519Argfs)</i>	BC (68)	Not
6031	HCBNRH	<i>BRCA1 c.1556delA (p.Lys519Argfs)</i> <i>BRCA1</i> <i>c.1556delA (p.Lys519Argfs)</i>	BC (66)	Not
6032	HCBNRH	<i>BRCA1 c.3228_3229delAG (p.Gly1077Alafs)</i> <i>BRCA1</i> <i>c.3228_3229delAG</i> (<i>p.Gly1077Alafs</i>)	OC (55)	Not
6207	HCBNRH	<i>BRCA1 c.697_698delGT (p.Val233Asnfs)</i> <i>BRCA1</i> <i>c.697_698delGT</i> (<i>p.Val233Asnfs</i>)	BC (47)	Not
8085	HCBNRH	<i>BRCA1 c.3228_3229delAG (p.Gly1077Alafs)</i> <i>BRCA1</i> <i>c.3228_3229delAG (p.Gly1077Alafs)</i>	BC (55), CC (66)	Not
11,717	HCBNRH	<i>BRCA1 c.1556delA (p.Lys519Argfs)</i> <i>BRCA1</i> <i>c.1556delA</i> (<i>p.Lys519Argfs</i>)	BC(42,57)	Not
12,470	HCBNRH	<i>BRCA1 c.3178G > T (p.Glu1060Ter)</i>	BC (39)	Not
13,023	HCBNRH	<i>BRCA2 c.5217_5223delTTTAAGT</i> (<i>p.Tyr1739Terfs</i>)	BC (59)	Not
15,529	HCBNRH	<i>BRCA2 c.4821_4823delTGAins</i>	BC (48)	Not
22,325	HCBNRH	<i>BRCA1 c.5047G > T</i> (<i>p.Glu1683Ter</i>)	BC (45)	Not
1,100,948	UM	<i>BRCA2 c.6591_6592delITG</i> (<i>p.Glu2198Asnfs</i>)	BC (44)	<i>BRCA2 c.9382C > T</i> (<i>p.Arg3128Ter</i>)
12,010,643	UM	<i>BRCA2 c.7360delA</i> (<i>p.Ile2454Phefs</i>)	BC (56)	<i>MUTYH c.1178G > A</i> (<i>p.Gly393Asp</i>)
75,443	UM	<i>BRCA2 c.5909C > A</i> (<i>p.Ser1970Ter</i>)	BC (55)	Not
88,295	UM	<i>BRCA2 c.7977-1G > C</i>	BC (44)	Not
64,949	UM	<i>BRCA2 c.5909C > A</i> (<i>p.Ser1970Ter</i>)	BC (55)	Not
67,723	UM		BC (46)	Not

Table 1 Summary of the 48 prospective BC or gynecological cancer patients included in the study (Continued)

Patient_ID	Institution	Familial path_ BRCA1 or path_ BRCA2 variant Familial path_ BRCA1 or path_ BRCA2 variant	ICD9 diagnosis (age)	Pathogenic variant identified in the current study
84,510	UM	<i>BRCA2</i> c.4866delA p.(Arg1622Serfs*14)	BC (67)	Not
13,007,862	UM	<i>BRCA2</i> c.5946delT (p.Ser1982Argfs)	BC (31)	Not
9,009,462	UM	<i>BRCA2</i> c.5909C > A (p.Ser1970Ter)	BC (67)	Not
900,178	UM	<i>BRCA2</i> c.6535_6536insA (p.Val2179Aspfs)	BC (49,77)	Not
10,005,829	UM	<i>BRCA2</i> c.9541_9554del p.(Met318CysfsTer13)	BC (38)	Not
10,007,016	UM	<i>BRCA2</i> c.632-1G > A	BC (51)	Not
10,003,959	UM	<i>BRCA2</i> c.6275_6276delTT (p.Leu2092Profs)	BC (55)	Not
12,852	UM	<i>BRCA2</i> c.1929delG (p.Arg645Glu fs)	BC (56)	Not
12,001,161	UM	<i>BRCA2</i> c.7958 T > C (p.Leu2653Pro)	BC (67)	Not
13,017,067	UM	<i>BRCA2</i> c.755_758delACAG (p.Asp252Val fs)	BC (74)	Not
688	UM	<i>BRCA2</i> c.1929delG (p.Arg645Glu fs)	BC (32)	Not
40,540	UM	<i>BRCA2</i> c.8535_8538delAGAG p.(Glu2846LysfsTer16)	BC (69)	Not
9,001,644	UM	<i>BRCA2</i> c.4965C > G (p.Tyr1655Ter)	BC (39, 45)	Not
89,205	UM	<i>BRCA2</i> c.5946delT (p.Ser1982Argfs)	BC (77)	Not
10,002,068	UM	<i>BRCA2</i> del exons 14–16	BC (37)	Not
10,004,590	UM	<i>BRCA2</i> c.2672dupT	BC (67,67)	Not
40,286	UM	<i>BRCA2</i> c.7069_7070delCT p.(Leu2357Val fsTer2)	BC (36,53)	Not
76,618	UM	<i>BRCA2</i> c.4478_4481delAAAG (p.Glu1493Val fs)	BC (51)	Not
12,015,576	UM	<i>BRCA2</i> c.9382C > T (p.Arg3128Ter)	BC (45)	Not
61,420	UM	<i>BRCA2</i> c.5350_5351delAA p.(Asn1784HisfsTer2)	BC (59)	Not
960,579	UM	<i>BRCA2</i> c.2808_2811del4 (p.Ala938Profs)	BC (39)	Not
14,965	UM	<i>BRCA2</i> c.5682C > G p.(Tyr1894Ter)	BC (59)	Not
20,468	UM	<i>BRCA2</i> c.6275_6276delTT (p.Leu2092Profs)	BC (38)	Not
56,193	UM	<i>BRCA2</i> c.7884dupA (p.Trp2629Metfs)	BC (79)	Not

HCBNRH Hereditary Cancer Biobank from the Norwegian Radium Hospital (Norway), *UM* University of Manchester (United Kingdom), *ICD9 diagnosis* International Classification of Diseases, 9th Revision, *OC* Ovary cancer, *BC* Breast cancer, *EC* Endometrial cancer, *MTHM* Malignant neoplasm of thymus, heart, and mediastinum, *CC* Colon cancer, *CVC* Cervical cancer, *Considered pathogenic based in its nature (nonsense and frameshift), *VUS* Variants of unknown significance, *NM* for *ATM* NM_000051, *BRCA1* NM_007294.3, *BRCA2* NM_000059.3, *MSH6* NM_001281492, *MUTYH* NM_012222

five out of six predictions suggested a potentially damaging effect (Fig. 2).

Discrepancies in protein-related predictions were even more pronounced for the variants in *APC*, *AXIN2*,

RAD51B, *DVL2*, *RAD51D*, *CDH1* and *MSH2* c.2164G > A. In contrast, none of the six prediction tools showed deleterious effects for the detected variants in the *AXIN2*, *ATM*, *RAD51B* and *MAP3K1* genes (*AXIN2*


Table 2 RNA splicing-dedicated in silico analyses for the VUS identified in our study (Continued)

Patient ID	Genomic position (GRCh37)	Gene	Exon	Nucleotide change (cNomen)	Predicted protein change (pNomen)	dbSNPsID	Non-Finnish European population frequency*	Reference splice site-dedicated analyses	MES scores				SFSL scores				Cryptic splice site-dedicated analyses	Local MES scores	ESR-dedicated analyses	ΔHzei	ΔVP			
									WT		Var		WT		Var							Potential local splice effect	WT	Var
									WT	Var	WT	Var	WT	Var	WT	Var								
960, 579	chr_5_56 177843 _C_G	MAP3K1	14	c.28 16C> G	p.Ser939 Cys	45,556, 841	0.0221	447	12.0 063	12.0 0063	0	100	100	0	-	-	-0.4 86881	-16.1	0					
1,000, 459	chr13_28 537449 _ACTT_A	CDX2	3	c.742 _744 del	p.Lys248de IAAG	553, 066, 746	0.000 1682	55	11.0 7045	11.0 7045	0	87.0 4307	87.0 4307	0	-	-	-2.4 6964	-100.08	-					
1,100, 948	chr_17_7133187 _A_G	DVL2	5	c.59 6T> C	p.Met199 Thr	372, 715, 697	6.01e-05	-61	6.34 467	6.34 467	0	80.0 4452	80.0 4452	0	-	-	0.0 509 416	-1.77	0.54					
10,002, 068	chr_17_63526198 _C_T	AXIN2	11	c.24 28G> A	p.Asp810 Asn	140, 344, 858	1.5e-05	23	11.0 6727	11.0 6727	0	87.0 3948	87.0 3948	0	-	-	-1.2 2987	-14.33	-					
10,005, 829	chr_14_69061228 _G_A	RAD51B	11	c.10 63G> A	p.Ala355 Thr	61,758, 785	0.007 1658	27	11.8	11.8	0	80.2	80.2	0	-	-	-1.2 4035	-50.64	-					
1,100, 093640	chr8_9093640 _C_T	NBN	3	c.28 3G> A	p.Asp95	61,753, 720	0.00 30459	-38	10.0 7663	10.0 7663	0	94.0 6711	94.0 6711	0	-	-	0.31 8238	24.6	0.03					
8155132	chr_11_10 8155132 _G_A	ATM	26	c.39 25G> A	p.Ala1309 Thr	149, 711, 770	0.000 9147	-29	9.9 8517	9.9 8517	0	84.0 8076	84.0 8076	0	-	-	0.67 6556	32.96	0.04					
12,001, 161	chr_14_68353893 _A_G	RAD51B	7	c.72 8A> G	p.Lys243 Arg	34,594, 234	0.01 0682	-29	9.09184 9184	9.09184 9184	0	78.0 9497	78.0 9497	0	0.9	7.9	-1.4 8785	-40.0 54	-0.19					
12,015, 576	chr19_15291551 _C_G	NOTCH3	19	c.30 83G> C	p.Trip1028 Ser	rs14682 9488	na	-60	11.0 1124	11.0 1124	0	82.0 5954	82.0 5954	0	-	-	0.3 00 115	-6.4	0.1					
11,717	chr1_45797881 _C_T	MUTYH	10	c.88 1G> A	p.Cys294 Tyr	r8792 54257	na	-44	6.0 31089	6.0 31089	0	72.0 818	72.0 818	0	-	-	1.0 94 96	-7.06	0.04					
17, 161	chr_11_108139187 _T_A	ATM	18	c.26 89T> A	p.Phe897Ile	147, 122,522	4.5e-05	51	9.0 8979	9.0 8979	0	93.0 4253	93.0 4253	0	-	-	0.5 54 269	87.95	0.01					
22	chr_12_133241897 _A_G	POLE	21	c.24 59T> C	p.Met820 Thr	767, 460, 640	0	-10	6.0 58677	6.0 58677	0	77.0 9039	77.0 9039	0	-	-	1.2 8743	-2.13	0.06					

Table 2 RNA splicing-dedicated in silico analyses for the VUS identified in our study (Continued)

Patient ID	Genomic position (GRCh37)	Gene	Exon	Nucleotide change (cNomen)	Predicted protein change (pNomen)	dbSNPsID	Non-Finnish European population frequency*	Reference splice site-dedicated analyses	MES scores		SSFL scores		Cryptic splice site-dedicated analyses	ESR-dedicated analyses
									WT	Var	WT	Var		
chr_14_68 352672 _A_G	RAD 518	6	c.53 9A> G	p.Tyr1 80Cys	28, 910, 275	0.00 45906	-34	5'	9.	9.	83.	83.	Potential local splice effect	0.8 81 539
									54919	54919	7411	7411		
chr_5_5 6155672 _A_G	MAP 3K1	3	c.76 4A> G	p.Asn 255Ser	56,069, 227	0.0269	-71	5'	7.	7.	78.	78.	New Acceptor Site?	-1.1 8661
									52484	484	4708	4708		
6207	chr_17_63 530163 _C_T	AXIN2	c.22 72G> A	p.Ala 758 Thr	145, 007, 501	0.003 9861	35	3'	6.	6.	86.	86.	-0.9 42 617	0.12 09
									34671	671	1925	1925		
6475	chr_17_33 433488 _G_A	RAD 51D	c.49 3C> T	p.Arg 165 Trp	544, 654, 228	6.94 e-05	13	3'	8.	8.	85.	85.	-2.5 5724	-22. 32
									20686	686	1161	1161		

na not available; *Non-Finnish European population based on ExAC database; NM for APC: NM_000038; ATM: NM_000051; AXIN2: NM_004655; BRCA1: NM_007300; CDH1: NM_004360; CDX2: NM_001265; DVL2: NM_004422; MAP3K1: NM_005921; MSH2: NM_000251; MRE11A: NM_005591; MUTYH: NM_012222; NBN: NM_002485; NOTCH3: NM_000435; POLE: NM_006231; PTEN: NM_000314; RAD51B: NM_133509; RAD51D: NM_002878. In order to predict their biological impact, RNA splicing-dedicated bioinformatics analyses were performed as described under Materials and Methods. Results shown in bold were considered as predictive of a potential variant-induced negative biological effect. MES MaxEntScan, SSFL Splice Site Finder-Like, nt Nucleotide, 3' or 5'ss 3' splice site or 5' splice site, ESR Exonic splicing regulators




c.2272G > A, *ATM* c.2689 T > A, *RAD51B* c.539A > G and c.1063G > A and *MAP3K1* c.764A > G) (Fig. 2).

Splicing-dedicated in silico analysis and minigene splicing assays

Out of the 26 unique VUS, two (*APC* c.721G > A and *MAP3K1* c.764A > G) were bioinformatically predicted as the most likely to affect RNA splicing, either by potentially creating a new splice site or by altering putative exonic splicing regulatory elements, respectively (Table 2). Given that RNA data was not available for *APC* c.721G > A, we set out to experimentally evaluate the impact on RNA splicing produced by this variant, by performing a cell-based minigene splicing assay. As shown in Fig. 3, we observed that c.721G > A did not affect the splicing pattern of *APC* exon 7 in our system. These results are reminiscent of those recently obtained for *MAP3K1* c.764A > G by using a similar splicing assay, in which the variant did not cause an alteration in the minigene’s splicing pattern (Dominguez-Valentin et al. *under submission*). It would be important in both

cases to validate the minigene results by analyzing RNA from the variant carriers/patients as compared to those from healthy controls. However, we do not have such material in our biobank.

To our knowledge, the only other VUS from our list for which RNA data is available is *MSH2* c.815C > T (p.Ala272Val). Previous results from different minigene assays revealed that, albeit located outside the splice sites, *MSH2* c.815C > T induces partial skipping of exon 5 [28]. These results agree, at least in part, with those obtained by analyzing RNA from a LS patient carrying this same variant [29]. Indeed, the latter study revealed aberrantly spliced *MSH2* transcripts associated with the presence of c.815C > T, but where the severity of the splicing defect was not addressed at the time. Of note, here we identified *MSH2* c.815C > T together with another VUS (*DVL2* c.596 T > C) and a *path_BRCA2* c.9382C > T (different from the familial *path_BRCA2*) in a patient diagnosed with ductal carcinoma at 44 years of age (Patient 1,100,948) (Table 1).



Discussion

Among prospectively detected BC or gynecological cancer phenocopies in the *path_BRCA1/2* families, we found that 4/48 have pathogenic variants in high-penetrance cancer genes: two BC- and one CRC-associated gene (*ATM*, *BRCA2* and *MSH6*, respectively). Our findings are in line with a previous study, which detected a likely pathogenic variant in a gene other than *BRCA1/2* in a BC patient, i.e. *MSH6* c.3848_3862del (p.(Ile1283_Tyr1287del) [30]. In addition, we found the *MUTYH* c.1178G > A (p.Gly393Asp) variant in a BC case, which is one of the most common *path_MUTYH* variants. Pathogenic *MUTYH* variants may cause a recessively inherited colon cancer syndrome. Whether or not individuals who are heterozygous for *MUTYH* mutations may be at risk for cancer is debated [31]. Among the five cases found to carry pathogenic variants, 2/13 were identified from families with *path_BRCA1* and 3/35 with *path_BRCA2* variants.

Our results are in concordance with the recently published NGS panel studies, which have demonstrated that

besides high-risk genes, like *BRCA1/2* and MMR genes, other genes may also contribute to familial cancer predisposition, thus providing a broader picture on the genetic heterogeneity of cancer syndromes [25, 32, 33]. In this regard, a molecular diagnosis yield of approximately 9% to identify a pathogenic or likely pathogenic variant in BC has been reported, and with yields of 13% in ovarian and 15% in colon/stomach cancer cases [25]. On the other hand, family history is currently used to identify high risk patients. However, the use of family history fails to identify women without close female relatives who are carriers of pathogenic variants [9].

Despite the potential of NGS to identify genetic causes among families that tested negative for pathogenic variants in high-risk genes using traditional methods [25, 32, 33], a high number of VUS are also detected and constitute a major challenge in oncogenetics [34]. In this study, we subjected 26 VUS to RNA splicing and protein in silico evaluations, and the bioinformatics predictions indicated that two VUS (*APC* c.721G > A and *MAP3K1* c.764A > G) were likely to affect RNA splicing. Our

results from minigene splicing assays suggest, however, that this is not the case. Complementary analysis of patients' RNA will be important to verify the impact on splicing of these variants in vivo. Of note, none of the six protein in silico prediction tools showed a deleterious effect for the *MAP3K1* c.764A > G missense variant and inconsistencies were found for the *APC* c.721G > A variant.

Bioinformatics prediction tools are widely used to aid the biological and clinical interpretation of sequence variants, although it is well recognized that they have their limitations. Co-segregation studies for further evaluation will be key for understanding whether some of the VUS detected in this work may have a causal effect. Some of the VUS may in the future be reclassified as deleterious or benign, but in the meantime, they cannot be used to make clinical decisions [30].

A polygenic model involving a combination of multiple genomic risk factors, including the effect of low- or moderate- penetrance susceptibility alleles may explain the increased BC risk in women who tested negative for family's *path_BRCA1/2* variants [5]. In addition, heterozygous whole gene deletions (WGD) and intragenic microdeletions have been reported to account for a significant proportion of pathogenic variants underlying cancer predisposition syndromes, although WGD were not a common mechanism in any of the three high-risk BC genes, *BRCA1*, *BRCA2* and *TP53* [35].

The clinical utility of gene panels such as the one used in this study is not yet fully established and the appropriate routes for clinical deployment of such tests remain under discussion [36]. So far, the large patient datasets generated by NGS panels may be used to explore the specific penetrance of the genes included in these panels, and to assess the performance and implications of the use of NGS in clinical diagnostics [34].

Conclusions

In kindreds carrying *path_BRCA1/2* variants, testing only for the already known *path_BRCA1/2* variants in the family may not be sufficient to exclude increased risk neither for BC nor for ovarian cancer or other cancers in the healthy female relatives. Our findings suggest that all women in BC or breast/ovarian cancer kindreds would benefit from being offered genetic testing irrespective of which causative genetic variants have been demonstrated in their relatives. In addition, we found a number of VUS in genes other than *BRCA1/2* i.e. *AXIN2*, *APC*, *DVL2*, *MAP3K1*, *RAD51B*, *NBN*, *POLE*, *CDH1*, *CDX2*, *MRE11A*, *MUTYH*, *NOTCH3*, *PTEN* and *RAD51D*. All these may be suspected of being associated with cancer in the families studied and may be considered as candidates for being included in future gene

panel testing to better understand why some families present aggregation of cancer cases.

Additional files

Additional file 1: The concentration in a 10 ml PCR was 1xThermopol Reaction Buffer with 2 mM MgSO₄, 0.3 μM "reverse" primers, 0.15 μM "forward" primer, 0.1 μM, 6-Carboxyfluorescein-GC clamp primer, 600 μM dNTP, 100 μg Bovine Serum Albumine (Sigma-Aldrich, Oslo, Norway) and 0.75 U Taq DNA polymerase. Plates were sealed with two strips of electrical tape (Clas Ohlson, Oslo, Norway). The temperature cycling was repeated 35 times; 94 °C for 30 s, annealing temperature held for 30 s and extension at 72 °C for 60 s (Eppendorf Mastercycler ep gradient S (Eppendorf, Hamburg, Germany)). **Table S1.** primers used to amplify PCR product to be analysed by cycling temperature capillary electrophoresis. (DOCX 16 kb)

Additional file 2: Primers used in the pCAS2 minigene splicing assay. (DOCX 14 kb)

Abbreviations

ACMG: American College of Medical Genetics and Genomics; BC: Breast cancer; BIC: Breast Cancer Information Core Database; CRC: Colorectal cancer; ENIGMA: Evidence-based Network for the Interpretation of Germline Mutant Alleles; ESR: Exonic splicing regulatory elements; HGMD: Human Gene Mutation Database; InSIGHT: International Society of Gastrointestinal Hereditary Tumors Database; LOVD: Leiden Open Variation Database; LS: Lynch syndrome; MAF: Minor allele frequency; MES: MaxEntScan; NGS: Next generation sequencing; *path_BRCA1/2*: Pathogenic (disease-causing) variant of the *BRCA1* or the *BRCA2* genes; SNPs: Single nucleotide polymorphisms; SNV: Single-nucleotide variants; SSFL: SSF-like; VUS: Variants of unknown significance; WGD: Whole gene deletions; WT: Wild type

Acknowledgements

We thank the included families for their contribution to this study.

Funding

This work was supported by the Radium Hospital Foundation (Oslo, Norway), Helse Sør-Øst (Norway), the French Association Recherche contre le Cancer (ARC), the Groupement des Entreprises Françaises dans la Lutte contre le Cancer (Gefluc), the Association Nationale de la Recherche et de la Technologie (ANRT, CIFRE PhD fellowship to H.T.) and by the OpenHealth Institute.

Availability of data and materials

All data generated or analyzed during this study are included in the manuscript.

Authors' contributions

All authors have taken part in the different steps of the study: MDV, DGRÉ, PM and EH designed the study, AM, HT performed in silico splicing predictions and the minigene assays, POE performed validation experiments, MM, AN and EHF performed in silico protein predictions, SN, DV performed the sequence analysis. MDV drafted the manuscript and all have read, revised and approved the manuscript.

Ethics approval and consent to participate

Ethical approval for the prospective study was granted from the Norwegian Data Inspectorate and Ethical Review Board (ref 2015/2382). All examined patients had signed an informed consent for their participation in the study.

Consent for publication

Not Applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Tumor Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway. ²Department of Genetic Medicine, The University of Manchester, Manchester Academic Health Science Centre, St. Mary's Hospital, Manchester, UK. ³Genesis Prevention Centre, University Hospital of South Manchester, Southmoor Road, Wythenshawe, UK. ⁴Inserm-U1245, UNIROUEN, Normandie Univ, Normandy Centre for Genomic and Personalized Medicine, Rouen, France. ⁵Interactive Biosoftware, Rouen, France. ⁶Medizinische Klinik und Poliklinik IV, Campus Innenstadt, Klinikum der Universität München, Ziemssenstr. 1, Munich, Germany. ⁷MGZ—Medizinisch Genetisches Zentrum, Munich, Germany. ⁸Department of Human Medicine, Universität Witten/Herdecke, Witten, Germany. ⁹Department of Medical Genetics, Oslo University Hospital, Oslo, Norway. ¹⁰Department of Informatics, University of Oslo, Oslo, Norway. ¹¹Institute of Cancer Genetics and Informatics, Oslo University Hospital, Oslo, Norway.

Received: 2 October 2017 Accepted: 10 January 2018

Published online: 15 January 2018

References

- Mavaddat N, Peock S, Frost D, Ellis S, Platte R, Fineberg E, Evans DG, Izatt L, Eeles RA, Adlard J, et al. Cancer risks for BRCA1 and BRCA2 mutation carriers: results from prospective analysis of EMBRACE. *J Natl Cancer Inst*. 2013;105(11):812–22.
- King MC, Marks JH, Mandell JB, Grp NYBCS. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science*. 2003;302(5645):643–6.
- Fackenthal JD, Olopade OI. Breast cancer risk associated with BRCA1 and BRCA2 in diverse populations. *Nat Rev Cancer*. 2007;7(12):937–48.
- Antoniou A, Pharoah PD, Narod S, Risch HA, Eyfjord JE, Hopper JL, Loman N, Olsson H, Johannsson O, Borg A, et al. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet*. 2003;72(5):1117–30.
- Obermeier K, Sachsenweger J, Friedl TW, Pospiech H, Winqvist R, Wiesmuller L. Heterozygous PALB2 c.1592delT mutation channels DNA double-strand break repair into error-prone pathways in breast cancer patients. *Oncogene*. 2016;35(29):3796–806.
- Aloraifi F, McCartan D, McDevitt T, Green AJ, Bracken A, Geraghty J. Protein-truncating variants in moderate-risk breast cancer susceptibility genes: a meta-analysis of high-risk case-control screening studies. *Cancer Genet*. 2015;208(9):455–63.
- Harismendy O, Schwab RB, Alakus H, Yost SE, Matsui H, Hasteh F, Wallace AM, Park HL, Madlensky L, Parker B, et al. Evaluation of ultra-deep targeted sequencing for personalized breast cancer care. *Breast Cancer Res*. 2013;15(6):R115.
- Evans DGR, Ingham SL, Buchan I, Woodward ER, Byers H, Howell A, Maher ER, Newman WG, Laloo F. Increased rate of Phenocopies in all age groups in BRCA1/BRCA2 mutation kindred, but increased prospective breast cancer risk is confined to BRCA2 mutation carriers. *Cancer Epidebm Biomar*. 2013;22(12):2269–76.
- Moller P, Hagen AI, Apold J, Maehle L, Clark N, Fiane B, Lovslett K, Hovig E, Vabo A. Genetic epidemiology of BRCA mutations—family history detects less than 50% of the mutation carriers. *Eur J Cancer*. 2007;43(11):1713–7.
- Moller P, Stormorken A, Holmen MM, Hagen AI, Vabo A, Maehle L. The clinical utility of genetic testing in breast cancer kindreds: a prospective study in families without a demonstrable BRCA mutation. *Breast Cancer Res Treat*. 2014;144(3):607–14.
- Moller P, Evans G, Haites N, Vasen H, Reis MM, Anderson E, Apold J, Hodgson S, Eccles D, Olsson H, et al. Guidelines for follow-up of women at high risk for inherited breast cancer: consensus statement from the biomed 2 demonstration Programme on inherited breast cancer. *Dis Markers*. 1999;15(1–3):207–11.
- Dominguez-Valentin M, Nakken S, Tubeuf H, Vodak D, Ekstrom PO, Nissen AM, Morak M, Holinski-Feder E, Martins A, Moller P, et al. Potentially pathogenic germline CHEK2 c.319+2T>A among multiple early-onset cancer families. *Fam Cancer*. 2017. <https://doi.org/10.1007/s10689-017-0011-0>. [Epub ahead of print].
- Li L, Chen HC, Liu LX. Sequence alignment algorithm in similarity measurement. *Int Forum Info Technol Appl Proc*. 2009;1:453–456. <https://doi.org/10.1109/lfita.2009.119>.
- Borras E, Pineda M, Blanco I, Jewett EM, Wang F, Teule A, Caldes T, Urioste M, Martinez-Bouzas C, Brunet J, et al. MLH1 founder mutations with moderate penetrance in Spanish lynch syndrome families. *Cancer Res*. 2010;70(19):7379–91.
- Ekstrom PO, Warren DJ, Thilly WG. Separation principles of cycling temperature capillary electrophoresis. *Electrophoresis*. 2012;33(7):1162–8.
- Hinselwood DC, Abrahamsen TW, Ekstrom PO. BRAF mutation detection and identification by cycling temperature capillary electrophoresis. *Electrophoresis*. 2005;26(13):2553–61.
- Ekstrom PO, Khrapko K, Li-Sucholeiki XC, Hunter IW, Thilly WG. Analysis of mutational spectra by denaturing capillary electrophoresis. *Nat Protoc*. 2008;3(7):1153–66.
- Houdayer C, Caux-Moncoutier V, Krieger S, Barrois M, Bonnet F, Bourdon V, Bronner M, Buisson M, Coulet F, Gaildrat P, et al. Guidelines for splicing analysis in molecular diagnosis derived from a set of 327 combined in silico/in vitro studies on BRCA1 and BRCA2 variants. *Hum Mutat*. 2012;33(8):1228–38.
- Palles C, Cazier JB, Howarth KM, Domingo E, Jones AM, Broderick P, Kemp Z, Spain SL, Guarino E, Salguero I, et al. Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet*. 2013;45(2):136–44.
- Ekstrom PO, Nakken S, Johansen M, Hovig E. Automated amplicon design suitable for analysis of DNA variants by melting techniques. *BMC Res Notes*. 2015;8:667.
- den Dunnen JT, Antonarakis SE. Mutation nomenclature extensions and suggestions to describe complex mutations: a discussion. *Hum Mutat*. 2000;15(1):7–12.
- Antoniou AC, Kuchenbaecker KB, Soucy P, Beesley J, Chen XQ, McGuffog L, Lee A, Barrowdale D, Healey S, Sinilnikova OM, et al. Common variants at 12p11, 12q24, 9p21, 9q31.2 and in ZNF365 are associated with breast cancer risk for BRCA1 and/or BRCA2 mutation carriers. *Breast Cancer Research*. 2012;14(1):1–18.
- Di Giacomo D, Gaildrat P, Abuli A, Abdat J, Frebourg T, Tosi M, Martins A. Functional analysis of a large set of BRCA2 exon 7 variants highlights the predictive value of hexamer scores in detecting alterations of exonic splicing regulatory elements. *Hum Mutat*. 2013;34(11):1547–57.
- Erkelenz S, Hillebrand F, Widera M, Theiss S, Faysaz A, Degrandi D, Pfeffer K, Schaal H. Balanced splicing at the tat-specific HIV-1 3' splice site is critical for HIV-1 replication. *Retrovirology*. 2015;12:29.
- Susswein LR, Marshall ML, Nusbaum R, Vogel Postula KJ, Weissman SM, Yackowski L, Vaccari EM, Bissonnette J, Booker JK, Cremona ML, et al. Pathogenic and likely pathogenic variant prevalence among the first 10,000 patients referred for next-generation cancer panel testing. *Genet Med*. 2016;18(8):823–32.
- Soukarieh O, Gaildrat P, Hamieh M, Drouet A, Baert-Desurmont S, Frebourg T, Tosi M, Martins A. Exonic Splicing Mutations Are More Prevalent than Currently Estimated and Can Be Predicted by Using In Silico Tools. *Plos Genet*. 2016;12(1):1–26.
- Gaildrat P, Killian A, Martins A, Tournier I, Frebourg T, Tosi M. Use of splicing reporter minigene assay to evaluate the effect on splicing of unclassified genetic variants. *Methods Mol Biol*. 2010;653:249–57.
- Tournier I, Vezain M, Martins A, Charbonnier F, Baert-Desurmont S, Olschwang S, Wang Q, Buisine MP, Soret J, Tazi J, et al. A large fraction of unclassified variants of the mismatch repair genes MLH1 and MSH2 is associated with splicing defects. *Hum Mutat*. 2008;29(12):1412–24.
- Sjursen W, Haukanes BI, Grindedal EM, Aarset H, Stormorken A, Engebretsen LF, Jonsrud C, Bjornevoll I, Andresen PA, Ariansen S, et al. Current clinical criteria for lynch syndrome are not sensitive enough to identify MSH6 mutation carriers. *J Med Genet*. 2010;47(9):579–85.
- Pinto P, Paulo P, Santos C, Rocha P, Pinto C, Veiga I, Pinheiro M, Peixoto A, Teixeira MR. Implementation of next-generation sequencing for molecular diagnosis of hereditary breast and ovarian cancer highlights its genetic heterogeneity. *Breast Cancer Res Treat*. 2016;159(2):245–56.
- Hegde M, Ferber M, Mao R, Samowitz W, Ganguly A. Working Group of the American College of medical G, genomics laboratory quality assurance C: ACMG technical standards and guidelines for genetic testing for inherited colorectal cancer (lynch syndrome, familial

- adenomatous polyposis, and MYH-associated polyposis). *Genet Med.* 2014;16(1):101–16.
32. Tung N, Lin NU, Kidd J, Allen BA, Singh N, Wenstrup RJ, Hartman AR, Winer EP, Garber JE. Frequency of Germline mutations in 25 cancer susceptibility genes in a sequential series of patients with breast cancer. *J Clin Oncol.* 2016;34(13):1460–8.
 33. Castera L, Krieger S, Rousselin A, Legros A, Baumann JJ, Bruet O, Brault B, Fouillet R, Goardon N, Letac O, et al. Next-generation sequencing for the diagnosis of hereditary breast and ovarian cancer using genomic capture targeting multiple candidate genes. *Eur J Hum Genet.* 2014;22(11):1305–13.
 34. Kamps R, Brandao RD, Bosch BJ, Paulussen AD, Xanthoulea S, Blok MJ, Romano A. Next-Generation Sequencing in Oncology: Genetic Diagnosis, Risk Prediction and Cancer Classification. *Int J Mol Sci.* 2017;18(2):1–57.
 35. Smith MJ, Urquhart JE, Harkness EF, Miles EK, Bowers NL, Byers HJ, Bulman M, Gokhale C, Wallace AJ, Newman WG, et al. The contribution of whole gene deletions and large rearrangements to the mutation Spectrum in inherited tumor predisposing syndromes. *Hum Mutat.* 2015;
 36. Lincoln SE, Kobayashi Y, Anderson MJ, Yang S, Desmond AJ, Mills MA, Nilsen GB, Jacobs KB, Monzon FA, Kurian AW, et al. A systematic comparison of traditional and multigene panel testing for hereditary breast and ovarian cancer genes in more than 1000 patients. *J Mol Diagn.* 2015;17(5):533–44.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

