

First and Second Fundamental Solutions of the Time-Fractional Telegraph Equation of Order $2\alpha^*$

M. Ferreira^{†,‡} M.M. Rodrigues[‡] N. Vieira[‡]

[†] School of Technology and Management,
Polytechnic Institute of Leiria
P-2411-901, Leiria, Portugal.

[‡] CIDMA - Center for Research and Development in Mathematics and Applications
Department of Mathematics, University of Aveiro
Campus Universitário de Santiago, 3810-193 Aveiro, Portugal.

E-mails: milton.ferreira@ipleiria.pt, mrodrigues@ua.pt, nloureirovieira@gmail.com

Abstract

In this work we obtain the first and second fundamental solutions of the multidimensional time-fractional equation of order 2α , $\alpha \in]0, 1]$, where the two time-fractional derivatives are in the Caputo sense. We obtain representations of the fundamental solutions in terms of Hankel transform, double Mellin-Barnes integral, and H-functions of two variables. As an application, the fundamental solutions are used to solve a Cauchy problem, and to study telegraph process with Brownian time.

Keywords: Time-fractional telegraph equation; Telegraph Dirac operator; First and second fundamental solutions; Caputo fractional derivative; Multivariate Mittag-Leffler function; H-function of two variables.

MSC 2010: 30G35; 35R11, 26A33, 35A08, 35C15, 33E12.

1 Introduction

The telegraph equation is used as an alternative to the diffusion equation, since it has the potential to describe both diffusive and wave-like phenomena, due to the simultaneous presence of first and second order time derivatives. For example, in the case of the transport of energetic charged particle in turbulent magnetic fields such as low-energy cosmic rays in the solar wind, the diffusion equation can not be used to describe the transport for early times because it leads to a non-zero probability density everywhere, which would correspond to an infinite propagation speed. Using the telegraph equation in this case we get a more realistic model for the early phase transport because it combines diffusion with a finite propagation speed (see [18]). Telegraph equations have also an extraordinary importance in electrodynamics (the scalar Maxwell equations are of this type), in the theory of damped vibrations, and in probability because they are connected with finite velocity random motions (see [12, 16]).

One of the first works studying the time-fractional telegraph equation is the paper of Cascaval et al. (see [4]). Here, the authors discussed some properties of the time-fractional telegraph equation in $\mathbb{R} \times \mathbb{R}^+$ such as the well-posedness and the asymptotic behavior of the solutions, by using the Riemann-Liouville approach. In [15], Orsingher and Beghin obtained the fundamental solution of the time-fractional telegraph equation of order 2α in $\mathbb{R} \times \mathbb{R}^+$ and gave a representation of their inverses in terms of stable densities. For the special case $\alpha = \frac{1}{2}$, the authors showed that the fundamental solution is the probability density of a telegraph process with Brownian

*The final version is published in *AIP Conference Proceedings - ICNPAA 2018*, **2046**, (2018), Article No. 020079, 9 pp. It is available via the website: <https://aip.scitation.org/doi/abs/10.1063/1.5081599>

time. In [3] it was discussed the solution of a general space-time fractional telegraph equation by means of the Laplace and Fourier transforms in the variables $x \in \mathbb{R}$ and $t \in \mathbb{R}^+$, respectively. In [19] it was obtained the solutions of the space-time fractional telegraph equation in $\mathbb{R} \times \mathbb{R}^+$ in terms of Mittag-Leffler functions, using an operational approach. In [14], Mamchuev considered the inhomogeneous time-fractional telegraph equation with Caputo derivatives, and obtained a general representation of regular solution in rectangular domain in terms of fundamental solution and appropriate Green functions. Regarding the multidimensional case, in [5] the authors discussed and derived the solution of the time-fractional telegraph equation in $\mathbb{R}^n \times \mathbb{R}^+$ with three kinds of nonhomogeneous boundary conditions, by the method of separation of variables. Very recently, in [9, 10] the authors found several representations of the fundamental solution of the time-fractional telegraph and telegraph Dirac equations in $\mathbb{R}^n \times \mathbb{R}^+$, in terms of integrals, special functions, and series.

The aim of this paper is to representations for the first and second fundamental solutions of the time-fractional telegraph equation of order 2α in terms of Hankel transforms, double Mellin-Barnes integral and H-functions of two variables. Moreover, the first fundamental solution is used in the law of telegraph process with Brownian time. This connection is motivated by the fact that the iterated Brownian motion and telegraph process with Brownian time are governed by time-fractional telegraph equations (see [15]).

The structure of the paper reads as follows: in the Preliminaries section we recall some basic facts about some special functions and fractional calculus, which are necessary for the development of the work. In the following section we obtain the first and second fundamental solutions of the time-fractional telegraph equation

$$({}^C\partial_t^{2\alpha} + a {}^C\partial_t^\alpha - c^2 \Delta_x) u(x, t) = 0$$

where $x \in \mathbb{R}^n$, $t > 0$, $0 < \alpha \leq 1$, $a \geq 0$, $c > 0$, Δ_x is the Laplace operator in space, and the two time-fractional derivatives of orders α and 2α are in the Caputo sense, with $\alpha \in]0, 1]$. This section ends with an application to the resolution of a Cauchy problem. In the last section we present an application of our results to the law of a telegraph process with Brownian time.

2 Preliminaries

Here we recall the main tools concerning fractional derivatives and special functions that will be used in our work. We start by recalling the definition of the multivariate Mittag-Leffler function (see [13]).

Definition 2 *The multivariate Mittag-Leffler function $E_{(a_1, \dots, a_n), b}(z_1, \dots, z_n)$ of n complex variables $z_1, \dots, z_n \in \mathbb{C}$ with complex parameters $a_1, \dots, a_n, b \in \mathbb{C}$ (with positive real parts) is defined by*

$$E_{(a_1, \dots, a_n), b}(z_1, \dots, z_n) = \sum_{k=0}^{+\infty} \sum_{\substack{l_1 + \dots + l_n = k \\ l_1, \dots, l_n \geq 0}} \binom{k}{l_1, \dots, l_n} \frac{\prod_{i=1}^n z_i^{l_i}}{\Gamma(b + \sum_{i=1}^n a_i l_i)}, \quad (1)$$

where the multinomial coefficients are given by

$$\binom{k}{l_1, \dots, l_n} := \frac{k!}{l_1! \times \dots \times l_n!}.$$

When $n = 2$ we obtain the bivariate Mittag-Leffler function, which can be written as

$$E_{(a_1, a_2), b}(z_1, z_2) = \sum_{l_1=0}^{+\infty} \sum_{l_2=0}^{+\infty} \frac{(l_1 + l_2)!}{l_1! l_2!} \frac{z_1^{l_1} z_2^{l_2}}{\Gamma(b + a_1 l_1 + a_2 l_2)}. \quad (2)$$

From (2) we can deduce, after straightforward calculations, an addition formula for the bivariate Mittag-Leffler function.

Lemma 2 *Let $z_1, z_2 \in \mathbb{C}$, and $a_1, a_2, b \in \mathbb{C}$ (with positive real parts). Then it holds*

$$E_{(a_1, a_2), b}(z_1, z_2) = \frac{1}{\Gamma(b)} + z_1 E_{(a_1, a_2), b+a_1}(z_1, z_2) + z_2 E_{(a_1, a_2), b+a_2}(z_1, z_2). \quad (3)$$

For general properties of the Mittag-Leffler function see [11, 13]. Now we recall the definition of the H-function of two complex variables.

Definition 2.3 (see [1]) *The H-function of two complex variables is defined via a double Mellin-Barnes integral of the form*

$$H_{p_1, q_1; p_2, q_2; p_3, q_3}^{0, n_1; m_2, n_2; m_3, n_3} \left[\begin{array}{l} x \\ y \end{array} \middle| \begin{array}{l} (a_j; \alpha_j, A_j)_{1, p_1}; (c_j, \gamma_j)_{1, p_2}; (e_j, E_j)_{1, p_3} \\ (b_j; \beta_j, B_j)_{1, q_1}; (d_j, \delta_j)_{1, q_2}; (f_j, F_j)_{1, q_3} \end{array} \right] = \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_2} \int_{\mathcal{L}_1} \phi(s, w) \phi_1(s) \phi_2(w) x^s y^w ds dw,$$

where

$$\begin{aligned} \phi(s, w) &= \frac{\prod_{i=1}^{n_1} \Gamma(1 - a_i + \alpha_i s + A_i w)}{\prod_{i=n_1+1}^{p_1} \Gamma(a_i - \alpha_i s - A_i w) \prod_{j=1}^{q_1} \Gamma(1 - b_j + \beta_j s + B_j w)}, \\ \phi_1(s) &= \frac{\prod_{j=1}^{m_2} \Gamma(d_j - \delta_j s) \prod_{i=1}^{n_2} \Gamma(1 - c_i + \gamma_i s)}{\prod_{j=m_2+1}^{q_2} \Gamma(1 - d_j + \delta_j s) \prod_{i=n_2+1}^{p_2} \Gamma(c_i - \gamma_i s)}, \\ \phi_2(w) &= \frac{\prod_{j=1}^{m_3} \Gamma(f_j - F_j w) \prod_{i=1}^{n_3} \Gamma(1 - e_i + E_i w)}{\prod_{j=m_3+1}^{q_3} \Gamma(1 - f_j + F_j w) \prod_{i=n_3+1}^{p_3} \Gamma(e_i - E_i w)}, \end{aligned}$$

and where an empty product is interpreted as 1, $x, y \in \mathbb{C}$, $m_i, n_i, p_i, q_i \in \mathbb{Z}$ such that $0 \leq m_i \leq q_i$, $0 \leq n_i \leq p_i$ ($i = 1, 2, 3$); $a_i, b_j, c_i, d_j, e_i, f_j \in \mathbb{C}$, $\alpha_i, A_i, \beta_j, B_j, \gamma_i, \delta_j, E_i, F_j \in \mathbb{R}^+$ and the sequence of parameters (a_j) , (b_j) , (c_j) , (d_j) , (e_j) and (f_j) are restricted that none of the poles of the integrand coincide. The contour \mathcal{L}_1 in the complex s -plane, and the contour \mathcal{L}_2 in the complex w -plane, are of Mellin-Barnes type with indentations, if necessary, to ensure that they separate one set of poles from the other.

In [1] the author proved that if

$$\begin{aligned} \Omega_1 &= \sum_{i=1}^{n_1} \alpha_i - \sum_{i=n_1+1}^{p_1} \alpha_i - \sum_{i=1}^{q_1} \beta_i + \sum_{i=1}^{m_2} \delta_i - \sum_{j=m_2+1}^{q_2} \delta_j + \sum_{i=1}^{n_2} \gamma_i - \sum_{i=n_2+1}^{p_2} \gamma_i \geq 0, \\ \Omega_2 &= \sum_{i=1}^{n_1} A_i - \sum_{i=n_1+1}^{p_1} A_i - \sum_{i=1}^{q_1} B_i + \sum_{i=1}^{m_3} F_i - \sum_{j=m_3+1}^{q_3} F_j + \sum_{i=1}^{n_3} E_i - \sum_{i=n_3+1}^{p_3} E_i \geq 0, \end{aligned} \quad (4)$$

and with the points $x = 0$ and $y = 0$ being tacitly excluded, the double Mellin-Barnes integral converges absolutely inside the sector given by

$$|\arg(x)| < \frac{\pi}{2} \Omega_1, \quad |\arg(y)| < \frac{\pi}{2} \Omega_2.$$

For $E_{(a_1, a_2), b}$ we have the following two results (see [9]).

Lemma 2.4 *The bivariate Mittag-Leffler function $E_{(a_1, a_2), b}$ has the following representation in the form of double Mellin-Barnes integral*

$$E_{(a_1, a_2), b}(z_1, z_2) = \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(-s) \Gamma(-w)}{\Gamma(b + a_1 w + a_2 s)} z_2^s z_1^w dw ds.$$

Corollary 2.5 *The bivariate Mittag-Leffler function $E_{(a_1, a_2), b}$ has the following representation in the form of H-function of two variables*

$$E_{(a_1, a_2), b}(z_1, z_2) = H_{1,1; 0,1; 0,1}^{0,1; 1,0; 1,0} \left[\begin{array}{l} z_2 \\ z_1 \end{array} \middle| \begin{array}{l} (0, 1, 1); \text{---}; \text{---} \\ (1 - b; a_2, a_1); (0, 1); (0, 1) \end{array} \right]. \quad (5)$$

In (5) and throughout the paper a horizontal line in the H-function means the absence of parameters. Now we recall the definition of the spaces C_α , $\alpha \in \mathbb{R}$, and C_α^m , $m \in \mathbb{N}$, given in [13].

Definition 2.6 (see [13]) A real or complex-valued function $f(t)$, $t > 0$, is said to be in the space C_α , $\alpha \in \mathbb{R}$, if there exists a real number $p > \alpha$ such that $f(t) = t^p f_1(t)$ for some function $f_1 \in C[0, \infty)$.

It is easy to see that C_α is a vector space and the set of spaces C_α is ordered by inclusion according to $C_\alpha \subseteq C_\beta \Leftrightarrow \alpha \geq \beta$.

Definition 2.7 (see [13]) A function $f(t)$, $t > 0$, is said to be in the space C_α^m , $m \in \mathbb{N}$, if and only if $f^{(m)} \in C_\alpha$.

Let D_t^γ denotes the Caputo fractional derivative of order $\gamma > 0$ defined by:

$$D_t^\gamma u(t) = \begin{cases} \frac{1}{\Gamma(m-\gamma)} \int_0^t (t-s)^{-\gamma+m-1} u^{(m)}(s) ds, & m-1 < \gamma < m \\ u^{(m)}(t), & \gamma = m \end{cases} \quad (6)$$

where $u^{(m)} := \frac{d^m u}{dt^m}$, $m \in \mathbb{N}$ (see [2]). The next theorem will be used in our analysis and allow us to solve general linear fractional differential equations with constant coefficients and Caputo derivatives.

Theorem 2.8 (see [13, Thm 4.1]) Let $\mu > \mu_1 > \dots > \mu_n \geq 0$, $m_i - 1 < \mu_i \leq m_i$, $m_i \in \mathbb{N}$, $\lambda_i \in \mathbb{R}$, $i = 1, \dots, n$. Consider the initial value problem

$$\begin{cases} D_t^\mu y(t) - \sum_{i=1}^n \lambda_i D_t^{\mu_i} y(t) = g(t) \\ y^{(k)}(0) = c_k \in \mathbb{R}, \quad k = 0, \dots, m-1, \quad m-1 < \mu \leq m, \end{cases} \quad (7)$$

where g is assumed to lie in C_{-1} if $\mu \in \mathbb{N}_0$ or in C_{-1}^1 if $\mu \notin \mathbb{N}_0$. Then (7) has a unique solution in the space C_{-1}^m of the form

$$y(t) = \int_0^t s^{\mu-1} E_{(\mu-\mu_1, \dots, \mu-\mu_n), \mu}(\lambda_1 s^{\mu-\mu_1}, \dots, \lambda_n s^{\mu-\mu_n}) g(t-s) ds + \sum_{k=0}^{m-1} c_k u_k(t), \quad t \geq 0,$$

with, $k = 0, \dots, m-1$

$$u_k(t) = \frac{t^k}{k!} + \sum_{i=l_k+1}^n \lambda_i t^{k+\mu-\mu_i} E_{(\mu-\mu_1, \dots, \mu-\mu_n), k+1+\mu-\mu_i}(\lambda_1 t^{\mu-\mu_1}, \dots, \lambda_n t^{\mu-\mu_n}).$$

The natural numbers l_k , $k = 0, \dots, m-1$ are determined from the conditions $m_{l_k} \geq k+1$ and $m_{l_k+1} \leq k$. In the case $m_i \leq k$, $i = 0, \dots, m-1$, we set $l_k := 0$, and of $m_i \geq k+1$, $i = 0, \dots, m-1$, then $l_k := n$.

3 First and second fundamental solution of the time-fractional telegraph equation of order 2α

In this section we obtain the first and second fundamental solution of a particular case of the previous equation, where $\beta = 2\alpha$ and $0 < \alpha \leq 1$, i.e., we look for a function $G^\alpha(x, t)$ that satisfies the following Cauchy problem

$$\begin{cases} ({}^C\partial_t^{2\alpha} + a {}^C\partial_t^\alpha - c^2 \Delta_x) G^\alpha(x, t) = 0 \\ G^\alpha(x, 0) = C_0 \delta(x) \\ \frac{\partial G^\alpha}{\partial t}(x, 0) = C_1 \delta(x) \end{cases}, \quad (8)$$

where $x \in \mathbb{R}^n$, $t > 0$, $a \geq 0$, $c > 0$, $C_0, C_1 \in \mathbb{R}$, $\delta(x) = \prod_{j=1}^n \delta(x_j)$ is the distributional Dirac delta function in \mathbb{R}^n . Applying the Fourier transform in \mathbb{R}^n to the Cauchy problem we get the following initial-value problem

$$\begin{cases} ({}^C\partial_t^{2\alpha} + a {}^C\partial_t^\alpha + c^2 |\kappa|^2) \widehat{G}^\alpha(\kappa, t) = 0 \\ \widehat{G}^\alpha(\kappa, 0) = C_0 \\ \frac{\partial \widehat{G}^\alpha}{\partial t}(\kappa, 0) = C_1 \end{cases}. \quad (9)$$

To solve the problem (9), we apply Theorem 2.8 with $\lambda_1 = -a$, $\lambda_2 = -c^2 |\kappa|^2$, $\mu = 2\alpha$, $\mu_1 = \alpha$, $\mu_2 = 0$, $n = 2$, $m = 2$, $g(t) = 0$, and $k = 1$, obtaining the following solution

$$\widehat{G}^\alpha(\kappa, t) = C_0 \widehat{u}_0(\kappa, t) + C_1 \widehat{u}_1(\kappa, t), \quad (10)$$

where

$$\widehat{u}_0(\kappa, t) = 1 - c^2 |\kappa|^2 t^{2\alpha} E_{(\alpha, 2\alpha), 1+2\alpha}(-a t^\alpha, -c^2 |\kappa|^2 t^{2\alpha}),$$

$$\widehat{u}_1(\kappa, t) = t - a t^{1+\alpha} E_{(\alpha, 2\alpha), 2+\alpha}(-a t^\alpha, -c^2 |\kappa|^2 t^{2\alpha}) - c^2 |\kappa|^2 t^{1+2\alpha} E_{(\alpha, 2\alpha), 2+2\alpha}(-a t^\alpha, -c^2 |\kappa|^2 t^{2\alpha}).$$

Taking into account Lemma 2.2 we have the following alternative representation of \widehat{u}_0 and \widehat{u}_1

$$\widehat{u}_0(\kappa, t) = E_{(\alpha, 2\alpha), 1}(-a t^\alpha, -c^2 |\kappa|^2 t^{2\alpha}) + a t^\alpha E_{(\alpha, 2\alpha), 1+\alpha}(-a t^\alpha, -c^2 |\kappa|^2 t^{2\alpha}), \quad (11)$$

$$\widehat{u}_1(\kappa, t) = t E_{(\alpha, 2\alpha), 2}(-a t^\alpha, -c^2 |\kappa|^2 t^{2\alpha}). \quad (12)$$

From (2), we have the following representation of \widehat{u}_0 and \widehat{u}_1 in the form of a double series

$$\begin{aligned} \widehat{u}_0(\kappa, t) &= \sum_{p=0}^{+\infty} \sum_{q=0}^{+\infty} \frac{\Gamma(1+p+q)}{\Gamma(1+\alpha p+2\alpha q) p! q!} (-a t^\alpha)^p (-c^2 |\kappa|^2 t^{2\alpha})^q \\ &\quad + a t^\alpha \sum_{p=0}^{+\infty} \sum_{q=0}^{+\infty} \frac{\Gamma(1+p+q)}{\Gamma(1+\alpha+\alpha p+2\alpha q) p! q!} (-a t^\alpha)^p (-c^2 |\kappa|^2 t^{2\alpha})^q, \end{aligned} \quad (13)$$

$$\widehat{u}_1(\kappa, t) = t \sum_{p=0}^{+\infty} \sum_{q=0}^{+\infty} \frac{\Gamma(1+p+q)}{\Gamma(2+\alpha p+2\alpha q) p! q!} (-a t^\alpha)^p (-c^2 |\kappa|^2 t^{2\alpha})^q. \quad (14)$$

From Lemma 2.4 we have the following representations of \widehat{u}_0 and \widehat{u}_1 in the form of a double Mellin-Barnes integral

$$\begin{aligned} \widehat{u}_0(\kappa, t) &= \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(-s) \Gamma(-w)}{\Gamma(1+\alpha w+2\alpha s)} (a t^\alpha)^w (c^2 |\kappa|^2 t^{2\alpha})^s dw ds \\ &\quad + \frac{a t^\alpha}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(-s) \Gamma(-w)}{\Gamma(1+\alpha+\alpha w+2\alpha s)} (a t^\alpha)^w (c^2 |\kappa|^2 t^{2\alpha})^s dw ds, \end{aligned} \quad (15)$$

$$\widehat{u}_1(\kappa, t) = \frac{t}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(-s) \Gamma(-w)}{\Gamma(2+\alpha w+2\alpha s)} (a t^\alpha)^w (c^2 |\kappa|^2 t^{2\alpha})^s dw ds. \quad (16)$$

From Corollary 2.5 we can write \widehat{u}_0 and \widehat{u}_1 in the form of H-functions of two variables

$$\begin{aligned} \widehat{u}_0(\kappa, t) &= H_{1,1;0,1;0,1}^{0,1;1,0;1,0} \left[\begin{array}{c} c^2 |\kappa|^2 t^{2\alpha} \\ a t^\alpha \end{array} \middle| \begin{array}{c} (0; 1, 1); \text{---}; \text{---} \\ (0; 2\alpha, \alpha); (0, 1); (0, 1) \end{array} \right] \\ &\quad + a t^\alpha H_{1,1;0,1;0,1}^{0,1;1,0;1,0} \left[\begin{array}{c} c^2 |\kappa|^2 t^{2\alpha} \\ a t^\alpha \end{array} \middle| \begin{array}{c} (0; 1, 1); \text{---}; \text{---} \\ (-\alpha; 2\alpha, \alpha); (0, 1); (0, 1) \end{array} \right], \end{aligned} \quad (17)$$

$$\widehat{u}_1(\kappa, t) = t H_{1,1;0,1;0,1}^{0,1;1,0;1,0} \left[\begin{array}{c} c^2 |\kappa|^2 t^{2\alpha} \\ a t^\alpha \end{array} \middle| \begin{array}{c} (0; 1, 1); \text{---}; \text{---} \\ (-1; 2\alpha, \alpha); (0, 1); (0, 1) \end{array} \right]. \quad (18)$$

Inverting the Fourier transform in (10) we get

$$G^\alpha(x, t) = C_0 u_0(x, t) + C_1 u_1(x, t). \quad (19)$$

We will denote by $G_1^\alpha(x, t) = u_0(x, t)$ the first fundamental solution of the first equation of (8) that satisfies the initial conditions $G_1^\alpha(x, 0) = \delta(x)$ and $\frac{\partial G_1^\alpha}{\partial t}(x, 0) = 0$. Additionally, we denote by $G_2^\alpha(x, t) = u_1(x, t)$ the second fundamental solution of the first equation of (8) that satisfies the initial conditions $G_2^\alpha(x, 0) = 0$ and $\frac{\partial G_2^\alpha}{\partial t}(x, 0) = \delta(x)$. For obtaining the explicit expressions for G_1^α and G_2^α we recall the following formula (see [17])

$$\frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{-i\kappa \cdot x} \phi(|\kappa|) d\kappa = \frac{|x|^{1-\frac{n}{2}}}{(2\pi)^{\frac{n}{2}}} \int_0^{+\infty} \phi(\tau) \tau^{\frac{n}{2}} J_{\frac{n}{2}-1}(\tau|x|) d\tau, \quad (20)$$

where J_ν represents the Bessel function of first kind with index ν , and the right hand side can be seen as a Hankel transform. We need also the following integral formula (see Formula (7) in [6, p. 22])

$$\int_0^{+\infty} z^\mu J_\nu(z y)^{\frac{1}{2}} dz = \frac{2^{\mu+\frac{1}{2}} \Gamma\left(\frac{\mu}{2} + \frac{\nu}{2} + \frac{3}{4}\right)}{y^{\mu+1} \Gamma\left(\frac{\nu}{2} - \frac{\mu}{2} + \frac{1}{4}\right)} \quad (21)$$

which is valid under the condition $-\operatorname{Re}(\nu) - \frac{3}{2} < \operatorname{Re}(\mu) < -\frac{1}{2}$. Since the expressions for G_1^α are longer than the ones for G_2^α , we are going to present only the calculations for G_2^α . Applying the inverse Fourier transform to (12) and using (20), we get

$$G_2^\alpha(x, t) = \frac{t|x|^{1-\frac{n}{2}}}{(2\pi)^{\frac{n}{2}}} \int_0^{+\infty} \tau^{\frac{n}{2}} E_{(\alpha, 2\alpha), 2}(-a t^\alpha, -c^2 \tau^2 t^{2\alpha}) J_{\frac{n}{2}-1}(\tau|x|) d\tau. \quad (22)$$

Making use of Lemma 2.4, and interchanging the integrals due to the convergence, we obtain

$$G_2^\alpha(x, t) = \frac{t|x|^{1-\frac{n}{2}}}{(2\pi)^{\frac{n}{2}}} \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(-s) \Gamma(-w)}{\Gamma(2+\alpha w+2\alpha s)} (a t^\alpha)^w (c^2 t^{2\alpha})^s \times \int_0^{+\infty} \tau^{\frac{n}{2}+2s} J_{\frac{n}{2}-1}(\tau|x|) d\tau dw ds. \quad (23)$$

Finally, using (21) and Corollary 2.5, we obtain, under the condition $-\frac{n}{2} < \Re(s) < -\frac{n}{4}$, the representation of the second fundamental solution in terms of double Mellin-Barnes integral and H-function of two variables:

$$G_2^\alpha(x, t) = \frac{t}{\pi^{\frac{n}{2}} |x|^n} \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma\left(\frac{n}{2}+s\right) \Gamma(-w)}{\Gamma(2+\alpha w+2\alpha s)} (a t^\alpha)^w \left(\frac{4c^2 t^{2\alpha}}{|x|^2}\right)^s dw ds \quad (24)$$

$$= \frac{t}{\pi^{\frac{n}{2}} |x|^n} H_{1,1; 1,0; 0,1}^{0,1; 0,1; 1,0} \left[\begin{array}{c} 4c^2 t^{2\alpha} |x|^{-2} \\ a t^\alpha \end{array} \middle| \begin{array}{c} (0; 1, 1); \left(1 - \frac{n}{2}, 1\right); - \\ (-1; 2\alpha, \alpha); -; (0, 1) \end{array} \right], \quad (25)$$

where the double Mellin-Barnes integral is convergent because the conditions (4) are fulfilled. For G_1^α we obtain, analogously, under the condition $-\frac{n}{2} < \Re(s) < -\frac{n}{4}$, the following representations:

$$G_1^\alpha(x, t) = \frac{|x|^{1-\frac{n}{2}}}{(2\pi)^{\frac{n}{2}}} \int_0^{+\infty} \tau^{\frac{n}{2}} E_{(\alpha, 2\alpha), 1}(-a t^\alpha, -c^2 \tau^2 t^{2\alpha}) J_{\frac{n}{2}-1}(\tau|x|) d\tau + \frac{a t^\alpha |x|^{1-\frac{n}{2}}}{(2\pi)^{\frac{n}{2}}} \int_0^{+\infty} \tau^{\frac{n}{2}} E_{(\alpha, 2\alpha), 1+\alpha}(-a t^\alpha, -c^2 \tau^2 t^{2\alpha}) J_{\frac{n}{2}-1}(\tau|x|) d\tau \quad (26)$$

$$= \frac{1}{\pi^{\frac{n}{2}} |x|^n} \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma\left(\frac{n}{2}+s\right) \Gamma(-w)}{\Gamma(1+\alpha w+2\alpha s)} (a t^\alpha)^w \left(\frac{4c^2 t^{2\alpha}}{|x|^2}\right)^s dw ds + \frac{a t^\alpha}{\pi^{\frac{n}{2}} |x|^n} \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma\left(\frac{n}{2}+s\right) \Gamma(-w)}{\Gamma(1+\alpha+\alpha w+2\alpha s)} (a t^\alpha)^w \left(\frac{4c^2 t^{2\alpha}}{|x|^2}\right)^s dw ds \quad (27)$$

$$\begin{aligned}
&= \frac{1}{\pi^{\frac{n}{2}} |x|^n} H_{1,1;1,0;0,1}^{0,1;0,1;1,0} \left[\begin{array}{c} 4c^2 t^{2\alpha} |x|^{-2} \\ a t^\alpha \end{array} \middle| \begin{array}{c} (0; 1, 1); (1 - \frac{n}{2}, 1); \text{---} \\ (0; 2\alpha, \alpha); \text{---}; (0, 1) \end{array} \right] \\
&+ \frac{a t^\alpha}{\pi^{\frac{n}{2}} |x|^n} H_{1,1;1,0;0,1}^{0,1;0,1;1,0} \left[\begin{array}{c} 4c^2 t^{2\alpha} |x|^{-2} \\ a t^\alpha \end{array} \middle| \begin{array}{c} (0; 1, 1); (1 - \frac{n}{2}, 1); \text{---} \\ (-\alpha; 2\alpha, \alpha); \text{---}; (0, 1) \end{array} \right]. \tag{28}
\end{aligned}$$

We end this section solving a fractional Cauchy problem.

Theorem 3.1 *Let $x \in \mathbb{R}^n$, $t > 0$, $0 < \alpha \leq 1$, $a \geq 0$, and $c > 0$, then the fractional Cauchy problem*

$$\begin{cases} ({}^C\partial_t^{2\alpha} + a {}^C\partial_t^\alpha - c^2 \Delta_x) h(x, t) = 0 \\ h(x, 0) = f_1(x) \\ \frac{\partial h}{\partial t}(x, 0) = f_2(x), \end{cases}$$

is solvable, and its solution has the form

$$h(x, t) = \int_{\mathbb{R}^n} G_1^\alpha(x - z, t) f_1(z) dz + \int_{\mathbb{R}^n} G_2^\alpha(x - z, t) f_2(z) dz, \tag{29}$$

where G_1^α and G_2^α are the first and second fundamental solutions given by (28) and (25), respectively, and provided that the integrals in the right-hand side of (29) are convergent.

Remark 3.2 *Theorem 3.1, when $a = 0$ the solution (29) coincides with the one presented in [12, Cor.6.4] (with $\beta = 2\alpha$). Moreover, considering $a = 0$ the first fundamental solution G_1^α coincides with the expression presented in [7] for the fundamental solution of the time-fractional diffusion-wave operator (with $\beta = 2\alpha$).*

4 Telegraph process with Brownian time

Now we consider in (8) $n = 1$, $a = 2\lambda \geq 0$, $C_0 = 1$ and $C_1 = 0$, i.e., we consider the following time-fractional telegraph equation of order 2α

$$({}^C\partial_t^{2\alpha} + 2\lambda {}^C\partial_t^\alpha - c^2 \partial_{xx}^2) u(x, t) = 0, \tag{30}$$

where $x \in \mathbb{R}$, $t > 0$, $c > 0$, and subject to the initial condition $u(x, 0) = \delta(x)$ for $0 < \alpha \leq \frac{1}{2}$, while, for $\frac{1}{2} < \alpha \leq 1$, besides the previous condition, also $u_t(x, 0) = 0$ is assumed. Equation (30) was already studied in [15], where the authors presented only an integral representation for the Fourier transform of the fundamental solution. Moreover, for $\alpha = \frac{1}{2}$ it was obtained an integral representation of the fundamental solution based on the Fourier inversion transform (see Theorem 4.2 in [15]).

Physically, if we consider $\alpha = \frac{1}{2}$ in (30), we obtain a heat equation with damping term which depends on all values of u in $[0, t]$ and assigning an overwhelming weight to those close to t (see [15]). The damping effect of the fractional derivative reverberates on the distribution u , where the solution of the heat equation (governing term) is perturbed by the telegraph distribution (which represents the impact of the fractional derivative). Moreover, since the fundamental solution u of (30) reduces, in this particular case, to $G_1^{\frac{1}{2}}$ (see (19)), it can be understood as the distribution of a particle moving back and forth the real line with velocities $\pm c$ for a random

time interval. Making $n = 1$, $a = 2\lambda$ and $\alpha = \frac{1}{2}$ in (26), (27) and (28) we get the following representations $G_1^{\frac{1}{2}}$:

$$\begin{aligned}
G_1^{\frac{1}{2}}(x, t) &= \frac{1}{\pi} \int_0^{+\infty} E_{(\frac{1}{2}, 1), 1}(-2\lambda\sqrt{t}, -c^2\tau^2 t) \cos(\tau|x|) d\tau \\
&\quad + \frac{2\lambda\sqrt{t}}{\pi} \int_0^{+\infty} E_{(\frac{1}{2}, 1), \frac{3}{2}}(-2\lambda\sqrt{t}, -c^2\tau^2 t) \cos(\tau|x|) d\tau \\
&= \frac{1}{\sqrt{\pi}|x|} \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(\frac{1}{2}+s) \Gamma(-w)}{\Gamma(1+\frac{w}{2}+s)} (2\lambda\sqrt{t})^w \left(\frac{4c^2 t}{|x|^2}\right)^s dw ds \\
&\quad + \frac{2\lambda\sqrt{t}}{\sqrt{\pi}|x|} \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_1} \int_{\mathcal{L}_2} \frac{\Gamma(1+s+w) \Gamma(\frac{1}{2}+s) \Gamma(-w)}{\Gamma(\frac{3}{2}+\frac{w}{2}+s)} (2\lambda\sqrt{t})^w \left(\frac{4c^2 t}{|x|^2}\right)^s dw ds \\
&= \frac{1}{\sqrt{\pi}|x|} H_{1,1;1,0;0,1}^{0,1;0,1;1,0} \left[\begin{array}{c} 4c^2 t |x|^{-2} \quad \left| \quad (0; 1, 1); (\frac{1}{2}, 1); - \\ 2\lambda\sqrt{t} \quad \left| \quad (0; 1, \frac{1}{2}); -; (0, 1) \end{array} \right. \right] \\
&\quad + \frac{2\lambda\sqrt{t}}{\sqrt{\pi}|x|} H_{1,1;1,0;0,1}^{0,1;0,1;1,0} \left[\begin{array}{c} 4c^2 t |x|^{-2} \quad \left| \quad (0; 1, 1); (\frac{1}{2}, 1); - \\ 2\lambda\sqrt{t} \quad \left| \quad (-\frac{1}{2}; 1, \frac{1}{2}); -; (0, 1) \end{array} \right. \right].
\end{aligned}$$

Acknowledgement

The work of M. Ferreira, M.M. Rodrigues and N. Vieira was supported by Portuguese funds through the CIDMA - Center for Research and Development in Mathematics and Applications, and the Portuguese Foundation for Science and Technology (“FCT–Fundação para a Ciência e a Tecnologia”), within project UID/MAT/0416/2013.

N. Vieira was also supported by FCT via the FCT Researcher Program 2014 (Ref: IF/00271/2014).

References

- [1] R.G. Buschman, *H-functions of two variables I*, Indian J. Math., **20**, (1978), 132–153.
- [2] M. Caputo, *Linear Models of Dissipation Whose Q Is Almost Frequency Independent-II*, Geophys. J. R. Astron. Soc., **13**-No.5, (1967), 529–539.
- [3] R.F. Camargo, A.O. Chiacchio and E.C. de Oliveira, *Differentiation to fractional orders and the fractional telegraph equation*, J. Math. Phys., **49**-No.3, (2008), **Article ID** 033505, 12p.
- [4] R.C. Cascaval, E.C. Eckstein, L.C. Frota and J.A. Goldstein, *Fractional telegraph equations*, J. Math. Anal. Appl., **276**-No.1, (2002), 145–159.
- [5] J. Chen, F. Liu, V. Anh, *Analytical solution for the time-fractional telegraph equation by the method of separating variables*, J. Math. Anal. Appl., **338**-No.2, (2008), 1364–1377.
- [6] A. Erdélyi, W. Magnus, F. Oberhettinger and G. Tricomi, *Tables of integral transforms-Vol.II*, Bateman Manuscript Project, California Institute of Technology, McGraw-Hill Book Company, New York-Toronto-London, 1954.
- [7] M. Ferreira and N. Vieira, *Fundamental solutions of the time fractional diffusion-wave and parabolic Dirac operators*, J. Math. Anal. Appl., **447**-No.1, (2017), 329–353.
- [8] M. Ferreira, M.M. Rodrigues and N. Vieira, *First and second fundamental solutions of the time-fractional telegraph equation with Laplace or Dirac operators*, Adv. Appl. Clifford Algebr., **28**-No.2, (2018), 28–42.
- [9] M. Ferreira, M.M. Rodrigues and N. Vieira, *Fundamental solution of the multi-dimensional time fractional telegraph equation*, Fract. Calc. Appl. Anal., Fract. Calc. Appl. Anal., **20**-No.4, (2017), 868–894.

- [10] M. Ferreira, M.M. Rodrigues and N. Vieira, *Fundamental solution of the time-fractional telegraph Dirac operator*, Math. Meth. in Appl. Sc., **40**-No.18, (2017), 7033–7050.
- [11] H.J. Haubold, A.M. Mathai and R.K. Saxena, *Mittag-Leffler functions and their applications*, J. Appl. Math., (2011), **Article ID** 298628, 51 p.
- [12] A. Kilbas, H.M. Srivastava and J.J. Trujillo, *Theory and applications of fractional differential equations*, North-Holland Mathematics Studies-Vol.204, Elsevier, Amsterdam, 2006.
- [13] Y. Luchko and R. Gorenflo, *An operational method for solving fractional differential equations with the Caputo derivatives* Acta Math. Vietnam., **24**-No.2, (1999), 207–233.
- [14] M.O. Mamchuev, *Solutions of the main boundary value problems for the time-fractional telegraph equation by the Green function method*, Fract. Calc. Appl. Anal., **20**-No.1, (2017), 190–211.
- [15] E. Orsingher and L. Beghin, *Time-fractional telegraph equations and telegraph processes with Brownian time*, Probab. Theory Relat. Fields, **128**-No.1, (2004), 141–160.
- [16] I. Podlubny, *Fractional Differential Equations. An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*, Mathematics in Science and Engineering 198, Academic Press, San Diego, 1999.
- [17] S.G. Samko, A.A. Kilbas and O.I. Marichev, *Fractional integrals and derivatives: theory and applications*, Gordon and Breach, New York, NY, 1993.
- [18] R.C. Tautz and I. Lerche, *Application of the three-dimensional telegraph equation to cosmic-ray transport*, Res. Astron. Astrophys., **16**(10) (2016), 162.
- [19] S. Yakubovich and M.M. Rodrigues, *Fundamental solutions of the fractional two-parameter telegraph equation*, Integral Transforms Spec. Funct., **23**-No.7, (2012), 509–519.