

Accepted Manuscript

On interval dynamic logic: Introducing quasi-action lattices

Regivan Santiago, Benjamín Bedregal, Alex Madeira, Manuel A. Martins

PII: S0167-6423(19)30010-3
DOI: <https://doi.org/10.1016/j.scico.2019.01.007>
Reference: SCICO 2267

To appear in: *Science of Computer Programming*

Received date: 29 June 2017
Revised date: 16 January 2019
Accepted date: 17 January 2019



Please cite this article in press as: R. Santiago et al., On interval dynamic logic: Introducing quasi-action lattices, *Sci. Comput. Program.* (2019), <https://doi.org/10.1016/j.scico.2019.01.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

On Interval Dynamic Logic: *Introducing Quasi-action Lattices*

Regivan Santiago¹, Benjamín Bedregal¹, Alexandre Madeira^{2,3} and Manuel A. Martins²

¹ Univ. Federal do Rio Grande do Norte — UFRN
Dep. de Informática e Matemática Aplicada — DIMAp
Group for Logic, Language, Information, Theory and Applications — LoLITA
Natal-RN-Brazil

² CIDMA - Dep. Mathematics, Univ. Aveiro, Portugal

³ Dep. Informática Univ. Minho, Portugal

Abstract. In this paper we discuss the incompatibility between the notions of validity and impreciseness in the context of Dynamic Logics. To achieve that we consider the Łukasiewicz action lattice and its interval counterpart, we show how some validities fail in the context of intervals. In order to capture the properties of action lattices that remain valid for intervals we propose a new structure called *Quasi-action Lattices* which generalizes *action lattices* and is able to model both: The Łukasiewicz action lattice, L , and its interval counterpart, \hat{L} .

The notion of graded satisfaction relation is extended to quasi-action lattices. We demonstrate that, in the case of intervals, the relation of graded satisfaction is correct (c.f. Theorem 3) with respect to the graded satisfaction relation on the Łukasiewicz action lattice. Although this theorem guarantees that satisfiability is preserved on intervals, we show that validity is not. We propose, then, to weaken the notion of validity on action lattices to *designated validity* on quasi-action lattices. In this context, Theorem 4 guarantees that the dynamic formulæ which are *valid with respect to L* will be *designated valid with respect to \hat{L}* .

1 Introduction

Dynamic Logics (DL) are extensions of modal logic. They are recognised as the most adequate logics to reason about computational systems in an assertional way [15]. In its origin, DL was introduced by V. Pratt [31] as a modal logic suitable to represent and reason about Hoare triples. Since then, DL assumed a central role in program verification. Today, not only in response to the explosion of programming and specification languages, but also in the emerging heterogeneous nature that a program can assume, a wide family of DL were defined to be applied to more general complex behaviors. This ranges from the standard versions for sequential imperative programmes (e.g. [15]) to other versions tailored for new computing paradigms, either probabilistic systems, following the original work of D. Kozen [20], hybrid systems with the differential dynamic logic

by Platzer [30] or even quantum versions due to Baltag and Smets [1]. Within this variety of dynamic logics, in [23], a method for a systematic construction of many-valued dynamic logics was studied. The method is parameterized by an action lattice that supports both the computational paradigm and the truth space, combining in just one structure the underlying Kleene algebra for the computations and the residuated lattice for proposition assertions.

Logics with many-valued semantics are applied in a variety of fields such as: Decision Making, Image Processing, Clustering, etc. One of such logics, which is a very important logic, is that of Lukasiewicz [9], which semantics is based on the residuated lattice $\mathbf{L} = \langle [0, 1], \rightarrow, 0, 1 \rangle$ — where: $a \rightarrow b = \min(1, 1 - a + b)$. In this logic, the truth-values may be thought of as arising from normalized measurements of bounded physical observables, just as boolean truth-values arise from yes/no-observables [27, §1.6]. The space of valueness $[0, 1]$ models exact measurements which presupposes an uncertainty and are not encoded by the elements of $[0, 1]$. In order to capture and deal with such uncertainty the indicated approach is to use the space of closed intervals $\mathbb{I}([0, 1]) = \{[a, b] \mid [a, b] \subseteq [0, 1]\}$ instead of $[0, 1]$. The idea is that an interval $[a, b]$ represents any real number inside it with a quality of $b - a$.

In [34] Santiago et. al. showed how interval can be used to model computational systems with imprecise transitions through an interval Kleene algebra. In this paper, we show that such algebra cannot be extended to an action lattice and hence be a space of truth values for axioms of DL. To overcome that, we extend the notion of action lattices to a new algebraic structure called *Quasi-action lattices*. This structure models both: The Lukasiewicz action lattice \mathbf{L} and its interval counterpart $\widehat{\mathbf{L}}$.

The notion of *satisfiability* is extended to Quasi-action lattices and Theorem 3 shows that satisfiability on the action lattice \mathbf{L} is represented by satisfiability on $\widehat{\mathbf{L}}$. The notion of representation is discussed in Section 3.

Although satisfiability is preserved, we show in Section 5 that validity is not. We propose to extend the notion of *validity on action lattices* to *designated validity on Quasi-action lattices*. We show that every DL formula which is valid with respect to the action lattice \mathbf{L} is designated valid with respect to $\widehat{\mathbf{L}}$.

Outline of the paper. Section 2 introduces a Multi-valued Dynamic Logic following the ‘dynamization’ method proposed by Madeira et. al [23]. Section 3 makes an overview of interval arithmetic and gives the ‘intervalization’ of the Lukasiewicz action lattice: \mathbf{L} . It proves that this structure is a *Kleene algebra*. In Section 3.2, we discuss some pitfalls in defining action lattices over intervals. Section 4 introduces the notion of *Quasi-action Lattices* which is the abstraction of interval Lukasiewicz action lattice: $\widehat{\mathbf{L}}$. It proposes the notion of satisfiability on Quasi-action lattices and connects satisfiability with respect to \mathbf{L} with satisfiability with respect to $\widehat{\mathbf{L}}$. Section 5 shows *the impact of imprecision on validity*, proposes the weakening of validity on action lattices to designated validity on quasi-action lattices and proves that every valid formula with respect to \mathbf{L} is designated valid with respect to $\widehat{\mathbf{L}}$. Finally, in Section 6, we present some final remarks.

2 Ł-Fuzzy Dynamic Logic

Many-valued versions of Modal Logics have been discussed in the literature, the purposed logics vary in the focus where the many-valueness is presented: In accessibility relations, in propositions interpretation or in both. The latter is the case of the works [13, 14] of M. Fitting suggesting a logic with many-valueness evaluated in finite Heyting algebras. Later, it was deeply investigated by F. Bou et al in [7], who adopted the more generic truth support of finite integral commutative residuated lattices.

The literature is not so rich with respect to Many-Valued Dynamic Logics. J. Hughes et al introduced in [18] a propositional dynamic logic over the continuum truth $(0, 1)$ -lattice with the standard fuzzy residues. However, from the perspective of dynamic logic, this formalism is quite restrictive, since it leaves behind both transitive closure and non deterministic choice. In the context of rational decision theory, C. Liao [22] also introduced another different many-valued dynamic logic w.r.t. the specific continuum truth $(0, 1)$ -lattice.

A systematic method to build Multi-valued Dynamic Logics was then introduced in [23, 24]. This method is parameterized by an action lattice [19], an algebraic structure that provides a generic support for computational space (as a Kleene algebra) and for truth space (as residuated lattice). The logic introduced in this section is based on this work and can be captured as an instantiation of this method.

2.1 The Łukasiewicz action lattice

The role of the Łukasiewicz residuated lattice, i.e., the algebraic structure

$$\mathbf{L} = ([0, 1], \max, \odot, 0, 1, \rightarrow)$$

with

- $x \rightarrow y = \min(1, 1 - x + y)$ and
- $x \odot y = \max(0, y + x - 1)$,

is taken as the standard fuzzy truth space [36]. Moreover, as stated above, we are looking for a structure suitable to support a fuzzy computational model. Whenever the \max and the \odot operators are used to model the choice and composition of atomic actions, we need to consider a Kleene operator to model the recursive iteration of programs. These operators constitute the components of an action lattice [19], the structure taken in [23, 24] as a generic parameter for a multi-valued logic definition. More precisely:

Definition 1. An action lattice is a tuple

$$\mathbf{A} = (U, +, :, 0, 1, *, \rightarrow, \cdot)$$

where U is a set, 0 and 1 are constants in U , $*$ is a unary operation over U and $+, :, \rightarrow$ and \cdot are binary operations over U satisfying the axioms enumerated in Figure 1, where the relation \leq is induced by “+”: $a \leq b$ iff $a + b = b$.

$$\begin{array}{ll}
a + (b + c) = (a + b) + c & (1) \\
a + b = b + a & (2) \\
a + a = a & (3) \\
a + 0 = 0 + a = a & (4) \\
a; (b; c) = (a; b); c & (5) \\
a; 1 = 1; a = a & (6) \\
a; (b + c) = (a; b) + (a; c) & (7) \\
(a + b); c = (a; c) + (b; c) & (8) \\
a; 0 = 0; a = 0 & (9) \\
1 + a + (a^*; a^*) \leq a^* & (10) \\
a; x \leq x \Rightarrow a^*; x \leq x & (11) \\
x; a \leq x \Rightarrow x; a^* \leq x & (12) \\
a; x \leq b \Leftrightarrow x \leq a \rightarrow b & (13) \\
a \rightarrow b \leq a \rightarrow (b + c) & (14) \\
(x \rightarrow x)^* = x \rightarrow x & (15) \\
a \cdot (b \cdot c) = (a \cdot b) \cdot c & (16) \\
a \cdot b = b \cdot a & (17) \\
a \cdot a = a & (18) \\
a + (a \cdot b) = a & (19) \\
a \cdot (a + b) = a & (20)
\end{array}$$

Fig. 1. Axiomatisation of action lattices (from [19])

Note that, by (19) and (20), the natural order \leq can be equivalently defined by $a \leq b$ iff $a \cdot b = a$. Observe that by restricting the definition of \mathbf{A} to the structure $K(\mathbf{A}) = (U, +, ;, 0, 1, *)$ axiomatized by (1)–(12) we obtain the definition of a Kleene algebra [10, 21]. In the context of this work, this will be called the underlying Kleene algebra of \mathbf{A} . Moreover, by considering structure $(U, +, ;, 0, 1, \rightarrow, *)$ axiomatized by (1)–(15) we obtain the definition of (left-residuated) action algebra [32].

For the illustration of the structure with several examples and properties we suggest [23]. Just as example, we can consider a discrete 3-valued lattice underling the 3-valued logic:

Example 1 (3 - linear three-value lattice). The explicit introduction of a denotation for unknown gives rise to the following linear lattice of three elements

$$\mathbf{3} = (\{\top, u, \perp\}, \vee, \wedge, \perp, \top, *, \rightarrow, \wedge)$$

where

$$\begin{array}{c|ccc}
\vee & \perp & u & \top \\
\hline
\perp & \perp & u & \top \\
u & u & u & \top \\
\top & \top & \top & \top \\
\hline
\wedge & \perp & u & \top \\
\hline
\perp & \perp & \perp & \perp \\
u & \perp & u & u \\
\top & \perp & u & \top \\
\hline
\rightarrow & \perp & u & \top \\
\hline
\perp & \top & \top & \top \\
u & \perp & \top & \top \\
\top & \perp & u & \top \\
\hline
* & \perp & \top & \top \\
\hline
\perp & \perp & \top & \top \\
u & \perp & \top & \top \\
\top & \perp & u & \top \\
\hline
\top & \top & \top & \top
\end{array}$$

It is easy to observe that, as a consequence of axiom (10), whenever 1 is the greatest element, we have that $x^* = 1$, for all x .

Hence we have all the ingredients to introduce the Łukasiewicz arithmetic lattice, a structure that plays the main role in the theory developed in the following:

Definition 2 (L - the Łukasiewicz arithmetic lattice). *The Łukasiewicz arithmetic lattice is the structure:*

$$L = ([0, 1], \max, \odot, 0, 1, *, \rightarrow, \min)$$

where

- $x \rightarrow y = \min(1, 1 - x + y)$,
- $x \odot y = \max(0, y + x - 1)$ and
- $x^* = 1$.

Lemma 1. *L is an action lattice.*

Proof. The verification of properties (1)-(4), as and well (16)-(20) can be straightforwardly verified since L is a lattice. Since $x^* = 1$, property (10) trivially holds; moreover, assuming (4), we have also (11), (12) and (15). The remaining properties are directly entailed by the definitions of “ \rightarrow ” and “ \odot ”.

2.2 The L-Fuzzy Dynamic Logic

Signatures for the dynamic logic intended to be interpreted in L, \mathbf{LDL} , are exactly the same as the ones of Propositional Dynamic Logic: Signatures are pairs (Π, Prop) of disjoint sets of **atomic programs** Π and of **propositions symbols** Prop .

Formulae of \mathbf{LDL} consist of the positive fragment of Propositional Dynamic Logic: The set of Π -programs, denoted by $\mathbf{Prg}(\Pi)$, consists of all expressions generated by:


$$\pi ::= \pi_0 \mid \pi; \pi \mid \pi + \pi \mid \pi^*$$

for $\pi_0 \in \Pi$. Given a signature (Π, Prop) , we specify the set of \mathbf{LDL} -formulae for (Π, Prop) , denoted by $\mathbf{Fm}^{\mathbf{LDL}}(\Pi, \text{Prop})$, by the following grammar:

$$\alpha ::= \top \mid \perp \mid p \mid \alpha \vee \alpha \mid \alpha \wedge \alpha \mid \alpha \rightarrow \alpha \mid \alpha \leftrightarrow \alpha \mid \langle \pi \rangle \alpha \mid [\pi] \alpha$$

for $p \in \text{Prop}$ and $\pi \in \mathbf{Prg}(\Pi)$.

Semantics. As expected, the interpretation of atomic programs are Kripke structures with weighted transitions. For instance, the atomic programs $\Pi = \{\pi, \pi'\}$ can be realized by the following structures



where the tags mention the uncertainty level of each state transitions. These weighted state transition systems are usually represented by the underlying adjacency matrices. We use the notation $A_\pi(w, w')$ to mean the fuzziness degree at transition (w, w') of program π ; e.g. in the case of (21) we have $A_\pi(s_1, s_2) = \frac{\sqrt{2}}{3}$.

$$\mathcal{A}_\pi = \begin{bmatrix} 0 & \frac{\sqrt{2}}{3} \\ 0 & 0.7 \end{bmatrix} \quad \mathcal{A}_{\pi'} = \begin{bmatrix} 0 & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{3}}{2} & 0.5 \end{bmatrix}$$

Moreover, we need a mathematical framework to interpret more complex programs, i.e. regular expressions of atomic programs, e.g. “ $\pi + \pi'$ ”. In other words, we need to consider a computational space for LDL where the programs are interpreted. Based on the classic matricial constructions over Kleene algebras (see [10, 21]) we consider the following structure

$$\mathbb{M}_n(\mathbf{L}) = (M_n(\mathbf{L}), \max, \odot, 0, 1, *) \quad (22)$$

considering its components as follows:

1. $M_n(\mathbf{L})$ is the space of $(n \times n)$ -matrices over \mathbf{L} ; i.e. with elements in \mathbf{L} .
2. for any $A, B \in M_n(\mathbf{L})$, define $M = \max(A, B)$ by $M_{ij} = \max(A_{ij}, B_{ij})$, $i, j \leq n$.
3. for any $A, B \in M_n(\mathbf{L})$, define $M = A \odot B$ by taking

$$\max(A_{i1} \odot B_{1j}, \max(A_{i2} \odot B_{2j}, (\dots, \max(A_{in} \odot B_{nj}) \dots))$$

4. the matricial 1 and 0 are the $(n \times n)$ -matrices defined by $1_{i,j} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$

and $0_{i,j} = 0$, for any $i, j \leq n$.

5. for any $M = [a] \in \mathbb{M}_1(\mathbf{L})$, $M^* = [a^*]$;

for any $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in M_n(\mathbf{L})$, $n > 1$, where A and D are square matrices, define

$$M^* = \left[\begin{array}{c|c} F^* & F^* \odot (B \odot D^*) \\ \hline (D^* \odot C) \odot F^* & \max(D^*, (D^* \odot (C \odot (F^* \odot (B \odot D^*)))) \end{array} \right]$$

where $F = \max(A, B \odot (D^* \odot C))$. Note that this construction is recursively defined from the base case (where $n = 2$) where the operations of the base action lattice \mathbf{A} are used.

A classic result (e.g. [10, 21]) establishes that Kleene algebras are closed under formation of matrices.

Fact 1 *The structure $\mathbb{M}_n(\mathbf{L}) = (M_n(\mathbf{L}), \max, \odot, 0, 1, *)$ defined above is a Kleene algebra.*

This justifies the adoption of $\mathbb{M}_n(\mathbf{L})$ as a well behaved computational space for LDL .

Definition 3. *LDL -models for a set of propositions Prop and programs Π , denoted by $\text{Mod}^{\text{LDL}}(\Pi, \text{Prop})$, consist of tuples:*

$$\mathfrak{M} = (W, V, (\mathcal{A}_\pi)_{\pi \in \Pi})$$

where W is a finite set (of states), $V : \text{Prop} \times W \rightarrow [0, 1]$ is a function, and $\mathcal{A}_\pi \in \mathbb{M}_n(\mathbf{L})$, with n standing for the cardinality of W .

The interpretation of programs in these models belongs to the space of the matrices over the underlying Kleene algebra of \mathbf{L} . Each matrix represents the effect of a program executing from any point of the model. Formally, the interpretation of a program $\pi \in \mathbf{Prog}(II)$ in a model $\mathfrak{M} \in \text{Mod}^{\mathbf{L}\mathcal{D}\mathcal{L}}(II, \text{Prop})$ is recursively defined, from the set of atomic programs II , as follows:


$$\mathcal{A}_{\pi;\pi'} = \mathcal{A}_{\pi} \odot \mathcal{A}_{\pi'}, \mathcal{A}_{\pi+\pi'} = \max(\mathcal{A}_{\pi}, \mathcal{A}_{\pi'}) \text{ and } \mathcal{A}_{\pi^*} = \mathcal{A}_{\pi}^*.$$

together with the constants interpretations $\mathcal{A}_1 = \mathbf{1}$ and $\mathcal{A}_0 = \mathbf{0}$.

Returning to our running example, we are able to calculate the interpretation of the program $\pi + \pi'$ by making:

$$\mathcal{A}_{\pi+\pi'} = \max(\mathcal{A}_{\pi}, \mathcal{A}_{\pi'}) = \max\left(\begin{bmatrix} 0 & \frac{\sqrt{2}}{3} \\ 0 & 0.7 \end{bmatrix}, \begin{bmatrix} 0 & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{3}}{2} & 0.5 \end{bmatrix}\right) = \begin{bmatrix} 0 & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{3}}{2} & 0.7 \end{bmatrix} \quad (23)$$

that represents the following weighted transition system:



By considering the interpretation of the propositions $\text{Prop} = \{p, q\}$ as $V(p, s_1) = 0.1$, $V(q, s_1) = 0.5$, $V(p, s_2) = \frac{\pi}{4}$ and $V(q, s_2) = 0.75$ we have a complete description of a concrete $(\{\pi, \pi'\}, \{p, q\})$ -model $\mathcal{A} = (\{s_1, s_2\}, V, (\mathcal{A}_p)_{p \in \{\pi, \pi'\}})$.

Satisfaction.

As mentioned above, the carrier of \mathbf{L} corresponds to the space of truth degrees for $\mathbf{L}\mathcal{D}\mathcal{L}$. In what follows we define what is the graded satisfaction relation for a model $\mathfrak{M} \in \text{Mod}^{\mathbf{L}\mathcal{D}\mathcal{L}}(II, \text{Prop})$.

Definition 4. *The graded satisfaction relation for a model $\mathfrak{M} \in \text{Mod}^{\mathbf{L}\mathcal{D}\mathcal{L}}(II, \text{Prop})$, is a function:*

$$\models_{\mathbf{L}} : W \times \text{Fm}^{\mathbf{L}\mathcal{D}\mathcal{L}}(II, \text{Prop}) \rightarrow [0, 1]$$

recursively defined as follows:

- $(w \models_{\mathbf{L}} \top) = 1$
- $(w \models_{\mathbf{L}} \perp) = 0$
- $(w \models_{\mathbf{L}} p) = V(p, w)$, for any $p \in \text{Prop}$
- $(w \models_{\mathbf{L}} \rho \wedge \rho') = \min\{(w \models_{\mathbf{L}} \rho), (w \models_{\mathbf{L}} \rho')\}$
- $(w \models_{\mathbf{L}} \rho \vee \rho') = \max\{(w \models_{\mathbf{L}} \rho), (w \models_{\mathbf{L}} \rho')\}$
- $(w \models_{\mathbf{L}} \rho \rightarrow \rho') = (w \models_{\mathbf{L}} \rho) \rightarrow (w \models_{\mathbf{L}} \rho')$
- $(w \models_{\mathbf{L}} \langle \pi \rangle \rho) = \max\{\mathcal{A}_{\pi}(w, w') \odot (w' \models_{\mathbf{L}} \rho) \mid w' \in W\}$

$$- (w \models_{\mathbb{L}} [\pi]\rho) = \min \{ \mathcal{A}_{\pi}(w, w') \rightarrow (w' \models_{\mathbb{L}} \rho) \mid w' \in W \}$$

Example 2. In order to illustrate the definition, the calculation of the truth degree for the formula “ $\langle \pi + \pi' \rangle (p \rightarrow q)$ ” in the proposed model \mathcal{A} can be achieved as follows:

$$\begin{aligned} (s_1 \models_{\mathbb{L}} \langle \pi + \pi' \rangle (p \rightarrow q)) &= \max(0 \odot (0.1 \rightarrow 0.5), \frac{\sqrt{2}}{2} \odot (0.75 \rightarrow \frac{\pi}{4})) \\ &= \frac{\sqrt{2}}{2} \odot (0.75 \rightarrow \frac{\pi}{4}) \\ &= \frac{\sqrt{2}}{2} \odot \min(1, 1 - 0.75 + \frac{\pi}{4}) \\ &= \frac{\sqrt{2}}{2} \end{aligned} \quad (24)$$

Therefore, we conclude with a degree of certainty $\frac{\sqrt{2}}{2}$ that, after executing “ $\pi + \pi'$ ” from the state s_1 , we have $p \rightarrow q$.

Assuming this semantics for $\mathbb{L}\mathcal{D}\mathcal{L}$, the investigation of calculus for this logic is the natural step to proceed. The next theorem is a first step on this direction, it shows the validity of some propositional dynamic axioms [16] in $\mathbb{L}\mathcal{D}\mathcal{L}$:

Theorem 1. *The following are valid formulae in the logic $\mathbb{L}\mathcal{D}\mathcal{L}$:*

1. $\langle \pi \rangle (\rho \vee \rho') \leftrightarrow \langle \pi \rangle \rho \vee \langle \pi \rangle \rho'$
2. $\langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho'$
3. $\langle \pi + \pi' \rangle \rho \leftrightarrow \langle \pi \rangle \rho \vee \langle \pi' \rangle \rho$
4. $\langle \pi; \pi' \rangle \rho \leftrightarrow \langle \pi \rangle \langle \pi' \rangle \rho$
5. $\langle \pi \rangle \perp \leftrightarrow \perp$
6. $\langle \pi \rangle \rho \rightarrow \langle \pi^* \rangle \rho$
7. $\langle \pi^* \rangle \rho \leftrightarrow \langle \pi^*; \pi^* \rangle \rho$
8. $\langle \pi^* \rangle \rho \leftrightarrow \langle \pi^{**} \rangle \rho$
9. $\langle \pi^* \rangle \rho \leftrightarrow \rho \vee \langle \pi \rangle \langle \pi^* \rangle \rho$
10. $[\pi + \pi'] \rho \leftrightarrow [\pi] \rho \wedge [\pi'] \rho$
11. $[\pi] (\rho \wedge \rho') \leftrightarrow [\pi] \rho \wedge [\pi] \rho'$

Proof. As stated above, the logic $\mathbb{L}\mathcal{D}\mathcal{L}$ can be built as an instantiation of the generic method of dynamization [23]. The same reference reports, at this abstract level, a systematic study of the validity of the previous axioms in dynamizations parametric to generic action lattices. Hence, because of the properties of the action lattice \mathbb{L} , all these proofs can be constructed by instantiation from this generic study. Indeed, in order to illustrate the principles involved, we explicitly extract here the proof for the validity of formula 1.

Before we proceed, observe that if $x = 1$, then (13) can be written as: “ $a; 1 \leq b \Leftrightarrow 1 \leq a \rightarrow b$ ”; i.e. $a \leq b \Leftrightarrow a \rightarrow b = 1$. Therefore, for every $w \in W$ and formulae $\rho, \rho' \in \text{Fm}^{\mathbb{L}\mathcal{D}\mathcal{L}}(\Pi, \text{Prop})$:

$$(w \models_{\mathbb{L}} \rho \rightarrow \rho') = 1 \text{ iff } (w \models_{\mathbb{L}} \rho) \leq (w \models_{\mathbb{L}} \rho')$$

Hence,

$$\begin{aligned}
& (w \models_{\mathbf{L}} \langle \pi \rangle (\rho \vee \rho')) \\
= & \quad \{ \text{defn of } \models_{\mathbf{L}} \} \\
& \max \{ \mathcal{A}_{\pi}(w, w') \odot (w' \models_{\mathbf{L}} \rho \vee \rho') \mid w' \in W \} \\
= & \quad \{ \text{defn of } \models_{\mathbf{L}} \} \\
& \max \{ \mathcal{A}_{\pi}(w, w') \odot \max \{ (w' \models_{\mathbf{L}} \rho), (w' \models_{\mathbf{L}} \rho') \} \} \\
= & \quad \{ (7) \} \\
& \max \{ \mathcal{A}_{\pi}(w, w') \odot (w' \models_{\mathbf{L}} \rho), (\mathcal{A}_{\pi}(w, w') \odot (w' \models_{\mathbf{L}} \rho')) \mid w' \in W \} \\
= & \quad \{ \text{by (1) and (2)} \} \\
& \max \{ \max \{ \mathcal{A}_{\pi}(w, w') \odot (w' \models_{\mathbf{L}} \rho) \mid w' \in W \}, \max \{ \mathcal{A}_{\pi}(w, w') \odot (w' \models_{\mathbf{L}} \rho') \mid w' \in W \} \} \\
= & \quad \{ \text{defn of } \models_{\mathbf{L}} \} \\
& \max \{ (w \models_{\mathbf{L}} \langle \pi \rangle \rho), (w \models_{\mathbf{L}} \langle \pi \rangle \rho') \} \\
= & \quad \{ \text{defn of } \models_{\mathbf{L}} \} \\
& (w \models_{\mathbf{L}} \langle \pi \rangle \rho \vee \langle \pi \rangle \rho')
\end{aligned}$$

Therefore, $\langle \pi \rangle (\rho \vee \rho') \leftrightarrow \langle \pi \rangle \rho \vee \langle \pi \rangle \rho'$ is valid.

3 \mathbf{L} -Interval algebra

The space of values $[0, 1]$ models exact measurements/truth values which is far from the real-world. In fact, any measurement presupposes an uncertainty which is not encoded by the elements of $[0, 1]$. Another situation arises whenever an expert is unable to supply an exact membership of an object in a fuzzy set, in this case it can be provided a closed subinterval of $[0, 1]$ as an expression of the inability to supply an exact answer [8]. Therefore, assuming the **Lukasiewicz arithmetic lattice** $\mathbf{L} = \langle [0, 1], \max, \odot, 0, 1, *, \rightarrow, \min \rangle$ as a natural space of measurements/truth values [27, §1.6] it is reasonable to investigate its interval counterpart. But what would such interval counterpart be? Before we proceed to answer this question, let us recall some definitions and facts about the interval counterpart of real numbers algebra: $\langle \mathbb{R}; +, -, /, \times, 1, 0 \rangle$.

In the 50's Ramon Moore [25, 26] and Teruo Sunaga [35] proposed the so called **interval arithmetics**. Interval arithmetics is a set of operations on the set of all closed intervals $[a, b] \subseteq \mathbb{R}$. They defined the arithmetic in the following way:

1. $[a, b] + [c, d] = [a + c, b + d]$
2. $-[c, d] = [-d, -c]$
3. $[a, b] \cdot [c, d] = [\min P, \max P]$; where $P = \{a \cdot c, a \cdot d, b \cdot c, b \cdot d\}$

4. $[a, b]^{-1} = [1/b, 1/a]$; provided that $0 \notin [a, b]$
5. $[a, b] - [c, d] = [a - d, b - c]$
6. $[a, b]/[c, d] = [a, b] \cdot [c, d]^{-1}$

Observe what happens with each operation:

1. If $x \in [a, b]$ and $y \in [c, d]$, then $(x + y) \in [a, b] + [c, d]$,
2. If $x \in [a, b]$ and $y \in [c, d]$, then $(x \cdot y) \in [a, b] \cdot [c, d]$,
3. If $x \in [a, b]$ and $y \in [c, d]$, then $(x/y) \in [a, b]/[c, d]$, and
4. If $x \in [a, b]$, then $(-x) \in -[a, b]$.

The arithmetic on intervals reveals two desired properties: (a) **Correctness** and (b) **Optimality**.

“*Correctness*. . . when an expression is evaluated using intervals, it yields an interval containing all results of pointwise evaluations based on point values that are elements of the argument intervals.

...

“*Optimality*. By optimality, we mean that the computed floating-point interval is not wider than necessary.”

Hickey *et.al*[17, p.1040]

The term *Correctness* connects n -ary interval operations F with n -ary real operations f and means that if F is *correct with respect to* f , then we can enfold any exact value $r \in \mathbb{R}$ in a closed interval $[a, b]$, such that $r \in [a, b]$, and then simply operate with such “envelopes” by using F , because the resulting interval $F([a, b])$ will enfold the desired result $f(r)$. Formally, a function F is *correct with respect to a real function* f whenever:

$$r \in [a, b] \Rightarrow f(r) \in F([a, b]) \quad (25)$$

In practice, exact values are replaced by intervals which are operated with correct interval functions. Intervals enfold the exact values and provide a measure of impreciseness through their widths.

Santiago *et. al* [2, 33] investigated the notion of Correctness. Instead of correctness the authors used the term **representation**, since interval expressions could be faced not just as machine representations of an exact calculation, but also as an instance of a “mathematical representation of real numbers”⁴. Beyond correctness, interval arithmetic is also optimum; namely the resulting intervals contain only the values of real operations. We could say that the proposed algebra of intervals is the **best interval representation for the arithmetic of real numbers**.

One side-effect of this process (called *intervalization*) is the loss of algebraic properties. The resulting structure is not an euclidean field; for example “ $X - X$ ” is not always equal to $[0, 0]$. As we can see in the following, like real numbers, some properties of \mathbb{L} are lost when we consider intervals of $[0, 1]$.

⁴ This idea is confirmed in some Representation Theorems of Euclidean continuous functions.

3.1 On the Interval Lukasiewicz Lattice

The Lukasiewicz arithmetic lattice $\mathbf{L} = \langle [0, 1], \max, \odot, 0, 1, *, \rightarrow, \min \rangle$ contains non-finitely representable elements; e.g. irrational numbers. In a similar way we can think of an interval algebra for \mathbf{L} . A piece of such algebra was introduced by Bedregal and Santiago in [4]. There, the authors proposed a correct interval implication for “ \rightarrow ”. In what follows, we propose the interval counterpart for $\mathbf{L} = \langle [0, 1], \max, \odot, 0, 1, *, \rightarrow, \min \rangle$ in such a way that the resulting operations are correct and optimal, i.e. they are **best interval representations**.

Definition 5. Consider the real unit interval $U = [0, 1] \subseteq \mathbb{R}$ and the set $\mathbb{U} = \{[a, b] \mid 0 \leq a \leq b \leq 1\}$ of subintervals of U . For any interval $X \in \mathbb{U}$, \underline{X} is the minimum of X and \overline{X} the maximum of X ; i.e. $X = [\underline{X}, \overline{X}]$. Given two intervals $X, Y \in \mathbb{U}$, let be the following partial orders on \mathbb{U} :

(i) The *product or Kulisch-Miranker order* :

$$X \leq Y \Leftrightarrow \underline{X} \leq \underline{Y} \wedge \overline{X} \leq \overline{Y}; \quad (26)$$

(ii) The *set inclusion order*: for all $X, Y \in \mathbb{U}$,

$$X \subseteq Y \Leftrightarrow \underline{Y} \leq \underline{X} \wedge \overline{X} \leq \overline{Y}. \quad (27)$$

Definition 6 ([33]). An interval $X \in \mathbb{U}$ is a **representation** of any real number $\alpha \in X$. Considering two interval representations X and Y for a real number α , X is said to be an interval representation of α better than Y , if $X \subseteq Y$. This notion can also be naturally extended for n -tuples of intervals. A function $F : \mathbb{U}^n \rightarrow \mathbb{U}$ is said to be an **interval representation** of a real function $f : [0, 1]^n \rightarrow [0, 1]$ if, for each $\mathbf{X} \in \mathbb{U}^n$ and $\mathbf{x} \in \mathbf{X}$, $f(\mathbf{x}) \in F(\mathbf{X})$. F is also said to be correct with respect to f . An interval function $F : \mathbb{U}^n \rightarrow \mathbb{U}$ is said to be an interval representation of a real function $f : [0, 1]^n \rightarrow [0, 1]$ better than an interval function $G : \mathbb{U}^n \rightarrow \mathbb{U}$, if $F(\mathbf{X}) \subseteq G(\mathbf{X})$, for each $\mathbf{X} \in \mathbb{U}^n$. The **best interval representation** of a real function $f : [0, 1]^n \rightarrow [0, 1]$ is the interval function $\hat{f} : \mathbb{U}^n \rightarrow \mathbb{U}$, defined by

$$\hat{f}(\mathbf{X}) = [\inf\{f(\mathbf{x}) \mid \mathbf{x} \in \mathbf{X}\}, \sup\{f(\mathbf{x}) \mid \mathbf{x} \in \mathbf{X}\}]. \quad (28)$$

In what follows we show the best (possible) interval representation for $\mathbf{L} = \langle [0, 1], \max, \odot, 0, 1, *, \rightarrow, \min \rangle$. Before we proceed, we provide some required definitions and facts about some fuzzy connectives and their best interval representation:

Let (L, \leq_L, \perp, \top) be a bounded lattice. A t-norm on L is an \leq_L -increasing function $T : L^2 \rightarrow L$ that is commutative, associative and has \top as neutral element [28]. Dually, a t-conorm is a commutative, associative and \leq_L -increasing function $\mathcal{S} : L^2 \rightarrow L$ which has \perp as neutral element [28]. A function $I : L^2 \rightarrow L$ which is \leq_L -decreasing in the first variable, \leq_L -increasing in the second variable, $I(\perp, \perp) = I(\top, \top) = \top$ and $I(\top, \perp) = \perp$ is called implication of L [29].

- (Fact1)** If T is a t-norm on the bounded lattice $([0, 1], \leq, 0, 1)$ then \widehat{T} is a t-norm on the bounded lattice $(\mathbb{U}, \leq, [0, 0], [1, 1])$ such that, $\widehat{T}(X, Y) = [T(\underline{X}, \underline{Y}), T(\overline{X}, \overline{Y})]$ [5, 12].
- (Fact2)** If S is a t-conorm on the bounded lattice $([0, 1], \leq, 0, 1)$ then \widehat{S} is a t-conorm on the bounded lattice $(\mathbb{U}, \leq, [0, 0], [1, 1])$ such that, $\widehat{S}(X, Y) = [S(\underline{X}, \underline{Y}), S(\overline{X}, \overline{Y})]$ [6, 12].
- (Fact3)** If I is an implication on the bounded lattice $([0, 1], \leq, 0, 1)$ then \widehat{I} is an implication on the bounded lattice $(\mathbb{U}, \leq, [0, 0], [1, 1])$ such that, $\widehat{I}(X, Y) = [I(\overline{X}, \underline{Y}), I(\underline{X}, \overline{Y})]$ [4].

Definition 7. Given $X, Y \in \mathbb{U}$.

1. $Max(X, Y) = [\max(\underline{X}, \underline{Y}), \max(\overline{X}, \overline{Y})]$
2. $Min(X, Y) = [\min(\underline{X}, \underline{Y}), \min(\overline{X}, \overline{Y})]$
3. $X \odot Y = [(\underline{X} \odot \underline{Y}), (\overline{X} \odot \overline{Y})] = [\max(0, \underline{X} + \underline{Y} - 1), \max(0, \overline{X} + \overline{Y} - 1)]$
4. $X \Rightarrow Y = [(\overline{X} \rightarrow \underline{Y}), (\underline{X} \rightarrow \overline{Y})] = [\min(1, 1 - \overline{X} + \underline{Y}), \min(1, 1 - \underline{X} + \overline{Y})]$
5. $X^{\circledast} = [\underline{X}^*, \overline{X}^*] = [1, 1]$.

Proposition 1. All these interval operations are the best interval representations of the operations in L ; i.e. $Max = \widehat{\max}$, $Min = \widehat{\min}$, $\odot = \widehat{\odot}$, $\Rightarrow = \widehat{\Rightarrow}$, and $\circledast = \widehat{\circledast}$.

Proof. The operations “min” and “ \odot ” are T-norms on $[0, 1]$ and the best interval representation of T-norms, according to **(Fact1)**, is given by $\widehat{T}(X, Y) = [T(\underline{X}, \underline{Y}), T(\overline{X}, \overline{Y})]$. The “max” operation is a T-conorm on $[0, 1]$ and the best interval representations of T-conorms, according to **(Fact2)**, is given by $\widehat{S}(X, Y) = [S(\underline{X}, \underline{Y}), S(\overline{X}, \overline{Y})]$. According to **(Fact3)** $\Rightarrow = \widehat{\Rightarrow}$. Finally, it is trivial that $\circledast = \widehat{\circledast}$.

Proposition 2. $X \leq Y$ iff $Max(X, Y) = Y$.

Proof. $X \leq Y$ iff $\underline{X} \leq \underline{Y}$ and $\overline{X} \leq \overline{Y}$ iff $\max(\underline{X}, \underline{Y}) = \underline{Y}$ and $\max(\overline{X}, \overline{Y}) = \overline{Y}$ iff $Max(X, Y) = Y$.

Theorem 2. The structure $K(\widehat{L}) = \langle \mathbb{U}, \widehat{\max}, \widehat{\odot}, [0, 0], [1, 1], \widehat{\circledast} \rangle$ is a Kleene algebra.


Proof. $\widehat{\max}$ trivially satisfies equations (1)-(4). It is also clear that $\widehat{\odot}$ satisfies equations (5)-(6). Equation (7) is also satisfied, since given $X, Y, Z \in \mathbb{U}$, $X \odot \widehat{\max}(Y, Z) = [\underline{X} \odot \max(\underline{Y}, \underline{Z}), \overline{X} \odot \max(\overline{Y}, \overline{Z})] = [\max(\underline{X} \odot \underline{Y}, \underline{X} \odot \underline{Z}), \max(\overline{X} \odot \overline{Y}, \overline{X} \odot \overline{Z})] = \widehat{\max}(X \odot Y, X \odot Z)$. Equation (8) comes from equation (7) and the commutativity of $\widehat{\odot}$. Equation (9) is also satisfied, since given $X \in \mathbb{U}$, $X \odot [0, 0] = [\underline{X} \odot 0, \overline{X} \odot 0] = [0, 0] = [0 \odot \underline{X}, 0 \odot \overline{X}] = [0, 0] \odot X$. Since for every $A \in \mathbb{U}$, $A^* = [1, 1]$, and $[1, 1]$ is the top element in \mathbb{U} , then inequation (10) is trivially satisfied. Again, since $A^* = [1, 1]$, then $A^* \odot X = X$, therefore implication (11) is satisfied. A similar argument applies to implication (12).

Since $K(\widehat{\mathbf{L}})$ is a Kleene algebra, we can canonically construct, as in (22), the space of matrices $\mathbb{M}_n(K(\widehat{\mathbf{L}}))$ (which is also a Kleene algebra).

Observation: According to Proposition 1 every operation of the Kleene algebra $K(\widehat{\mathbf{L}})$ is the best interval representation of the respective operation of $K(\mathbf{L})$. Therefore, we can say that $K(\widehat{\mathbf{L}})$ and $\mathbb{M}_n(K(\widehat{\mathbf{L}}))$ are, respectively, **the best interval representation of the Kleene algebras \mathbf{L} and $\mathbb{M}_n(K(\mathbf{L}))$** .

Notation: In order to simplify the notation, we use the same symbols for the operations of Łukasiewicz Kleene algebras whenever the context is clear; namely: $K(\mathbf{L}) = \langle [0, 1], \max, \odot, 0, 1, * \rangle$, its interval representation $K(\widehat{\mathbf{L}}) = (\mathbb{U}, \max, \odot, 0, 1, *)$, and the corresponding spaces of matrices: $\mathbb{M}_n(K(\mathbf{L})) = (M_n(K(\mathbf{L})), \max, \odot, 0, 1, *)$ and $\mathbb{M}_n(K(\widehat{\mathbf{L}})) = (M_n(K(\widehat{\mathbf{L}})), \max, \odot, 0, 1, *)$.

The next automata is an interval representation of (21)




Their interval matrices are:⁵

$$\mathcal{A}_\pi = \begin{bmatrix} (0, 0) & (0.4, 0.5) \\ (0, 0) & (0.7, 0.7) \end{bmatrix} \quad \mathcal{A}_{\pi'} = \begin{bmatrix} (0, 0) & (0.6, 0.8) \\ (0.7, 0.9) & (0.5, 0.5) \end{bmatrix}$$

The interpretation of the program $\mathcal{A}_{\pi+\pi'}$ is

$$\max \left(\begin{bmatrix} (0, 0) & (0.4, 0.5) \\ (0, 0) & (0.7, 0.7) \end{bmatrix}, \begin{bmatrix} (0, 0) & (0.6, 0.8) \\ (0.7, 0.9) & (0.5, 0.7) \end{bmatrix} \right) = \begin{bmatrix} (0, 0) & (0.6, 0.8) \\ (0.7, 0.9) & (0.7, 0.7) \end{bmatrix}$$

which represents the following weighted transition system:



Although we have defined operation “ \Rightarrow ” on intervals, the relation between them and action lattices will be exposed in the next section.

⁵ Here we use (a, b) instead of $[a, b]$ in order to provide a clear reading.

3.2 Pitfalls in defining action lattices over intervals

Before we proceed, it must be clear why do we use intervals. Intervals are used in a variety of situations when it is not possible to use exact values. If the exact values can be used, then it does not make sense to use intervals.

Although it is possible to use a near exact value to represent a desired point (e.g. 3.14 would be used to represent π), the information about impreciseness is not codified by such exact value. Intervals provide such kind of information and the quality of such representation can be measured by the width of the interval: the tighter is the interval the better is the representation.

Sometimes intervals are the only representation available to work with; e.g. (1) some magnetic resonance machines provide intervals for non-exact values (2) some applications in fuzzy systems provide intervals as inexact membership degree or as the abstraction of several membership degrees provided by different experts.

In any case, intervals are the entities provided instead of exact values. To deal with intervals a **price must be paid**; namely: not all properties of the space containing the exact values are preserved in the interval space. For example, in the case of real numbers, interval representations in general do not satisfy the property: $X - X = 0$.

As we will see the same happens with the interval representation of the action lattice \mathbb{L} . All properties stated in Figure 1 are satisfied by \mathbb{L} , whereas some are not by its interval representation $\hat{\mathbb{L}}$. Since these properties are connected with Dynamic Logics, there will be impacts of interval representation on the logical axioms. Some of these impacts are discussed below:

Observe that in the Łukasiewicz action lattice, \mathbb{L} , the equation “ $x \rightarrow x = 1$ ” is satisfied while this is not true in its interval representation: $\hat{\mathbb{L}}$. *But this is a crucial feature of $\hat{\mathbb{L}}$!* Take the following example: $[0.5, 0.6] \Rightarrow [0.5, 0.6] = [0.6 \rightarrow 0.5, 0.5 \rightarrow 0.6] = [0.9, 1] \neq [1, 1]$. Although $1 \in [0.9, 1]$, what is happening here? Suppose that the interval $[0.5, 0.6]$ is the tightest machine interval which represents the non-finitely representable exact values in \mathbb{L} : $\frac{\pi}{6} = 0.523598775\dots$ and $\frac{\frac{\pi}{6} + 0.6}{2} = 0.5617993875\dots$. Then, in order to calculate the implications: $\frac{\pi}{6} \rightarrow \frac{\pi}{6}$, $\frac{\frac{\pi}{6} + 0.6}{2} \rightarrow \frac{\frac{\pi}{6} + 0.6}{2}$, $\frac{\pi}{6} \rightarrow \frac{\frac{\pi}{6} + 0.6}{2}$ and $\frac{\frac{\pi}{6} + 0.6}{2} \rightarrow \frac{\pi}{6}$, the only way is to calculate: $[0.5, 0.6] \Rightarrow [0.5, 0.6]$. In this case, $\frac{\pi}{6} \rightarrow \frac{\pi}{6} = 1$, $\frac{\frac{\pi}{6} + 0.6}{2} \rightarrow \frac{\frac{\pi}{6} + 0.6}{2} = 1$, $\frac{\pi}{6} \rightarrow \frac{\frac{\pi}{6} + 0.6}{2} = 1$, $\frac{\frac{\pi}{6} + 0.6}{2} \rightarrow \frac{\pi}{6} = 0.9617993875\dots$, and $[0.5, 0.6] \Rightarrow [0.5, 0.6] = [0.6 \rightarrow 0.5, 0.5 \rightarrow 0.6] = [0.9, 1]$. Therefore, all the previous exact implications are contained in the “imprecise implication” “[0.5, 0.6] \Rightarrow [0.5, 0.6]”. In other words, unless an interval X has the form $[a, a]$, it does not make sense to impose $X \Rightarrow X = [1, 1]$, since the same interval can be used to represent two different exact values. Therefore, the known logical laws of Dynamic Logic must be reviewed or the notion of validity must be extended.

The price to be paid for using intervals does not stop here, in what follows we show that the structure $\hat{\mathbb{L}} = \langle \mathbb{U}, \max, \odot, 0, 1, *, \Rightarrow, \min \rangle$ is not an action lattice. This means that to propose a Dynamic Logic which deals with interval values some properties of action lattices must be *generalized*.

Proposition 3.

1. Equation (13) $A \odot X \leq B \Leftrightarrow X \leq A \Rightarrow B$ (Left-residuation) fails.
2. Equation (15) $(X \Rightarrow X)^* = X \Rightarrow X$ fails.
3. $X \Rightarrow X \leq (X \Rightarrow X)^*$.

Proof.

1. Make $X = [1, 1]$, then $A \odot X \leq B \Leftrightarrow X \leq A \Rightarrow B$ becomes $A \leq B \Leftrightarrow A \Rightarrow B = [1, 1]$ which is not true, since although $Y \leq Y$, $Y \Rightarrow Y$ is not always equal to $[1, 1]$; make $Y = [0.5, 0.6]$.
2. Make $X = [0.5, 0.6]$. By definition $(X \Rightarrow X)^* = [1, 1]$, but $X \Rightarrow X = [0.9, 1]$.
3. Since $\underline{X \Rightarrow X} = \underline{X} \rightarrow \underline{X} \leq 1$ and $\overline{X \Rightarrow X} = \overline{X} \rightarrow \overline{X} = 1$, then $X \Rightarrow X \leq [1, 1] = (X \Rightarrow X)^*$.

Proposition 4. Equations (16)-(20) are satisfied.

Proof. Since the structure $\langle \mathbb{U}, Max, Min \rangle$ is a lattice [3, 8, 11], then it satisfies Equations (16)-(20).

4 Quasi-action Lattices and Interval-valued Propositions

In this section we provide an algebraic structure which generalizes *action lattices* and is enough to model those lattices and their interval counterpart.

Definition 8. Let

$$\mathbf{A} = (U, +, ;, 0, 1, *, \rightarrow, \cdot)$$

be an algebra where U is a set, $0, 1 \in U$ are constants, $*$ is a unary operation on U and $+, ;, \rightarrow$ and \cdot are binary operations on U . Let $\Delta = \{x \in U : (x \rightarrow x)^* = x \rightarrow x\}$ be called the set of total elements of \mathbf{A} . \mathbf{A} is called **quasi-action lattice** whenever it satisfies the axioms: (1) – (12), (14), and (16) – (20) in figure 1 and the axioms:

$$a; x \leq b \Leftrightarrow x \leq a \rightarrow b; \text{ for every } a \in \Delta. \quad (13')$$

$$0, 1 \in \Delta \text{ and } x + y, x; y, x^*, x \rightarrow y, x \cdot y \in \Delta \text{ whenever } x, y \in \Delta. \quad (29)$$

$$a \rightarrow b = 1 \Leftrightarrow \exists z \in \Delta, a \leq z \leq b \quad (30)$$

Example 3. Every action lattice, \mathbf{A} , is a quasi-action lattice, since $U = \Delta$ and it satisfies (13'). Proposition 6 shows the main Quasi-action lattice studied here.

Proposition 5. $\langle \Delta, +, ;, 0, 1, *, \rightarrow, \cdot \rangle$ is an action lattice.

Proof. Equations (1) – (12), (14), and (16) – (20) are satisfied because \mathbf{A} is a quasi-action lattice. Equation (13) is guaranteed by equations (29) and (13). Finally, equation (15) is satisfied by definition of Δ .

In what follows we define a relation which captures the order property (OP) which is usually satisfied by some fuzzy implications.

Definition 9. Given a quasi-action lattice and $a, b \in U$, we define the relation “ \triangleleft ” as follows:

$$a \triangleleft b \Leftrightarrow a \rightarrow b = 1 \quad (23)$$

Lemma 2.

1. $a \triangleleft b \Rightarrow a \leq b$.
2. $a \triangleleft b$ and $b \triangleleft a \Rightarrow a = b$.
3. $a \triangleleft b$ and $b \triangleleft c \Rightarrow a \triangleleft c$.
4. $a \triangleleft a$ iff $a \in \Delta$;
5. $a \triangleleft b \leq c \Rightarrow a \triangleleft c$.
6. $a \leq b \triangleleft c \Rightarrow a \triangleleft c$.
7. $a \triangleleft b \Rightarrow \exists c, a \triangleleft c \triangleleft b$.
8. If $a \in \Delta$, then $\forall b \in U, a \triangleleft a + b$.

Proof.

- | | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ol style="list-style-type: none"> 1. Suppose $a \triangleleft b$, by definition there is $c \in \Delta$ s.t. $a \leq c \leq b$, therefore $a \leq b$. 2. Straightforward from 1. 3. Straightforward from 1. 4. (\Rightarrow) If $a \triangleleft a$, then by (23) $a \rightarrow a = 1$. So, $(a \rightarrow a)^* = 1^* = 1$, $(a \rightarrow a)^* = a \rightarrow a$ and $a \in \Delta$. (\Leftarrow) If $a \in \Delta$, since $a \leq a \leq a$, by definition, (30) and (23), $a \triangleleft a$. | <ol style="list-style-type: none"> 5. Trivial. 6. Trivial. 7. Trivial. 8. Since $a \leq a \leq a + b$, then $a \triangleleft a + b$. |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Lemma 3.

1. $a \rightarrow 1 = 1$.
2. $0 \rightarrow a = 1$.
3. $a \leq 1 \rightarrow a$.
4. $a; a \rightarrow b \leq b$; for $a \in \Delta$.
5. $1 \rightarrow a = a$.
6. $1 = a; b \Leftrightarrow a = b = 1$.
7. $(a \rightarrow b); (b \rightarrow a) = 1 \Rightarrow a = b$.
8. If $a \triangleleft b$ and $c \triangleleft d$, then $a + c \triangleleft b + d$.
9. If $a \triangleleft b$ and $c \triangleleft d$, then $a \cdot c \triangleleft b \cdot d$.

Proof.

1. Since $1 \in \Delta$ and $a \leq 1 \leq 1$, then by axiom (30), $a \rightarrow 1 = 1$.
2. Analogous.
3. $a \leq a$ iff $1; a \leq a$ iff, by axiom (13'), $a \leq 1 \rightarrow a$.
4. Straightforward from (13') and reflexivity of “ \leq ”.

5. Straightforward from Lemma 3.4⁶, for $a = 1$, together with Lemma 3.3.
6. (\Rightarrow) Since $K(\mathbf{A}) = (U, +, ;, 0, 1, *)$ is a Kleene algebra, the “;” is increasing monotonic in both components. Therefore, $1 \leq a; b \Rightarrow 1 \leq a; 1 \Rightarrow 1 \leq a \Rightarrow 1 = a$. The same for b . Therefore, $a = b = 1$. The converse, (\Leftarrow), is trivial.
7. $(a \rightarrow b; b \rightarrow a) = 1$. Then, by Lemma 3.6, $(a \rightarrow b) = 1$ and $(b \rightarrow a) = 1$. By axiom (23) $(a \leq b)$ and $(b \leq a)$. By Lemma 2.1, $a \leq b$ and $b \leq a$. Hence $a = b$.
8. If $a \leq b$ and $c \leq d$, then there are $z, z' \in \Delta$ such that $a \leq z \leq b$ and $c \leq z' \leq d$. Therefore, by [23, (33) p.7] $a + c \leq z + z' \leq b + d$. Since $z + z' \in \Delta$, then $a + c \leq b + d$.
9. Analogous applying [23, (35) p.7].

Lemma 4.

1. The set $\Delta = \{X \in \mathbb{U} : (X \Rightarrow X)^\circledast = X \Rightarrow X\}$ is equal to $\{X \in \mathbb{U} : \underline{X} = \overline{X}\}$.
2. For every $A \in \Delta$ and $X, B \in \mathbb{U}$, $A \odot X \leq B \Leftrightarrow X \leq A \Rightarrow B$.

Proof.

1. Given $X \in \Delta$, $[1, 1] \stackrel{def}{=} (X \Rightarrow X)^\circledast = X \Rightarrow X$. Therefore, $[\overline{X} \rightarrow \underline{X}, \underline{X} \rightarrow \overline{X}] = [1, 1]$. Hence, $\overline{X} \rightarrow \underline{X} = 1$ and $\underline{X} \rightarrow \overline{X} = 1$. By (13) $\overline{X} \leq \underline{X}$ and $\underline{X} \leq \overline{X}$ and so $\overline{X} = \underline{X}$. The converse is trivial.
2. Given $A \in \Delta$ and $X, B \in \mathbb{U}$, $A = [a, a]$, $A \odot X \leq B \Leftrightarrow [a \odot \underline{X}, a \odot \overline{X}] \leq [B, B] \Leftrightarrow a \odot \underline{X} \leq B$ and $a \odot \overline{X} \leq B \Leftrightarrow \underline{X} \leq a \rightarrow B$ and $\overline{X} \leq a \rightarrow B \Leftrightarrow X \leq A \Rightarrow B$.

Proposition 6. $\widehat{L} = \langle \mathbb{U}, \widehat{\max}, \widehat{\odot}, [0, 0], [1, 1], \widehat{\ast}, \widehat{\Rightarrow}, \widehat{\min} \rangle$ is a quasi-action lattice.

Proof. According to **Theorem 2**, \widehat{L} satisfies axioms (1) – (12). Equations (16) – (20) are trivially satisfied. Since L is an action lattice, for every intervals: $[\underline{a}, \overline{a}]$, $[\underline{b}, \overline{b}]$ and $[\underline{c}, \overline{c}]$, $\overline{a} \rightarrow \underline{b} \leq \overline{a} \rightarrow \max(\underline{b}, \underline{c})$ and $\underline{a} \rightarrow \overline{b} \leq \underline{a} \rightarrow \max(\overline{b}, \overline{c})$. Hence, $[\overline{a} \rightarrow \underline{b}, \underline{a} \rightarrow \overline{b}] \leq [\overline{a} \rightarrow \max(\underline{b}, \underline{c}), \underline{a} \rightarrow \max(\overline{b}, \overline{c})]$ and $[\underline{a}, \overline{a}] \Rightarrow [\underline{b}, \overline{b}] \leq [\underline{a}, \overline{a}] \Rightarrow \max([\underline{b}, \overline{b}], [\underline{c}, \overline{c}])$; satisfying (14). Lemma 4.2 guarantees (13’). (29) is guaranteed by Lemma 4.1. For (30) suppose $[\underline{a}, \overline{a}] \Rightarrow [\underline{b}, \overline{b}] = [1, 1]$, then $\overline{a} \rightarrow \underline{b} = 1$ iff $\overline{a} \leq \underline{b}$. Making $z = [\overline{a}, \overline{a}]$, we have $[\underline{a}, \overline{a}] \leq z \leq [\underline{b}, \overline{b}]$. If $\exists [z, z] \in \Delta$ s.t. $[\underline{a}, \overline{a}] \leq [z, z] \leq [\underline{b}, \overline{b}]$, then $\overline{a} \leq z \leq \underline{b}$. Therefore, $\overline{a} \rightarrow \underline{b} = 1$ and $[\underline{a}, \overline{a}] \Rightarrow [\underline{b}, \overline{b}] = [1, 1]$.

Given a Quasi-action lattice, \mathbf{A} , signatures for Dynamic Logic intended to be interpreted in \mathbf{A} (\mathbf{ADL}) are exactly the same as those of Propositional Dynamic Logic. They are pairs, $(\mathbf{\Pi}, \mathbf{Prop})$, of disjoint sets of **atomic programs**, $\mathbf{\Pi}$, and of **proposition symbols**, \mathbf{Prop} . Formulae of \mathbf{ADL} consist of the positive fragment of Propositional Dynamic Logic: The set of $\mathbf{\Pi}$ -programs, denoted by $\mathbf{Prg}(\mathbf{\Pi})$, consists of all expressions specified by the following BNF:

$$\pi ::= \pi_0 \mid \pi; \pi \mid \pi + \pi \mid \pi^*$$

⁶ Lemma 3.4 means the 4th item of Lemma 3.

for $\pi_0 \in \mathbf{\Pi}$. The \mathbf{ADL} -formulae for $(\mathbf{\Pi}, \mathbf{Prop})$, denoted by $Fm^{\mathbf{ADL}}(\mathbf{\Pi}, \mathbf{Prop})$, are those given by BNF:

$$\alpha ::= \top \mid \perp \mid p \mid \alpha \vee \alpha \mid \alpha \wedge \alpha \mid \alpha \rightarrow \alpha \mid \alpha \leftrightarrow \alpha \mid \langle \pi \rangle \alpha \mid [\pi] \alpha$$

for $p \in \mathbf{Prop}$ and $\pi \in \mathbf{Prg}(\mathbf{\Pi})$. \mathbf{ADL} -models, \mathfrak{M} , for a set of propositions in \mathbf{Prop} and programs in $\mathbf{\Pi}$, denoted by: $\mathfrak{M} \in \mathbf{Mod}^{\mathbf{ADL}}(\mathbf{\Pi}, \mathbf{Prop})$, are tuples:

$$\mathfrak{M} = (W, V, (\mathcal{A}_\pi)_{\pi \in \mathbf{\Pi}})$$

in which W is a finite set (of states), $V : \mathbf{Prop} \times W \rightarrow U$ is a function called *valuation* and \mathcal{A}_π is a U -valued binary relation indexed by program π .

Definition 10. *The graded satisfaction relation for a model $\mathfrak{M} \in \mathbf{Mod}^{\mathbf{ADL}}(\mathbf{\Pi}, \mathbf{Prop})$ is the function:*

$$\models_U : W \times Fm^{\mathbf{ADL}}(\mathbf{\Pi}, \mathbf{Prop}) \rightarrow U$$

recursively defined as follows:

- $(w \models_U \top) = 1$;
- $(w \models_U \perp) = 0$;
- $(w \models_U p) = V(p, w)$, for any $p \in \mathbf{Prop}$;
- $(w \models_U \rho \wedge \rho') = (w \models_U \rho) \cdot (w \models_U \rho')$;
- $(w \models_U \rho \vee \rho') = (w \models_U \rho) + (w \models_U \rho')$;
- $(w \models_U \rho \rightarrow \rho') = (w \models_U \rho) \rightarrow (w \models_U \rho')$;
- $(w \models_U \rho \leftrightarrow \rho') = [(w \models_U \rho) \rightarrow (w \models_U \rho')] ; [(w \models_U \rho') \rightarrow (w \models_U \rho)]$;
- $(w \models_U \langle \pi \rangle \rho) = \sum_{w' \in W} (\mathcal{A}_\pi(w, w') ; (w' \models_U \rho))$; ⁷
- $(w \models_U [\pi] \rho) = \prod_{w' \in W} (\mathcal{A}_\pi(w, w') \rightarrow (w' \models_U \rho))$.

Lemma 5. *Let \mathbf{A} be a quasi-action lattice, then:*

1. $(w \models_U \alpha \rightarrow \alpha') = 1$ iff $(w \models_U \alpha) \leq (w \models_U \alpha')$.
2. If $(w \models_U \alpha \leftrightarrow \alpha') = 1$, then $(w \models_U \alpha) = (w \models_U \alpha')$.


Proof.

1. $(w \models_U \alpha \rightarrow \alpha') = 1$ iff $[(w \models_U \alpha) \rightarrow (w \models_U \alpha')] = 1$ iff $(w \models_U \alpha) \leq (w \models_U \alpha')$.
2. Suppose that $(w \models_U \alpha \leftrightarrow \alpha') = 1$, then $[(w \models_U \alpha) \rightarrow (w \models_U \alpha')] ; [(w \models_U \alpha') \rightarrow (w \models_U \alpha)] = 1$. By Lemma 3.7, $(w \models_U \alpha) = (w \models_U \alpha')$.

⁷ Since “+” and “.” are associative, we can generalize them to n -ary operations and we resort to notation “ \sum ” for iterated version of “+” and to “ \prod ” for the iterated version of “.”.

Now, let us see the situation in the context of intervals. Observe how the notion of interval representation behaves as expected.


For example, considering the propositions: $\text{Prop} = \{p, q\}$, the transition system:



and the interpretation, $V : \text{Prop} \rightarrow [0, 1]$, again, as $V(p, s_1) = 0.1$, $V(q, s_1) = 0.5$, $V(p, s_2) = \frac{\pi}{4}$ and $V(q, s_2) = 0.75$, then

$$(s_1 \models_{\mathbb{L}} \langle \pi + \pi' \rangle (p \rightarrow q)) = \frac{\sqrt{2}}{2}$$

For the interval representation of (31):



Considering the interpretation on the quasi action lattice $\hat{\mathbb{L}}$, $\mathbf{Rep}(V) : \text{Prop} \rightarrow \mathbb{U}$ as: $\mathbf{Rep}(V)(p, s_1) = [0.1, 0.1]$, $\mathbf{Rep}(V)(q, s_1) = [0.5, 0.5]$, $\mathbf{Rep}(V)(p, s_2) = [0.7, 0.8]$ and $\mathbf{Rep}(V)(q, s_2) = [0.75, 0.75]$ we would have:

$$\begin{aligned} (s_1 \models_{\hat{\mathbb{L}}} \langle \pi + \pi' \rangle (p \rightarrow q)) &= \max([0, 0] \odot ([0.1, 0.1] \Rightarrow [0.5, 0.5]), [0.6, 0.8] \odot ([0.5, 0.5] \Rightarrow [0.7, 0.8])) \\ &= \max([0, 0], [0.6, 0.8] \odot [0.5 \rightarrow 0.7, 0.5 \rightarrow 0.8]) \\ &= [0.6, 0.8] \odot [1, 1] \\ &= [0.6, 0.8]. \end{aligned}$$

In other words, considering the valuation $V : \text{Prop} \times W \rightarrow [0, 1]$, on the set of exact values $[0, 1]$, and the valuation $\mathbf{Rep}(V) : \text{Prop} \times W \rightarrow \mathbb{U}$, on the set of intervals, it is expected that they relate in the following way:

- (a) $V(p, s_1) \in \mathbf{Rep}(V)(p, s_1)$;
- (b) $V(p, s_2) \in \mathbf{Rep}(V)(p, s_2)$;
- (c) $V(q, s_1) \in \mathbf{Rep}(V)(q, s_1)$;
- (d) $V(q, s_2) \in \mathbf{Rep}(V)(q, s_2)$;

i.e. the interval valuation $\mathbf{Rep}(V)$ represents the exact valuation V . Further, it is also expected that satisfiability relation defined on exact values relate to the satisfiability relation defined on intervals in the following way:

$$[s_1 \models_{\mathbb{L}} \langle \pi + \pi' \rangle (p \rightarrow q)] \in [s_1 \models_{\hat{\mathbb{L}}} \langle \pi + \pi' \rangle (p \rightarrow q)].$$

So, it is expected that if intervals are used to represent exact truth values then the satisfiability relation on intervals should be correct; namely it gives a truth value (an interval) which contain the exact truth-value of the satisfiability relation on exact values.

The next theorem guarantees the above situation in a general setting:

Theorem 3. *Let $\Theta = \{p_1, \dots, p_n\} \subseteq \text{Prop} \cup \{\perp, \top\}$; $w, w' \in W$; $A_\pi(w, w') \in \mathbb{M}_n(L)$ and $\mathbf{Rep}(A_\pi)(w, w') \in \mathbb{M}_n(K(\widehat{L}))$. For every formula $\alpha_\Theta \in \text{Fm}^{L\mathcal{DL}}$ built up recursively from Θ , $(w \models_L \alpha_\Theta) \in (w \models_{\widehat{L}} \alpha_\Theta)$, whenever $V(p_i, w) \in \mathbf{Rep}(V)(p_i, w)$ and $A_\pi(w, w') \in \mathbf{Rep}(A_\pi)(w, w')$.⁸*

Proof. The proof is by induction on the structure of α_Θ .

$$(B-1) \quad (w \models_L \top) = 1 \in [1, 1] = (w \models_{\widehat{L}} \top);$$

$$(B-2) \quad (w \models_L p_i) \stackrel{\text{def}}{=} V(p_i, w) \stackrel{\text{hyp}}{\in} \mathbf{Rep}(V)(p_i, w) \stackrel{\text{def}}{=} (w \models_{\widehat{L}} \top);$$

(I) Suppose $x = (w \models_L \rho) \in (w \models_{\widehat{L}} \rho) = [a, b]$ and $y = (w \models_L \rho') \in (w \models_{\widehat{L}} \rho') = [c, d]$, then:

$$(I-1) \quad (w \models_L \rho \wedge \rho') \stackrel{\text{def}}{=} \min(x, y) \in [\min(a, c), \min(b, d)] \stackrel{\text{def}}{=} \min([a, b], [c, d]) \stackrel{\text{def}}{=} \min(w \models_{\widehat{L}} \rho, w \models_{\widehat{L}} \rho') \stackrel{\text{def}}{=} w \models_{\widehat{L}} \rho \wedge \rho'.$$

$$(I-2) \quad \text{The proof is analogous for } (w \models_L \rho \vee \rho') \in (w \models_{\widehat{L}} \rho \vee \rho').$$

$$(I-3) \quad b \rightarrow c \leq x \rightarrow y \leq a \rightarrow d \Leftrightarrow x \rightarrow y \in [b \rightarrow c, a \rightarrow d] \stackrel{\text{def}}{\Leftrightarrow} x \rightarrow y \in [a, b] \Rightarrow [c, d] \Leftrightarrow (w \models_L \rho \rightarrow \rho') \in (w \models_{\widehat{L}} \rho \rightarrow \rho').$$

$$(I-4) \quad \text{By (I-3)} \quad (w \models_L \rho \rightarrow \rho') \in (w \models_{\widehat{L}} \rho \rightarrow \rho') \text{ and } (w \models_L \rho' \rightarrow \rho) \in (w \models_{\widehat{L}} \rho' \rightarrow \rho). \text{ By Proposition 1 } (w \models_L \rho \rightarrow \rho') \odot (w \models_L \rho' \rightarrow \rho) \in (w \models_{\widehat{L}} \rho \rightarrow \rho') \odot (w \models_{\widehat{L}} \rho' \rightarrow \rho).$$

$$(I-5) \quad \text{Let } w' \in W. \text{ Assuming that } (w' \models_{\widehat{L}} \rho) = [e, f], \mathbf{Rep}(A_\pi)(w, w') = [g, h], u = (w' \models_L \rho) \in [e, f]. \text{ By hypothesis, } v = A_\pi(w, w') \in [g, h]. \text{ By Proposition 1, } v \odot u \in [g, h] \odot [e, f]. \text{ Hence } (A_\pi(w, w') \odot (w' \models_L \rho)) \in (\mathbf{Rep}(A_\pi)(w, w') \odot (w' \models_{\widehat{L}} \rho)); \text{ for any } w, w' \in W. \text{ Therefore, } \max \{(A_\pi(w, w') \odot (w' \models_L \rho)) : w' \in W\} \in \max \{\mathbf{Rep}(A_\pi)(w, w') \odot (w' \models_{\widehat{L}} \rho) : w' \in W\}. \text{ By definition, } (w \models_L \langle \pi \rangle \rho) \in (w \models_{\widehat{L}} \langle \pi \rangle \rho).$$

$$(I-6) \quad \text{The proof is analogous for } (w \models_L [\pi]\rho) \in (w \models_{\widehat{L}} [\pi]\rho).$$

5 Satisfiability vs Validity

Although satisfiability is preserved by intervalization (Theorem 3) *validity is not*. In fact, this section shows that the *introduction of impreciseness on logical values affects validity* and as a consequence this concept must be weakened.

⁸ Observe that $\mathbf{Rep}(V)(p_i, w)$ and $\mathbf{Rep}(A_\pi)(w, w')$ are intervals.

Suppose we state, as usually, that valid formulæ are those which are always interpreted in quasi-action lattices as “1”. This is always true for the case in which $\Delta = U$. This is the case of the formula:

$$\langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho'. \quad ^9 \quad (32)$$

In general this is not true if Δ is a proper subset of U . The proof of such validity relies on the following property:

$$(x \leq y) \rightarrow (a; x \leq a; y), \quad (33)$$

see [23, Lemma 1 (36), p. 7]. Proposition 5 guarantees that this is true for $x, y, a \in \Delta$. In order to extend this property to the whole quasi-action lattice we should require:

$$(x < y) \rightarrow (a; x < a; y) \quad (34)$$

which is not always true.

In fact, this is not true for the interval model proposed here. Observe that in such quasi-action lattice the subset “ $U - \Delta$ ” is the set of non-degenerate intervals; and hence the set of imprecise values. Take for example: $[0.3, 0.5], [0.6, 0.8] \in U - \Delta$, in this case $[0.3, 0.5] < [0.6, 0.8]$, but $[0.4, 0.7] \odot [0.3, 0.5] = [0, 0.2] \not\leq [0, 0.5] = [0.4, 0.7] \odot [0.6, 0.8]$. This is due to the correctness of “ \odot ” which propagates the uncertainty on $[0.4, 0.7]$ and does not “respect” the relation “ $<$ ” — instead of “[0.4, 0.7]” try any degenerate interval; e.g. $[0.4, 0.4]$. From the abstract viewpoint, whenever Δ is a proper subset of U , the nature of objects $a \notin \Delta$ affects the relation “ $<$ ” via the Kleene operation “ $;$ ”.

Since the operation “ $;$ ” plays a fundamental role on the definition of modalities, we are obligated to generalize the notion of validity. *Instead of validity we propose **designated validity***; intuitively:

“A formula ‘ ρ ’ will be **designated valid** whenever for every quasi-action lattice \mathbf{A} , the value of ‘ $w \models_U \rho$ ’, for every $w \in W$, belongs to a ‘suitable’ proper subset $D \subset U$ such that $1 \in D$ and $0 \notin D$.”

In what follows we define the set of designated values for *any* quasi-action lattice followed by the notion of **designated model** and **designated validity**.

Definition 11. *Let \mathbf{A} be a quasi-action lattice. The **set of designated elements** of \mathbf{A} is the set:*

$$D = \{x \in U : \forall y \in U, x < y \text{ implies } y = 1\}. \quad (35)$$

*Given a quasi-action lattice, \mathbf{A} , the set of its designated elements, D , and a formula $\rho \in Fm^{\mathbf{A}\mathcal{D}\mathcal{L}}(\mathbf{II}, \mathbf{Prop})$, \mathbf{A} is a **designated model** for ρ or ρ is **designated valid in \mathbf{A}** , $\models_{\mathbf{A}}^d \rho$, if for every $w \in W, (w \models_U \rho) \in D$. ρ is **designated valid** if for every quasi-action lattice, \mathbf{A} , $\models_{\mathbf{A}}^d \rho$.*

⁹ Remember that $\Delta = U$ means that \mathbf{A} is an action lattice. The validity of (32) is proven in [23, Lemma 4.2].

The next lemma shows that the designated elements in \mathbb{U} are precisely those intervals whose maximum element is “1”; i.e. they contain the truth “1”, or in other words they are the truth “1” together with some imprecision.

Lemma 6. *The subset of designated elements of \mathbb{U} is precisely the set of intervals containing “1”; namely:*

$$D = \{[x, 1] \in \mathbb{U} : x \in [0, 1]\}. \quad (36)$$

Proof. According to Equation (35), the set of designated elements of \mathbb{U} is the set: $D = \{X \in \mathbb{U} : \forall Y \in \mathbb{U}, X \triangleleft Y \text{ implies } Y = [1, 1]\}$.

1. Given $X \in D$, suppose $\overline{X} < 1$, then $\underline{X} < \overline{X} < \frac{\overline{X}+1}{2} < 1$ and $X \Rightarrow [\frac{\overline{X}+1}{2}, \frac{\overline{X}+1}{2}] = [1, 1]$. Therefore, $X < [\frac{\overline{X}+1}{2}, \frac{\overline{X}+1}{2}]$. But, since $X \in D$ (by hypothesis) and $\frac{\overline{X}+1}{2} \neq [1, 1]$, we have a contradiction. Hence $X = [\underline{X}, 1]$.
2. Given $[x, 1] \in \mathbb{U}$ and $Y \in \mathbb{U}$, if $[x, 1] \triangleleft [\underline{Y}, \overline{Y}]$, then $[x, 1] \Rightarrow [\underline{Y}, \overline{Y}] = [1 \rightarrow \underline{Y}, x \rightarrow \overline{Y}] = [1, 1]$; i.e. $1 \rightarrow \underline{Y} = 1$ and $x \rightarrow \overline{Y} = 1$. Since $1 \rightarrow \underline{Y} = \underline{Y}$, then $\underline{Y} = 1$. Hence $[\underline{Y}, \overline{Y}] = [1, 1]$. Therefore, $[x, 1] \in D$.

This lemma is very significant. It points out what we have written before: *A designated value on intervals is the “absolute truth”, 1, together with some information of impreciseness*; i.e. they are logical values of the form: $[x, 1]$. The next theorem guarantees that $\widehat{\mathbb{L}}$ is a designated model for every formula of Theorem 1.

Theorem 4. *$\widehat{\mathbb{L}}$ is a designated model for every valid formula α in the logic $ED\mathcal{L}$.*

Proof. We prove for the second formula of Theorem 1:

Since “ $\langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho'$ ” is a valid formula with respect to action lattices, then for every $w \in W$,

$$(w \models_{\mathbb{L}} \langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho') = 1.$$

By Theorem 3,

$$(w \models_{\mathbb{L}} \langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho') \in (w \models_{\widehat{\mathbb{L}}} \langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho').$$

Therefore, $(w \models_{\widehat{\mathbb{L}}} \langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho')$ is an interval of the form $[x, 1]$. Hence $\models_{\widehat{\mathbb{L}}}^d \langle \pi \rangle (\rho \wedge \rho') \rightarrow \langle \pi \rangle \rho \wedge \langle \pi \rangle \rho'$.

Since this argument is general, it holds for every formula of Theorem 1.

6 Final Remarks

In this paper we have shown how the notion of validity on Łukasiewicz Dynamic Logic must be reviewed when the notion of impreciseness (through intervals) need to be taken into account for the specification/verification of computational systems. There are two ways to achieve that (Section 3.2): To review the laws of Dynamic Logics or extend the notion of validity. We chose the second way and proposed the notion of *designated validity* which is a weakening of validity. In the case of intervals, the result is: *Every law of Dynamic Logic is preserved under the notion of Designated Validity*. Therefore, the quasi-action lattice \hat{L} is a structure where there is some harmony between impreciseness and a kind of validity.

However, we could not demonstrate that all designated valid formulae with respect to \hat{L} is also valid with respect to L . If we prove this, it means that this specific Dynamic Logic is not affected when we deal with imprecision, what is required is to enlarge the concept of validity. This is a subject of a current investigation.

Acknowledgements. This paper is a result of the cooperation between UFRN (Brazil) and Aveiro University (Portugal) supported by Marie Curie Foundation. R. Santiago and B. Bedregal are supported by Marie Curie project PIRSES-GA-2012-318986 GetFun funded by EU-FP7 and by the *Brazilian National Council for Scientific and Technological Development* (CNPq, Portuguese: *Conselho Nacional de Desenvolvimento Científico e Tecnológico*) under the Projects 304597/2015-5 and 307681/2012-2. This work is also financed by the ERDF – European Regional Development Fund through the Operational Programme for Competitiveness and Internationalisation - COMPETE 2020 and by National Funds through the Portuguese funding agency, FCT - *Fundação para a Ciência e a Tecnologia* within project POCI-01-0145-FEDER- 016692. A. Madeira and M. Martins are also supported by the FCT BPD individual grant SFRH/BPD/103004/2014 and UID/MAT/04106/2019 at CIDMA, respectively.

References

1. BALTAG, A., AND SMETS, S. Quantum logic as a dynamic logic. *Synthese* 179, 2 (2011), 285–306.
2. BEDREGAL, B., AND SANTIAGO, R. Some continuity notions for interval functions and representation. *Computational and Applied Mathematics* 32, 3 (2013), 435–446.
3. BEDREGAL, B. R. C., BUSTINCE, H., PALMEIRA, E. S., DIMURO, G. P., AND FERNÁNDEZ, J. Generalized interval-valued OWA operators with interval weights derived from interval-valued overlap functions. *Int. J. Approx. Reasoning* 90 (2017), 1–16.
4. BEDREGAL, B. R. C., AND SANTIAGO, R. H. N. Interval representations, Łukasiewicz implicators and Smets–Magrez axioms. *Information Sciences* 221 (2013), 192 – 200.

5. BEDREGAL, B. R. C., AND TAKAHASHI, A. The best interval representations of t-norms and automorphisms. Fuzzy Sets and Systems 157, 24 (2006), 3220 – 3230.
6. BEDREGAL, B. R. C., AND TAKAHASHI, A. Interval valued versions of t-conorms, fuzzy negations and fuzzy implications. In 2006 IEEE International Conference on Fuzzy Systems (2006), pp. 1981–1987.
7. BOU, F., ESTEVA, F., GODO, L., AND RODRÍGUEZ, R. O. On the minimum many-valued modal logic over a finite residuated lattice. J. Log. Comput. 21, 5 (2011), 739–790.
8. BUSTINCE, H., BARRENECHEA, E., PAGOLA, M., FERNANDEZ, J., XU, Z., BEDREGAL, B., MONTERO, J., HAGRAS, H., HERRERA, F., AND DE BAETS, B. A historical account of types of fuzzy sets and their relationships. IEEE Transactions on Fuzzy Systems 24, 1 (Feb 2016), 179–194.
9. CIGNOLI, R., D’OTTAVIANO, I., AND MUNDICI, D. Algebraic Foundations of Many-Valued Reasoning. Trends in Logic. Springer Netherlands, 1999.
10. CONWAY, J. H. Regular Algebra and Finite Machines. Printed in GB by William Clowes & Sons Ltd, 1971.
11. CORNELIS, C., DESCHRIJVER, G., AND KERRE, E. E. Advances and challenges in interval-valued fuzzy logic. Fuzzy Sets and Systems 157, 5 (2006), 622–627.
12. DIMURO, G. P., BEDREGAL, B. R. C., SANTIAGO, R. H. N., AND REISER, R. H. S. Interval additive generators of interval t-norms and interval t-conorms. Inf. Sci. 181, 18 (2011), 3898–3916.
13. FITTING, M. Many-valued modal logics. Fundam. Inform. 15, 3-4 (1991), 235–254.
14. FITTING, M. Many-valued model logics II. Fundam. Inform. 17, 1-2 (1992), 55–73.
15. HAREL, D., KOZEN, D., AND TIURYN, J. Dynamic Logic. MIT Press, Cambridge, MA, USA, 2000.
16. HAREL, D., KOZEN, D., AND TIURYN, J. Dynamic Logic. MIT Press, 2000.
17. HICKEY, T., JU, Q., AND VAN EMDEN, M. H. Interval arithmetic: From principles to implementation. J. ACM 48, 5 (Sept. 2001), 1038–1068.
18. HUGHES, J., ESTERLINE, A. C., AND KIMIAGHALAM, B. Means-end relations and a measure of efficacy. Journal of Logic, Language and Information 15, 1-2 (2006), 83–108.
19. KOZEN, D. On action algebras. manuscript in: Logic and Flow of Information, Amsterdam, 1991.
20. KOZEN, D. A probabilistic PDL. J. Comput. Syst. Sci. 30, 2 (1985), 162–178.
21. KOZEN, D. A completeness theorem for Kleene algebras and the algebra of regular events. Inf. Comput. 110, 2 (1994), 366–390.
22. LIAU, C. Many-valued dynamic logic for qualitative decision theory. In New Directions in Rough Sets, Data Mining, and Granular-Soft Computing, 7th International Workshop, RSFDGrC ’99, Yamaguchi, Japan, November 9–11, 1999, Proceedings (1999), N. Zhong, A. Skowron, and S. Ohsuga, Eds., vol. 1711 of Lecture Notes in Computer Science, Springer, pp. 294–303.
23. MADEIRA, A., NEVES, R., AND MARTINS, M. A. An exercise on the generation of many-valued dynamic logics. J. Log. Algebr. Meth. Program. 85, 5 (2016), 1011–1037.
24. MADEIRA, A., NEVES, R., MARTINS, M. A., AND BARBOSA, L. S. A dynamic logic for every season. In Formal Methods: Foundations and Applications - 17th Brazilian Symposium, SBMF 2014, Maceió, AL, Brazil, September 29–October 1, 2014. Proceedings (2014), C. Braga and N. Martí-Oliet, Eds., vol. 8941 of Lecture Notes in Computer Science, Springer, pp. 130–145.

25. MOORE, R. E. Interval Arithmetic and Automatic Error Analysis in Digital Computing. Ph.D. dissertation, Department of Mathematics, Stanford University, Stanford, CA, USA, Nov. 1962. Also published as Applied Mathematics and Statistics Laboratories Technical Report No. 25.
26. MOORE, R. E., AND YANG, C. T. Interval analysis I. Technical Document LMSD-285875, Lockheed Missiles and Space Division, Sunnyvale, CA, USA, 1959.
27. MUNDICI, D. Advanced Łukasiewicz calculus and MV-algebras. Trends in Logic. Springer Netherlands, 2011.
28. PALMEIRA, E. S., AND BEDREGAL, B. R. C. Extension of fuzzy logic operators defined on bounded lattices via retractions. Computers & Mathematics with Applications 63, 6 (2012), 1026–1038.
29. PALMEIRA, E. S., BEDREGAL, B. R. C., FERNÁNDEZ, J., AND JURIO, A. On the extension of lattice-valued implications via retractions. Fuzzy Sets and Systems 240 (2014), 66–85.
30. PLATZER, A. Logical Analysis of Hybrid Systems - Proving Theorems for Complex Dynamics. Springer, 2010.
31. PRATT, V. R. Semantical considerations on floyd-hoare logic. In 17th Annual Symposium on Foundations of Computer Science, Houston, Texas, USA, 25-27 October 1976 (1976), IEEE Computer Society, pp. 109–121.
32. PRATT, V. R. Action logic and pure induction. In JELIA (1990), vol. 478 of Lecture Notes in Computer Science, Springer, pp. 97–120.
33. SANTIAGO, R. H. N., BEDREGAL, B. R. C., AND ACÍÓLY, B. M. Formal aspects of correctness and optimality of interval computations. Formal Aspects of Computing 18, 2 (2006), 231–243.
34. SANTIAGO, R. H. N., BEDREGAL, B. R. C., MADEIRA, A., AND MARTINS, M. A. On interval dynamic logic. In Formal Methods: Foundations and Applications - 19th Brazilian Symposium, SBMF 2016, Natal, Brazil, November 23-25, 2016, Proceedings (2016), L. Ribeiro and T. Lecomte, Eds., vol. 10090 of Lecture Notes in Computer Science, pp. 129–144.
35. SUNAGA, T. Theory of an interval algebra and its application to numerical analysis [reprint of res. assoc. appl. geom. mem. 2 (1958), 29–46]. Japan J. Indust. Appl. Math. 26, 2-3 (10 2009), 125–143.
36. XU, Y., RUAN, D., QIN, K., AND LIU, J. Lattice-Valued Logic: An Alternative Approach to Treat Fuzziness and Incomparability. Studies in Fuzziness and Soft Computing, Springer Berlin Heidelberg, 2012.