



Fractional differential equations and Volterra–Stieltjes integral equations of the second kind

Avyt Asanov¹ · Ricardo Almeida² · Agnieszka B. Malinowska³

Received: 28 December 2018 / Revised: 15 June 2019 / Accepted: 24 September 2019
© The Author(s) 2019

Abstract

In this paper, we construct a method to find approximate solutions to fractional differential equations involving fractional derivatives with respect to another function. The method is based on an equivalence relation between the fractional differential equation and the Volterra–Stieltjes integral equation of the second kind. The generalized midpoint rule is applied to solve numerically the integral equation and an estimation for the error is given. Results of numerical experiments demonstrate that satisfactory and reliable results could be obtained by the proposed method.

Keywords Fractional differential equation · Volterra–Stieltjes integral equation · Generalized midpoint rule

Mathematics Subject Classification 26A33 · 45D05 · 34K28 · 65R20

1 Introduction

Fractional differential equations are a generalization of ordinary differential equations, where integer-order derivatives are replaced by fractional derivatives. Over the last decade, these equations have attracted a lot of attention from researchers from different areas, since frac-

Communicated by José Tenreiro Machado.

✉ Agnieszka B. Malinowska
a.malinowska@pb.edu.pl

Avyt Asanov
avyt.asanov@manas.edu.kg

Ricardo Almeida
ricardo.almeida@ua.pt

¹ Department of Mathematics, Kyrgyz-Turkish Manas University, Customs of the Kyrgyz Republic, 720038 Bishkek, Kyrgyzstan

² Department of Mathematics, Center for Research and Development in Mathematics and Applications (CIDMA), University of Aveiro, 3810–193 Aveiro, Portugal

³ Faculty of Computer Science, Białystok University of Technology, 15–351 Białystok, Poland

tional derivatives provide an excellent tool for the description of memory and hereditary properties of various materials and processes (Atanackovic and Stankovic 2009; Carpinteri and Mainardi 2014; Demir et al. 2012; Kulish and Lage 2002; Meerschaert 2011; Podlubny 1999; Rezazadeh et al. 2018; Tariq et al. 2018; Vazquez 2005). Several types of fractional derivatives have been suggested to describe more accurately real-world phenomena, each one with their own advantages and disadvantages (Djida et al. 2017; Kilbas et al. 2006; Osman 2017; Osman et al. 2019; Podlubny 1999; Rezazadeh et al. 2019). A more general unifying perspective to the subject was proposed in Agrawal (2010), Klimek and Lupa (2013), Malinowska et al. (2015), by considering fractional operators depending on general kernels. In this work, we follow the special case of this approach that was developed in Almeida (2017a, b), Almeida et al. (2018, 2019), Garra et al. (2019), Kilbas et al. (2006), Yang and Machado (2017). Namely, we focus on nonlinear fractional differential equations involving a Caputo-type fractional derivative with respect to another function, called ψ -Caputo derivative. These types of equations have been successfully used to model the world population growth (Almeida 2017a, b) and gross domestic product (Almeida 2017b). On the other hand, it is well known that fractional differential equations often have to be solved numerically (Arqub and Maayah 2018; Arqub and Al-Smadi 2018a, b; Diethelm 2010; Ford and Connolly 2006; Lubich 1985; Morgado et al. 2013; Sousa and Oliveira 2019). Therefore, knowing the usefulness of the ψ -Caputo fractional derivative, an important issue is to discuss the numerical methods for differential equations with this derivative.

Motivated by the above discussion, in this paper, we provide a numerical scheme to solve fractional differential equations with the ψ -Caputo derivative. The main idea is to rewrite the considered equation as the Volterra–Stieltjes integral equation and then apply the generalized midpoint rule that was developed by Asanov et al. (2011b). Regarding integral equations, the best general reference is the handbook by Polyanin and Manzhirov (2008). Results on nonclassical Volterra integral equations of the first kind can be found in Apartsyn (2003). In Asanov (1998), problems of regularization, uniqueness and existence of solutions for Volterra integral and operator equations of the first kind are studied. Some properties of Volterra and Volterra–Stieltjes integral operators are given in Bukhgeim (1999), and Banas and Regan (2005). In Banas et al. (2000) and Federson and Bianconi (2001), quadratic integral equations of Urysohn–Stieltjes type and their applications are investigated. Various numerical solution methods for integral equations are presented in Asanov et al. (2011a, b), Asanov and Abdujabbarov (2011), Asanov et al. (2016), Delves and Walsh (1974), Federson et al. (2002). In particular, the generalized trapezoid rule and the generalized midpoint rule to evaluate the Stieltjes integral approximately by employing the notion of derivative of a function by means of a strictly increasing function (Asanov et al. 2011a, b; Asanov 2001), the generalized trapezoid rule for linear Volterra–Stieltjes integral equations of the second kind (Asanov et al. 2016), and the generalized midpoint rule for linear Fredholm–Stieltjes integral equations of the second kind (Asanov and Abdujabbarov 2011).

The paper is organized as follows. First, in Sect. 2, we review some necessary concepts and results on fractional calculus. In Sect. 3, we state an initial value problem with the ψ -Caputo fractional derivative of order $\alpha > 1$ and prove several results concerning this problem which will be needed in the forthcoming sections. Then, in Sect. 4, using the generalized midpoint rule, we exhibit a numerical procedure to solve the Volterra–Stieltjes integral equation that corresponds to the given fractional differential equation. An upper bound formula for the error in our approximation is derived. Results of numerical experiments are presented in Sect. 5. We verify the accuracy and analyse the stability of the numerical scheme. Section 6 provides conclusions and suggestions for future work and thus completes this work.

2 Preliminaries on fractional calculus

Let $\alpha > 0$ be a real number and $x : [a, b] \rightarrow \mathbb{R}$ a function. Given another function $\psi \in C^1[a, b]$ such that ψ is increasing and $\psi'(t) \neq 0$, for all $t \in [a, b]$, the ψ -Riemann–Liouville fractional integral of x , of order α , is defined as

$$I_{a+}^{\alpha, \psi} x(t) = \frac{1}{\Gamma(\alpha)} \int_a^t \psi'(\tau) (\psi(t) - \psi(\tau))^{\alpha-1} x(\tau) d\tau,$$

where $\Gamma(\cdot)$ is the Gamma function, i.e., $\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$, $z > 0$. The ψ -Riemann–Liouville fractional derivative of x , of order α , is defined as

$$D_{a+}^{\alpha, \psi} x(t) = \left(\frac{1}{\psi'(t)} \frac{d}{dt} \right)^n I_{a+}^{n-\alpha, \psi} x(t),$$

where $n = [\alpha] + 1$. The other important definition, which will be used in this work, is the ψ -Caputo fractional derivative of x of order α :

$${}^C D_{a+}^{\alpha, \psi} x(t) = D_{a+}^{\alpha, \psi} \left[x(t) - \sum_{k=0}^{n-1} \frac{x_\psi^{[k]}(a)}{k!} (\psi(t) - \psi(a))^k \right],$$

where

$$n = [\alpha] + 1 \quad \text{for } \alpha \notin \mathbb{N}, \quad n = \alpha \quad \text{otherwise,}$$

and

$$x_\psi^{[k]}(t) = \left(\frac{1}{\psi'(t)} \frac{d}{dt} \right)^k x(t).$$

We see at once that, if $\alpha = m \in \mathbb{N}$, then the ψ -Caputo fractional derivative coincides with the ordinary derivative

$${}^C D_{a+}^{m, \psi} x(t) = x_\psi^{[m]}(t),$$

while if $\alpha \in \mathbb{N}$ and $x \in C^n[a, b]$, then

$${}^C D_{a+}^{\alpha, \psi} x(t) = \frac{1}{\Gamma(n-\alpha)} \int_a^t \psi'(\tau) (\psi(t) - \psi(\tau))^{n-\alpha-1} x_\psi^{[n]}(\tau) d\tau.$$

Important relations between the two fractional operators are as follows (Almeida et al. 2018):

1. If $x \in C[a, b]$, then

$${}^C D_{a+}^{\alpha, \psi} I_{a+}^{\alpha, \psi} x(t) = x(t).$$

2. If $x \in C^{n-1}[a, b]$, then

$$I_{a+}^{\alpha, \psi} {}^C D_{a+}^{\alpha, \psi} x(t) = x(t) - \sum_{k=0}^{n-1} \frac{x_\psi^{[k]}(a)}{k!} (\psi(t) - \psi(a))^k.$$

3 Fractional differential equations

In this section, we consider the following nonlinear fractional differential equation with the ψ -Caputo derivative:

$${}^C D_{a+}^{\alpha, \psi} x(t) = F(t, x(t)) + f(t), \quad t \in [a, b], \quad (1)$$

subject to the initial conditions

$$x(a) = x_a, \quad x_{\psi}^{[k]}(a) = x_a^k, \quad k = 1, \dots, n-1, \quad (2)$$

where

1. $1 < \alpha \notin \mathbb{N}$ and $n = [\alpha] + 1$,
2. x_a and x_a^k , for $k = 1, \dots, n-1$, are fixed reals,
3. $F : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and $F(t, 0) = 0$, for all $t \in [a, b]$,
4. $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Regarding the existence and uniqueness of solutions for nonlinear fractional differential equation with the ψ -Caputo derivative of type (1), we refer the reader to Almeida et al. (2018). For stability results, we suggest the work of Almeida et al. (2019).

We shall prove some preparatory results providing a basis for the later development of a numerical method for solving the initial value problem (1) and (2). First, let us recall that problem (1) and (2) can be rewritten as the Volterra–Stieltjes integral equation, i.e., a Volterra-type integral equation involving the Riemann–Stieltjes integral.

Theorem 1 Almeida et al. (2018) *A function $x \in C^{n-1}[a, b]$ is a solution to problem (1) and (2) if and only if x satisfies the following Volterra–Stieltjes integral equation:*

$$x(t) = I_{a+}^{\alpha, \psi} F(t, x(t)) + g(t), \quad t \in [a, b], \quad (3)$$

where

$$\begin{aligned} I_{a+}^{\alpha, \psi} F(t, x(t)) &= \frac{1}{\Gamma(\alpha)} \int_a^t (\psi(t) - \psi(\tau))^{\alpha-1} F(\tau, x(\tau)) d\psi(\tau), \\ g(t) &= \sum_{k=0}^{n-1} \frac{x_a^k}{k!} (\psi(t) - \psi(a))^k + I_{a+}^{\alpha, \psi} f(t), \quad t \in [a, b]. \end{aligned} \quad (4)$$

In what follows we assume that:

(H1) Function F is Lipschitz with respect to the second variable, that is, there exists a positive constant L such that

$$|F(t, x_1) - F(t, x_2)| \leq L|x_1 - x_2|, \quad \forall t \in [a, b], \quad \forall x_1, x_2 \in \mathbb{R}.$$

(H2) There exist nonnegative constants L_1 and $\beta \in (0, 1]$ such that

$$|F(t_1, x) - F(t_2, x)| \leq L_1|t_1 - t_2|^\beta |x|, \quad \forall t_1, t_2 \in [a, b], \quad \forall x \in \mathbb{R}.$$

Lemma 1 Let $\alpha > 1$ and $s_1, s_2 \in [c, d]$, where $0 \leq c < d$. Then, $|s_1^\alpha - s_2^\alpha| \leq c_0|s_1 - s_2|$, where $c_0 = \alpha d^{\alpha-1}$.

Proof The proof follows from the mean value theorem. Let $\sigma(t) = t^\alpha$, $t \in [c, d]$. Then,

$$|s_1^\alpha - s_2^\alpha| \leq \sup_{t \in (c, d)} |\sigma'(t)| \cdot |s_1 - s_2| \leq \alpha d^{\alpha-1} |s_1 - s_2|.$$

□

For a given function $\varphi \in C[a, b]$, let $\|\cdot\|_C$ denote the usual norm:

$$\|\varphi\|_C = \sup_{t \in [a, b]} |\varphi(t)|.$$

Lemma 2 Let $\alpha > 1$, $f \in C[a, b]$, and $x \in C[a, b]$ be the solution to Eq. (3). Then, under assumption (H1) it holds that

$$\|x\|_C \leq L_0 \|g\|_C, \quad (5)$$

where

$$L_0 = \exp \left\{ \frac{L(b-a)}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|\psi'\|_C \right\},$$

and g is defined by (4).

Proof Since $F(t, 0) = 0$, it follows that

$$|F(\tau, x(\tau))| = |F(\tau, x(\tau)) - F(\tau, 0)| \leq L|x(\tau)|.$$

Hence, by (3), we obtain

$$|x(t)| \leq \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|\psi'\|_C \int_a^t |x(s)| ds + \|g\|_C, \quad t \in [a, b].$$

Applying the Grönwall inequality we get (5). \square

Lemma 3 Let $\alpha \in (1, 2)$, $s_1, s_2 \in [c, d]$, where $0 \leq c < d$, $p > 1$ such that $p(2 - \alpha) < 1$, and $\frac{1}{p} + \frac{1}{q} = 1$. Then,

$$|s_1^{\alpha-1} - s_2^{\alpha-1}| \leq c_1 |s_1 - s_2|^{\frac{1}{q}},$$

where

$$c_1 = (\alpha - 1) \left(\frac{1}{1 - p(2 - \alpha)} \right)^{\frac{1}{p}} \left(d^{1-p(2-\alpha)} - c^{1-p(2-\alpha)} \right)^{\frac{1}{p}}.$$

Proof Let $s_1, s_2 \in [c, d]$, with $s_2 > s_1$. Then, by the Hölder inequality,

$$\begin{aligned} s_2^{\alpha-1} - s_1^{\alpha-1} &= (\alpha - 1) \int_{s_1}^{s_2} \frac{1}{\tau^{2-\alpha}} d\tau \leq (\alpha - 1) \left(\int_{s_1}^{s_2} \tau^{-p(2-\alpha)} d\tau \right)^{\frac{1}{p}} \left(\int_{s_1}^{s_2} d\tau \right)^{\frac{1}{q}} \\ &= c_1 |s_2 - s_1|^{\frac{1}{q}}. \end{aligned}$$

\square

Theorem 2 Let $\alpha > 1$ and $x \in C[a, b]$ be the solution to Eq. (3). Then, under assumption (H1), it holds that

$$|x(t_2) - x(t_1)| \leq c_2 |t_2 - t_1|, \quad \forall t_1, t_2 \in [a, b],$$

where

$$\begin{aligned} c_2 &= \frac{\|\psi'\|_C}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} [L_0 L \|g\|_C + \|f\|_C] \\ &\quad + \sum_{k=1}^{n-1} \frac{|x_a^k|}{(k-1)!} (\psi(b) - \psi(a))^{k-1} \|\psi'\|_C, \end{aligned}$$

and L_0 as defined in Lemma 2.

Proof Let $\alpha > 1$ and $t_2 > t_1$, where $t_1, t_2 \in [a, b]$. Then, by Eqs. (3) and (4) and Lemma 2, we have

$$\begin{aligned} |x(t_2) - x(t_1)| &\leq \frac{L_0 L \|g\|_C}{\Gamma(\alpha)} \int_a^{t_1} [(\psi(t_2) - \psi(\tau))^{\alpha-1} - (\psi(t_1) - \psi(\tau))^{\alpha-1}] d\psi(\tau) \\ &\quad + \frac{L L_0 \|g\|_C}{\Gamma(\alpha)} \int_{t_1}^{t_2} (\psi(t_2) - \psi(\tau))^{\alpha-1} d\psi(\tau) \\ &\quad + \sum_{k=1}^{n-1} \frac{|x_a^k|}{k!} [(\psi(t_2) - \psi(a))^k - (\psi(t_1) - \psi(a))^k] \\ &\quad + \frac{\|f\|_C}{\Gamma(\alpha)} \int_a^{t_1} [(\psi(t_2) - \psi(\tau))^{\alpha-1} - (\psi(t_1) - \psi(\tau))^{\alpha-1}] d\psi(\tau) \\ &\quad + \frac{\|f\|_C}{\Gamma(\alpha)} \int_{t_1}^{t_2} (\psi(t_2) - \psi(\tau))^{\alpha-1} d\psi(\tau). \end{aligned}$$

Computing the integrals and applying the mean value theorem, we get

$$\begin{aligned} |x(t_2) - x(t_1)| &\leq \frac{L_0 L \|g\|_C}{\alpha \Gamma(\alpha)} [(\psi(t_2) - \psi(a))^\alpha - (\psi(t_1) - \psi(a))^\alpha - (\psi(t_2) - \psi(t_1))^\alpha \\ &\quad + (\psi(t_2) - \psi(t_1))^\alpha] + \frac{\|f\|_C}{\alpha \Gamma(\alpha)} [(\psi(t_2) - \psi(a))^\alpha \\ &\quad - (\psi(t_1) - \psi(a))^\alpha - (\psi(t_2) - \psi(t_1))^\alpha + (\psi(t_2) - \psi(t_1))^\alpha] \\ &\quad + \sum_{k=1}^{n-1} \frac{|x_a^k|}{(k-1)!} (\psi(b) - \psi(a))^{k-1} \|\psi'\|_C (t_2 - t_1). \end{aligned}$$

Finally, by Lemma 1 (with $d = \psi(b) - \psi(a)$) and the mean value theorem, we obtain

$$\begin{aligned} |x(t_2) - x(t_1)| &\leq \frac{L_0 L \|g\|_C}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|\psi'\|_C (t_2 - t_1) \\ &\quad + \frac{\|f\|_C}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|\psi'\|_C (t_2 - t_1) \\ &\quad + \sum_{k=1}^{n-1} \frac{|x_a^k|}{(k-1)!} (\psi(b) - \psi(a))^{k-1} \|\psi'\|_C (t_2 - t_1). \end{aligned}$$

□

Theorem 3 Let $x \in C[a, b]$ be the solution to Eq. (3), $G = \{(t, \tau) : a \leq \tau \leq t \leq b\}$ and

$$k(t, \tau) = \frac{1}{\Gamma(\alpha)} (\psi(t) - \psi(\tau))^{\alpha-1} F(\tau, x(\tau)). \quad (6)$$

Under assumptions (H1) and (H2) the following hold:

1. If $\alpha > 2$ and $\beta = 1$, then

$$|k(t, \tau_1) - k(t, \tau_2)| \leq c_3 |\tau_1 - \tau_2|, \quad (7)$$

for all $(t, \tau_1), (t, \tau_2) \in G$, where

$$c_3 = \frac{1}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-2} \\ \times \left[(L_0 L_1 \|g\|_C + Lc_2) (\psi(b) - \psi(a)) + (\alpha - 1) L L_0 \|g\|_C \|\psi'\|_C \right].$$

2. If $\alpha \in (1, 2)$, $p > 1$ with $p(2 - \alpha) < 1$, $\frac{1}{p} + \frac{1}{q} = 1$, and $\gamma = \min\{\beta, \frac{1}{q}\}$, then

$$|k(t, \tau_1) - k(t, \tau_2)| \leq c_4 |\tau_1 - \tau_2|^\gamma, \quad (8)$$

for all $(t, \tau_1), (t, \tau_2) \in G$, where

– for $\beta \geq \frac{1}{q}$

$$c_4 = \frac{1}{\Gamma(\alpha)} \left[L_0 L_1 (\psi(b) - \psi(a))^{\alpha-1} (b-a)^{\beta-\frac{1}{q}} \|g\|_C \right. \\ \left. + Lc_2 (\psi(b) - \psi(a))^{\alpha-1} (b-a)^{1-\frac{1}{q}} \right. \\ \left. + L L_0 (\alpha - 1) \left(\frac{1}{1 - p(2 - \alpha)} \right)^{\frac{1}{p}} (\psi(b) - \psi(a))^{\frac{1}{p}(1-p(2-\alpha))} \|g\|_C \|\psi'(t)\|_C \right];$$

– for $\beta < \frac{1}{q}$

$$c_4 = \frac{1}{\Gamma(\alpha)} \left[L_0 L_1 (\psi(b) - \psi(a))^{\alpha-1} \|g\|_C \right. \\ \left. + Lc_2 (\psi(b) - \psi(a))^{\alpha-1} (b-a)^{1-\beta} + L L_0 (\alpha - 1) \left(\frac{1}{1 - p(2 - \alpha)} \right)^{\frac{1}{p}} \right. \\ \left. \times (\psi(b) - \psi(a))^{\frac{1}{p}(1-p(2-\alpha))} \|g\|_C \|\psi'\|_C (b-a)^{\frac{1}{q}-\beta} \right].$$

Proof Observe that, for all $(t, \tau_1), (t, \tau_2) \in G$, we have

$$k(t, \tau_2) - k(t, \tau_1) = \frac{1}{\Gamma(\alpha)} \left[(\psi(t) - \psi(\tau_2))^{\alpha-1} - (\psi(t) - \psi(\tau_1))^{\alpha-1} \right] F(\tau_2, x(\tau_2)) \\ + \frac{1}{\Gamma(\alpha)} (\psi(t) - \psi(\tau_1))^{\alpha-1} [F(\tau_2, x(\tau_2)) - F(\tau_1, x(\tau_2))] \\ + \frac{1}{\Gamma(\alpha)} (\psi(t) - \psi(\tau_1))^{\alpha-1} [F(\tau_1, x(\tau_2)) - F(\tau_1, x(\tau_1))]. \quad (9)$$

Applying Lemmas 1 and 2, and Theorem 2, we can assert that

$$\left| \frac{1}{\Gamma(\alpha)} (\psi(t) - \psi(\tau_1))^{\alpha-1} [F(\tau_2, x(\tau_2)) - F(\tau_1, x(\tau_2))] \right| \\ \leq \frac{L_0 L_1}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|g\|_C |\tau_1 - \tau_2|^\beta, \quad (10)$$

$$\left| \frac{1}{\Gamma(\alpha)} (\psi(t) - \psi(\tau_1))^{\alpha-1} [F(\tau_1, x(\tau_2)) - F(\tau_1, x(\tau_1))] \right| \\ \leq \frac{Lc_2}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} |\tau_1 - \tau_2|, \quad (11)$$

and

$$\begin{aligned} & \left| \frac{1}{\Gamma(\alpha)} [(\psi(t) - \psi(\tau_2))^{\alpha-1} - (\psi(t) - \psi(\tau_1))^{\alpha-1}] F(\tau_2, x(\tau_2)) \right| \\ & \leq \frac{LL_0(\alpha-1)}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-2} \|\psi'\|_C \|g\|_C |\tau_1 - \tau_2|, \quad \alpha > 2. \end{aligned} \quad (12)$$

If $\alpha > 2$ and $\beta = 1$, then taking into account (10)–(12), from (9) we obtain (7). On the other hand, if $\alpha \in (1, 2)$, $p > 1$, $p(2 - \alpha) < 1$, $\frac{1}{p} + \frac{1}{q} = 1$, then by Lemmas 2 and 3, and Theorem 2 it follows that

$$\begin{aligned} & \left| \frac{1}{\Gamma(\alpha)} [(\psi(t) - \psi(\tau_1))^{\alpha-1} - (\psi(t) - \psi(\tau_2))^{\alpha-1}] F(\tau_2, x(\tau_2)) \right| \\ & \leq \frac{LL_0}{\Gamma(\alpha)} (\alpha-1) \left(\frac{1}{1 - p(2 - \alpha)} \right)^{\frac{1}{p}} (\psi(b) - \psi(a))^{\frac{1}{p}(1 - p(2 - \alpha))} \|g\|_C \|\psi'\|_C |\tau_1 - \tau_2|^{\frac{1}{q}}. \end{aligned} \quad (13)$$

In this case, taking into account (10)–(11) and (13), from (9) we obtain (8). \square

4 Numerical method and error analysis

The purpose of this section is to construct a method to find approximate solutions to fractional initial value problems of type (1) and (2). We use an equivalence relation between problem (1), (2) and integral equation (3). Then, the approximation routine is based on the generalized midpoint rule that was first developed by Asanov et al. (2011b). They proposed an approximation of the Stieltjes integral with the use of the notion of the derivative of a function with respect to the strictly increasing function (see Asanov 2001). The generalized midpoint rule summarizes the midpoint rule (Kalitkin 1978), and here we use this method to solve numerically the integral equation (3).

For $n \in \mathbb{N}$, let

$$h = \frac{b-a}{2n}, \quad t_k = a + kh, \quad k = 0, 1, 2, \dots, 2n.$$

We substitute $t = t_k, k = 0, 1, 2, \dots, 2n$ into integral equation (3) and examine the following system of equations:

$$\begin{cases} x(t_0) = g(t_0), \quad t_0 = a \\ x(t_1) = \int_a^{t_1} k(t_1, \tau) d\psi(\tau) + g(t_1) \\ x(t_{2i}) = \int_a^{t_{2i}} k(t_{2i}, \tau) d\psi(\tau) + g(t_{2i}), \quad i = 1, \dots, n \\ x(t_{2j+1}) = \int_a^{t_{2j+1}} k(t_{2j+1}, \tau) d\psi(\tau) + g(t_{2j+1}), \quad j = 1, \dots, n-1, \end{cases} \quad (14)$$

where $k(t, \tau)$ is defined by (6). Using the generalized midpoint rule for integrals in system (14) we get

$$\left\{ \begin{aligned} \int_a^{t_{2i}} k(t_{2i}, \tau) d\psi(\tau) &= \sum_{m=1}^i k(t_{2i}, t_{2m-1}) [\psi(t_{2m}) - \psi(t_{2m-2})] \\ &+ \sum_{m=1}^i R_m^{(2n)}(x, i), \quad i = 1, \dots, n \\ \int_a^{t_1} k(t_1, \tau) d\psi(\tau) &= k(t_1, t_0) [\psi(t_1) - \psi(t_0)] + R_0^{(0)}(x) \\ \int_a^{t_{2j+1}} k(t_{2j+1}, \tau) d\psi(\tau) &= \int_a^{t_1} k(t_{2j+1}, s) d\psi(s) + \int_{t_1}^{t_{2j+1}} k(t_{2j+1}, \tau) d\psi(\tau) \\ &= k(t_{2j+1}, t_0) [\psi(t_1) - \psi(t_0)] + R_j^{(0)}(x) + \sum_{m=1}^j k(t_{2j+1}, t_{2m}) \\ &\times [\psi(t_{2m+1}) - \psi(t_{2m-1})] + \sum_{m=1}^j R_m^{(2n-1)}(x, j), \quad j = 1, \dots, n-1, \end{aligned} \right. \quad (15)$$

where

$$\left\{ \begin{aligned} R_0^{(0)}(x) &= \int_a^{t_1} [k(t_1, \tau) - k(t_1, t_0)] d\psi(\tau) \\ R_j^{(0)}(x) &= \int_a^{t_{2j+1}} [k(t_{2j+1}, \tau) - k(t_{2j+1}, t_0)] d\psi(\tau), \quad j = 1, \dots, n-1 \\ R_m^{(2n)}(x, i) &= \int_{t_{2m-2}}^{t_{2m}} [k(t_{2i}, \tau) - k(t_{2i}, t_{2m-1})] d\psi(\tau), \quad m = 1, \dots, i \\ R_m^{(2n-1)}(x, j) &= \int_{t_{2m-1}}^{t_{2m+1}} [k(t_{2j+1}, \tau) - k(t_{2j+1}, t_{2m})] d\psi(\tau), \quad m = 1, \dots, j. \end{aligned} \right. \quad (16)$$

If $\alpha > 2$ and $\beta = 1$, then on the strength of Theorem 3 for $R_m^{(2n)}(x, i)$ and $R_m^{(2n-1)}(x, j)$ we obtain the following estimates:

$$\left\{ \begin{aligned} |R_m^{(2n)}(x, i)| &\leq c_3 h [\psi(t_{2m}) - \psi(t_{2m-2})], \quad m = 1, \dots, i \\ |R_m^{(2n-1)}(x, j)| &\leq c_3 h [\psi(t_{2m+1}) - \psi(t_{2m-1})], \quad m = 1, \dots, j, \end{aligned} \right. \quad (17)$$

where $i = 1, 2, \dots, n, j = 1, 2, \dots, n-1$ (cf. Corollary 2 in Asanov et al. 2011b). Then taking into account (16) and (17), we have

$$\left\{ \begin{aligned} \left| \sum_{m=1}^i R_m^{(2n)}(x, i) \right| &\leq c_3 h [\psi(t_{2i}) - \psi(a)], \quad i = 1, \dots, n \\ |R_0^{(0)}(x)| &\leq c_3 h [\psi(t_1) - \psi(a)] \\ |R_j^{(0)}(x)| &\leq c_3 h [\psi(t_1) - \psi(a)] \\ \left| \sum_{m=1}^j R_m^{(2n-1)}(x, j) \right| &\leq c_3 h [\psi(t_{2j+1}) - \psi(t_1)], \quad j = 1, \dots, n-1. \end{aligned} \right. \quad (18)$$

If $\alpha \in (1, 2)$, $p > 1$ with $p(2 - \alpha) < 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $\gamma = \min\{\beta, \frac{1}{q}\}$, then on the strength of Theorem 3, for $R_m^{(2n)}(x, i)$ and $R_m^{(2n-1)}(x, j)$, we obtain the following estimates:

$$\left\{ \begin{aligned} |R_m^{(2n)}(x, i)| &\leq c_4 h^\gamma [\psi(t_{2m}) - \psi(t_{2m-2})], \quad m = 1, \dots, i \\ |R_m^{(2n-1)}(x, j)| &\leq c_4 h^\gamma [\psi(t_{2m+1}) - \psi(t_{2m-1})], \quad m = 1, \dots, j, \end{aligned} \right. \quad (19)$$

for $i = 1, 2, \dots, n, j = 1, 2, \dots, n - 1$ (cf. Corollary 2 in Asanov et al. 2011b). Then, taking into account (16) and (19), we have

$$\left\{ \begin{array}{l} \left| \sum_{m=1}^i R_m^{(2n)}(x, i) \right| \leq c_4 h^\gamma [\psi(t_{2i}) - \psi(a)], \quad i = 1, \dots, n \\ |R_0^{(0)}(x)| \leq c_4 h^\gamma [\psi(t_1) - \psi(a)] \\ |R_j^{(0)}(x)| \leq c_4 h^\gamma [\psi(t_1) - \psi(a)] \\ \left| \sum_{m=1}^j R_m^{(2n-1)}(x, j) \right| \leq c_4 h^\gamma [\psi(t_{2j+1}) - \psi(t_1)], \quad j = 1, \dots, n - 1. \end{array} \right. \quad (20)$$

Combining relations (6) and (15) we can rewrite (14) as

$$\left\{ \begin{array}{l} x(t_0) = g(t_0), \quad t_0 = a \\ x(t_1) = \frac{1}{\Gamma(\alpha)} (\psi(t_1) - \psi(t_0))^{\alpha-1} F(t_0, x(t_0)) [\psi(t_1) - \psi(t_0)] + g(t_1) + R_0^{(0)}(x) \\ x(t_{2i}) = \frac{1}{\Gamma(\alpha)} \sum_{m=1}^i (\psi(t_{2i}) - \psi(t_{2m-1}))^{\alpha-1} F(t_{2m-1}, x(t_{2m-1})) \\ \quad \times [\psi(t_{2m}) - \psi(t_{2m-2})] + g(t_{2i}) + \sum_{m=1}^i R_m^{(2n)}(x, i), \quad i = 1, \dots, n \\ x(t_{2j+1}) = \frac{1}{\Gamma(\alpha)} (\psi(t_{2j+1}) - \psi(t_0))^{\alpha-1} F(t_0, x(t_0)) (\psi(t_1) - \psi(t_0)) \\ \quad + \frac{1}{\Gamma(\alpha)} \sum_{m=1}^j (\psi(t_{2j+1}) - \psi(t_{2m}))^{\alpha-1} F(t_{2m}, x(t_{2m})) [\psi(t_{2m+1}) - \psi(t_{2m-1})] \\ \quad + g(t_{2j+1}) + R_j^{(0)}(x) + \sum_{m=1}^j R_m^{(2n-1)}(x, j), \quad j = 1, \dots, n - 1. \end{array} \right. \quad (21)$$

Omitting

$$\sum_{m=1}^i R_m^{(2n)}(x, i), \quad R_0^{(0)}(x), \quad R_j^{(0)}(x), \quad \sum_{m=1}^j R_m^{(2n-1)}(x)$$

in equations of system (21) and writing the sought solution x at the nodes t_k , we get the system of equations in terms of x_k :

$$\left\{ \begin{array}{l} x_0 = g(t_0), \quad t_0 = a \\ x_1 = \frac{1}{\Gamma(\alpha)} (\psi(t_1) - \psi(t_0))^{\alpha-1} F(t_0, x_0) [\psi(t_1) - \psi(t_0)] + g(t_1) \\ x_{2i} = \frac{1}{\Gamma(\alpha)} \sum_{m=1}^i (\psi(t_{2i}) - \psi(t_{2m-1}))^{\alpha-1} F(t_{2m-1}, x_{2m-1}) \\ \quad \times [\psi(t_{2m}) - \psi(t_{2m-2})] + g(t_{2i}), \quad i = 1, \dots, n \\ x_{2j+1} = \frac{1}{\Gamma(\alpha)} (\psi(t_{2j+1}) - \psi(t_0))^{\alpha-1} F(t_0, x_0) (\psi(t_1) - \psi(t_0)) \\ \quad + \frac{1}{\Gamma(\alpha)} \sum_{m=1}^j (\psi(t_{2j+1}) - \psi(t_{2m}))^{\alpha-1} F(t_{2m}, x_{2m}) [\psi(t_{2m+1}) - \psi(t_{2m-1})] \\ \quad + g(t_{2j+1}), \quad j = 1, \dots, n - 1, \end{array} \right. \quad (22)$$

where $x_0 = x(a)$ and $x_i \approx x(t_i)$, for $i = 1, 2, \dots, 2n$.

Now, we make the truncation error and convergent order analysis for the proposed numerical scheme. Let x be the solution to integral equation (3). As before, an approximation for $x(t_k)$ at a node t_k is denoted by x_k .

Theorem 4 Let $\alpha > 2$ and $\beta = 1$. Under assumptions (H1) and (H2) it holds that

$$|x(t_k) - x_k| \leq c_6 h, \quad k = 0, 1, \dots, 2n, \quad (23)$$

as $n \rightarrow \infty$, where

$$c_6 = c_3[\psi(b) - \psi(a)]e^{c_5(b-a)}, \quad c_5 = \frac{L}{\Gamma(\alpha)}(\psi(b) - \psi(a))^\alpha.$$

Proof Let the error be denoted by

$$z_k = x(t_k) - x_k, \quad k = 0, 1, \dots, 2n.$$

Taking into account (21) and (22), we have

$$\left\{ \begin{array}{l} z_0 = 0 \\ z_1 = R_0^{(0)}(x) \\ z_{2i} = \frac{1}{\Gamma(\alpha)} \sum_{m=1}^i (\psi(t_{2i}) - \psi(t_{2m-1}))^{\alpha-1} [F(t_{2m-1}, x(t_{2m-1})) \\ \quad - F(t_{2m-1}, x_{2m-1})](\psi(t_{2m}) - \psi(t_{2m-2})) + \sum_{m=1}^i R_m^{(2n)}(x, i), \quad i = 1, \dots, n \\ z_{2j+1} = \frac{1}{\Gamma(\alpha)} \sum_{m=1}^j (\psi(t_{2j+1}) - \psi(t_{2m}))^{\alpha-1} [F(t_{2m}, x(t_{2m})) \\ \quad - F(t_{2m}, x_{2m})](\psi(t_{2m+1}) - \psi(t_{2m-1})) \\ \quad + R_j^{(0)}(x) + \sum_{m=1}^j R_m^{(2n-1)}(x, j), \quad j = 1, \dots, n-1. \end{array} \right. \quad (24)$$

From this, by (18), we conclude that

$$\left\{ \begin{array}{l} |z_1| \leq R(h) \\ |z_k| \leq R(h) + c_5 h \sum_{j=1}^{k-1} |z_j|, \quad k = 2, \dots, 2n, \end{array} \right. \quad (25)$$

where

$$R(h) = c_3[\psi(b) - \psi(a)]h. \quad (26)$$

Now, let us consider the system

$$\epsilon_k = R(h) + c_5 h \sum_{j=1}^{k-1} |\epsilon_j|, \quad k = 2, \dots, 2n, \quad (27)$$

and $\epsilon_1 = R(h)$ as an initial condition. It is easily seen that $|z_k| \leq \epsilon_k$ for $k = 1, \dots, 2n$. Indeed, this can be verified by mathematical induction as follows: for $k = 1$ it is obvious. Let $|z_j| \leq \epsilon_j$ for $j = 1, \dots, k-1$. Then, by inequality (25), we get

$$|z_k| \leq R(h) + c_5 h \sum_{j=1}^{k-1} \epsilon_j = \epsilon_k.$$

Observe that $\epsilon_j, j = 1, \dots, 2n$, given by

$$\epsilon_j = R(h)[1 + c_5h]^{j-1},$$

satisfy system (27). In fact, using the inequality

$$(1 + \gamma)^{k-1} - 1 = \gamma \sum_{j=1}^{k-1} (1 + \gamma)^{j-1}, \quad k \geq 2,$$

and taking $\gamma = c_5h$ we obtain

$$\begin{aligned} R(h) + c_5h \sum_{j=1}^{k-1} \epsilon_j &= R(h) \left\{ 1 + c_5h \sum_{j=1}^{k-1} (1 + c_5h)^{j-1} \right\} \\ &= R(h) \{1 + [(1 + c_5h)^{k-1} - 1]\} = \epsilon_k, \quad k \geq 2. \end{aligned}$$

Consequently, we get the following estimates:

$$|z_k| \leq R(h)(1 + c_5h)^{k-1}, \quad k = 1, \dots, 2n. \quad (28)$$

Using the fact that $(1 + t)^{\frac{1}{t}}$ is increasing and approaches the number e as $t \rightarrow 0^+$ we get

$$(1 + c_5h)^{k-1} \leq (1 + c_5h)^{\frac{b-a}{h}} = [(1 + c_5h)^{\frac{1}{c_5h}}]^{c_5(b-a)} \leq e^{c_5(b-a)}$$

for $k \leq \frac{b-a}{h}$. This together with (26) and (28) gives (23). \square

Theorem 5 Let $\alpha \in (1, 2)$, $p > 1$ with $p(2 - \alpha) < 1$,

$$\frac{1}{p} + \frac{1}{q} = 1, \gamma = \min \left\{ \beta, \frac{1}{q} \right\}, \quad \gamma_0 = \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^\alpha < 1.$$

Under assumptions (H1) and (H2) it holds that

$$|x(t_k) - x_k| \leq c_7 h^\gamma, \quad k = 0, 1, \dots, 2n, \quad (29)$$

as $n \rightarrow \infty$, where

$$c_7 = \frac{c_4}{1 - \gamma_0} [\psi(b) - \psi(a)].$$

Proof Using the conditions of Theorem 5 and the estimates (20), from (24) we obtain following inequalities for z_k :

$$\left\{ \begin{aligned} |z_1| &\leq c_4 h^\gamma [\psi(b) - \psi(a)] \\ |z_{2i}| &\leq \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \sup_{k=1, \dots, 2i-1} |z_k| \sum_{m=1}^i [\psi(t_{2m}) - \psi(t_{2m-2})] \\ &\quad + c_4 h^\gamma [\psi(t_{2i}) - \psi(a)], \quad i = 1, \dots, n \\ |z_{2j+1}| &\leq \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \sup_{k=2, \dots, 2j} |z_k| \sum_{m=1}^j [\psi(t_{2m+1}) - \psi(t_{2m-1})] \\ &\quad + c_4 h^\gamma [\psi(t_{2j+1}) - \psi(t_1)], \quad j = 1, \dots, n-1. \end{aligned} \right. \quad (30)$$

Set $\|z\| = \sup_{k=0, \dots, 2n} |z_k|$. Then, by (30), we have

$$|z_{2i}| \leq \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|z\| [\psi(t_{2i}) - \psi(a)] + c_4 h^\gamma [\psi(t_{2i}) - \psi(a)], \quad i = 1, \dots, n,$$

and

$$|z_{2j+1}| \leq \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^{\alpha-1} \|z\| [\psi(t_{2j+1}) - \psi(a)] \\ + c_4 h^\gamma [\psi(t_{2j+1}) - \psi(t_1)], \quad j = 1, \dots, n-1.$$

Hence

$$\|z\| \leq \frac{L}{\Gamma(\alpha)} (\psi(b) - \psi(a))^\alpha \|z\| + c_4 h^\gamma [\psi(b) - \psi(a)],$$

what finishes the proof. \square

5 Numerical examples

In the following examples, to show the efficiency of the proposed numerical method, we approximate the solutions for some fractional differential equations of order $\alpha > 1$. In all examples, corresponding systems of algebraic equations of type (22) are solved using the command *fsolve* in Maple. In tables, we present the absolute error

$$E = \max_{i=0,1,\dots,2n} |x_i - x(t_i)|,$$

and the elapsed CPU time in seconds of the proposed numerical scheme (22).

Example 1 Consider the following fractional differential equation:

$${}^C D_{0+}^{5/2, \psi} x(t) = \frac{x^3(t)}{1+x^2(t)} - \frac{1}{2}, \quad t \in [0, 1], \quad (31)$$

with the initial conditions and kernel

$$x_0 = 1, \quad x_0^1 = x_0^2 = 0 \quad \text{and} \quad \psi(t) = t + t\sqrt{t}, \quad (32)$$

respectively. Therefore, in integral equation (3) we have $\alpha = \frac{5}{2}$ and

$$F(t, x) = \frac{x^3}{1+x^2}, \quad g(t) = 1 - \frac{1}{2\Gamma(\frac{7}{2})} \left(t + t\sqrt{t} \right)^{\frac{5}{2}}.$$

In this case, conditions (H1) and (H2) are satisfied for $L = 9/8$, $L_1 = 0$, and $\beta = 1$. Note that, the exact solution to (31) and (32) is $x(t) = 1$, $t \in [0, 1]$. The exact and numerical solutions for $n = 20, 40, 60$, as well as the evolution of the error, are shown in Fig. 1. The maximum of absolute error for different values of n and the elapsed CPU time in seconds are displayed in Table 1. As expected, as n increases, the error decreases.

Example 2 Consider the following fractional differential equation:

$${}^C D_{0+}^{3/2, \psi} x(t) = \frac{x^5(t)}{(1+x^2(t))^2} - \frac{243}{100}, \quad t \in [0, 1], \quad (33)$$

with the initial conditions and kernel

$$x_0 = 3, \quad x_0^1 = 0 \quad \text{and} \quad \psi(t) = 2t + t^{\frac{4}{3}}, \quad (34)$$

respectively. Therefore, in integral equation (3) we have $\alpha = \frac{3}{2}$ and

$$F(t, x) = \frac{x^5}{(1+x^2)^2}, \quad g(t) = 3 - \frac{243}{100\Gamma(\frac{5}{2})} \left(2t + t^{\frac{4}{3}} \right)^{\frac{3}{2}}.$$




Fig. 1 Comparison of the exact and numerical solutions (left), and errors (right) in Example 1

Table 1 Maximum of the absolute error and elapsed CPU time in seconds for the numerical scheme (Example 1)

n	20	40	60
E	0.000419022	0.000101614	0.000044441
Time	0.375	3.046	10.000




Fig. 2 Comparison of the exact and numerical solutions (left), and errors (right) in Example 2

In this case, conditions (H1) and (H2) are met for $L = 125/108$, $L_1 = 0$, and $\beta = 1$. Note that the exact solution to (33) and (34) is $x(t) = 3$, $t \in [0, 1]$. The exact and numerical solutions for $n = 20, 40, 60$, as well as the evolution of the error, are shown in Fig. 2. The maximum of absolute error for different values of n and the elapsed CPU time in seconds are displayed in Table 2.

Example 3 Consider the following fractional differential equation:

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5],$$

Table 2 Maximum of the absolute error and elapsed CPU time in seconds for the numerical scheme (Example 2)

n	20	40	60
E	0.172337940	0.059299298	0.031882101
Time	0.453	3.515	11.718




Fig. 3 Comparison of the exact and numerical solutions (left), and errors (right) in Example 3

Table 3 Maximum of the absolute error and elapsed CPU time in seconds for the numerical scheme (Example 3)

n	20	40	60
E	0.002572167258	0.000904114	0.000490134258
Time	0.046	0.203	0.515

with the initial conditions and kernel

$$x_0 = 0, \quad x_0^1 = 0, \quad \psi(t) = \sqrt{t+1},$$

respectively. Therefore, in integral equation (3) we have $\alpha = \frac{3}{2}$ and

$$F(t, x) = x, \quad g(t) = \frac{(\sqrt{t+1} - 1)^{\frac{3}{2}}}{\Gamma\left(\frac{5}{2}\right)}.$$

Once ${}^C D_{0+}^{3/2, \psi} E_{3/2}(\psi(t) - \psi(0))^{3/2} = (\psi(t) - \psi(0))^{3/2}$, where $E_{3/2}(\cdot)$ denotes the Mittag-Leffler function of order $\alpha = 3/2$, we conclude that the solution to this problem is $x(t) = E_{3/2}(\psi(t) - \psi(0))^{3/2} - 1$, $t \in [0, 5]$. The exact and numerical solutions for $n = 20, 40, 60$, as well as the evolution of the error, are shown in Fig. 3. The maximum of absolute error for different values of n and the elapsed CPU time in seconds are displayed in Table 3.

In the next examples, we analyse the stability of the proposed numerical method.

Example 4 Consider the two fractional differential equations with initial conditions:

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = 0, \quad x_0^1 = 0 \quad (35)$$




Fig. 4 Comparison of functions X and E , for $\mu = 0.1$ (left) and $\mu = 0.2$ (right) in Example 4

Table 4 Approximate values of $\bar{x}(5)$, where \bar{x} is the solution to (36), for different values of μ and n in Example 4

$\bar{x}(5)$	$n = 20$	$n = 40$	$n = 60$
$\mu = 0.1$	2.232897331	2.230742939	2.230209768
$\mu = 0.2$	2.527240325	2.524599595	2.523947221
$\mu = 0.3$	2.821583318	2.818456250	2.817684672

and

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = \mu, \quad x_0^1 = 0. \quad (36)$$

The kernel is given by $\psi(t) = \sqrt{t+1}$ and μ is a real number. Let x and \bar{x} be solutions of (35) and (36), respectively. In this case, $F(t, x) = x$ for both equations and function g is given by

$$g(t) = \frac{(\sqrt{t+1} - 1)^{\frac{3}{2}}}{\Gamma\left(\frac{5}{2}\right)} \quad \text{and} \quad g(t) = \mu + \frac{(\sqrt{t+1} - 1)^{\frac{3}{2}}}{\Gamma\left(\frac{5}{2}\right)},$$

respectively. By the Grönwall inequality (cf. Almeida et al. 2019; Sousa and Oliveira 2019), we conclude that

$$|x(t) - \bar{x}(t)| \leq |\mu| E_{3/2} \left((\sqrt{t+1} - 1)^{3/2} \right), \quad t \in [0, 5].$$

Consider the two functions

$$X : t \mapsto |x(t) - \bar{x}(t)| \quad \text{and} \quad E : t \mapsto |\mu| E_{3/2} \left((\sqrt{t+1} - 1)^{3/2} \right).$$

In Fig. 4, we present the plots for two values of μ . It can be observed that the proposed numerical scheme preserves the underlying structural stability of the initial value problem, with respect to small perturbation of the initial conditions.

In Table 4, we display approximate values of $\bar{x}(5)$ for different values of μ and n .




Fig. 5 Numerical solutions of Eq. (37) for $n = 40$ and with respect to perturbations of parameters: μ_1 (left), μ_2 (center) and μ_3 (right). (Example 5)

Example 5 Let us now consider the following fractional differential equation:

$${}^C D_{0+}^{3/2+\mu_3, \psi} x(t) = \sin((\mu_1 - 1)x(t)) + x(t) + \mu_1, \quad t \in [0, 5], \quad (37)$$

with initial conditions

$$x_0 = \mu_2, \quad x_0^1 = 0,$$

where $\mu_1, \mu_2, \mu_3 \in \mathbb{R}$ are three parameters. Let $\psi(t) = \sqrt{t+1}$. In this case, we have

$$F(t, x) = \sin((\mu_1 - 1)x) + x, \quad g(t) = \mu_2 + \frac{\mu_1(\sqrt{t+1} - 1)^{\frac{3}{2}+\mu_3}}{\Gamma\left(\frac{5}{2} + \mu_3\right)}.$$

To analyse the stability of the numerical method, we consider perturbations of parameters μ_1, μ_2, μ_3 . First, we fix $\mu_2 = 0 = \mu_3$ and we analyse the numerical solutions of the two following equations:

$${}^C D_{0+}^{3/2, \psi} x(t) = \sin((\mu_1 - 1)x(t)) + x(t) + \mu_1, \quad t \in [0, 5], \quad x_0 = 0, \quad x_0^1 = 0, \quad (38)$$

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = 0, \quad x_0^1 = 0, \quad (39)$$

Table 5 Numerical results for equations (38) and (39), with $n = 40$, in Example 5

$\mu_1 - 1$	$\max_i x(t_i) - \bar{x}(t_i) $
-0.2	0.503263941
-0.1	0.260315080
0.1	0.278317099
0.2	0.574119985

Table 6 Numerical results for equations (40) and (41), with $n = 40$, in Example 5

μ_2	$\max_i x(t_i) - \bar{x}(t_i) $
0.1	0.3018276927
0.2	0.587713310
0.3	0.881569965
0.4	1.175426619

Table 7 Numerical results for Eqs. (42) and (43), with $n = 40$, in Example 5

μ_3	$\max_i x(t_i) - \bar{x}(t_i) $
0.1	0.103558302
0.2	0.207116085
0.3	0.309937744
0.4	0.411352999

for $\mu_1 \in \{0.8, 0.9, 1, 1.1, 1.2\}$. Let x and \bar{x} be solutions of (38) and (39), respectively. Figure 5 (left) shows x for different values of μ_1 and $n = 40$. In Table 5, we display $\max_i |x(t_i) - \bar{x}(t_i)|$. Next, we assume that $\mu_1 = 1$, $\mu_3 = 0$, and consider numerical solutions to equations:

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = \mu_2, \quad x_0^1 = 0, \quad (40)$$

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = 0, \quad x_0^1 = 0, \quad (41)$$

for $\mu_2 \in \{0, 0.1, 0.2, 0.3, 0.4\}$. Let x and \bar{x} be solutions of (40) and (41), respectively. Numerical results, for different values of μ_2 , are presented in Fig. 5 (center) and Table 6.

Finally, we fix $\mu_1 = 1$, $\mu_2 = 0$, and analyse numerical solutions to equations:

$${}^C D_{0+}^{3/2+\mu_3, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = 0, \quad x_0^1 = 0, \quad (42)$$

$${}^C D_{0+}^{3/2, \psi} x(t) = x(t) + 1, \quad t \in [0, 5], \quad x_0 = 0, \quad x_0^1 = 0, \quad (43)$$

for $\mu_3 \in \{0, 0.1, 0.2, 0.3, 0.4\}$. Let x and \bar{x} be solutions of (42) and (43), respectively. Numerical results, for different values of μ_3 , are presented in Fig. 5 (right) and Table 7.

6 Conclusions

Recently, the Caputo derivative with respect to a kernel function ψ was proposed and applied to some real-world processes (Almeida 2017a,b; Voyiadjis and Sumelka 2019). As usual, an important issue is to develop numerical methods for fractional differential equations with

this new type of derivative. In this paper, such a numerical scheme is presented and its error bound is investigated. The idea is based on an equivalence relation between the fractional differential equation with ψ -Caputo derivative and the Volterra–Stieltjes integral equation. To the latter equation, the generalized midpoint rule is applied. The numerical simulations show that the proposed numerical scheme preserves the underlying structural stability of the initial value problem, with respect to small perturbation of the initial data, and satisfactory and reliable results could be obtained. It means that the approximation routine presented in (Asanov et al. 2011b) for Stieltjes integral can also be successfully applied to solve fractional differential equations, where the fractional derivative operator depends on an increasing function. Nevertheless, as mentioned in Introduction, there exist various numerical methods for integral equations. Therefore, we expect that some of those methods could be adopted for fractional differential equations with ψ -Caputo derivative. This important issue will be considered in a forthcoming paper.

Acknowledgements R. Almeida is supported by Portuguese funds through the CIDMA -Center for Research and Development in Mathematics and Applications, and the Portuguese Foundation for Science and Technology (FCT-Fundação para a Ciência e a Tecnologia), within project UID/MAT/04106/2013. A. B. Malinowska is supported by the Bialystok University of Technology Grant S/WI/1/2016.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Agrawal OP (2010) Generalized variational problems and Euler–Lagrange equations. *Comput Math Appl* 59(5):1852–1864
- Almeida R (2017a) A Caputo fractional derivative of a function with respect to another function. *Commun Nonlinear Sci Numer Simul* 44:460–481
- Almeida R (2017b) What is the best fractional derivative to fit data? *Appl Anal Discrete Math* 11:358–368
- Almeida R, Malinowska AB, Monteiro MTT (2018) Fractional differential equations with a Caputo derivative with respect to a kernel function and their applications. *Math Methods Appl Sci* 41:336–352
- Almeida R, Malinowska AB, Odziejewicz T (2019) On systems of fractional differential equations with the ψ -Caputo derivative and their applications. *Math Methods Appl Sci*. <https://doi.org/10.1002/mma.5678>
- Apartsyn AS (2003) Nonclassical linear Volterra equations of the first kind. VSP, Utrecht
- Arqub OA, Al-Smadi M (2018a) Atangana–Baleanu fractional approach to the solutions of Bagley–Torvik and Painlevé equations in Hilbert space. *Chaos Solitons Fractals* 117:161–167
- Arqub OA, Al-Smadi M (2018b) Numerical algorithm for solving time-fractional partial integrodifferential equations subject to initial and Dirichlet boundary conditions. *Numer Methods Partial Differ Equ* 34:1577–1597
- Arqub OA, Maayah B (2018) Numerical solutions of integrodifferential equations of Fredholm operator type in the sense of the Atangana–Baleanu fractional operator. *Chaos Solitons Fractals* 117:117–124
- Asanov A (1998) Regularization, uniqueness and existence of solutions of Volterra equations of the first kind. VSP, Utrecht
- Asanov A (2001) The derivative of a function by means of an increasing function. *Manas J Eng* 1:18–64 (in Russian)
- Asanov A, Abdujabbarov MM (2011) Solving linear Fredholm–Stieltjes integral equations of the second kind by using the generalized midpoint rule. *J Math Syst Sci* 5:459–463
- Asanov A, Chelik MH, Chalish A (2011a) Approximating the Stieltjes integral by using the generalized trapezoid rule. *Matematiches* 66(2):13–21
- Asanov A, Chelik MN, Abdujabbarov MM (2011b) Approximating the Stieltjes integral using the generalized midpoint rule. *Matematika* 27(2):139–148
- Asanov A, Hazar E, Eroz M, Matanova K, Abdyldeaeva E (2016) Approximate solution of Volterra–Stieltjes linear integral equations of the second kind with the generalized trapezoid rule. *Adv Math Phys* 2016:1–6

- Atanackovic TM, Stankovic B (2009) Generalized wave equation in nonlocal elasticity. *Acta Mech* 208(1–2):1–10
- Banas J, Regan OO (2005) Volterra–Stieltjes integral operators. *Math Comput Model Dyn Syst* 1(2–3):335–344
- Banas J, Rodrigues JR, Sadarangani K (2000) On a class of Urysohn–Stieltjes quadratic integral equations and their applications. *J Comput Appl Math* 113(1–2):35–50
- Bukhgeim AL (1999) Volterra equations and inverse problems. VSP, Utrecht
- Carpinteri A, Mainardi F (2014) Fractals and fractional calculus in continuum mechanics, vol 378. Springer, Wien
- Delves LM, Walsh J (1974) Numerical solution of integral equations. Oxford University Press, Oxford
- Demir DD, Bildik N, Sinir GB (2012) Application of fractional calculus in the dynamics of beams. *Bound Value Probl* 2012(135):1–13
- Diethelm K (2010) The analysis of fractional differential equations: an application-oriented exposition using differential operators of Caputo type. Lecture notes in mathematics. Springer, Berlin
- Djida JD, Atangana A, Area I (2017) Numerical computation of a fractional derivative with non-local and non-singular kernel. *Math Model Nat Phenom* 12(3):4–13
- Federson M, Bianconi R (2001) Linear Volterra–Stieltjes integral equations in the sense of the Kurzweil–Henstock integral. *Arch Math* 37(4):307–328
- Federson M, Bianconi R, Barbanti L (2002) Linear Volterra integral equations. *Acta Math Appl Sin* 18(4):553–560
- Ford NJ, Connolly JA (2006) Comparison of numerical methods for fractional differential equations. *Commun Pure Appl Anal* 5(2):289–306
- Garra R, Giusti A, Mainardi F (2018) The fractional Dodson diffusion equation: a new approach. *Ricerche Mat* 67(2):899–909
- Kalitkin NN (1978) Calculus of approximations. Nauka, Moscow (in Russian)
- Kilbas AA, Srivastava HM, Trujillo JJ (2006) Theory and applications of fractional differential equations. North-Holland Mathematics Studies, vol 204. Elsevier Science B.V., Amsterdam
- Klimek M, Lupa M (2013) Reflection symmetric formulation of generalized fractional variational calculus. *Fract Calc Appl Anal* 16(1):243–261
- Kulish VV, Lage JL (2002) Application of fractional calculus to fluid mechanics. *J Fluids Eng* 124(3):803–806
- Lubich C (1985) Fractional linear multistep methods for Abel–Volterra integral equations of the second kind. *Math Comput* 45:463–469
- Malinowska AB, Odziejewicz T, Torres DFM (2015) Advanced methods in the fractional calculus of variations, Springer briefs in applied sciences and technology. Springer, Cham
- Meerschaert MM (2011) Fractional calculus, anomalous diffusion, and probability. Fractional dynamics. World Scientific Publishing Co. Pte. Ltd., Singapore, pp 265–284
- Morgado ML, Ford NJ, Lima PM (2013) Analysis and numerical methods for fractional differential equations with delay. *J Comput Appl Math* 252:159–168
- Osman MS (2017) Multiwave solutions of time-fractional $(2 + 1)$ -dimensional Nizhnik–Novikov–Veselov equations. *Pramana J Phys* 88(4):67
- Osman MS, Rezazadeh H, Eslami M (2019) Traveling wave solutions for $(3+1)$ dimensional conformable fractional Zakharov–Kuznetsov equation with power law nonlinearity. *Nonlinear Eng* 8(1):559–567
- Podlubny I (1999) Fractional differential equations. Academic, San Diego
- Polyanin AD, Manzhirov AV (2008) Handbook of integral equations, 2nd edn. Hall/CRC Press, Boca Raton
- Rezazadeh H, Osman MS, Eslami M, Ekici M, Sonmezoglu A, Asma M, Biswas A (2018) Mitigating Internet bottleneck with fractional temporal evolution of optical solitons having quadratic-cubic nonlinearity. *Optik* 164:84–92
- Rezazadeh H, Osman MS, Eslami M, Mirzazadeh M, Zhou Q, Badri SA, Korkmaz A (2019) Hyperbolic rational solutions to a variety of conformable fractional Boussinesq-like equations. *Nonlinear Eng* 8(1):224–230
- Sousa JVC, Oliveira EC (2019) A Gronwall inequality and the Cauchy-type problem by means of ψ -Hilfer operator. *Differ Equ Appl* 11:87–106
- Tariq KU, Younis M, Rezazadeh H, Rizvi STR, Osman MS (2018) Optical solitons with quadratic-cubic nonlinearity and fractional temporal evolution. *Mod Phys Lett B* 32(26):1850317
- Vazquez L (2005) A fruitful interplay: from nonlocality to fractional calculus, nonlinear waves: classical and quantum aspects. *NATO Sci Ser II Math Phys Chem* 153:129–133
- Voyiadis GZ, Sumelka W (2019) Brain modelling in the framework of anisotropic hyperelasticity with time fractional damage evolution governed by the Caputo–Almeida fractional derivative. *J Mech Behav Biomed Mater* 89:209–216

Yang XJ, Machado JA (2017) A new fractional operator of variable order: application in the description of anomalous diffusion. *Phys A* 481:276–283

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.