


## RESEARCH

## Open Access



# Diurnal regulation of *SDG2* and *JMJ14* by circadian clock oscillators orchestrates histone modification rhythms in *Arabidopsis*

Qingxin Song<sup>1,2,3†</sup>, Tien-Yu Huang<sup>1,2†</sup>, Helen H. Yu<sup>1,2</sup>, Atsumi Ando<sup>1,2</sup>, Paloma Mas<sup>4</sup>, Misook Ha<sup>5\*</sup> and Z. Jeffrey Chen<sup>1,2,3\*</sup> 

## Abstract

**Background:** Circadian rhythms modulate growth and development in all organisms through interlocking transcriptional-translational feedback loops. The transcriptional loop involves chromatin modifications of central circadian oscillators in mammals and plants. However, the molecular basis for rhythmic epigenetic modifications and circadian regulation is poorly understood.

**Results:** Here we report a feedback relationship between diurnal regulation of circadian clock genes and histone modifications in *Arabidopsis*. On one hand, the circadian oscillators *CCA1* and *LHY* regulate diurnal expression of genes coding for the eraser (*JMJ14*) directly and writer (*SDG2*) indirectly for H3K4me3 modification, leading to rhythmic H3K4me3 changes in target genes. On the other hand, expression of circadian oscillator genes including *CCA1* and *LHY* is associated with H3K4me3 levels and decreased in the *sdg2* mutant but increased in the *jmj14* mutant. At the genome-wide level, diurnal rhythms of H3K4me3 and another histone mark H3K9ac are associated with diurnal regulation of 20–30% of the expressed genes. While the majority (86%) of H3K4me3 and H3K9ac target genes overlap, only 13% of morning-phased and 22% of evening-phased genes had both H3K4me3 and H3K9ac peaks, suggesting specific roles of different histone modifications in diurnal gene expression.

**Conclusions:** Circadian clock genes promote diurnal regulation of *SDG2* and *JMJ14* expression, which in turn regulate rhythmic histone modification dynamics for the clock and its output genes. This reciprocal regulatory module between chromatin modifiers and circadian clock oscillators orchestrates diurnal gene expression that governs plant growth and development.

**Keywords:** Diurnal, Histone modification, *Arabidopsis*

## Background

Circadian clocks anticipate diurnal cycles to coordinate physiological and developmental processes in all organisms from cyanobacteria, flies, to humans and plants [1–3], which are achieved through feedback transcriptional-translational regulatory modules [4–7]. In the transcriptional module, chromatin modifications regulate central circadian oscillators in mammals [8–10] and

plants [7, 11–14]. In mouse liver, there are time-dependent patterns of transcription factor binding, RNA polymerase II recruitment, RNA abundance, and chromatin states, which account for 22% of mRNA cycling genes, suggesting a transcriptional mechanism for regulating the mammal circadian network [8].

In *Arabidopsis*, the central transcriptional loop involves two morning-phased Myb-like transcription factors, CIRCADIAN CLOCK ASSOCIATED 1 (*CCA1*) and LATE ELONGATED HYPOCOTYL (*LHY*) [15, 16], which repress the evening-phased regulator, *timing of CAB expression 1 (TOC1)*, by directly binding to its promoter [17, 18]. In turn, *TOC1* indirectly promotes the expression of *CCA1* and *LHY* by inhibiting the

\* Correspondence: [misook.ha@gmail.com](mailto:misook.ha@gmail.com); [zjchen@austin.utexas.edu](mailto:zjchen@austin.utexas.edu)

<sup>†</sup>Qingxin Song and Tien-Yu Huang contributed equally to this work.

<sup>5</sup>Samsung Advanced Institute of Technology, Samsung Electronics Corporation, Suwon 443-803, South Korea

<sup>1</sup>Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX 78712, USA

Full list of author information is available at the end of the article



repressive *PSEUDO-RESPONSE REGULATOR* (*PRR*) genes or directly represses expression of *CCA1* and *LHY* by binding their promoters [19, 20]. Integrating other clock regulators including *GI* and *ELF4–ELF3–LUX* complex with the central *CCA1/LHY/TOC1* loop provides a classic feedback regulatory mechanism for the circadian clock in *Arabidopsis* [21, 22], which includes a number of core molecular oscillators that constitute interlocking transcriptional-translational negative feedback loops (see reviews in [2, 4, 7, 23, 24]).

Using candidate gene approaches, several studies have indicated links between histone modifications and expression of circadian clock genes. For example, induction of the morning-phased *TOC1* expression wave form is related to histone acetylation, which is repressed by *CCA1* and coincident with histone deacetylase activities and repressive chromatin status at dawn [12]. Rhythmic expression of *CCA1* and *LHY* and their reciprocal regulator *TOC1* is associated with H3K4me3 and H3K9/14 ac levels in the translational start sites of these genes [11, 13], but correlated negatively with H3K36me2 levels [11]. Moreover, changes in diurnal transcript levels correlate with H3K9ac, H3K27ac, and H3S28p levels between end-of-day and end-of-night [25]. Inhibiting acetylation and H3K4me3 abolishes rhythmic expression of circadian oscillators, while blocking H3K4me3 leads to increased levels of clock-repressor binding, suggesting a transitional role for the H3K4me3 mark in modulating clock gene expression from activation to repression [14]. *JMJ5*, a histone demethylase in humans and plants, is co-regulated with evening-phased clock regulators and promotes expression of clock genes at dawn [26]. However, neither the basis for histone mark rhythms nor the relationship between histone modifications and circadian clock regulators is clearly defined. Here we report dynamic interactions between chromatin modification of circadian clock gene expression and diurnal regulation of histone modifications, which regulate diurnal gene expression networks in *Arabidopsis*.


## Results

### Diurnal regulation of histone methyltransferase and demethylase by clock genes

The association of rhythmic histone modifications with expression of individual circadian clock genes in *Arabidopsis thaliana* [14] and with altered expression of the core clock genes in *Arabidopsis* hybrids and allopolyploids [27] may suggest a general role of histone modifications in diurnal regulation of the circadian clock and its output genes. The diurnal rhythms are likely established through diurnal transcriptional regulation of corresponding genes or diurnal oscillation of histone-modifying enzymes. But the relationship between circadian clock regulators and histone-modifying factors is

largely unknown. In *Arabidopsis*, five histone methyltransferases (*ATX1*, *SDG2*, *SDG4*, *SDG25*, and *SDG26*) and three histone demethylases (*JMJ14*, *JMJ15*, and *JMJ18*) are reported to be responsible for adding and removing H3K4me3, respectively [28–35]. Among them, *SDG2* and *JMJ14* are associated with dramatic changes of genome-wide H3K4me3 levels in their respective mutants [36–38], suggesting that these two genes may be primary targets of the circadian clock regulation. To test this, we examined expression patterns of *SDG2* and *JMJ14* subjected to diurnal (24 h, 16 h light/8 h dark) and circadian conditions (48 h, constant light) in the wild type (*Col-0*). *SDG2* showed rhythmic expression peaks under diurnal but not constant light conditions (Additional file 1: Figure S1a). However, *JMJ14* showed rhythmic expression patterns under both diurnal and constant light conditions (Additional file 1: Figure S1b). Thus, expression patterns of *SDG2* and *JMJ14* are diurnal and circadian, respectively. To further explore the roles of clock genes in *SDG2* and *JMJ14* regulation, we examined *SDG2* and *JMJ14* expression changes under diurnal (24 h, 16 h light/8 h dark, LD) conditions in the wild type (*Ws*) compared to the *cca1 lhy* double mutant (*Ws* background) and in the *CCA1* overexpression line (*CCA1-OX*, *Col-0* background) relative to *Col-0*. Under diurnal condition, the expression of *SDG2* peaked in the morning (ZT28, ZT0 or 24 = dawn) in the wild type and was significantly downregulated in the *cca1 lhy* mutant but upregulated in the *CCA1-OX* line (Fig. 1a). Expression of *SDG2* in the WT started to increase from ZT24 (dawn), and overexpressing *CCA1* may have a “burst” effect at ZT24 and accelerated the *SDG2* expression peak (Fig. 1a), with a caveat of a potential genotypic effect (*Ws* vs. *Col*).

In contrast to *SDG2*, *JMJ14* expression peaked at night (ZT40) and reached the valley after dawn (ZT28) in the WT and was significantly upregulated in the *cca1 lhy* mutant and downregulated in the *CCA1-OX* line under diurnal condition (Fig. 1b). The data suggest that *CCA1/LHY* exerts temporal effects on positive and negative correlation, respectively, with *SDG2* and *JMJ14* expression levels. Although both *SDG2* and *JMJ14* have one *CCA1*-binding site (CBS) in their promoter sequences, *SDG2* and *JMJ14* were not reported as target genes of *CCA1* according to the published dataset of chromatin immunoprecipitation sequencing (ChIP-seq), which was performed using *CCA1p::CCA1-GFP* transgenic plants at ZT2 and ZT14 [39]. To determine whether *CCA1* can bind promoters of *SDG2* and *JMJ14* at different time points, we performed ChIP-qPCR using antibodies against *CCA1* at ZT0 when *CCA* expression peaks. ChIP-qPCR showed significant enrichment of *CCA1* in the promoters of *JMJ14* and *TOC1* (normalized to *UBQ10*), but not in the promoter of *SDG2* (Fig. 1c).



**Fig. 1** The clock regulates expression of histone methyltransferase (*SDG2*) and demethylase (*JMJ14*) genes. **a, b** Relative expression levels (R.E.L.) of *SDG2* (**a**) and *JMJ14* (**b**) in the wild type (WT, *Ws*), *cca1 lhy* mutant (*Ws*) (upper panel), and WT (*Col-0*) and *CCA1-OX* (*Col-0*) (lower panel) plants under a diurnal cycle (16 h light/8 h dark). Black and white bars indicate dark (ZT16, 20, and 24) and light (ZT28, 32, 36, and 40), respectively. **c** ChIP-qPCR showed the fold enrichment (Y-axis) of *CCA1*-binding fractions in the promoters of *SDG2* and *JMJ14* using rosette tissues from 3-week-old *A. thaliana* at ZT0. The *TOC1* promoter containing evening elements was used as the positive control. *UBQ10* was used as a nonbinding control for normalization. Error bars in **a–c** indicate standard deviation of biological replicates ( $n = 3$ ) with an asterisk showing the statistical significance level ( $P < 0.05$ , Student *t* test). **d** Binding profiles of H3K4me3 at the TSS  $\pm 1.5$  kb in the wild type and *cca1 lhy* mutant with two biological replicates. **e** Examples showing decreased binding of H3K4me3 to the genes *AT1G21270* (upper panel) and *AT3G05880* (lower panel) in *cca1 lhy* mutant compared with the WT (*Ws*)

Although we could not exclude a possibility of non-specific binding of *CCA1*-antibodies, this result indicates that *CCA1/LHY* may directly bind to the *JMJ14* promoter and regulates its expression, as other members (*JMJ30* and *JMJ5*) of the gene family that are co-regulated with evening-phased genes [26, 40], but may indirectly regulate *SDG2* expression through other factors such as *CCA1*-mediated genes or in the *CCA1* complex. Alternatively, regulation of *SDG2* by *CCA1* could occur at a different time of the day. In addition, other H3K4 methyltransferase *ATX1* and H3K4 demethylase *JMJ15* were also down- and upregulated, respectively, in the *cca1 lhy* mutant (Additional file 1: Figure S1c, d). The data suggest that *CCA1/LHY* can upregulate expression of H3K4me3 writers (*SDG2* and *ATX1*) and downregulate H3K4me3 erasers (*JMJ14* and *JMJ15*), which would result in the overall decreased distributions of H3K4me3 in the *cca1 lhy* mutant.

Using ChIP-seq analysis with antibodies against H3K4me3, we analyzed H3K4me3 distribution patterns in the rosette leaves of wild type (*Ws*) and *cca1 lhy*


mutant plants at dawn (ZT0) under diurnal conditions, when *CCA1/LHY* expression level peaked [41]. Compared to the wild type, H3K4me3 levels near the transcription start sites of genic regions were damped in the *cca1 lhy* mutant (Fig. 1d). A total of 880 genes showed decreased levels of H3K4me3 peaks (one-way ANOVA,  $P < 0.05$ ) (Additional file 2: Table S1), including two examples shown in Fig. 1e. A previous study has identified *CCA1* target genes by ChIP-seq using 2-week-old seedlings grown under 12 h light/12 h dark for 12 days and then transferred to constant light for 2 days [39]. The overlapping fraction is small between the *CCA1* target genes (5%, 78/1433) and those of reduced H3K4me3 levels in the *cca1 lhy* mutant ( $P > 0.5$ , hypergeometric test). The results indicate that *CCA1* affects overall H3K4me3 accumulation levels but does not directly participate in the establishment of H3K4me3 marks for *CCA1* target genes. However, different growth conditions and tissue stages in these two studies may also contribute to the small overlapping fraction between the *CCA1* target genes and genes with reduced H3K4me3 levels in the *cca1 lhy* mutant.


### Disrupted histone modifications lead to altered expressions of clock genes

To test how histone modifications affect expression of circadian clock genes, we analyzed H3K4me3 level changes of the clock genes in the *sdg2* (10-day-old seedlings under long-day conditions) or *jmj14* (3-week-old rosette leaves 16 h light/8 h dark) mutant using the published ChIP-seq datasets [42, 43]. Consistent with the previous report [14], disruption of *SDG2* resulted in a dramatic decrease of H3K4me3 accumulation levels in most clock genes examined, including *CCA1* and *LHY* (Fig. 2a), *TOC1*, *PRR5*, *PRR7*, *PRR9*, *GI*, *ELF3*, *ELF4*, and *LUX* (Additional file 1: Figure S2). ChIP-qPCR using rosette leaves at ZT0, ZT12, and ZT24 further confirmed a significant reduction of H3K4me3 levels near the 5' region of *CCA1* and *LHY* in the *sdg2* mutant (Additional file 1: Figure S3a). Consequently, expression levels of *CCA1* and *LHY* were dramatically reduced in the *sdg2* mutant (Fig. 2b). In contrast to the results in the *sdg2* mutant, H3K4me3 accumulation levels of *CCA1*, *LHY*, and several other clock genes were slightly increased in the *jmj14* mutant (Fig. 2c and Additional file 1: Figure S2). This moderate effect may be related to the genetic redundancy of JMJ demethylase family including six other members (*JMJ15*, *JMJ16*, *JMJ17*, *JMJ18*, *JMJ19*, and *JMJ30*) in *Arabidopsis* [32, 34, 40]. ChIP-qPCR confirmed a significant increase of H3K4me3 levels near the 5' region of *CCA1* and *LHY* in the *jmj14* mutant (Additional file 1: Figure S3b), which correlated with increased expression levels of those clock genes (Fig. 2d).

### Dynamic diurnal rhythms of histone modifications in *Arabidopsis*

In animals and plants, histone marks like H3K4me3 and H3K9ac are associated with active transcription and can sometimes be part of transcriptional activation [44, 45]. Clock-mediated diurnal expression of histone-modifying genes suggest rhythmic distributions of histone marks. To examine diurnal rhythms of histone modifications, we used ChIP-seq to examine diurnal rhythms of genome-wide H3K4me3 levels every 3 h in a 24-h period using rosette leaves of *A. thaliana* grown under the long-day cycle (16 h light/8 h dark). For comparative studies, we also performed ChIP-seq experiments using another active histone mark H3K9ac. We identified a total of 15,151 H3K4me3 and 16,966 H3K9ac high-confidence peaks ( $P_{\text{adj}} < 0.01$ , MACS), corresponding to 14,818 and 15,422 unique genes, respectively (Additional file 3: Table S2). The distribution of H3K4me3 and H3K9ac in both evening and morning phases may suggest that these marks are also required for general transcriptional regulation [46]. Of those peaks, 4420 (29%) H3K4me3 and 3493 (21%) H3K9ac peaks exhibited diurnal rhythms at a statistically significant level ( $P_{\text{adj}} < 0.01$ , JTK\_CYCLE using Kendall's tau correlation and Bonferroni correction to adjust  $P$  value) (Fig. 3a and Additional file 4: Table S3). Those oscillating H3K4me3 and H3K9ac peaks overlapped 4367 and 3276 genes, respectively. This distribution rhythm is not likely affected by histone H3, and only 21 (0.5%) H3K4me3 and 32 (0.9%) H3K9ac rhythmic peaks exhibited histone H3 distribution rhythms at a statistically significant level ( $P_{\text{adj}} < 0.01$ , JTK\_CYCLE) (Additional file 1: Figure S3c, d). Another study identified 657 and 1495





genes with oscillating H3K4me3 and H3K9ac peaks, respectively, using two time points (end-of-day and end-of-night) in the short-day condition [25]. Between two datasets, 38% (250/657) and 28% (418/1495) of the genes with rhythmic H3K4me3 and H3K9ac peaks also showed oscillating H3K4me3 and H3K9ac rhythms in our study ( $P < 1e-38$ , hypergeometric test). The difference between these two studies could be related to the differences in the time points (nine vs. two) examined and in the growth conditions (long-day vs short-day) used between two experiments.

The phase distribution analysis indicated genome-wide binding rhythms of H3K4me3 with a major peak at ZT12 and a minor peak at ZT6, whereas the major peak of H3K9ac binding rhythms was delayed towards ZT15 and the minor peak was moved forward to ZT3 (Additional file 1: Figure S4a). This is consistent with total levels of H3K4me3 and H3K9ac as determined by

the Western blot analysis (Additional file 1: Figure S4b). The ChIP-seq data further showed a diurnal rhythm of the total H3K4me3 levels with the peak at ZT12, whereas the total H3K9ac levels did not show obvious peaks during the day and night (Additional file 1: Figure S4b). The inconsistency between diurnal H3K9ac distribution and global H3K9ac content may be related to inconspicuous H3K9ac distribution variation in the time points examined in total H3K9ac levels (ZT0, 6, 12, 18, 24). Alternatively, non-rhythmic H3K9ac distribution may account for the majority of overall H3K9ac variation because of its role in other aspects of plant growth and development [47, 48].

The ChIP-seq data were further classified into the morning-phased (between ZT0 and ZT6) and evening-phased (between ZT9 and ZT18) peaks using k-medoids clustering [49] (Fig. 3a and Additional file 4: Table S3). A majority (86%, 14,029/16,391) of H3K4me3- or H3K9ac-enriched genes had both H3K4me3 and H3K9ac peaks,

and only a small proportion of genes was associated with only one histone mark (Fig. 3b). This suggests a coordinated mechanism for H3K4me3 and H3K9ac to regulate gene expression [45]. However, among morning-phased genes, only 13% (313/2335) were associated with both morning-phased H3K4me3 and H3K9ac peaks. Among evening-phased genes, 696 genes (22%) were related to both evening-phased H3K4me3 and H3K9ac peaks (Fig. 3b). The proportion of genes associated with both histone marks in morning-phased (13%) or evening-phased (22%) peaks was significantly lower than that enriched with either H3K4me3 or H3K9ac peaks ( $P < 2e^{-100}$ , hypergeometric test).

Gene ontology (GO) analysis suggested that the morning- and evening-phased H3K4me3 and H3K9ac marks were enriched in genes with different gene ontology (GO) groups (Fig. 3c). The morning-phased genes are enriched with photosynthetic activities and responses to a variety of stresses, while the evening-phased genes are enriched with DNA and RNA metabolism and protein ubiquitination and folding (Fig. 3c). These data implied a specificity for diurnal H3K4me3 and H3K9ac rhythms in the expression of genes involved in different biological networks. A total of 22% (958/4367) and 28% (1208/4367) genes with rhythmic H3K4me3 peaks in the *sdg2* and *jmj14* mutants, respectively, showed altered H3K4me3 levels (Fig. 3d), which was 1.5-fold higher than those genes (12%, 1252/10,503 and 17%, 1810/10,503) with non-rhythmic H3K4me3 peaks in the mutants, respectively, displaying altered H3K4me3 levels ( $P < 1e^{-10}$ , Fisher's exact test). The data suggest that the genes with rhythmic histone modifications are more sensitive to the loss of histone methyltransferase and demethylase activities than other genes.


To examine the correlation between oscillating histone modifications and gene expression, we analyzed expression patterns of genes using published data of expression profiling in *Arabidopsis* under diurnal (16 h light/8 h dark) and circadian (constant light) conditions [41]. For the genes with both diurnal and circadian expression patterns, 14% (612/4377) had both rhythmic H3K4me3 and H3K9ac marks (Additional file 4: Table S3). However, the fraction of both rhythmic H3K4me3 and H3K9ac marks was much lower for the genes with circadian but not diurnal expression patterns (4%, 150/4134) or for the genes with diurnal but not circadian expression patterns (4%, 139/3121). These results suggest different impact of histone-modifying enzymes in diurnal and/or circadian expression patterns. JMJ proteins are circadian-mediated probably through coregulation with the clock components [26, 40], while *SGD2* expression is indirectly regulated by *CCA1* (Fig. 1c and Additional file 1: Figure S1). At the genome-wide level, ~48% (2104/4367) of the genes with oscillating H3K4me3 and ~47% (1552/3276) of the genes with oscillating H3K9ac displayed rhythmic

expression with distinct morning-phased and evening-phased patterns [41] (Fig. 3e), indicating a strong correlation between diurnal histone modifications and rhythmic gene expression.

#### Roles of histone modifications in core clock genes

Further analysis of the ChIP-seq data showed rhythmic patterns of either or both H3K4me3 and H3K9ac marks for all core clock genes examined, including *CCA1*, *LHY*, *TOC1*, *PRR5*, *PRR7*, *GI*, *ELF3*, and *LUX* (Fig. 4a–d and Additional file 1: Figure S5 and S6). This confirms previous reports about the correlation between oscillating chromatin modifications and rhythmic expressions of key clock genes, including *LHY*, *CCA1*, *TOC1*, *PRRs*, and *LUX* [11–14]. Most core clock genes possessed both marks; the morning-expressed genes like *CCA1* had H3K4me3 and H3K9ac peaks at dawn (ZT0–ZT3) (Fig. 4a), while the evening-expressed genes like *TOC1* had H3K4me3 and H3K9ac peaks at noon (ZT12) (Fig. 4c). Expectedly, the transcript levels of *CCA1* and *TOC1* correlated strongly with H3K4me3 and H3K9ac accumulation on a diurnal basis (Fig. 4b, d). This suggests a synchronization between circadian expression and chromatin rhythms.


A possibility for expression rhythms and chromatin modifications is that the core circadian clock genes may interact with chromatin-modifying enzymes to alter histone modifications and regulate target gene expression. To test this, we analyzed binding targets of *CCA1* and *TOC1* from the published ChIP-seq datasets [20, 39], and examined enrichment levels of H3K4me3 and H3K9ac in the target genes. As *CCA1* and *LHY* often repress expression of evening-phased genes by binding their promoters [39], the majority of *CCA1* targets showed rhythmic expression patterns with an evening phase. The results showed that approximately 33% and 20% of the *CCA1* targets had evening-phased binding patterns for H3K4me3 and H3K9ac ( $P < 1e^{-50}$ , hypergeometric test), respectively, while fewer than 7% of *CCA1* targets exhibited morning-phased profiles for H3K4me3 and H3K9ac (Fig. 4e). This finding is consistent with the expression patterns of those *CCA1* target genes in the evening phase (Additional file 1: Figure S7), although phased distributions of morning and evening genes could change under long- or short-day conditions [41, 50]. In contrast, 23% and 13% of *TOC1* target genes correlated with H3K4me3 and H3K9ac marks in the morning phase (Fig. 4f) ( $P < 1e^{-5}$ , hypergeometric test), while a smaller proportion of target genes correlated with those marks in the evening. This is consistent with *TOC1* being a regulator of the morning-phased genes [19, 20] (Additional file 1: Figure S7). To further test the roles of *SDG2* and *JMJ14* in circadian-mediated gene expression, we randomly selected three *CCA1* target genes



and examined enrichment of H3K4me3 in genic region of these genes in wild type and *sdg2* and *jmj14* mutants at ZT0, 12, and 24. Enrichment of H3K4me3 in all three *CCA1* target genes was significantly decreased in the *sdg2* mutant at similar levels in the time points tested but increased in the *jmj14* mutant in most time points (Additional file 1: Figure S8). The effect in a few time points was moderate in the *jmj14*, probably due to genetic redundancy, as observed for the clock genes (Fig. 2c and Additional file 1: Figure S3). Notably, the H3K4me3 levels in the *sdg2* mutant are reduced at all time points tested (Additional file 1: Figure S8a), consistent with that *SDG2* is not directly regulated by *CCA1* (Fig. 1c). *JMJ14*, however, showed increased H3K4me3 levels with relative rhythms (Additional file 1: Figure S8b), consistent with a role for *JMJ* proteins in co-regulating with the circadian clock and its target genes [26, 32]. These data indicate that *SDG2* and *JMJ14* are involved in regulation of rhythmic deposition of H3K4me3 in *CCA1* target genes.

### Discussion and conclusions

We provided a comprehensive genome-wide landscape of histone modification dynamics in *Arabidopsis*, which has supported diurnal regulation of histone modifications on several individual circadian clock genes as previously reported [11–14, 25, 26]. The current data support a model that integrates clock rhythms with histone modifications to establish rhythmic activities of gene expression, growth, and development (Fig. 5). Circadian clock genes *CCA1* and *LHY* correlate indirectly with the upregulation of the histone methyltransferase gene *SDG2* and directly with repression of the histone demethylase gene *JMJ14* to coordinate histone modification levels in the downstream genes of the circadian output network. Coincidentally, recent studies showed REVEILLE8 interacted with NIGHT LIGHT-INDUCIBLE AND CLOCKREGULATED proteins (LNKs) [51–53] to regulate RNA Pol II and H3K4me3 occupancy on *PRR5* and *TOC1* loci to control their expression [54], further confirming the complex



interaction between circadian oscillator, histone modifications, and basal transcriptional machinery. It is predictable that the dynamic histone modifications could change the chromatin structures of genes and affect the accessibility of clock regulators and/or RNA Polymerase II, contributing to rhythmic expression of the target genes [7, 8]. On the other hand, histone methyltransferases and demethylases directly regulate the expression of core clock genes through adding or removing histone modifications. This interactive and reciprocal regulation between circadian clock oscillators and chromatin modifiers orchestrates diurnal gene expression that controls growth and development in plants and animals [6, 55].

A large proportion of histone modification sites exhibit rhythmic patterns, while different histone modifications (H3K4me3 and H3K9ac) have distinct phases for their target genes, which is consistent with the report in mammals [8]. The data collectively indicate a conserved correlation between histone modifications and circadian clock regulation in plants and mammals, as well as a specific distribution of different histone modifications in morning-phased and evening-phased genes. This could be related to distinct peaks of H3K4me3 and H3K9ac in the morning and evening phases, respectively, which correspond to expression of morning- and evening-

phased genes (Fig. 3). We also noticed both H3K4me3 and H3K9ac showed lower levels at the night (ZT18-24) than during the day (Fig. 3a). One possibility is that total RNA transcription is lower at night than during the day, resulting in lower levels of H3K4me3 and H3K9ac modifications at night. Alternatively, activation of those genes at the night may not require both H3K4me3 and H3K9ac modifications. Other histone modifications could also be involved in evening-phased genes. Further examination of additional histone marks will provide a comprehensive view for diurnal rhythms of histone modifications and gene expression.

For H3K4me3, the rhythm for regulating circadian-mediated gene expression is associated with expression of the H3K4me3 methyltransferase gene *SDG2*, which is expressed at significantly higher levels during the day than at night; on the other hand, the expression level of the H3K4me3 demethylase gene (*JMJ14*) is low during the day but high at night. This provides a basis for transcriptional regulation of rhythmic histone modifications. Indeed, both global H3K4me3 and morning-phased H3K4me3 levels peak during the day (ZT12) (Additional file 1: Figure S4), suggesting that H3K4me3 is largely established during the day. However, the global H3K4me3 peak is delayed 8 h, compared to the expression peak of H3K4me3 methyltransferase *SDG2* (ZT4). This could be related to the protein and mRNA that accumulate at different levels in a diurnal cycle. For example, CCA1 binds to the *JMJ14* promoter at ZT0 in our ChIP-qPCR study but *JMJ14* is not a target gene at ZT2 or ZT14 in the ChIP-seq analysis [39]. CCA1 protein peaks 3 h later than mRNA in maize [56]. The decrease of *JMJ14* transcripts during the day period may result in the presence of H3K4me3 for a longer time. In addition, the dynamic histone modifications depend on maximal activity of histone-modifying enzymes that may change over time. Notably, other members of H3K4me3 writers (e.g., ATX1) and erasers (e.g., JMJ15) may also contribute to expression rhythms of circadian-mediated regulatory networks.

How histone methyltransferases and histone demethylases are recruited to their target genes remains unknown. One possibility is that histone methyltransferases such as ATX1 can be recruited by the transcriptional machinery through interaction with the Ser5-phosphorylated CTD of Pol II for transcription initiation and elongation [46]. Alternatively, histone modifications could be established through the interaction of sequence-specific transcription factors with histone methyltransferases/demethylases to regulate target gene expression [57, 58]. The targeting and binding time of transcription regulators for specific sequences could affect the time of H3K4me3 deposition. A recent study reported that lysine-specific demethylase 1 (LSD1)-like histone demethylases (LDL1 and LDL2) can interact with CCA1/LHY to repress *TOC1* expression



[59]. Moreover, LDL1 and LDL2 interact with HDA6 to co-regulate *TOC1* expression by histone demethylation and deacetylation. It is probable that this type of histone complex can include JM14 and other factors to regulate expression of clock genes. It is equally plausible that histone methyltransferases and acetyltransferases could be in the same complex to carry out the opposite direction of circadian-mediated gene expression. It will be interesting to further test how specific transcription factors including circadian clock regulators recruit other histone methyltransferases and demethylases to establish rhythmic histone modifications that exert growth and developmental regulation in plants and animals.

## Methods

### Plant materials and growth conditions

*A. thaliana* ecotypes Col-0 and Ws were used in this study. *cca1 lhy* mutant in Ws background and *CCA1-OX* transgenic lines in Col-0 background were generated as previously described [27]. The mutants *sdg2* (SALK-021008) and *jmj14* (SALK\_135712) were obtained from Arabidopsis Biological Resource Center (ABRC). The primer sequences for genotyping are listed in Additional file 5: Table S4. For diurnal conditions, plants were grown under the light/dark (L/D) cycle of 16 h/L and 8 h/D at 22 °C.

### RNA extraction and qRT-PCR

Total RNA was isolated from aerial rosette leaves of ~3-week-old plants using Plant RNA Reagent (Thermo Fisher Scientific, Waltham, Massachusetts). After digestion by RNase-Free DNase (Promega, Madison, Wisconsin), total RNA (1 µg) was used to produce first-strand cDNA with the Omniscript RT Kit (Qiagen, Valencia, California). The cDNA was used as the template for qRT-PCR using FastStart Universal SYBR Green Master in a LightCycler<sup>®</sup>96 System (Roche, Indianapolis, Indiana). The relative expression level was quantified using the internal control *ACT7* (*AT5G09810*). Three biological replicates were performed for each experiment, and three technical replicates were used for each biological replicate in qRT-PCR analysis. Student's *t* test and calculation of error bars were performed to determine the significance level in each comparison using three biological replicates. The primer sequences are listed in Additional file 5: Table S4.

### Histone extraction and Western blot analysis

Histone extraction was performed as described previously [60]. An aliquot (1 g) of aerial rosette leaves from ~3-week-old plants were ground to powder with liquid nitrogen and then suspended in NIB buffer (250 mM sucrose, 60 mM KCl, 15 mM NaCl, 5 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>, 15 mM PIPES, pH 6.8, 0.8% Triton X-100, 1 mM

PMSF, and 1x cocktail). The homogenized solution was filtered through four layers of cheesecloth; the supernatant was centrifuged at 3000g for 20 min at 4 °C. Pellets were re-suspended in 800 µl 0.4 N H<sub>2</sub>SO<sub>4</sub> and incubated with a rotator overnight at 4 °C. After centrifugation at 16,000g at 4 °C for 10 min, the supernatant was transferred into a new tube by adding 264 µl trichloroacetic acid. Samples were incubated overnight at 4 °C and centrifuged again at 16,000g at 4 °C for 10 min to precipitate histone proteins. After washing by acetone, proteins were dissolved in water for Western blot analysis. The antibodies were anti-histone H3 (Abcam; Ab1791, 1:5000 dilution), anti-H3K4me3 (Abcam; Ab8580, 1:5000 dilution), and anti-H3K9Ac (Abcam; Ab10812, 1:5000 dilution).

### Chromatin immunoprecipitation (ChIP) and ChIP-seq

Chromatin immunoprecipitation was performed following the published protocol [61]. Briefly, 2 g of aerial rosette tissues from ~3-week-old *A. thaliana* (Col-0) were cross-linked in 1% formaldehyde, and the chromatin fractions were isolated and purified. Antibodies against histone H3 (Abcam; ab1791), H3K4me3 (Abcam; Ab8580), H3K9ac (Abcam; Ab10812), or CCA1 (Abiocode, R1234-3) were added to the chromatin extracts (with a 1:1000 dilution) and incubated overnight at 4 °C with gentle rotation. Immunoprecipitated DNA was extracted using Protein A/G Magnetic Beads (Thermo Fisher Scientific, Waltham, Massachusetts). Immunoprecipitated DNA (IP DNA) and input DNA (without IP but with incubation of beads) were subjected to reverse cross-linking and purified using QIAquick PCR Purification Kit (Qiagen, Valencia, California). For ChIP-seq, libraries with three biological replicates were constructed from IP DNA and non-IP DNA using NEBNext<sup>®</sup> Ultra™ II DNA Library Prep Kit for Illumina (NEB, Ipswich, Massachusetts). For ChIP-qPCR, immunoprecipitated DNA was used as the template for qPCR using FastStart Universal SYBR Green Master in a LightCycler<sup>®</sup>96 System (Roche, Indianapolis, Indiana). The relative enrichment was quantified using the internal control *AT2G26560*. Three biological replicates were performed for each experiment with three technical replicates for each biological replicate. Student's *t* test was used to determine the significance level in each comparison.

### ChIP-seq data analysis

ChIP-seq dataset generated in this study and the published ChIP-seq datasets in *sdg2* and *jmj14* mutants were analyzed as follows. After removing adapters, low-quality reads, and PCR duplicates, the filtered pair-end reads were aligned to *A. thaliana* reference genome (TAIR10) using Bowtie2 (Version 2.3.0) with parameters "bowtie2 --no-mixed --no-discordant" [62]. Only uniquely and concordantly mapped reads were used for further analysis. In order to adjust different

sequencing depths among different samples, the uniquely mapped reads were “down-sampled” to the lowest read number of samples, as previously described [56]. Enriched peaks were identified using Model-based Analysis of ChIP-Seq (MACS, version 2.1.0) with the option “macs2 callpeak -g dm -q 0.01” [63]. Peaks were used for further analysis only if they were present in two biological replicates. To make a master-peak list for all time points, the peaks obtained from each time point were merged. When different samples were compared, the read coverage was normalized to 1× sequencing depth to eliminate bias of different insertion lengths in pair-end reads. The genes overlapping with peaks in the gene body or ± 1-kb regions were defined as peak-overlapped genes. The peaks were compared between the wild type and *cca1 lhy* mutant using one-way ANOVA, and their significance was determined with  $P < 0.05$ . As no biological replicate was used from the ChIP-seq datasets in *sdg2* and *jmj14* mutants, we used 1.5-fold changes to determine differential peaks between the wild type and mutants. The non-parametric algorithm JTK\_CYCLE was used to detect rhythmic peaks and identify the phase of each peak with  $P_{\text{adj}} < 0.01$  [64]. Gene Ontology (GO) analysis was performed using DAVID [65].

## Additional files

- Additional file 1:** Supplementary Figure S1–S8. (PS 3131 kb)
- Additional file 2: Table S1.** List of peaks with different H3K4me3 enrichment between WT and *cca1 lhy* mutant. (XLSX 101 kb)
- Additional file 3: Table S2.** List of all identified H3K4me3 and H3K9ac peaks with coverage levels at all time points. (XLSX 11750 kb)
- Additional file 4: Table S3.** The list of diurnal H3K4me3 and H3K9ac peaks with coverage levels at all time points. (XLSX 2978 kb)
- Additional file 5: Table S4.** Primers used in this study. (DOCX 14 kb)

## Acknowledgements

We thank Dr. Hong Qiao (The University of Texas at Austin) and her laboratory for the technical assistance in Western blot analysis and the staff members at Genome Sequencing and Analysis Facility for sequencing.

## Authors' contributions

QS, MH, and ZJC conceived and designed the research. QS, TH, HHY, and AA performed the experiments and analyzed the data. PM analyzed the data. QS, TYH, MH, and ZJC wrote the manuscript. All authors read and approved the final manuscript.

## Funding

The research is supported by grants to Z.J.C. from the National Institute of General Medical Sciences (GM109076) and Samsung Electronics Co., Ltd. (UTA12-000733, 2012–2014).

## Availability of data and materials

All raw sequencing data have been deposited in NCBI Nucleotide and Sequence Read Archive (SRA) (<https://trace.ncbi.nlm.nih.gov/Traces/sra/?study=SRP152291>) [66]. Most analyses were performed using publicly available software, and the codes used for statistical analysis are available upon request.

## Ethics approval and consent to participate

Not applicable.

## Consent for publication

Not applicable.

## Competing interests

All the authors declare that they have no competing interests.

## Author details

<sup>1</sup>Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX 78712, USA. <sup>2</sup>Department of Integrative Biology, The University of Texas at Austin, Austin, TX 78712, USA. <sup>3</sup>State Key Laboratory of Crop Genetics and Germplasm Enhancement, Nanjing Agricultural University, Nanjing 210095, China. <sup>4</sup>Center for Research in Agricultural Genomics (CRAG), Consortium CSIC-IRTA-UAB-UB, Campus UAB, Bellaterra, 08193 Barcelona, Spain. <sup>5</sup>Samsung Advanced Institute of Technology, Samsung Electronics Corporation, Suwon 443-803, South Korea.

Received: 29 November 2018 Accepted: 29 July 2019

Published online: 20 August 2019

## References

- Bass J, Takahashi JS. Circadian integration of metabolism and energetics. *Science*. 2010;330(6009):1349–54.
- Greenham K, McClung CR. Integrating circadian dynamics with physiological processes in plants. *Nat Rev Genet*. 2015;16(10):598–610.
- Wijnen H, Young MW. Interplay of circadian clocks and metabolic rhythms. *Annu Rev Genet*. 2006;40:409–48.
- Nohales MA, Kay SA. Molecular mechanisms at the core of the plant circadian oscillator. *Nat Struct Mol Biol*. 2016;23(12):1061–9.
- Turek FW. Circadian clocks: tips from the tip of the iceberg. *Nature*. 2008;456(7224):881–3.
- Takahashi JS. Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet*. 2017;18(3):164–79.
- Seo PJ, Mas P. Multiple layers of posttranslational regulation refine circadian clock activity in *Arabidopsis*. *Plant Cell*. 2014;26(1):79–87.
- Koike N, Yoo SH, Huang HC, Kumar V, Lee C, Kim TK, Takahashi JS. Transcriptional architecture and chromatin landscape of the core circadian clock in mammals. *Science*. 2012;338(6105):349–54.
- Etchegaray JP, Lee C, Wade PA, Reppert SM. Rhythmic histone acetylation underlies transcription in the mammalian circadian clock. *Nature*. 2003;421(6919):177–82.
- Ripperger JA, Schibler U. Rhythmic CLOCK-BMAL1 binding to multiple E-box motifs drives circadian Dbp transcription and chromatin transitions. *Nat Genet*. 2006;38(3):369–74.
- Song HR, Noh YS. Rhythmic oscillation of histone acetylation and methylation at the *Arabidopsis* central clock loci. *Mol Cells*. 2012;34(3):279–87.
- Perales M, Mas P. A functional link between rhythmic changes in chromatin structure and the *Arabidopsis* biological clock. *Plant Cell*. 2007;19(7):2111–23.
- Hemmes H, Henriques R, Jang IC, Kim S, Chua NH. Circadian clock regulates dynamic chromatin modifications associated with *Arabidopsis* *CCA1/LHY* and *TOC1* transcriptional rhythms. *Plant Cell Physiol*. 2012;53(12):2016–29.
- Malapeira J, Khaitova LC, Mas P. Ordered changes in histone modifications at the core of the *Arabidopsis* circadian clock. *Proc Natl Acad Sci U S A*. 2012;109(52):21540–5.
- Mizoguchi T, Wheatley K, Hanzawa Y, Wright L, Mizoguchi M, Song HR, Carre IA, Coupland G. LHY and CCA1 are partially redundant genes required to maintain circadian rhythms in *Arabidopsis*. *Dev Cell*. 2002;2(5):629–41.
- Wang ZY, Tobin EM. Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. *Cell*. 1998;93(7):1207–17.
- Alabadi D, Oyama T, Yanovsky MJ, Harmon FG, Mas P, Kay SA. Reciprocal regulation between TOC1 and LHY/CCA1 within the *Arabidopsis* circadian clock. *Science*. 2001;293(5531):880–3.
- Harmer SL, Kay SA. Positive and negative factors confer phase-specific circadian regulation of transcription in *Arabidopsis*. *Plant Cell*. 2005;17(7):1926–40.

19. Gendron JM, Pruneda-Paz JL, Doherty CJ, Gross AM, Kang SE, Kay SA. Arabidopsis circadian clock protein, TOC1, is a DNA-binding transcription factor. *Proc Natl Acad Sci U S A*. 2012;109(8):3167–72.
20. Huang W, Perez-Garcia P, Pokhilko A, Millar AJ, Antoshechkin I, Riechmann JL, Mas P. Mapping the core of the Arabidopsis circadian clock defines the network structure of the oscillator. *Science*. 2012;336(6077):75–9.
21. Pokhilko A, Mas P, Millar AJ. Modelling the widespread effects of TOC1 signalling on the plant circadian clock and its outputs. *BMC Syst Biol*. 2013; 7:23.
22. Dalchau N, Hubbard KE, Robertson FC, Hotta CT, Briggs HM, Stan GB, Goncalves JM, Webb AA. Correct biological timing in Arabidopsis requires multiple light-signaling pathways. *Proc Natl Acad Sci U S A*. 2010;107(29): 13171–6.
23. Harmer SL. The circadian system in higher plants. *Annu Rev Plant Biol*. 2009; 60:357–77.
24. Staiger D, Shin J, Johansson M, Davis SJ. The circadian clock goes genomic. *Genome Biol*. 2013;14(6):208.
25. Baerenfaller K, Shu H, Hirsch-Hoffmann M, Futterer J, Opitz L, Rehrauer H, Hennig L, Grissem W. Diurnal changes in the histone H3 signature H3K9ac[H3K27ac]H3S28p are associated with diurnal gene expression in Arabidopsis. *Plant Cell Environ*. 2016;39(11):2557–69.
26. Jones MA, Covington MF, DiTacchio L, Vollmers C, Panda S, Harmer SL. Jumonji domain protein JMJ5 functions in both the plant and human circadian systems. *Proc Natl Acad Sci U S A*. 2010;107(50):21623–8.
27. Ni Z, Kim ED, Ha M, Lackey E, Liu J, Zhang Y, Sun Q, Chen ZJ. Altered circadian rhythms regulate growth vigour in hybrids and allopolyploids. *Nature*. 2009;457(7227):327–31.
28. Berr A, McCallum EJ, Menard R, Meyer D, Fuchs J, Dong A, Shen WH. Arabidopsis SET DOMAIN GROUP2 is required for H3K4 trimethylation and is crucial for both sporophyte and gametophyte development. *Plant Cell*. 2010;22(10):3232–48.
29. Tamada Y, Yun JY, Woo SC, Amasino RM. ARABIDOPSIS TRITHORAX-RELATED7 is required for methylation of lysine 4 of histone H3 and for transcriptional activation of FLOWERING LOCUS C. *Plant Cell*. 2009;21(10): 3257–69.
30. Berr A, Shafiq S, Pinon V, Dong A, Shen WH. The trxG family histone methyltransferase SET DOMAIN GROUP 26 promotes flowering via a distinctive genetic pathway. *Plant J*. 2015;81(2):316–28.
31. Cartagena JA, Matsunaga S, Seki M, Kurihara D, Yokoyama M, Shinozaki K, Fujimoto S, Azumi Y, Uchiyama S, Fukui K. The Arabidopsis SDG4 contributes to the regulation of pollen tube growth by methylation of histone H3 lysines 4 and 36 in mature pollen. *Dev Biol*. 2008;315(2):355–68.
32. Lu F, Cui X, Zhang S, Liu C, Cao X. JMJ14 is an H3K4 demethylase regulating flowering time in Arabidopsis. *Cell Res*. 2010;20(3):387–90.
33. Yang H, Mo H, Fan D, Cao Y, Cui S, Ma L. Overexpression of a histone H3K4 demethylase, JMJ15, accelerates flowering time in Arabidopsis. *Plant Cell Rep*. 2012;31(7):1297–308.
34. Yang H, Han Z, Cao Y, Fan D, Li H, Mo H, Feng Y, Liu L, Wang Z, Yue Y, et al. A companion cell-dominant and developmentally regulated H3K4 demethylase controls flowering time in Arabidopsis via the repression of FLC expression. *PLoS Genet*. 2012;8(4):e1002664.
35. Chen Q, Chen X, Wang Q, Zhang F, Lou Z, Zhang Q, Zhou DX. Structural basis of a histone H3 lysine 4 demethylase required for stem elongation in rice. *PLoS Genet*. 2013;9(1):e1003239.
36. Guo L, Yu Y, Law JA, Zhang X. SET DOMAIN GROUP2 is the major histone H3 lysine [corrected] 4 trimethyltransferase in Arabidopsis. *Proc Natl Acad Sci U S A*. 2010;107(43):18557–62.
37. Berr A, McCallum EJ, Alioua A, Heintz D, Heitz T, Shen WH. Arabidopsis histone methyltransferase SET DOMAIN GROUP8 mediates induction of the jasmonate/ethylene pathway genes in plant defense response to necrotrophic fungi. *Plant Physiol*. 2010;154(3):1403–14.
38. Le Masson I, Jauvin V, Bouteiller N, Rivard M, Elmayan T, Vaucheret H. Mutations in the Arabidopsis H3K4me2/3 demethylase JMJ14 suppress posttranscriptional gene silencing by decreasing transgene transcription. *Plant Cell*. 2012;24(9):3603–12.
39. Nagel DH, Doherty CJ, Pruneda-Paz JL, Schmitz RJ, Ecker JR, Kay SA. Genome-wide identification of CCA1 targets uncovers an expanded clock network in Arabidopsis. *Proc Natl Acad Sci U S A*. 2015;112(34):E4802–10.
40. Lu SX, Knowles SM, Webb CJ, Celaya RB, Cha C, Siu JP, Tobin EM. The Jumonji C domain-containing protein JMJ30 regulates period length in the Arabidopsis circadian clock. *Plant Physiol*. 2011;155(2):906–15.
41. Mockler TC, Michael TP, Priest HD, Shen R, Sullivan CM, Givan SA, McEntee C, Kay SA, Chory J. The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis. *Cold Spring Harb Symp Quant Biol*. 2007;72:353–63.
42. Greenberg MV, Deleris A, Hale CJ, Liu A, Feng S, Jacobsen SE. Interplay between active chromatin marks and RNA-directed DNA methylation in Arabidopsis thaliana. *PLoS Genet*. 2013;9(11):e1003946.
43. Chen LQ, Luo JH, Cui ZH, Xue M, Wang L, Zhang XY, Pawlowski WP, He Y. ATX3, ATX4, and ATX5 encode putative H3K4 methyltransferases and are critical for plant development. *Plant Physiol*. 2017;174(3):1795–806.
44. Jenuwein T, Allis CD. Translating the histone code. *Science*. 2001;293(5532): 1074–80.
45. Ha M, Ng DW, Li WH, Chen ZJ. Coordinated histone modifications are associated with gene expression variation within and between species. *Genome Res*. 2011;21(4):590–8.
46. Ding Y, Avramova Z, Fromm M. Two distinct roles of ARABIDOPSIS HOMOLOG OF TRITHORAX1 (ATX1) at promoters and within transcribed regions of ATX1-regulated genes. *Plant Cell*. 2011;23(1):350–63.
47. Tian L, Fong MP, Wang JJ, Wei NE, Jiang H, Doerge RW, Chen ZJ. Reversible histone acetylation and deacetylation mediate genome-wide, promoter-dependent and locus-specific changes in gene expression during plant development. *Genetics*. 2005;169(1):337–45.
48. Tian L, Chen ZJ. Blocking histone deacetylation in Arabidopsis induces pleiotropic effects on plant gene regulation and development. *Proc Natl Acad Sci U S A*. 2001;98(1):200–5.
49. Park HS, Jun CH. A simple and fast algorithm for K-medoids clustering. *Expert Syst Appl*. 2009;36(2):3336–41.
50. Para A, Farre EM, Imaizumi T, Pruneda-Paz JL, Harmon FG, Kay SA. PRR3 is a vascular regulator of TOC1 stability in the Arabidopsis circadian clock. *Plant Cell*. 2007;19(11):3462–73.
51. Xie Q, Wang P, Liu X, Yuan L, Wang L, Zhang C, Li Y, Xing H, Zhi L, Yue Z, et al. LNK1 and LNK2 are transcriptional coactivators in the Arabidopsis circadian oscillator. *Plant Cell*. 2014;26(7):2843–57.
52. Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, Salemi MR, Phinney BS, Harmer SL. REVEILLE8 and PSEUDO-RESPONSE REGULATOR5 form a negative feedback loop within the Arabidopsis circadian clock. *PLoS Genet*. 2011;7(3): e1001350.
53. Farinas B, Mas P. Functional implication of the MYB transcription factor RVE8/LCL5 in the circadian control of histone acetylation. *Plant J*. 2011;66(2):318–29.
54. Ma Y, Gil S, Grasser KD, Mas P. Targeted recruitment of the basal transcriptional machinery by LNK clock components controls the circadian rhythms of nascent RNAs in Arabidopsis. *Plant Cell*. 2018;30(4):907–24.
55. Chen ZJ, Mas P. Interactive roles of chromatin regulation and circadian clock function in plants. *Genome Biol*. 2019;20(1):62.
56. Ko DK, Rohozinski D, Song Q, Taylor SH, Juenger TE, Harmon FG, Chen ZJ. Temporal shift of circadian-mediated gene expression and carbon fixation contributes to biomass heterosis in maize hybrids. *PLoS Genet*. 2016;12(7): e1006197.
57. Ning YQ, Ma ZY, Huang HW, Mo H, Zhao TT, Li L, Cai T, Chen S, Ma L, He XJ. Two novel NAC transcription factors regulate gene expression and flowering time by associating with the histone demethylase JMJ14. *Nucleic Acids Res*. 2015;43(3):1469–84.
58. Zhang S, Zhou B, Kang Y, Cui X, Liu A, Deleris A, Greenberg MV, Cui X, Qiu Q, Lu F, et al. C-terminal domains of a histone demethylase interact with a pair of transcription factors and mediate specific chromatin association. *Cell Discov*. 2015;1.
59. Hung FY, Chen FF, Li C, Chen C, Lai YC, Chen JH, Cui Y, Wu K. The Arabidopsis LDL1/2-HDA6 histone modification complex is functionally associated with CCA1/LHY in regulation of circadian clock genes. *Nucleic Acids Res*. 2018;46(20):10669–81.
60. Zhang F, Qi B, Wang L, Zhao B, Rode S, Riggan ND, Ecker JR, Qiao H. EIN2-dependent regulation of acetylation of histone H3K14 and non-canonical histone H3K23 in ethylene signalling. *Nat Commun*. 2016;7:13018.
61. Yamaguchi N, Winter CM, Wu MF, Kwon CS, William DA, Wagner D. PROTOCOLS: chromatin immunoprecipitation from Arabidopsis tissues. *Arabidopsis Book*. 2014;12:e0170.
62. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*. 2012;9(4):357–U354.
63. Zhang Y, Liu T, Meyer CA, Eeckhoutte J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, et al. Model-based analysis of ChIP-Seq (MACS). *Genome Biol*. 2008;9(9):R137.

64. Hughes ME, Hogenesch JB, Kornacker K. JTK\_CYCLE: an efficient nonparametric algorithm for detecting rhythmic components in genome-scale data sets. *J Biol Rhythm*. 2010;25(5):372–80.
65. Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW, Lane HC, Lempicki RA. The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol*. 2007;8(9):R183.
66. Song Q, Huang TY, Yu HH, Ando A, Mas P, Ha M, Chen ZJ. Diurnal regulation of SDG2 and JMJ14 by circadian clock oscillators orchestrates histone modification rhythms in Arabidopsis. Datasets. Sequence read archive. <https://trace.ncbi.nlm.nih.gov/Traces/sra/?study=SRP152291> (2019).

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

