# FIRST RECORD OF THE MAPLE GALLWASP, PEDIASPIS ACERIS (GMELIN, 1790) (CYNIPIDAE: PEDIASPIDINI) FROM TURKEY, WITH A NEW HOST RECORD

## Yusuf Katılmış\* and Musa Azmaz\*\*

- $^*$  Pamukkale University, Faculty of Arts & Sciences, Department of Biology, Denizli, TURKEY.
- \*\* Pamukkale University, Acıpayam Vocational High School, Department of Veterinary, Laboratory & Veterinary Health Program, Acıpayam, Denizli, TURKEY. E-mail: musazmaz@gmail.com

[Katılmış, Y. & Azmaz, M. 2017. First record of the maple gallwasp, *Pediaspis aceris* (Gmelin, 1790) (Cynipidae: Pediaspidini) from Turkey, with a new host record. Munis Entomology & Zoology, 12 (1): 285-287]

ABSTRACT: In this study, we report a new record that *Pediaspis aceris* (Gmelin, 1790) belonging to the maple gallwasp (Pediaspidini), on *Acer hyrcanum* Fisch. & Mey. (Aceraceae) in Isparta. *P. aceris* is a new record for Cynipidae fauna of Turkey in tribe, genus and species levels. At the same time, *A. hyrcanum* is a new host record for this species. Geographical distribution, host and photos of the gall are given.

KEY WORDS: Cynipidae, Pediaspidini, new record, new host, Turkey

Although Cynipidae belonging to Cynipoidea is the largest family with roughly 1400 described species (Ronquist et al., 2015), Pediaspidini belonging to Cynipidae has only 2 species belonging to 2 genera (*Himalocynips*, *Pediaspis*). Both genera are monotypic. *Himalocynips vigintilis* Yoshimoto, 1970 is only known from Nepal (Eastern Palearctic) and there is no data about its biology (Liljeblad & Ronquist, 1998). *P. aceris* as a European species is significant because it is the only gall wasp except Cynipini to show both sexual and asexual generations, a life cycle trait that has either evolved independently in the Pediaspidini, or indicates a sister group relationship between Cynipini and Pediaspidini. Moreover, both generations of *P. aceris* induce gall on maple (*Acermonspessulanum* L., *A. platanoides* L., *A. pseudoplatanus* L. and *A. opalus* Miller) (Melika, 2006).

Cynipidae fauna of Turkey is represented by seven taxa (Aulacideini, Aylacini, Ceroptresini, Cynipini, Diastrophini, Diplolepidini, Synergini). Approximately 150 species belonging to these taxa were totally recorded from Turkey according to faunistic studies in last decade (Kemal & Koçak, 2010; Kıyak & Katılmış, 2010; Katılmış & Kıyak, 2008, 2009a,b, 2010, 2011a,b,c,d, 2012a,b; Mutun & Dinç, 2011, 2015; Mete & Demirsoy, 2012; Çetin et al., 2014; Dinç et al., 2014; Mutun et al., 2014; Azmaz & Katılmış, 2015; Katılmış & Azmaz, 2015).

## MATERIALS AND METHODS

All gall specimens on host plant were collected from Isparta in 2010 and 2015. After photos of the gall specimens were taken, data about locality and host plant were recorded. Galls were kept in laboratory conditions and checked weekly for emerged wasps. Adult gall wasps were subsequently fixed in ethanol. The adults were then air-dried, fixed on cards (5x10 mm), and pinned. The terminology used to describe gall wasp morphology follows Melika (2006). The gall specimens and adults were deposited in the Pamukkale University, Faculty of Arts & Sciences, Department of Biology, Entomology Laboratory, Denizli, Turkey.

## RESULTS

## Tribe Pediaspidini Ashmead, 1903 Genus *Pediaspis* Tischbein, 1852 Species *Pediaspis aceris* (Gmelin, 1790)

**Material examined:** ISPARTA, Eğirdir, above Yukarıgökdere village, Davraz mountain, Kasnak Oak Nature Protection Area, 37°44′ N, 30°49′ E, 1540 m, collected 02.VI.2010, 11.VI.2015, 29, 10°, on *A. hyrcanum* (Fig. 1).

**Host:** Acer hyrcanum is a new host record as a maple tree.

**Distribution:** Known from Austria, Belgium, France, Germany, Georgia, Hungary, Iberian Peninsula, Iran, Romania, Russia, Switzerland, Ukraine (Melika, 2006).

### DISCUSSION

As faunistic studies about gall wasps in Turkey include mainly oak gall wasps (Cynipini) on oaks, *P. aceris* that known only from Western Palearctic as a European species, has not been unfortunately recorded from Turkey so far. As a result of this study, tribe Pediaspidini was recorded as a new taxon for Cynipidae fauna of Turkey. Furthermore, a new host record (*A. hyrcanum*) was provide for *P. aceris*. We predict that species richness of the Turkish fauna will increasingly continue with faunistic studies.

### ACKNOWLEDGEMENT

We would like to thank Dr. Mehmet Çiçek, that provided support for the identification of the plant species.

## LITERATURE CITED

- Azmaz, M. & Katılmış, Y. 2015. Four new records of Oak Gall Wasp (Hymenoptera: Cynipidae, Cynipini) from Turkey. Munis Entomology & Zoology, 10: 201-204.
- Çetin, G., Orman, E. & Polat, Z. 2014. First record of the Oriental chestnut gall wasp, Dryocosmus kuriphilus Yasumatsu (Hymenoptera: Cynipidae) in Turkey. Bitki Koruma Bülteni, 54: 303-309.
- Dinc, S., Mutun, S. & Melika, G. 2014. A new species of Andricus Hartig oak gallwasp from Turkey (Hymenoptera: Cynipidae, Cynipini). North-West. J. Zool., 10: 122-127.
- Katılmış, Y. & Azmaz, M. 2015. Investigation on the inquilines (Hymenoptera: Cynipidae, Synergini) of oak galls from inner western Anatolia, Turkey. Turk. J. Zool., 39: 168-173.
- Katılmış, Y. & Kıyak, S. 2008. Checklist of Cynipidae of Turkey, with a new genus record. J. Nat. Hist., 42: 2161-2167. Katılmış, Y. & Kıyak, S. 2009a. Oak gallwasp Genus Andricus (Hymenoptera, Cynipidae) new records from Turkey.
- Zool. Middle East, 48: 108-110.

  Katılmış, Y. & Kıyak, S. 2009b. The oak gallwasp Aphelonyx persica: a new record from Turkey, with some new host records. Phytoparasitica, 37: 95-97.
- Katılmış, Y. & Kıyak, S. 2010. Distribution, Phenology and Effects of *Diplolepis* spp. (Hymenoptera: Cynipidae) on *Rosa camina* in the Inland Western Anatolian. J. Entomol. Res. Soc., 12: 31-36.
- Katılmış, Y. & Kıyak, S. 2011a. Oak gallwasps (Hymenoptera: Cynipidae) fauna of inner western Anatolia. Munis Entomology & Zoology, 6: 735-757.
- Katılmış, Y. & Kıyak, S. 2011b. Herb gallwasp *Neaylax salviae* (Giraud, 1859) (Hymenoptera, Aylacini) new record from Turkey. Turk. J. Zool., 35: 707-710.
- Katılmış, Y. & Kıyak, S. 2011c. New records of herb gallwasps (Hymenoptera, Cynipidae, Aylacini) from Turkey. North-West. J. Zool., 7: 17-19.
- Katılmış, Y. & Kıyak, S. 2011d. First records of inquiline wasps of the genus Ceroptres Hartig, 1840 (Hymenoptera, Cynipidae, Synergini) from Turkey. J. Entomol. Res. Soc., 13: 65-69.
- Katılmış, Y. & Kıyak, S. 2012a. Further study on Periclistus brandtii (Ratzeburg, 1831) (Hymenoptera, Cynipidae) from Turkey. Entomol. News, 122: 51-54.
- Katılmış, Y. & Kıyak, S. 2012b. Two new records of inquiline wasp of the genus Saphonecrus Dalla Torre and Kieffer, 1910 (Hymenoptera: Cynipidae: Synergini) from Turkey, their associated galls and hosts. Munis Entomology & Zoology, 7: 1141-1144.
- Kemal, M. & Koçak, A. Ö. 2010. Winter trips to south Van Lake and the cynipid galls on oaks (Hymenopteera, Cynipidae). Cesa News, 56: 1-66.
- Kıyak, S. & Katılmış, Y. 2010. Oak gallwasp Dryocosmus mayri (Hymenoptera, Cynipidae)- new record from Turkey. J. Entomol. Res. Soc., 12: 67-70.
- Liljeblad, J. & Ronquist, F. 1998. A phylogenetic analysis of higher-level gall wasps relationships (Hymenoptera: Cynipidae). Systematic Entomology, 28: 229-252.

Melika, G. 2006. Gall Wasps of Ukraine. Cynipidae. Vestnik zoologii, 21: 1-644.

Mete, Ö. & Demirsoy, A. 2012. A Preliminary Study on The Gallwasp Fauna of Kemaliye (Erzincan, Turkey) and A New Record for Turkey. Hacettepe J. Biol. & Chem., Special Issue, 351-363.

Mutun, S. & Dinc, S. 2011. Contributions to the Gallwasp (Hymenoptera: Cynipidae) Fauna of Turkey with One New Record. J. Appl. Biol. Sci., 5: 83-85.

Mutun, S. & Dinc, S. 2015. Twelve oak gall wasp species (Hymenoptera, Cynipidae) new to the Turkish fauna. Turk. J.

Muttin, S. & Dift, S. 2015. Iweive oak gair wasp species (Hymenoptera, Cympolacy, Nov. 13 and 12 and 12 and 13 and 13 and 13 and 14 and 14 and 14 and 15 and 16 a



Figure 1. Pediaspis aceris sexual galls on leaf (a) and on fruit (b) (a scale bar 1 cm).