

## Subjective wellbeing, sense of humor and psychological health in hemodialysis patients

Luis Manuel Mota Sousa<sup>1</sup>, Ana Vanessa Antunes<sup>2</sup>, Cristina Maria Alves Marques-Vieira<sup>3</sup>, Paulo César Lopes Silva<sup>4</sup>, Olga Maria Martins de Sousa Valentim<sup>5</sup>, Helena Maria Guerreiro José<sup>6</sup>

<sup>1</sup> Ortopedia, Hospital Curry Cabral del Centro Hospitalar Lisboa Central, Lisboa. Portugal y Escuela Superior de Salud Atlántica, Oeiras. Portugal

<sup>2</sup> Escuela Superior de Salud Egas Moniz, Almada. Portugal

<sup>3</sup> Instituto de Ciências de la Salud, Universidade Católica Portuguesa, Lisboa. Portugal e Investigador CIIS. Portugal

<sup>4</sup> Unidad de Terapia Intensiva, Hospital José Joaquim Fernandes, Unidad Local de Salud del Baixo Alentejo, Beja. Portugal

<sup>5</sup> Escuela Superior de Salud Atlántica, Oeiras. Portugal

<sup>6</sup> Universidad de Algarve, Escuela Superior de Salud y UICISA-E, Coimbra. Portugal

### Abstract

**Objective:** To investigate the relationship between satisfaction with life in general, sense of humor, and anxiety, depression and stress with subjective happiness in chronic kidney disease patients undergoing hemodialysis.

**Material and Methods:** A cross-sectional and correlational study was developed in two units of the Diaverum dialysis clinic and one hospital unit, with 183 chronic kidney disease patients undergoing hemodialysis. An instrument was used to characterize the sociodemographic and clinical profile of the sample (age, gender, nationality, education, occupation, marital status, dialysis sessions length, presence of hypertension and diabetes): the subjective happiness scale; the satisfaction with life in general; depression, anxiety and stress scale 21; and multidimensional sense of humor scale. Inferential procedures included Spearman correlation coefficients, and multiple linear regression adjusting to age, marital status, professional activity and diabetes.

**Results:** Subjective Happiness was positively correlated with satisfaction with life in general, and the three dimensions of Sense of Humor. Nevertheless, subjective happiness was negatively correlated with stress / anxiety and depression. Satisfaction with life in general, humor production and social use of humor, and attitude towards humor had a positive relationship with subjective happiness. However, depression had a negative relationship with subjective happiness.

**Conclusions:** Higher levels of subjective happiness were associated with higher levels of satisfaction with life in general, and sense of humor, however they were also associated with lower levels of depression in patients undergoing hemodialysis.

**KEYWORDS:** renal dialysis; happiness; wit and humor as topic; depression.

### Introduction

Hemodialysis (HD) is the main treatment for patients in end-stage renal disease, however, it presents a high risk for adverse patient outcomes<sup>1</sup>. HD affects the life of Chronic kidney disease (CKD) patients and it can have negative effects on general health and well-being including mental, social, physical, emotional, spiritual, and subjective aspects. Furthermore, it can have a negative influence on physical performance, mental

#### Correspondencia:

Luis Manuel Mota Sousa

Universidade de Évora, Escola Superior de Enfermagem de São João de Deus.

Largo do Senhor da Pobreza, 7000-811 Évora. Portugal

E-mail: [luismmosousa@gmail.com](mailto:luismmosousa@gmail.com)

status, and social relationships<sup>2</sup>. Anxiety, depression and stress are frequent conditions in CKD patients, essentially during the transition from the pre-dialysis period to renal replacement therapy. Nevertheless, it tends to decrease after beginning dialysis<sup>3</sup>.

The World Health Organization (WHO) points out that subjective wellbeing is considered within the research community as an indicator of Quality of Life evaluation<sup>4</sup>. The subjective wellbeing is characterized by emotional responses, satisfaction domains, and the judgment about global satisfaction with life. The subjective wellbeing components are pleasant affect (e.g. joy, contentment, pride, affection, and happiness), unpleasant affect (e.g. guilt and shame, anxiety and worry, anger, stress, and depression), life satisfaction (e.g. desire to change, satisfaction with current life, past, and future), and a satisfaction domain (e.g. work, family, leisure, health, finances, and self)<sup>5</sup>.

In the CKD patients the subjective wellbeing (measured by personal wellbeing index) is positively related with subjective happiness, positive affect, and quality of life and is negatively associated with negative affect. Subjective happiness, negative affect, and the physical component of quality of life are associated to the subjective wellbeing<sup>6</sup>. Moreover, higher levels of happiness are associated with higher levels of satisfaction with life in general (personal wellbeing index)<sup>7</sup>, and also with higher levels of sense of humor<sup>8</sup>.

This study intends to disclose if CKD patients' subjective well-being, sense of humor, and anxiety, depression and stress are significant predictors of their subjective happiness, further influencing patients' disease management strategies and health status.

The objective was to investigate the relationship between subjective well-being, sense of humor, and anxiety, depression and stress with subjective happiness in CKD patients undergoing hemodialysis.

## Material and Methods

### Study design

A cross-sectional and correlational study, developed in two units of the Diaverum Dialysis Clinic and one hospital unit in the Lisbon region, Portugal, with CKD patients undergoing HD between May and June 2015.

### Subjects and setting

The inclusion criteria defined were: people aged 18 years or over undergoing HD routinely for at least six months. Exclusion criteria were: people with active psychiatric illness, cognitive impairment, and non-compensated visual or hearing impairment. Information regarding these aspects was confirmed in each person's clinical records. 253 CKD patients met the eligibility criteria (139 in Clinic 1, 114 in Clinic 2, and 15 in the hospital unit). The sample calculation was set out with a Confidence Interval of 95% and sample error of 5%. Afterwards, a random selection without replacement was made. A randomized sample of 183 people undergoing HD was selected from the dialysis clinics, 93 of clinic 1, 78 of clinic two and 12 of the hospital unit.

### Procedures

Interviews were performed during the HD session by five trained nurses. The main researcher previously met with these nurses to explain the objectives and how to collect the data. Afterwards a written roadmap was provided to assist in completing the data collection instruments.

Various instruments were used to characterize the socio demographic and clinical profile of the sample (age, gender, nationality, education, occupation, marital status, dialysis sessions length, presence of hypertension and diabetes), the subjective happiness scale (SHS)<sup>7</sup>; the satisfaction with life in general (SWLG) was measured by the Personal Wellbeing Index<sup>9</sup>; Depression, Anxiety and Stress Scale 21 (DASS – 21)<sup>10</sup>; and Multidimensional Sense of Humor Scale (MSHS)<sup>8</sup>.

### Outcomes measurement

The SHS consists of four items, in which participants are asked to self-characterize themselves compared to their peers (items two and three) and classify their levels of happiness and unhappiness (items one and four). The last item score is reversed. The SHS uses a visual analogue scale with seven positions based on two antagonistic statements which express the level of happiness or lack of it. The application of the Portuguese scale to CKD patients shows a single factor with an internal reliability of Cronbach's  $\alpha$  of 0.90<sup>7</sup>. Satisfaction with life in general or the personal wellbeing index is made of seven domains for the overall measure of life-satisfaction (satisfaction with standard of living, health, personal development, personal relationships, sense of security, connection to the community, and security for the future). For each domain respondents were asked to classify their satisfaction within a scale from 0 (extremely dissatisfied) to 10 (very satisfied) with a neutral intermediate position. The

personal wellbeing index was measured on a scale ranging from 0 to 100 (maximum percentage of the scale). The application of the Portuguese scale to CKD patients shows the existence of a single factor, with internal reliability of Cronbach's  $\alpha$  of 0.82<sup>9</sup>. DASS-21 was organized into three subscales: depression, anxiety and stress, each consisting of seven items. The Portuguese version of DASS 21 in CKD patients presents a bi-dimensional structure, "Depression" and "Stress/Anxiety", with Cronbach's  $\alpha$  values of 0.86 for "Depression" and 0.88 for "Stress/Anxiety"<sup>10</sup>.

The MSHS structure was globally consistent with studies conducted in other samples, presenting satisfactory internal consistency values. It is considered a valid instrument for characterizing an individual's "state of humor" and to describe its different dimensions. The Portuguese version of MSHS showed a structure with three factors, "Humor Production and Social Use of Humor", "Adaptive Humor and Appreciation Humor" and "Attitude Towards Humor", with Cronbach's  $\alpha$  values of 0.93, 0.90 and 0.83 respectively<sup>8</sup>.

### Ethical approval

Approval was received from the ethics committee of Diaverum (Approval N° 1/2015) as from the ethics committee of the Hospital Center (Approval No 175/2015). All participants signed the informed consent form after being informed about the guarantee of confidentiality of their data and the right of withdrawal without risk to oneself. Informed consent was therefore obtained from those who met the inclusion criteria and agreed to participate.

### Data analysis

Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS), version 23.0. Descriptive statistics are reported as frequency, percentage, mean, and standard deviations were calculated using two-tailed Student's t-test for pairwise comparisons, one-way analysis of variance (ANOVA) for multiple comparisons, and two-way ANOVA for multiple comparisons involving two independent variables. ANOVA analyses were subjected to Bonferroni's post-hoc test. Inferential procedures included Spearman correlation coefficients, and multiple linear regression with control of age, marital status, professional activity and diabetes. The significance level was set at  $p < 0.05$ .

## Results

The sample consisted of 183 patients diagnosed with CKD, with an average age of 59.17 ( $\pm 14.64$ ) years, mostly men (59.6%). The nationality of the patients was distributed as follows: most patients are Portuguese (78.7%), followed by Cape Verdean (13.7%), Sao Tomean (3.3%), Angolan (2.2%), Guinean (1.6%), and Bulgarian (0.5%). Concerning literacy level, 3.3% were illiterate, 41.1% had the 4<sup>th</sup> grade, 18.9% the 6<sup>th</sup> grade, 15% the 9<sup>th</sup> grade, 12.2% the 12<sup>th</sup> grade and 9.4% university education. In this sample 28% of the patients were single, 53.8% married, 11.5% widowed and 6.6% divorced. Concerning occupation, 76% were retired while 24% had a regular job. Regarding health data, patients underwent hemodialysis for 70.09 ( $\pm 54.2$ ) months, 61.9% had high blood pressure and 25.8% diabetes.

CKD patients under 63 years of age had higher levels of subjective happiness ( $p = 0.013$ ) than older patients. Portuguese CKD patients had the lowest score for subjective happiness ( $p = 0.013$ ), compared with the remaining population (Cape Verdean; São Toméan; Angolan, Guinean and Bulgarian). Retired patients had lower levels of subjective happiness ( $p = 0.002$ ) than those who have regular jobs. Widowed and divorced patients had lower levels of subjective happiness ( $p = 0.033$ ) in comparison to married and single patients (Bonferroni correction). Finally, CKD patients with diabetes had a lower level of subjective happiness ( $p = 0.012$ ). **Table 1** shows both socio demographic and clinical factors related with CKD, that are associated with subjective happiness.

Mean score of subjective happiness was 19.9 ( $\pm 6.1$ ), satisfaction with life in general/personal wellbeing index has a mean score of 64.2% ( $\pm 18.5\%$ ). Mean scores for the other variables are as following: stress/anxiety 6.9 ( $\pm 6.5$ ); depression 4.3 ( $\pm 4.4$ ); humor production and social use of humor 37.7 ( $\pm 11.2$ ); adaptive humor and appreciation humor 27.1 ( $\pm 6.1$ ) and attitude towards humor with 24.7 ( $\pm 5.1$ ) (**Table 2**).

**Table 3** shows that the subjective happiness was positively correlated with professional activity ( $\rho = 0.256$ ,  $p < 0.01$ ); satisfaction with life in general/personal wellbeing index ( $\rho = 0.627$ ,  $p < 0.001$ ), humor production and social use of humor ( $\rho = 0.476$ ,  $p < 0.001$ ); adaptive humor and appreciation humor ( $\rho = 0.387$ ,  $p < 0.001$ ); and attitude towards humor ( $\rho = 0.364$ ,  $p < 0.001$ ). However, it presented a low negative correlation with stress/anxiety ( $\rho = -0.239$ ,  $p < 0.01$ ) and depression ( $\rho = -0.391$ ,  $p < 0.001$ ). Higher scores on subjective happiness were associated

**Table 1.** Socio demographic and clinical factors associated to subjective happiness.

	Subjective Happiness	p-value
<b>Gender</b>		
Male	19.7±5.8	p=0.614
Female	20.2±6.6	
<b>Age</b>		
Under 63 years	21.1±6.2	p=0.013
More than 63 years	18.9±5.7	
<b>Nationality</b>		
Portuguese	19.3±6.2	p=0.013
Other	22.1±5.5	
<b>Professional activity</b>		
Retired	19.1±6.1	p=0.002
Active	22.4±5.5	
<b>Marital Status</b>		
Single	20.9±5.8	p=0,033
Married	20.1±6.1	
Other (widowed and divorced)	17.5±6.1	
<b>Arterial hypertension</b>		
No	20.5±5.8	p=0.332
Yes 19.9±6.2		
<b>Diabetes</b>		
No	20.5±5.9	p=0.012
Yes	17.9±6.3	
<b>Hemodialysis time</b>		
Less than five years	19.3±5.8	p=0.297
More than five years	20.3±6.4	

**Table 2.** Mean and standard deviation for the different variables.

	Range	Minimum	Maximum	Mean	Standard deviation
1. Subjective Happiness	4-28	4	28	19.9	6.1
2. Satisfaction with Life in General / Personal Wellbeing Index	0-100	6.3	100	64.2	18.5
3. Stress / Anxiety	0-42	0	34	6.9	6.5
4. Depression	0-21	0	18	4.3	4.4
5. Humor Production and Social Use of Humor	11-55	11	55	37.7	11.2
6. Adaptive Humor and Appreciation Humor	7-35	10	35	27.1	6.1
7. Attitude Towards Humor	6-30	6	30	24.7	5.1

with higher levels of satisfaction with life in general/personal wellbeing index, humor production and social use of humor, adaptive humor and appreciation humor and attitude towards humor, though lower levels of subjective happiness were associated with higher levels of stress/anxiety and depression.

Professional activity was negatively correlated with depression ( $\rho=-0.222$ ,  $p<0.001$ ), satisfaction with life in general/personal wellbeing index was positively correlated with humor production and social use of humor ( $\rho=0.353$ ,  $p<0.001$ ); adaptive humor and appreciation humor ( $\rho=0.270$ ,  $p<0.001$ ) and attitude towards humor ( $\rho=0.211$ ,  $p<0.001$ ). Nevertheless, it presented a negative correlation with stress/anxiety ( $\rho=-0.297$ ,  $p<0.001$ ) and depression ( $\rho=-0.440$ ,  $p<0.001$ ). Higher scores on satisfaction with life in general/personal wellbeing index were associated with higher levels of humor production and social use of humor, adaptive humor and appreciation humor, and attitude towards humor, and lower levels of stress/anxiety and depression.

Stress/anxiety showed a significant positive correlation with depression ( $\rho=0.670$ ,  $p<0.001$ ) and attitude towards humor ( $\rho=0.232$ ,  $p<0.01$ ).

Depression was negatively correlated with humor production, with social use of humor ( $\rho=-0.164$ ,  $p<0.05$ ) and with attitude towards humor ( $\rho=-0.240$ ,  $p<0.01$ ). When depression values increased, humor production and social use of humor and attitude towards humor values decreased.

Humor production and social use of humor showed a significant positive correlation with adaptive humor and humor appreciation ( $\rho=0.718$ ,  $p<0.001$ ); and attitude towards humor ( $\rho=0.234$ ,  $p<0.001$ ). Lastly, adaptive humor and humor appreciation were positively correlated with attitude towards humor ( $\rho=0.338$ ,  $p<0.001$ ).

The model with professional activity, explains 8% of the subjective happiness variance. This control variable is the one that most contributes to the explanation of subjective happiness ( $\Delta R^2=0.115$ ,  $p<0.01$ ).

In the second step, the adjusted  $R^2$  for the model was 51.4% for professional activity, satisfaction with life in general/personal wellbeing index, stress/anxiety, depression, humor production and social use of humor, and attitude towards humor, being these the variables

**Table 3.** Regression for Subjective Happiness with other variables and correlations.

	1	2	3	4	5	6
1. Subjective Happiness						
2. Satisfaction with Life in General/Personal Wellbeing Index	0.627‡					
3. Stress / Anxiety	-0.239†	-0.297‡				
4. Depression	-0.391‡	-0.440‡	0.670‡			
5. Humor Production and Social Use of Humor	0.476‡	0.353‡	-0.095	-0.164*		
6. Adaptive Humor and Appreciation Humor	0.387‡	0.270‡	-0.123	-0.131	0.718‡	
7. Attitude Towards Humor	0.364‡	0.211‡	0.232†	-0.240†	0.234†	0.338‡

\*Significance p<0.05, †Significance p<0.01, ‡Significance p<0.001.

that best explain subjective happiness. The resulting R<sup>2</sup> were statistically significant at the p<0.00 and p<0.05 levels.

Professional activity, satisfaction with life in general/personal wellbeing index, humor production and social use of humor, and attitude towards humor had a positive effect on subjective happiness (respectively, β=0.142, p<0.05; β=0.430, p<0.001; β=0.239, p<0.05; β=0.165, p<0.01). Nevertheless, depression had a negative effect on subjective happiness (β=-0.178, p<0.05). These variables are the ones contributing the most to the explanation of subjective happiness (ΔR<sup>2</sup>=0.431, p<0.001).

### Discussion

Our study was aimed at examining the relationship between subjective happiness and satisfaction with life in general/personal wellbeing index, stress/anxiety and depression, sense of humor and socio-demographic characteristics, as well as, verifying the relationship between these variables and subjective happiness.

The findings are aligned with the literature on the effects of HD on CKD patients' life, and on the psychological stressors that may decrease wellbeing and quality of life, therefore contributing to less positive clinical outcomes<sup>11</sup>.

**Table 4.** Multiple linear regression adjusted for variables associated with the Subjective Happiness.

Predictors	Subjective Happiness			
	Step 1		Step 2	
	β	t	β	T
Constant		14.354		0.609
1. Age	0.003	0.030	-0.021	-0.314
2. Nationality	-0.047	-0.565	0.049	0.777
3. Professional activity	0.221	2.655†	0.142	2.281*
4. Marital status	-0.111	-1.315	-0.067	-1.040
2. Diabetes	-0.142	-1.812	-0.099	-1.684
3. Satisfaction with Life in General /Personal Wellbeing Index			0.430	6.553‡
4. Stress / Anxiety			0.115	1.427
5. Depression			-0.178	-2.077*
6. Humor Production and Social Use of Humor			0.239	2.866*
7. Adaptive Humor and Appreciation Humor			0.005	0.067
8. Attitude Towards Humor			0.165	2.790†
F	4.259†		17.268‡	
Adj. R-Sq	0.088		0.514	
R-Sq Change	0.115		0.431	

\*Significance p<0.05, †Significance p<0.01, ‡Significance p<0.001.



The highest levels of happiness were found in younger CKD patients with regular employment, married or single, non-Portuguese nationality and without diabetes. These results also corroborate the findings of another study carried out in patients undergoing hemodialysis, namely, standard of living and satisfaction with life achievements carried higher importance in subjective wellbeing<sup>12</sup>.

Such results can also be partly explained by the aging process. Older people patients on hemodialysis have peculiar characteristics, not only because of the greater number of associated comorbidities, but because the aging process itself entails physical and psychological changes that directly affect quality and life. Other aspects must also be considered as high stress factors during hemodialysis sessions<sup>13</sup>, such as the vascular access, which includes the cannulation process, bleeding, infection, lack of flow, and the inexperience of the team. Having a job minimizes the impact of the treatment routine, it makes the individual feel useful and socially active, which explains higher levels of personal fulfillment and greater happiness. One study points out that by the end of 2015, approximately 36% of patients undergoing peritoneal dialysis had full-time work, contrasting with 22% of hemodialysis patients<sup>14</sup>.

Despite the impact of treatment on the personal routine, married patients find in their family nuclei an informal but more solid support network, which strengthens their emotional component and eventually their perception of happiness. On the other hand, widowers under hemodialysis experience more hopelessness and had lower level of general health<sup>15</sup>, probably because they have nobody to share their sorrows, maintaining their feelings on the burden of disease<sup>16</sup>. These patients consequently have greater difficulty in finding adaptive strategies that allow maintaining or increasing their levels of happiness.

Concerning nationality, the non-Portuguese patients of our sample are predominantly African. The higher levels of happiness found in this population may be explained by cultural issues, as these individuals live in larger family and social networks with expanded support. This idea is supported, in part, by a study that observed a level of low contentment, and medium to high levels of hedonic affection in a group of black Africans<sup>17</sup>.

Happiness can influence socioeconomic phenomena such as consumption activities, behavior at work, investment behavior and political behavior, presenting itself as a determining factor in person's quality of life<sup>18</sup>. In the

same way, happiness was associated with higher levels of satisfaction with life in general, humor production and social use of humor, adaptive humor and appreciation humor and attitude towards humor. Though lower levels of subjective happiness were associated with higher levels of stress/anxiety and depression. The psychological disorders are the most common problems in patients undergoing dialysis, such as depression and anxiety. Hemodialysis is considered a long-term treatment and has significant side effects on physical and mental well-being<sup>19</sup>. Stress is a common reaction of patients with CKD undergoing HD which appears as a way to deal with the anxiety associated with the threat of the disease. Since they are constantly alert, they are also more vulnerable and have more difficulty in keeping an emotional state of happiness. Such findings may have important clinical implications for these patients. Untreated mental problems increase mortality, decrease quality of life, and increase the risk of suicide<sup>19</sup>. Healthcare teams must consider the need for a periodic evaluation of the stressors and their effects on the health and quality of life of the patients, with the consequent development of strategies that offer these individuals interventions that propitiate an improvement in their quality of life.

Patients undergoing hemodialysis with higher levels of subjective happiness were found to be associated with higher levels of satisfaction with life in general<sup>9</sup>. CKD patients with higher levels of sense of humor were associated with higher levels of subjective happiness and satisfaction with life in general<sup>8</sup>.

Professional activity, satisfaction with life in general, humor production and social use of humor, and attitude towards humor had a positive effect on subjective happiness. Thus, depression had a negative effect on subjective happiness.

Fordyce's happiness program was effective in reducing depression and anxiety<sup>11,20</sup> and improving life of quality of patients undergoing hemodialysis<sup>20</sup>.

Keeping in mind the limitations and implications for practice of this study, we emphasize the need for further studies on nursing interventions under the promotion of happiness in the most diverse contexts of the practice of care for the person with chronic kidney disease. Its disclosure is imperative.

#### **Practical implications for nursing**

Our results suggest that CKD patients, professionally active, with lower levels of depression, with higher levels

of satisfaction with life in general/personal wellbeing index, and humor production and social use of humor, and attitude towards humor had higher levels of subjective happiness. We therefore recommend that nurses should actively participate in the process of identifying and minimizing stressors, thus favouring the treatment, quality of life and wellbeing of these patients.

Such results may also help future interventions related to the emotional wellbeing (subjective happiness) of CKD patients, particularly those aimed at improving nurse training and the identification of nursing diagnosis related to well-being and monitoring the effectiveness of nursing interventions. Nurses with this knowledge could be able to optimize their patients' health outcomes.

Depressive symptoms in CKD patients are associated with decreased wellbeing<sup>16</sup> and consequently decreased quality of life<sup>21-22</sup>.

In fact, with our data we can infer that when nurses implement interventions that increase the perception of subjective happiness, they can improve the patients' well-being and, consequently, their quality of life when patients undergoing hemodialysis however, further research is required. Several authors point out the Fordyce's happiness program<sup>11,20</sup>, the integration of laughter yoga<sup>23</sup>, the simulated laughter programme<sup>24</sup> or watching humor movies, according to the patients' preferences<sup>25</sup>, as important strategies to increase CKD patients' happiness.

### Limitations

Some limitations can be identified. Both units involved in this study are in the same geographical region influencing socio demographic characteristics, which prevents the generalization of conclusions. Also, the data collection environment (HD room) can work as a distraction factors for CKD patients. Finally, the small sample size might limit the external validity of the results.

From the results found we can conclude about the relationship between subjective well-being, sense of humor, and psychological health with subjective happiness in CKD patients. Satisfaction with life in general/personal wellbeing index, humor production and social use of humor, and attitude towards humor had a positive relationship with subjective happiness. However, depression had a negative relationship with subjective happiness.

Recibido: 26 septiembre 2018  
 Revisado: 4 enero 2019  
 Modificado: 20 febrero 2019  
 Aceptado: 25 febrero 2019

### References

1. Rivara MB, Chen CH, Nair A, Cobb D, Himmelfarb J, Mehrotra R. Indication for dialysis initiation and mortality in patients with chronic kidney failure: a retrospective cohort study. *Am J Kidney Dis.* 2017;69(1):41-50.
2. Seraji M, Shojaeizadeh D, Rakhshani F. Well-being in Hemodialysis Patients. *Iran J Public Health.* 2018;47(8):1222-3.
3. Bezerra CI, Silva BC, Elias RM. Decision-making process in the pre-dialysis CKD patients: do anxiety, stress and depression matter? *BMC Nephrol.* 2018;19(1):98.
4. Weinberg MK, Bennett PN, Cummins RA. Validation of the personal wellbeing index for people with end stage kidney disease. *Appl Res Qual Life.* 2016;11(4):1227-40.
5. Diener E, Suh EM, Lucas RE, Smith HL. Subjective well-being: Three decades of progress. *Psychol Bull.* 1999;125(2):276-302.
6. Sousa LM, Antunes AV, Baixinho CR, Severino SS, Marques-Vieira C, José HM. Subjective Wellbeing Assessment in People with Chronic Kidney Disease Undergoing Hemodialysis. *In T Rath (Ed.) Chronic Kidney Disease - from Pathophysiology to Clinical Improvements. InTech;* 2018:281-93 DOI: 10.5772/intechopen.71194.
7. Sousa LM, Marques-Vieira CM, Severino SS, Pozo-Rosado JL, José HM. Validation of the Subjective Happiness Scale in people with Chronic Kidney Disease. *Enferm Global.* 2017;16(3):60-70.
8. Sousa LMM, Marques-Vieira CM, Severino S, Pozo-Rosado J, Antunes AV, José HM. Validation of the multidimensional sense of humor scale in people with chronic kidney disease. *J Nurs Educ Pract.* 2018;8(3):72-9.

9. Sousa LMM, Marques-Vieira CMA, Severino SSP, Pozo Rosado JL, José HM. Validación del Índice de Bien-estar Personal en personas con enfermedad renal crónica. *Enferm Nefrol.* 2016;19(2):135-41.
10. Sousa LM, Marques-Vieira CM, Severino SS, Co-reo JL, Gomes JC, José HM. A Depression Anxiety Stress Scale em pessoas com doença renal crónica. *Rev Port Enferm Saúde Ment.* 2017;17:50-7.
11. Mehrabi Y, Ghazavi Z, Shahgholian N. Effect of fordyce's happiness program on stress, anxiety, and depression among the patients undergoing hemodialysis. *Iran J Nurs Midwifery Res.* 2017; 22(3):190-4.
12. Bennett PN, Weinberg MK, Bridgman T, Cummins RA. The happiness and subjective well being of people on haemodialysis. *J Ren Care.* 2015;41(3):156-61.
13. Casey JR, Hanson CS, Winkelmayr WC, Craig JC, Palmer S, Strippoli GF, et al. Patients' perspectives on hemodialysis vascular access: a systematic review of qualitative studies. *Am J Kidney Dis.* 2014;64(6):937-53.
14. Jin DC, Yun SR, Lee SW, Han SW, Kim W, Park J. Current characteristics of dialysis therapy in Korea: 2015 registry data focusing on elderly patients. *Kidney Res Clin Pract.* 2016;35(4):204-11.
15. Shakya D, Tuladhar J, Poudel S. Burden and Depression among Caregivers of Hemodialysis Patients. *Palliat Med Care.* 2017;4(1):1-6.
16. Ercan F, Demir S. Hopelessness and Quality of Life Levels in Hemodialysis Patients. *Gazi Med J.* 2018;29(3):169-174.
17. Brulé G, Veenhoven R. Geography of happiness: configurations of affective and cognitive appraisal of life across nations. *Int J Happiness Dev.* 2015;2(2):101-17.
18. Sulemana I. Are happier people more willing to make income sacrifices to protect the environment? *Soc Indic Res.* 2016;127(1):447-67.
19. El Filali A, Bentata Y, Ada N, Oneib B. Depression and anxiety disorders in chronic hemodialysis patients and their quality of life: a cross-sectional study about 106 cases in the northeast of Morocco. *Saudi J Kidney Dis Transplant.* 2017;28(2):341-8.
20. Tabatabaei MF, Raghbi M. Effect of happiness training in depression, anxiety, and quality of life among hemodialysis patients. *J Res Health.* 2017;7(4):935-43.
21. Belayev LY, Mor MK, Sevic MA, Shields AM, Rollman BL, Palevsky PM, et al. Longitudinal associations of depressive symptoms and pain with quality of life in patients receiving chronic hemodialysis. *Hemodial Int.* 2015;19(2):216-24.
22. Ottaviani AC, Betoni LC, Pavarini SC, Gramani Say K, Zazzetta MS, Orlandi FD. Association between anxiety and depression and quality of life of chronic renal patients on hemodialysis. *Texto Contexto Enferm.* 2016;25(3):e00650015.
23. Bennett PN, Parsons T, Ben-Moshe R, Neal M, Weinberg MK, Gilbert K, et al. Intradialytic Laughter Yoga therapy for haemodialysis patients: a pre-post intervention feasibility study. *BMC Complement Altern Med.* 2015;15(1):176.
24. Heo EH, Kim S, Park HJ, Kil SY. The effects of a simulated laughter programme on mood, cortisol levels, and health-related quality of life among haemodialysis patients. *Complement Ther Clin Pract.* 2016;25:1-7.
25. Sousa LM, Marques-Vieira C, Severino SS, Pozo-Rosado JL, José HM. Fatores explicativos da apreciação de filmes cómicos em pessoas com doença renal crónica. *Rev Inv Enferm.* 2016;S2(15):49-55.

Este artículo se distribuye bajo una Licencia Creative Commons Atribución-NoComercial 4.0 Internacional.

<https://creativecommons.org/licenses/by-nc/4.0/>

