

10-7-2019

Longitudinal Change in Common Impairments in Children with Cerebral Palsy from age 1.5 to 11 years

Lynn M Jeffries

Alyssa Laforme Fiss

Sarah Westcott McCoy

Lisa Avery

Follow this and additional works at: <https://ir.lib.uwo.ca/ptpub>

 Part of the [Physical Therapy Commons](#)

Citation of this paper:

Jeffries, Lynn M; Laforme Fiss, Alyssa; Westcott McCoy, Sarah; and Avery, Lisa, "Longitudinal Change in Common Impairments in Children with Cerebral Palsy from age 1.5 to 11 years" (2019). *Physical Therapy Publications*. 61.
<https://ir.lib.uwo.ca/ptpub/61>

1 **Title: Longitudinal Change in Common Impairments in Children with Cerebral Palsy from age**
2 **1.5 to 11 years**

3 **Short Title:** Change in Impairments

4
5 **Lynn M. Jeffries, PT, PhD, PCS**, Associate Professor, Department of Rehabilitation Sciences,
6 University of Oklahoma Health Sciences Center, Oklahoma City, OK 73117, [lynn-](mailto:lynn-jeffries@ouhsc.edu)
7 jeffries@ouhsc.edu

8 **Alyssa LaForme Fiss, PT, PhD, PCS**, Associate Professor, Department of Physical Therapy,
9 Mercer University, Atlanta, GA 30341, fiss_al@mercer.edu

10 **Sarah Westcott McCoy, PT, PhD, FAPTA**, Professor, Rehabilitation Medicine, University of
11 Washington, Seattle, WA 98195, westcs@uw.edu

12 **Lisa Avery, B.Eng, M.Sc**, Avery Information Services Ltd, Orillia, ON, CAN,
13 lisa@averyinformation.com

14
15

16 Conflicts of Interest: The author(s) declares no conflict of interest.

17

18 Correspondence: Lynn Jeffries, PT, PhD, PCS, Department of Rehabilitation Sciences, University
19 of Oklahoma Health Sciences Center, 1200 N Stonewall, Rm 3092, Oklahoma City, OK 73117,
20 lynn-jeffries@ouhsc.edu

21

22 Grant Support: The Move & PLAY Study was supported through funds from the Canadian
23 Institutes of Health Research (MOP 81107) and the US Department of Education, National
24 Institutes of Disability and Rehabilitation Research (H133G060254). The On Track Study was
25 funded by the Canadian Institutes of Health Research (MOP # 119276) and the Patient Centered
26 Outcomes Research Institute (Grant # 5321). The statements presented in this work are solely
27 the responsibility of the authors and do not necessarily represent the views of the Patient-
28 Centered Outcomes Research Institute (PCORI), its Board of Governors or Methodology
29 Committee.

1 **Acknowledgements**

2

3 We acknowledge additional Move & PLAY and On Track Study Team members including
4 academic researchers: Doreen Bartlett, Lisa Chiarello, Robert Palisano, Pitor Wilk, Peter
5 Rosenbaum, and Jan Willem Gorter; Canadian project coordinator Barb Galuppi; US project
6 coordinator Monica Smersh; and parent researchers: Lisa Diller, Paula Drew, Nancy Ford,
7 Marquitha Gilbert, tina hjernegaard, Kimberly Rayfield, and Barbara Sieck Taylor.

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1 **Abstract**

2 **Purpose**

3 This project aimed to determine if change occurs over time for impairments of balance, range
4 of motion (ROM), endurance, and strength of children with cerebral palsy (CP), by Gross Motor
5 Function Classification System (GMFCS) levels.

6 **Methods**

7 Measurements were completed in 77 children at two sessions (T1, T2) on average 5.8 years
8 apart. Mean ages were 2.9 years (SD .9) and 8.7 years (SD 1.1) at T1 and T2, respectively.

9 **Results**

10 Significant differences were noted from T1 to T2 for some children (GMFCS levels I, II, and III/IV:
11 balance increased, GMFCS levels I and II: strength increased, and GMFCS levels III/IV and V:
12 ROM decreased). Endurance scores were not different. Endurance scores did not change.

13 **Conclusions**

14 Longitudinal changes in most impairments occurred in children with CP. Monitoring and
15 targeted interventions should support each child's development.

16

17

18

19

20

1 Introduction

2 Children with cerebral palsy (CP) present with impairments of body function and structure.^{1,2}
3 Balance is considered a primary impairment since challenges in postural control in both static
4 and dynamic activities are frequently present at the time of diagnosis.³⁻⁵ As children with CP
5 age, secondary impairments often develop including restrictions in joint range of motion (ROM)
6 and decreased endurance and strength.^{2,6,7} A primary goal of therapy is to monitor the
7 development of impairments and focus intervention on the reduction of current impairments
8 and prevention of further secondary impairments.

9 Researchers have documented impairments in children with CP across all functional
10 motor ability levels of the Gross Motor Function Classification System (GMFCS)⁸ and as young as
11 18 months.^{2,6,9-11} However, most studies use cross-sectional methodology that provides results
12 related to a cohort of children at one point in time. Based on this cross-sectional data,
13 impairments in children with CP are present and differ by GMFCS levels.² A cross-sectional
14 study provides valuable information but does not provide a clear depiction of the development
15 of the impairments over time.

16 The opportunity occurred to examine longitudinally a sub-set of children with CP who
17 participated in two multi-site, international, prospective studies.^{12,13} Using the same clinical
18 measures, we had the ability to examine changes in impairments in a group of children with CP
19 following several years of development. The purposes of this study were to: 1) determine if a
20 change occurs over time for impairments of balance (Early Clinical Assessment of Balance
21 (ECAB)),¹⁷ range of motion (Spinal Alignment and Range of Motion Measure (SAROMM)),¹⁵

1 endurance for activity (Early Activity Scale for Endurance (EASE)),¹⁶ and strength (Functional
2 Strength Assessment (FSA))² in children with CP.

3 **Methods**

4 This study examined results from the Move & PLAY and On Track studies. Movement
5 and Participation in Life Activities of Young Children with Cerebral Palsy (Move & PLAY) aimed
6 to understand the child, family, and service delivery determinants that together explained the
7 motor abilities, self-care, and play of young children with CP.^{12,17} On Track: Monitoring
8 Development of Children with Cerebral Palsy and Gross Motor Delay aimed to develop
9 longitudinal developmental trajectories and reference percentiles for impairments, health
10 conditions, and participation variables for children with CP.¹³ This current analysis includes
11 children with CP who participated in both studies. Full study protocols have been reported
12 elsewhere.^{13,18} All participating institutions and recruitment sites with Institutional Review
13 Boards (IRBs) provided ethics approval. Parents or guardians provided informed consent and
14 children, as appropriate and in compliance with the specific IRB, provided assent for both
15 studies.

16 **Participants**

17 A convenience sample of 77 children with CP participated in both studies, from six
18 provinces across Canada, including British Columbia, Saskatchewan, Manitoba, Ontario, Nova
19 Scotia, and Newfoundland, and four regions of the United States, including areas within and
20 surrounding Georgia, Oklahoma, Pennsylvania, and Washington. Of this sample 52% of the
21 participants were from the United States. Participants were recruited through children's
22 rehabilitation centers in Canada and through physical therapists, occupational therapists,

1 physicians, and hospital systems in the United States. All children had a diagnosis of CP.
2 Children were excluded if their parents were unable to speak and understand English, French,
3 or Spanish.

4 **Measures**

5 *Gross Motor Function Classification System (GMFCS)*

6 The GMFCS is a five-point classification system used to describe gross motor function
7 ability including sitting, transfers, walking and wheeled mobility for children with CP. The child's
8 functional abilities, use of assistive technology, and need for care giver assistance differentiates
9 the levels.⁸ The GMFCS levels are divided into age bands to clearly describe gross motor
10 function as age. GMFCS content validity,⁸ construct validity, and inter-rater reliability have
11 previously been supported.¹⁹⁻²¹

12 *Early Clinical Assessment of Balance (ECAB)*

13 The ECAB provides an estimate of postural stability for children with CP across all
14 GMFCS levels.¹⁴ The assessor examines the child's head and trunk control, protective responses,
15 upright posture in sitting and standing, and postural adjustments during voluntary movements
16 in standing. The ECAB has known-groups validity for children with CP 1.5-12 years of age, with
17 average scores that differ between age groups and all GMFCS levels ($p < 0.001$),¹⁴ as well as
18 excellent inter-rater (ICC (2,1) = 0.99) and test-retest (ICC (2,1) = 0.99) reliability.²² The ECAB has
19 a minimal detectable change (MDC₉₅) of 10 points.¹⁴ The total ECAB score out of 100 was used
20 for analysis. The higher the score, the better the balance.

21 *Spinal Alignment and Range of Motion Measure (SAROMM)*

1 The SAROMM provides an overall estimate of *spinal alignment* and *ROM and muscle*
2 *extensibility* using standard physical therapy measurement techniques.¹⁵ The assessor scores 4
3 spinal alignment items using a 5-point ordinal score of 0 (“no alignment limitations with active
4 correction”) to 4 (“Fixed” – limitation is structural, static, not reducible and severe). For the
5 remaining extremity *ROM and muscle extensibility* items, the assessor scores items using a 5-
6 point ordinal score of 0 (“normal” - no restrictions of ROM on passive testing and no postures
7 typical of some children with CP) to 4 (“fixed” – limitation is structural, static, irreducible and is
8 severe), hence a lower score is better ROM.¹⁵ Researchers report good validity, inter-rater
9 reliability (ICC (2,1) = 0.89), and test-retest reliability (ICC (2,1) = 0.93) when used with children
10 with CP.¹⁵ For the SAROMM, total score the MDC₉₀ has been reported as 3.22²³ and MDC₉₅ as 9
11 points,¹⁵ and the minimal clinically important difference (MCID) is 4.53.²³ The mean of all item
12 scores for each child was used for analysis.

13 *4-Item Early Activity Scale for Endurance (EASE)*

14 The 4-item EASE includes four questions of parent perception of the child’s endurance
15 for activity.¹⁶ Questions are scored using a 5-point ordinal scale of 1(Never) to 5 (Always) and
16 include the child’s (1) physical activity related to peers, (2) physical energy level and their need
17 to take breaks, (3) frequency of breathing quickly and getting flushed during activity, and (4)
18 frequency of daily activities requiring a lot of physical energy. Higher scores indicate greater
19 endurance for activity. The EASE is moderately correlated (Spearman $r=0.41$, $p = 0.01$) with the
20 Six-Minute Walk Test¹⁶ and has acceptable inter-rater reliability (ICC (2,1) = 0.79).¹⁶ The EASE
21 does not have a calculated MDC. The mean EASE score was used for analysis.

22 *Functional Strength Assessment (FSA)*

1 The FSA includes an assessment of eight movements against gravity and resistance,
2 providing an estimate of the child’s strength in major muscle groups.² The assessor rates the
3 child’s strength using a 5-point ordinal scale ranging from 1 (only flicker of contraction or just
4 initiates movement against gravity) to 5 (full available range against gravity and strong
5 resistance) for major muscle groups (neck and trunk flexors and extensors, hip extensors, knee
6 extensors, and shoulder flexors).² The FSA has excellent inter-rater reliability (ICC (2,1) =
7 0.996);² however, the FSA does not have a calculated MDC. The mean FSA score is used for
8 analysis with a higher score indicating better strength.

9 The ECAB, SAROMM, EASE, and FSA forms and training protocols can be accessed
10 through the CanChild website ([https://www.canchild.ca/en/research-in-practice/current-
11 studies/on-track/on-track-measures](https://www.canchild.ca/en/research-in-practice/current-studies/on-track/on-track-measures)).

12 **Procedures**

13 Therapist assessors were physical therapists and occupational therapists, with at least
14 one year experience, from Canada and the United States who completed onsite training prior to
15 data collection. Therapist’s also completed videotaped criterion tests of the ECAB, SAROMM
16 and FSA measures and obtained greater than or equal to 80% item agreement with the study
17 investigators providing ‘gold standard’ responses.

18 Data from the first assessment of the Move & PLAY study (T1) collected between
19 summer 2007 and spring 2009, and data from the first assessment of the On Track study (T2)
20 collected between spring 2013 and summer 2014 were used for this analysis. Parents
21 completed the EASE and a demographic questionnaire. Therapist assessors completed the
22 ECAB, SAROMM, and FSA. The parent and therapist assessor independently completed the

1 GMFCS and a consensus rating was determined.²⁴ For this analysis, we used the GMFCS rating
2 from the On Track study (T2) because some children at T1 were under 2 years old and GMFCS
3 reliability is greater after 2 years of age.²⁰

4 **Data Analysis**

5 Data were analyzed using the Statistical Package for Social Sciences (Version 18).
6 Descriptive statistics of participants are in Table 1. Median and minimum/maximum range for
7 impairments (ECAB, SAROMM, EASE, and FSA) are presented in Table 2. Because of the small
8 sample size of children at GMFCS Level III, these children were combined with children at
9 GMFCS Level IV for group comparisons for the impairment measures. The Wilcoxon Signed
10 Ranked test was used for comparison of medians between T1 and T2 for GMFCS level groups
11 (Table 2). Comparisons of median scores for each variable across GMFCS level groups were
12 completed for data at both T1 and T2 using non-parametric, Kruskal-Wallis tests with post hoc
13 pairwise comparisons to determine if significant differences existed across GMFCS levels (Table
14 3). An alpha level of $p < 0.05$ was used to indicate significance for overall testing.

15 **Results**

16 Children were 1.5 to 4.6 years old (mean 2.9 yrs, SD=.9) at T1 and 6.3 to 11.1 years old
17 (mean 8.7 yrs, SD=1.1) at T2. Participants were 52% males and 79% white. The proportion of
18 children in each GMFCS level in our sample was: GMFCS I – 26.0% (N=20); GMFCS II – 29.9%
19 (N=23); GMFCS III – 9.1% (N=7); GMFCS IV – 13.0% (N=10); GMFCS V – 22.1% (N=17). Table 1
20 contains demographic information. On average, the time difference between T1 and T2 was 5.8
21 years (SD .6 yrs).

1 Comparing ECAB scores over time, significant improvements in balance were noted for
2 children in GMFCS levels I ($p < 0.001$), II ($p < 0.001$), and III/IV ($p = 0.008$), (Table 2). At both T1 and
3 T2 assessments, significant differences were noted on the ECAB across all comparisons with
4 children with more functional mobility demonstrating higher ECAB scores than children with
5 less functional mobility (Table 3).

6 Overtime SAROMM scores were significantly higher at T2, for children in GMFCS levels
7 III/IV ($p = 0.005$) and V ($p < 0.001$) indicating greater ROM restrictions (Table 2). At both T1 and T2
8 assessments, significant differences were noted on the SAROMM across all comparisons, except
9 between children in levels II and III/IV at T1 (Table 3).

10 No differences in EASE scores within GMFCS levels were noted over time (Table 2). On
11 the EASE, at both T1 and T2 assessments, significant differences were noted across all
12 comparisons, except between children in levels II and III/IV at both T1 and T2 (Table 3).

13 Comparing FSA scores over time, significant improvements in strength were noted for
14 children in GMFCS levels I ($p < 0.001$) and II ($p < 0.001$) (Table 2). At both T1 and T2 assessments,
15 significant differences were noted on the FSA across all comparisons, except between children
16 in levels II and III/IV at T1, and between children in levels I and II at T2 (Table 3).

17 **Discussion**

18 This study provided the opportunity to follow 77 children with CP longitudinally over a
19 multiyear period to explore how commonly identified impairments changed over time. Finding
20 for each construct are discussed below.

21 Significant differences in balance were noted from T1 to T2 for all children with CP
22 except those with GMFCS level V. As expected, children with more gross motor ability

1 demonstrated better balance than children with less gross motor ability. For the majority of
2 children, the amount of change in balance was greater than the minimal amount of change
3 required to differentiate a true change versus a change due to variability in performance.

4 As balance is correlated to gross motor ability²⁵ and gross motor skills continue to
5 develop as children with CP age²⁶ this improvement was expected. For children in level V, the
6 balance median score decreased. Gross motor skills of children at level V plateau on average at
7 2 years 7 months¹⁹ indicating gross motor skills are not significantly changing, therefore, one
8 could theorize that balance skills will also not change. This longitudinal exploration of balance in
9 children with CP provides a beginning analysis. Continued exploration is needed to determine if
10 differences in service focus and frequency influences a child's balance abilities.

11 Similar to the results of Ostensjo and colleagues,⁹ children with CP in our study
12 presented with some degree of ROM and spinal alignment restrictions regardless of GMFCS
13 level. As expected, children with higher functional mobility presented with fewer restrictions
14 compared to children with lower functional abilities. For the majority of children, the amount of
15 change in ROM scores was greater than the minimal amount of change required to differentiate
16 a true change and they had restrictions beyond what is typically considered a clinical change
17 that influences their daily activities.

18 When ROM and spinal alignment were measured over time, children at GMFCS levels I
19 and II did not present with a significant change. Based on this study's methodology, direct
20 correlations cannot be examined to determine what prevented the progression of joint
21 restrictions; however, one can hypothesize independent mobility likely facilitates joint and
22 spinal flexibility. Children with higher GMFCS levels rely on assistive devices for mobility

1 (walkers or wheelchairs) and spend a good deal of time seated or even lying down, therefore,
2 these static postures likely contribute to the development of secondary impairments in ROM.
3 This presents as an opportune window for intervention to prevent increases in joint restrictions
4 in children in levels III-V particularly. The challenge is what intervention should be used. Based
5 on a systematic review brief stretching has little to no effect either short-term or long-term on
6 improving joint mobility in persons with neurological conditions.²⁷ A heightened focus on
7 increasing targeted functional activities and more frequent changes in positions for those at
8 GMFCS Level V may assist to prevent further ROM restrictions. Within our data, however, we
9 do not know the details of interventions to prevent secondary impairments and how increasing
10 environmental modifications may alter the development of ROM restrictions.

11 As we expected endurance for activity was higher for children with higher functional
12 mobility. Median EASE scores decreased at each GMFCS level but were not significantly
13 different over time. What is unknown is if endurance is relatively constant over time for
14 children with CP in this age range, the tool is not sensitive enough for longitudinal change, or a
15 larger sample of children is needed to identify longitudinal change. We know from physical
16 activity literature that endurance typically decreases in adolescents without disabilities,²⁸ so
17 tracking children with CP into the future to identify if and when changes in endurance occur is
18 needed. Also since parents report no change in endurance, we hypothesis that maybe this is an
19 area of intervention therapists do not focus on. We recommend in the future therapists should
20 consider targeted interventions focused on endurance at all GMFCS levels and ages.

21 As expected, strength scores were higher for children with higher motor function. Over
22 time strength scores significantly improved for children in GMFCS levels I and II. This is

1 congruent with literature demonstrating children with CP have the ability to strengthen various
2 muscle groups.²⁹ Additionally the FSA scores for children in the other GMFCS levels increased
3 but were not significantly different across time. There are varying results in the literature
4 related to strengthening interventions for children with CP and the carry over to functional
5 motor skills.³⁰ Most literature focuses on children at GMFCS levels I-III since they more often
6 have the ability to demonstrate selective motor control. Given that a focus on services related
7 to secondary impairments increases over time, more details from therapists would be helpful to
8 determine effective strengthening interventions especially for children with lower functional
9 ability.

10 **As most children with CP receive therapy services during their early childhood,**
11 **therapists have the opportunity to support maturation during this critical period of growth**
12 **and skill development. Although, we were unable to determine the impact of therapy**
13 **services nor the child's natural evolution of skill mastery, most children with CP were**
14 **stronger, had better balance, and their endurance had not significantly declined after almost**
15 **six years. Based on these results and the knowledge that in adolescents and young**
16 **adulthood, individuals get heavier and flexibility and endurance decline,^{28, 31, 32} therapists**
17 **should encourage health and wellness programs that focus on strength and cardiopulmonary**
18 **fitness whether that be in a therapist directed or community program.**

19 **Limitations**

20 The primary limitation of this study is the small sample size. To acknowledge this, we
21 used non-parametric statistics for data analysis. **We did not correct for multiple analyses,**
22 **however as most p values were below <0.001 one could determine this was not a problem.**

1 We also combined data for children in GMFCS levels III and IV. The EASE and FSA measures do
2 not have MDC or MCID values, which would have provided additional information regarding
3 change over time. This study had a slightly lower proportion of children at GMFCS level I and a
4 greater proportion of children at level V compared to the Reid and colleagues³³ determination
5 of the GMFCS distribution of CP based on multiple international registries. Finally, a potential
6 sampling bias could be present as parents agreed to participate in two research studies over
7 time which could indicate they are closely linked to rehabilitation providers and potentially
8 engaged in intervention programs more than other children.

9 **Conclusion**

10 The results of this study indicate there are improvements in children with CP within
11 some impairment areas (balance and strength), however greater ROM restrictions and no
12 changes in endurance are noted over an extended time. These changes support the need for
13 physical therapists to monitor and focus interventions on primary and secondary impairments
14 in children with CP, given the hypothesis that each of these impairments can potentially impact
15 the children's ability to perform daily activities and participate in home, school and community
16 environments. Based on this longitudinal study, continued monitoring of impairments and
17 collaboration with families is important for the development of children with CP.

References

1. Rosenbaum P, Stewart D. The World Health Organization International Classification of Functioning, Disability, and Health: a model to guide clinical thinking, practice and research in the field of cerebral palsy. *Semin Pediatr Neurol*. 2004;11:5-10.
2. Jeffries LM, Fiss A, McCoy S, Bartlett DJ. Description of primary and secondary impairments in young children with cerebral palsy. *Ped Phys Ther*. 2016;28:7-14.
3. Woollacott M, Shumway-Cook A. Postural dysfunction during standing and walking in children with cerebral palsy: what are the underlying problems and what new therapies might improve balance? *Neural Plast*. 2005;12(2-3):211–219.
4. Liu W, Zaino C, Westcott McCoy S. Anticipatory postural adjustments in children with cerebral palsy and children with typical development during functional reaching: a center of pressure study. *Pediatr Phys Ther*. 2007;19:188–195.
5. Hsue B, Miller F, Su F. The dynamic balance of children with cerebral palsy and typical development during gait. Part I: Spatial relationship between COM and COP trajectories. *Gait and Posture*. 2009;29:465–470.
6. Nordmark E, Hagglund G, Lauge-Pederson H, Wagner P, Westbom L. Development of lower limb range of motion from early childhood to adolescence in cerebral palsy: A population-based study. *BMC Med*. 2009;7:65.
7. Bartlett DJ, Palisano RJ. Physical therapists' perceptions of factors influencing the acquisition of motor abilities of children with cerebral palsy: implications for clinical reasoning. *Phys Ther*. 2002;85:237-248.
8. Palisano R, Rosenbaum P, Bartlett D, Livingston M. Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol*. 2008;50(10):744-50.
9. Ostensjo S, Carlberg EB, Vollestad NK. Motor impairments in young children with cerebral palsy; relationship to gross motor function and everyday activities. *Dev Med Child Neurol*. 2004;46:580-589.
10. Ryan JM, Forde C, Hussey JM, et al. Comparison of patterns of physical activity and sedentary behavior between children with cerebral palsy and children with typical development. *Phys Ther*. 2015;95:1609-1616.
11. Bjornson KF, Belza B, Kartin D, et al. Ambulatory physical activity performance in youth with cerebral palsy and youth who are developing typically. *Phys Ther*. 2007;87:248–257.

12. Bartlett DJ, Chiarello LA, McCoy SW, et al. Determinants of gross motor function of young children with cerebral palsy: A prospective cohort study. *Dev Med Child Neurol*, 2014a;56:275-82.
13. McCoy SW, Bartlett D, Smersh M, Galuppi B, Hanna S. Collaboration Group: On Track Study Team. Monitoring development of children with cerebral palsy: the On Track study. Protocol of a longitudinal study of development and services. Available at: <https://www.canchild.ca/en/resources/294-monitoring-development-of-children-with-cerebralpalsy-the-on-track-study-protocol-of-a-longitudinal-study-of-development-and-services>. Accessed November 1, 2018.
14. McCoy SW, Bartlett DJ, Yocum A, et al. Development and validity of the Early Clinical Assessment of Balance for young children with cerebral palsy. *Dev Neurorehabil*. 2014;17:375-83.
15. Bartlett DJ, Purdie B. Testing of the Spinal Alignment and Range of Motion Measure: A discriminative measure of posture and flexibility for children with cerebral palsy. *Dev Med Child Neurol*. 2005;47:739-743.
16. McCoy S, Yocum A, Bartlett D, et al. Development of the Early Activity Scale for Endurance for children with cerebral palsy. *Ped Phys Ther*. 2012;24:232-240.
17. Chiarello LA, Palisano RJ, Bartlett DJ, McCoy SW. A multivariate model of determinants of change in gross-motor abilities and engagement in self-care and play in young children with cerebral palsy. *Phys Occup Ther Pediatr*. 2011;31:150-168.
18. Bartlett DJ, Chiarello LA, McCoy SW, et al. The measurement model of the Move & PLAY study: An example of comprehensive outcomes research in rehabilitation. *Phys Ther*. 2010;90:1-13.
19. Rosenbaum PL, Walter SD, Hanna SE, et al. Prognosis for gross motor function in cerebral palsy; creation of motor development curves. [JAMA](#). 2002;288:1357-1363.
20. Palisano RJ, Rosenbaum PL, Walter S, et al. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol*. 1997;39:214-223.
21. Wood, E, Rosenbaum P. The Gross Motor Function Classification System for cerebral palsy. A study of reliability and stability over time. *Dev Med Child Neurol*. 2000;42:292-296.
22. Randall K, Bartlett D, McCoy SW. Measuring postural stability in young children with cerebral palsy: A comparison of two instruments. *Ped Phys Ther*. 2014;26:332-337.

23. Chen C, Wu, KPH, Lie, W, Cheng HK, Shen H, Lin, K. Validity and clinimetric properties of the Spinal Alignment and Range of Motion Measure in children with cerebral palsy. *Dev Med Child Neurol*. 2013;55:745-750.
24. Bartlett DJ, Galuppi B, Palisano RJ, McCoy SW. Consensus classifications of gross motor, manual ability, and communication function classification systems between therapists and parents of children with cerebral palsy. *Dev Med Child Neurol*. 2016;58:98-99.
25. Liao HF, Hwan AW. Relations of balance function and gross motor ability for children with cerebral palsy. *Percept Motor Skills*. 2003;96:1173-1184.
26. Russell D, Rosenbaum P, Avery L, Lane M. Gross Motor Function Measure (GMFM-66 & GMFM-88) User's Manual. London, UK: Mac Keith Press; 2002.
27. Katalinic OM, Harvy LA, Herbert RD. Effectiveness of stretch for the treatment and prevention of contracture in people with neurological conditions: A systematic review. *Phys Ther*. 2011;91(1):11-24.
28. Durmith SC, Gigante DP, Domingues MR, Kohl HW. Physical activity change enduring adolescence: A systematic review and a pooled analysis. *Int J of Epi*. 2011;40(11):685-698.
29. Park EY, Kim WH. Meta-analysis of the effect of strengthening interventions in individuals with cerebral palsy. *Res Dev Disabil*. 2014;35(2):239-249.
30. Novak I, McIntyre S, Morgan C. A systematic review of interventions for children with cerebral palsy: State of the evidence. *Dev Med Child Neuro*. 2013;55:885-910.
31. Maher, CA, Williams, MT, Olds T, Lane AE. Physical and sedentary activity in adolescents with cerebral palsy. *Dev Med Child Neurol*. 2007; 49:450-457.
32. Day SM, Strauss DJ, Vachon PJ, Rosenbloom L, Shavelle, RM, Wu, YW. Growth patterns in a population of children and adolescents with cerebral palsy. *Dev Med Child Neurol*. 2007; 49:167-171.
33. Reid SM, Carlin JB, Reddihough DS. Using the Gross Motor Function Classification System to describe patterns of motor severity in cerebral palsy. *Dev Med Child Neurol*. 2011; 53:1007-1012.

Table 1. Children’s demographics and clinical characteristics

Child Demographics		
Gender (%)	Male	40 (52)
	Female	37 (48)
Ethnicity (%)	African-American	3 (4)
	Asian or Pacific Islander	3 (4)
	Hispanic/ Latino	1 (1)
	Native American	3 (4)
	White	61 (80)
	Other	4 (6)
	Data not available	2(1)
GMFCS Level (%)	I: Independent self-mobility	20 (26)
	II	23 (30)
	III	7 (9)
	IV	10 (13)
	V: Severe limitations in posture/self-mobility	17 (22)
Distribution of Involvement (%)	Quadriplegia	40 (52)
	Hemiplegia	19 (25)
	Diplegia	14 (18)
	Triplegia	3 (4)
	Data not available	1 (1)

Abbreviation: SD = standard deviation

Table 2: Balance, Range of Motion, Endurance, and Strength comparisons across time by Gross Motor Function Classification System (GMFCS) level

Early Clinical Assessment of Balance (ECAB)			
GMFCS level	Time 1 Median (min, max)	Time 2 Median (min, max)	Wilcoxon Signed Ranks Test
I	87.00 (43.50, 100.0)	100.00 (78.00, 100.00)	Z=3.62; p < 0.001
II	45.00 (18.00, 100.0)	86.0 (48.0, 100.00)	Z=3.90; p < 0.001
III/IV	21.00 (9.00, 40.50)	32.0 (6.00, 51.50)	Z=2.66; p = 0.008
V	5.00 (2.00, 15.00)	3.5.0 (0, 29.00)	Z=-0.57; p = 0.57
Spinal Alignment and Range of Motion Measure (SAROMM)			
I	.23 (.00, .85)	.31 (.08, .96)	Z=1.39; p = 0.17
II	.50 (.00, 1.42)	.85 (.08, 1.85)	Z=1.99; p = 0.05
III/IV	.54 (.12, 1.73)	1.54 (.08, 2.08)	Z=2.82; p = 0.005
V	1.38 (.65, 1.77)	2.23 (.96, 2.92)	Z=3.29; p < 0.001
Early Activity Scale for Endurance (EASE)			
I	4.25 (3.0, 5.0)	4.00 (2.25, 5.0)	Z=1.28; p = 0.20
II	3.25 (1.75, 5.0)	3.25 (1.75, 4.75)	Z=1.10; p = 0.27
III/IV	3.00 (1.25, 5.0)	2.50 (1.25, 4.25)	Z=1.09; p = 0.27
V	1.75 (1.0, 3.5)	1.25 (1.0, 3.25)	Z=0.75; p = 0.45
Functional Strength Assessment (FSA)			
I	3.5 (2.88, 4.00)	4.56 (2.88, 5.00)	Z=3.69; p < 0.001
II	3.00 (1.75, 4.00)	4.13 (2.5, 5.00)	Z=4.17; p < 0.001
III/IV	2.63 (1.5, 3.75)	3.13 (1.25, 4.25)	Z=1.89; p = 0.06
V	1.25 (.13, 2.5)	1.63 (1.00, 3.38)	Z=1.29; p = 0.20

Table 3: Balance, Range of Motion, Endurance, and Strength pairwise comparisons by Gross Motor Function Classification System (GMFCS) level

Early Clinical Assessment of Balance (ECAB)			
GMFCS level	Comparison GMFCS level	P-value T1	P-value T2
I	II	<.001*	0.003*
	III/IV	<.001*	<0.001*
	V	<.001*	<0.001*
II	III/IV	<.001*	<0.001*
	V	<.001*	<0.001*
III/IV	V	<.001*	<0.001*
Spinal Alignment and Range of Motion Measure (SAROMM)			
GMFCS level	Comparison GMFCS level	p-value T1	p-value T2
I	II	.004*	0.001*
	III/IV	.001*	<0.001*
	V	.000*	<0.001*
II	III/IV	.30	0.02*
	V	<.001*	<0.001*
III/IV	V	.002*	0.002*
Early Activity Scale for Endurance (EASE)			
GMFCS level	Comparison GMFCS level	p-value T1	p-value T2
I	II	.01*	0.01*
	III/IV	<.001*	0.001*
	V	<.001*	<0.001*
II	III/IV	.07	0.07
	V	<.001*	<0.001*
III/IV	V	.001*	0.003*
Functional Strength Assessment (FSA)			
GMFCS level	Comparison GMFCS level	p-value T1	p-value T2
I	II	.002*	0.15
	III/IV	<.001*	<0.001*
	V	<.001*	<0.001*
II	III/IV	.09	<0.001*
	V	<.001*	<0.001*
III/IV	V	<.001*	<0.001*

*P < 0.05