

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

The ecology of feral cats (*Felis catus*) on a New Zealand offshore island: Considerations for management

A thesis presented in partial fulfilment of the requirements for the degree of

Doctor of Philosophy

in

Zoology

at Massey University, Manawatu,

New Zealand.



Kathryn Emily Strang 2018

There is a predator within me One that craves to stalk you My neverending prey For with you I have The thrill of the hunt In the night and in the jungle Feel my roar vibrate through your bones And know me to be a beast Aggressively marking you as my territory Claws that dig into your fur Canines that linger in your warmth And this is the beginning Of our wild hunt

Poem by Cheyenne Raine (<u>www.rainepoetry.com</u>) and artwork by Lucy Newton.

Abstract

Invasive species have contributed to the global biodiversity crisis, with the majority of recentday terrestrial vertebrate extinctions implicating invasive predators. Domestic cats (*Felis catus*), particularly feral cats, are among the main culprits. In New Zealand, the continued decline of native species is largely due to invasive predators, which has led conservation efforts to focus on pest management. Feral cats are often a secondary focus for pest control, and their impacts within an ecosystem are not well understood. The main objective of this study was to produce a comprehensive study of feral cat ecology using a population of feral cats on Ponui Island, New Zealand, and highlight factors that should be considered for efficient control operations.

Predators distribute themselves and move within their environment in relation to prey availability and habitat type. These factors vary between locations, leading to differences in home range sizes that are difficult to predict. A relationship between home range size and population density was identified in the literature, which I used to predict changes in home range size with changing population density. I mapped the home ranges of eight cats for one year using location points from Very High Frequency (VHF) radio-telemetry and camera traps, then two males were removed from the population, seven months apart. The first removal resulted in an increase in the home range sizes of the remaining cats, whereas the second removal saw an invasion of four unmarked male cats. These results show that feral cats change their home ranges accordingly with changes in density, and removals (such as control operations) result in a change in feral cat ranging behaviour.

Feral cats are often live-trapped during control programmes to reduce capturing non-target animals. This technique is labour intensive because traps need to be checked frequently for welfare purposes. Describing feral cat activity patterns can dictate when live-traps should be set and checked. Camera traps have recently been used to estimate activity patterns, but have yet to be validated against accelerometry devices. Therefore, I compared the activity patterns obtained through camera trap data to that from collar-mounted accelerometry devices, and found a high correlation ($R^2 > 0.9$) between the two methods. The highest correlation was when activity from 600 or more videos was used. Feral cats were most active between sunset and sunrise, and live-traps should be set during these times to increase trapping efficiency and reduce bycatch.

The reproductive biology of invasive predators can be used to predict the recovery of populations following control operations, however there is very little information available for

feral cats. Therefore, I investigated the reproduction of feral cats in a stable, insular population. Using camera trap data, I found that females had high reproductive output, averaging three kittens per litter and two litters per year. However, the recruitment rate was low in this population; only 3-4% of kittens survived to one-year-old. There were two infanticide events observed; the first reported for solitary-living feral cats. Females moved shorter distances when they had young kittens (less than seven-weeks-old). These results show that feral cats can rapidly recover from control operations. Females are less likely to encounter traps when they have young kittens, suggesting there are optimal seasons to capture cats.

Population genetics is used in invasive biology to identify populations that are isolated and have limited immigration. Eradication of isolated populations will be easier and more cost-efficient, with low chances of reinvasion. Although offshore islands are geographically isolated, there is the chance of reinvasion that is assisted by humans. I investigated the population genetics of the feral cats on Ponui Island using genetic samples collected from adults and kittens, and also opportunistically studied parentage. Most of the fathers of the litters were the heaviest males, and the males that had the most home range overlap with queens. Population genetics showed that the cats have most likely been isolated with no recent introductions, suggesting the removal of cats from this island would be successful with low chances of reinvasion.

Feral cats can have large impacts on native species, but prey on rats that also have detrimental effects on wildlife. This led me to investigate the diet of feral cats using scat analysis on an island with native birdlife, and rodents at high densities. I examined season and sex differences on diet and the impacts of cats on native species. Feral cats consumed prey based on seasonal availability, with cats eating rats when rats were at their highest density. Females ate smaller prey more frequently than males, such as passerines. The cats on this island are not reliant on the rat population, and were found to eat many native species such as brown kiwi, morepork, and fantails.

The findings from this thesis can be applied to feral cat management to develop efficient control operations. The decision to control a population should be based on both dietary and genetic data to reduce possible cascading ecological effects from the predator removal and identify genetically isolated populations. Home range, activity, and diet data can be used to determine the control protocol, such as; trap spacing, the time traps should be set and checked, and if secondary or primary poisoning should be used and the seasons that cats would uptake the baits. Finally, knowledge of feral cat reproduction can be used to predict the recovery of the population.

vi

Acknowledgements

My supervisors, Isabel Castro, Murray Potter, and Nick Cave. Isabel, I guess this journey started because of you. I had been applying for PhDs overseas when you invited me on a trip to Ponui, got me talking to the Chamberlins about the feral cats, and convinced me to apply for funding at Massey. I see what you did there... I appreciate you encouraging me to take every opportunity that came my way, not only for my PhD. Often, I felt like it would have been easier to herd cats than what it was to get all of you in one room at the same time, but the great ideas talked about in these meetings was worth it. I appreciate you all taking on this project and I feel like everyone had a different expertise to offer that moulded this thesis into what it is.

Kristene Gedye, you were basically my genetics supervisor. Thank you for holding my hand throughout all of my lab work. For teaching me so much about genetics, coming up with new ways that we could extract DNA (in a simple and cheap way), and boosting my spirits when the PCR gods really didn't like me. I learnt a lot of things from you, and I really appreciate all of the new skills I have.

Steve Trewick, thank you for giving me a very quick crash course on population genetics. Chapter 5 would not have been the same without your input. Thank you for reading through the drafts to make sure I knew what I was talking about!

My sincere thanks goes to all of the volunteers and interns who helped me in the field. Aden and Sarah, thank you so much for all of your hard work. Your company out on Ponui was amazing, as well as binge watching Friends and Game of Thrones at night. I appreciate the many sleepless nights trapping, walking endless kilometres to find cats, sorting through the massive dataset that was our camera trap data, and sorting through cat scats and pitfalls in the lab. Cecilia, thank you for coming out and helping me re-bait traps and keeping me company in the field. Warren, thank you for sorting through cat scats. It really helped me with being able to finally sort through the mountain of scats that awaited me after finishing field work.

To the Rawhiti guys, Inge and John, Tom Dixon, Alex Wilson, and Alex Brighton; I have loved your company every catch week. I appreciated you all coming out to help me check traps during my first week trapping. Rana, thank you for lending me traps, running around the island after the pit tag reader, and for your quick hands when Tigger refused to go into the crush cage. Inge, you taught me so much during that first week of trapping. I never would have gained the confidence to anaesthetise feral cats without you being there that first time. John, you also ran around for

me checking traps, and even brought us hot chocolate all the way up to Road trap on a long, miserable night trapping. Maria, I owe you a lot. You enthusiastically took over the position from Inge of accompanying me out on trapping nights. Your vet knowledge is incredible, and I will forever appreciate you being on call for me, and for looking after Shadow.

To the Chamberlins, words cannot express my gratitude. None of this would have been possible without you guys. Thank you for welcoming me and sharing your amazing island and farm with me. Ponui is absolutely one of my favourite places in the world. Thank you for letting me borrow vehicles when I was trapping, for the many barge and boats trips, the impromptu trips to Rotoroa Island, teaching me about sheep farming, all the fruit from the orchard (especially the wheeny grapefruit!), collecting dead cats for me whenever you came across them on the farm, and most of all, your company.

Maurice Alley, thank you for teaching me so much about post-mortems, and for not laughing at me when I brought you bones and asked you what it died from. You are quite the investigator, and I enjoyed piecing together the puzzle with you for both the kiwi and cat post-mortems.

To all those people who helped in what may seem like small but were very important ways. Ellen Schoener, Kim Singleton and the Wilson family, thank you for honouring my unusual request to be sent some of your cat's hairs for the genetic sampling. Paul, Tracy, Shaun, Cleland, Sharon, and Sandra, thank you for making it easier to get things done for this PhD. For mending equipment, building antennas, and ordering endless amounts of batteries. The Ecology Group would be lost without you. To the Waikato DOC office in Rotorua and the Palmerston North City Council, thank you for lending me traps. Thanks goes to Phil and Katie for helping me identify the munched up feathers that I found in cat scats. Keren Dittmer, for letting me use your lab for all of my genetic work, and checking up on me when I was working late nights in the lab. Anastasia, for helping Kristene and I grind up the bone samples for the genetic work. Lee Shapiro, thank you for the initial advice on setting up the trapping programme for the feral cats on the island and for all of the advice on baiting and also collaring the cats. To the girls at the Feline Unit; thanks for the many coffees, the kitten cuddles, for lending me data, and letting me store my anaesthetic.

To the Ponui team, enjoy! Good luck for your postgrad studies, and thank you for all of the trips that we shared together on Ponui. Thanks especially to Tom and Alex who taught me so much about the island when I was just a newbie.

viii

To my friends, my office mates, and the Ecology Group, I could not think of better people to go through this with. Thank you for all of the support, the lunches at 12pm on the dot, the movie and Game of Thrones nights, and the parties when we really needed them. Nina, the weekly pints before pilates really saved me (for both my PhD and pilates)! Yvonne, the endless coffees at uni and de-briefs, and just lending an ear when it was a bad health week. Nicki, Katie, and Emily, you know everything that you have done for me; the surprise trips, Mean Girl movie nights, and the very weird but so much fun Spice Girl night, to name a few. Thank you for it all.

My family, thank you for all of the support. Thank you Mum and Dad for instilling hard-working values into me throughout my life, it certainly helped prepare me for the long days in the field, lab and office. Thanks for the petrol and food top-ups on my way through to either Ponui or back down to Palmy. And most of all, thanks for taking an interest in the feral cats on Ponui every time I called in.

My thanks goes to those who saw the value in my research on an underestimated invasive predator: Massey University Doctoral Scholarship, Julie Alley Bursary, JP Skipworth (Ecology) Scholarship, Ecology Bursary, NZVA Wildlife Society Marion Cunningham Memorial Fund, Freemasons Postgraduate Scholarship, Australasian Wildlife Management Society Postgraduate Student Award, Lovell & Berys Clark Scholarship, and Massey University Doctoral Completion Grant.

This thesis was not possible without the help of so many people, and a huge amount of effort went into this (summarised in the table below). Cats are not always easy to track, and unorthodox methods were used at times, such as the top of the barge. Results from this thesis have been presented at numerous conferences, which is summarised in Appendix A.



Days spent on Ponui from 2014-2018	375
No. of boat trips to Ponui	43
Kilometres driven between Palmerston North and Kawakawa Bay	42570 (559 hours)
Kilometres of transect walked	924
Sleepless nights trapping	24
No. of camera trap videos	174571
Hours of camera trap footage	1454 (60 days)
Hours spent sifting and IDing cat scats	325
PCRs run	2589
PCRs rerun (because the PCR gods failed me)	975
No. coffees drunk during PhD	~5200
No. muesli bars eaten in the field (they are a lifeline)	~774

Table of Contents

Abstract	v
Acknowledgements	vii
List of figures	xv
List of tables	xvii
List of plates	xix
Chapter 1 - Feral cats: What is known globally and what do we need to know in a New Z context?	ealand1
The contribution of invasive species to the global biodiversity crisis	3
Invasive predators in New Zealand	3
Predator-prey dynamics	4
Feral cats: A global issue	8
Feral cats in New Zealand	10
Invasive predator control and public perceptions	11
Research priorities to inform feral cat management	13
Home range	14
Activity	14
Reproduction	15
Genetics	16
Diet	17
Integrating knowledge on feral cat ecology for control operations	18
Thesis structure	19
References	22
Chapter 2 – Changes in feral cat home range in response to an experimental removal of Considerations for control operations.	f males: 33
Introduction	35
Methods	40
Study site	40
Baiting phase to attract cats to live-traps	42
Live-trapping	44
Tracking through radio-telemetry	50
Camera trapping	53
Home range calculation from telemetry and camera trap data	55
Estimation of habitat type in home range	57
Calculation of home range overlap	57

Removal of male cats	57
Statistical analysis	57
Results	59
Home range	59
Morphometrics	64
Removal of males	66
Home range estimation using two techniques	71
Discussion	72
Home range versus density relationship	72
Sex differences between home range size and overlap	73
Comparison of home range size between Ponui Island and mainland/offshore stud	ies 74
Home range estimates using different techniques	75
Management	76
References	79
Supplementary material	85
Chapter 3: Camera traps versus accelerometry: Do they tell the same story?	101
Introduction	103
Methods	106
Accelerometers	106
Camera traps	107
Analysis	108
Results	112
Feral cat activity	112
Comparison of camera trap data with accelerometry	117
Activity around live-traps	119
Camera trap hotspots	119
Differences in male and female camera trap hotspots	124
Discussion	127
Camera versus accelerometer comparison	127
Feral cat activity pattern	129
Feral cat activity around live-traps and camera trap hotspots	131
References	133
Supplementary material	141

Chapter 4 - Infanticide, recruitment, and kitten growth rates in a population of (<i>Felis catus</i>).	of island feral cats 143
Introduction	
Methods	
Detecting pregnancies	
Finding den sites and processing kittens	
Tracking queens and estimating core home range	
Colony cats	
Post-mortems	
Statistical analysis	
Results	
Seasonal pregnancies	
Litter size and kitten growth	
Movements	
Den sites	
Recruitment rate	
Causes of mortality	
Discussion	
Seasonal pregnancies and recruitment rate	
Growth rate differences between feral and colony kittens	
Movements of the sub-adult and queens	
Movements of kittens between dens	
Infanticide	
Management considerations	
References	
Supplementary material	
Chapter 5 – Population genetics of feral cats on Ponui Island and parentage o	of their kittens 189
Introduction	
Methods	
Sample collection	
DNA Extraction	
Genotyping	
Analysis	
Results	

Genetic diversity	207
Relationships between cats on Ponui Island and other populations	208
Parentage	212
Discussion	215
Population genetics and diversity within the Ponui Island feral cats	215
Parentage of Ponui Island kittens	217
Implications for management	219
References	221
Supplementary material	229
Chapter 6 – Feral cats on a rat-dominated offshore island: Season and sex differences and impact on brown kiwi	in diet 235
Introduction	237
Methods	243
Collection	243
Sorting and ID	243
Sexing scats	244
Determining sex ratio	246
Calculating calorie intake	246
Calculating number of kiwi eaten per year	247
Statistical analysis	247
Results	249
Diet composition	249
Kiwi	254
Seasonal variations in diet	256
Sex bias in scats	258
Discussion	261
References	267
Supplementary material	274
Chapter 7 – Major findings and integrating aspects of feral cat ecology into effective c	ontrol
	279
Synthesis	281
Types of control for feral cats, and past eradications	282
How representative are Ponui Island feral cats of other populations?	283
An integration of ecology and control	284
Deciding to control a feral cat population	285

Implementing a control programme	. 290
Predicting the recovery and reinvasion of a population	. 294
Future research and challenges arising from this thesis	. 295
Concluding remarks	. 297
References	.298
Appendix A: List of conference presentations and other works from this thesis.	.303
Appendix B: Information on previous studies collaring feral cats	.305
Appendix C: Endoparasites identified in scats and gut contents from Ponui Island feral cats	. 312
Appendix D: Sex ratio of feral cat populations	313

List of figures

Chapter 2

Figure 1: Relationship between home range size and population density of female and male feral cats.
Figure 2: A) Satellite photo of New Zealand showing the location of Ponui Island. B) Satellite photo of Ponui Island showing the division into three privately-owned farms
Figure 3: A) Heatmap showing the number of individual cats detected at camera trap locations from January 2014 - January 2015 around Southern Ponui Island. B) Map of the study site showing the set-up of live-capture traps during the study
Figure 4: Methods used for processing live-trapped cats
Figure 5: Satellite photo of South Ponui Island showing the two transects and the triangulation sampling locations for each transect
Figure 6: Satellite photo from Google Earth (2017) showing the triangulation of bearings taken from transect points
Figure 7: Satellite image showing the 28 camera traps placed across South Ponui Island from September 2015 - March 2017
Figure 8: Kernel density estimations (KDE) for the three female and five male cats monitored on Southern Ponui Island
Figure 9: Map showing the 100% MCP home ranges and overlap of the feral cats radio-collared and studied from April 2015 – March 2017 on Ponui Island
Figure 10: Relationship between head-body length and body weight of the feral cats trapped in South Ponui Island
Figure 11: Relationship between home range and density in all feral cat studies from Table 1, as well as information on the Ponui Island cat population from this study
Figure 12: Movements of Midnight, and of Tigger before and after Midnight's death (Removal 1) 68
Figure 13: Number of unmarked cats captured on cameras traps from month 1 to 6 following Removal 2 on South Ponui Island
Figure 14: Relationship between home range and population density in all feral cat studies from Table 1, as well as the cats from this study. Results, regression values, and equations are shown for A) Removal 1, and B) Removal 2
Figure 15: Correlation in home range size (ha) between A) grid, and B) ½ average distance (AD) camera trap techniques

Chapter 3

Figure 8: Camera trap videos (n = 322) of cats visiting cage traps, eating bait, or being captured in a cage live-trap compared to the overall activity pattern detected by accelerometry data (delta-g/hour \pm SE)..119

Chapter 4

Figure 1: Number of female cats that were pregnant and the first month that the pregnancy was detected from January 2014 - March 2017. 152
Figure 2: Growth rate of feral and colony raised cats (fed ad libitum) during the first year of life
Figure 3: Comparison of the range of adult body weights between colony raised adult (n = 49 males, 59 females) and feral adult cats (n = 8 m, 3 f)
Figure 4: Boxplot showing the distance moved (m) from the den during different reproductive states of two female cats
Figure 5: Locations of the sub-adult (Shadow) when she was between 6-12 months-old in South Ponui Island
Figure 6: Den site locations for three different litters
Figure 7: Estimated survival of feral kittens in their first year of life given as the percentage of kittens seen alive at two-weeks-old, weaning (16-weeks-old), and recruitment age (one-year-old) from the estimated number of kittens born (54-72 kittens) during January 2014 - March 2017163

Chapter 5

Figure 1: A) Location of Ponui Island (the main study site) and distances to populated areas and cities as potential sources of immigration into the feral cat population. B) Location of genetic samples used in this study
Figure 2: Electropherogram of two kittens from the same litter, Cleo TS and Cleo black, showing allele size (bp) for locus F53
Figure 3: Genetic clusters on microsatellite data (loci = 18) of individuals sampled from Ponui Island and four outgroup populations
Figure 4: Clustering based on microsatellite data (loci = 18) of 29 individuals sampled from Ponui Island only

Figure 5: Overlap in home range (100% MCP) between Cleo and Socks (females) and the potential fathers of their litters. 214

Chapter 6

 Figure 1: Composition of prey found in feral cat scats.
 250

 Figure 2: A comparison of the number of brown kiwi clutches that hatched (September 2013 – January 2016) from 16 males (n = 33 clutches) and the percentage of scats analysed from each month (total n = 434) that had kiwi present (n = 34).
 254

 Figure 3: Percentage of scats (n = 434) that contained each of the main prey types across different seasons.
 257

 Figure 4: Non-metric MDS ordination of the six prey types found in feral cat scats (n = 434) collected in different seasons.
 257

Figure 5: A) Percentage of scats that contained six different prey types in scats belonging to either male (n = 180) or female (n = 63) feral cats. **B)** Percentage of occurrence for different sized birds in male (n = 76) or female (n = 21) scats that contained birds. 260

List of tables

Chapter 2

Table 1: Modified table from Liberg et al. (2000) showing a summary of the home range and popula density of feral cats 39	ition 9
Table 2: Total number of nights live-trapping feral cats from April 2015 - January 2017 to radio-collar or download accelerometry data	cats ô
Table 3: Home range size (ha) estimated for each radio-collared cat using five different levels measurement. 60	s of 0
Table 4: Percentage of home range (100% MCP) for each individual that encompassed forest/scrub hat as opposed to pasture.	bitat 0
Table 5: Percentage of overlap in home range (100% MCP) between each paired combination of c 62	ats. 2
Table 6: Morphometrics, location data, and general information for each feral cat in the study	5
Table 7: Home range changes of Tigger for Removal 167	7

Chapter 3

 Table 1: Habitat and location features associated with the 28 camera trap locations used in this study that were tested as predictors for the number of individual cats and the number of cat videos captured at each location.

 111

Table 3: GLM with negative binomial probability distribution results for the overall model on factors driving the number of cat videos captured at camera trap locations.
 121

Table 6: Results of the final GLM, estimate, standard error (SE), z values, and significance levels for each category as a predictor of the number of individual cats captured at different camera trap locations. ...122

Table 7: Results of the GLM with Poisson probability distribution, estimate, standard error (SE), z values, and significance levels for each category as a predictor of the number of female cat videos captured at different camera trap locations.

 125

Table 8: Results of the GLM with Poisson probability distribution, estimate, standard error (SE), z values, and significance levels for each category as a predictor of the number of male cat videos captured at different camera trap locations.

 125

Chapter 4

Table 1: General information on the adult feral cats that were live-trapped in South Ponui Island.......153

 Table 2: Number of cats that were pregnant each month and the corresponding months the kittens would have been born.

 154

 Table 3: The average monthly home range size (ha) (100% MCP) for the two females followed during different reproductive phases.

 158

Table 5: Causes of mortality in seven kittens, one sub-adult, and three adults found on Ponui Island.. 164

 Table 6: Description of the behaviours by both the infanticidal male (Simba) and the queen (Cleo) for the male infanticide event.

 166

Chapter 5

Table 1: Information about the 36 cats sampled fo	genetic analysis19	97
---	--------------------	----

Table 2: The nineteen primer pairs used for genotyping, length of the repeating sequence, and the observed base pair (bp) range found for all Ponui Island and outgroup samples (n = 36)......203

 Table 4: Number of alleles matched up between the potential father and each kitten for Cleo's four litters and Socks' one litter.

 213

Chapter 6

 Table 1: Percentage of occurrence of food items of feral cats studied on New Zealand's mainland and offshore islands.
 240

Table 3: Estimated number of prey consumed each year by an individual Ponui Island cat, and calorie intake from each prey item per year.
 252

Table 5: The sex of the cats that deposited the scats and the location (in relation to a tra	il) where the scats
were collected	

Chapter 7

Table 1: Number of rats depredated per season each year on Ponui Island by the feral cat popula 28	tion 38
Table 2: Number of rats on Ponui Island and estimated rat density with no feral cat depredation.	
	38

List of plates

Plate 1: Still from a camera trap video of one of the study cats (Midnight) with a rat he had caught	i 1
Plate 2: Photo showing part of the study site taken from the Eastern ridge transect	33
Plate 3: Still from a camera trap video of Mittens captured on Ponui Island.	101
Plate 4: Photos of two of the kittens found in the study site; Cleo BW from Cleo's May 2016 Pumpkin from Socks' November 2015 litter.	litter, and 143
Plate 5: Photo of Cleo's May 2016 litter in their natal den	189
Plate 6: Photo of kereru (Hemiphaga novaeseelandiae) prey remains from feral cat depredation gully.	n in Kauri 235
Plate 7: Still from camera trap showing an encounter between an adult brown kiwi and a feral cat (Patches). 279