



University of Groningen

Kinetics of hydrolysis of 4-methoxyphenyl-2,2-dichloroethanoate in binary water-cosolvent mixtures; the role of solvent activity and solute-solute interactions

Rispens, T; Cabaleiro-Lago, C; Engberts, JBFN

Published in:
Organic & Biomolecular Chemistry

DOI:
[10.1039/b414593c](https://doi.org/10.1039/b414593c)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rispens, T., Cabaleiro-Lago, C., & Engberts, JBFN. (2005). Kinetics of hydrolysis of 4-methoxyphenyl-2,2-dichloroethanoate in binary water-cosolvent mixtures; the role of solvent activity and solute-solute interactions. *Organic & Biomolecular Chemistry*, 3(4), 597-602. <https://doi.org/10.1039/b414593c>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supporting Information for:

Kinetics of hydrolysis of 4-methoxyphenyl-2,2-dichloroethanoate in binary water-cosolvent mixtures; the role of solvent activity and solute-solute interactions

Theo Rispens,^{a,b} Celia Cabaleiro-Lago,^{a,c} and Jan B. F. N. Engberts,^{a,d,*}

- (a) Physical Organic Chemistry Unit, Strating Institute, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands.
(b) E-mail: t.rispens@chem.rug.nl
(c) E-mail: celiacl@uvigo.es
(d) E-mail: j.b.f.n.engberts@chem.rug.nl

Contents:

- Kinetic data for the neutral hydrolysis of *p*-methoxyphenyl-2,2-dichloroethanoate at 25.0 °C for:

Water/Cyanomethane Mixtures	Table 1	S2
Water/PEG400 Mixtures	Table 2	S2
Water/Tetrahydrofuran Mixtures	Table 3	S3
Water/Ethanol Mixtures	Table 5	S4
Water/1-Propanol Mixtures	Table 6	S4
Water/2-Propanol Mixtures	Table 7	S4
Water/1-Butanol Mixtures	Table 8	S5
Water/2-Methyl-2-propanol Mixtures	Table 9	S5

- Isobaric activation parameters for the neutral hydrolysis of *p*-methoxyphenyl-2,2-dichloroethanoate at 25.0 °C for mixtures of water with cyanomethane, PEG400 and tetrahydrofuran:

Table 4 S3

In pure water, the rate constant of the neutral hydrolysis of MPDE is $k_0 = 3.00 \times 10^{-3} \text{ s}^{-1}$ at 25.0 °C.

The accuracy of the kinetic data is 2-3% for cyanomethane, PEG400, and tetrahydrofuran; and 1-2% for the alcohols.

Table 1. Kinetic Data for Water/Cyanomethane Mixtures

[H ₂ O] (M)	<i>x</i> _{water}	ln(<i>k</i> / <i>k</i> ₀)	ln(<i>k</i> / <i>k</i> ₀) - 2 ln <i>a</i> _{water}
53.8	0.988	-0.261	-0.237
52.1	0.977	-0.545	-0.500
50.35	0.964	-0.844	-0.777
50.25	0.963	-0.925	-0.855
45.2	0.924	-1.94	-1.82
40	0.877	-3.07	-2.90
34.5	0.821	-4.09	-3.91
28.7	0.750	-5.01	-4.82
23.4	0.673	-5.70	-5.50

Table 2. Kinetic Data for Water/PEG400 Mixtures

[H ₂ O] (M)	<i>x</i> _{water}	ln(<i>k</i> / <i>k</i> ₀)	ln(<i>k</i> / <i>k</i> ₀) - 2 ln <i>a</i> _{water}
55.34	0.9999	-0.010	-0.006
52.31	0.9968	-0.301	-0.291
51.13	0.9955	-0.473	-0.461
49.24	0.9933	-0.720	-0.702
46.74	0.9902	-0.968	-0.941
44.7	0.9873	-1.24	-1.20
39.71	0.9793	-1.83	-1.77
37.28	0.9747	-2.18	-2.09
34.24	0.9681	-2.57	-2.45
30.21	0.9575	-3.31	-3.13

Table 3. Kinetic Data for Water/Tetrahydrofuran Mixtures

[H ₂ O] (M)	<i>x</i> _{water}	ln(<i>k</i> / <i>k</i> ₀)	ln(<i>k</i> / <i>k</i> ₀) - 2 ln <i>a</i> _{water}
52.2	0.986	-0.505	-0.481
49.0	0.971	-1.01	-0.960
45.7	0.954	-1.73	-1.66
42.1	0.933	-2.76	-2.66
39.9	0.918	-3.22	-3.11
34.0	0.875	-4.09	-3.96

Table 4. Isobaric activation parameters at 25.0 °C.

cosolvent	<i>x</i> _{water}	Δ [#] <i>G</i> (kJ/mol)	Δ [#] <i>H</i> (kJ/mol)	- <i>T</i> Δ [#] <i>S</i> (kJ/mol)
none	1	87.5	35.1	52.4
	0.922	92.5	35.0	57.5
	0.875	95.1	37.6	57.5
cyanomethane	0.817	97.8	57.4	40.3
	0.9873	90.5	37.7	52.8
	0.9747	92.9	42.0	50.9
PEG400	0.968	94.0	44.8	49.2
	0.971	90.0	40.5	49.5
	0.954	91.8	47.5	44.6
tetrahydrofuran	0.936	93.7	48.5	45.3
	0.918	95.5	44.5	51.0

The accuracy Δ[#]*G* of is 0.1 kJ/mol, that of Δ[#]*H* and -*T*Δ[#]*S* 1.5 kJ/mol.

Table 5. Kinetic Data for Water/Ethanol Mixtures

x_{water}	$\ln(k/k_0)$	$\ln(k/k_0) - 2 \ln x_{\text{water}}$
0.995	-0.0782	-0.0686
0.990	-0.153	-0.132
0.985	-0.238	-0.207
0.980	-0.318	-0.278
0.976	-0.385	-0.336
0.970	-0.490	-0.428

Table 6. Kinetic Data for Water/1-Propanol Mixtures

x_{water}	$\ln(k/k_0)$	$\ln(k/k_0) - 2 \ln x_{\text{water}}$
0.997	-0.0683	-0.0622
0.995	-0.127	-0.116
0.992	-0.189	-0.173
0.989	-0.266	-0.243
0.983	-0.394	-0.360

Table 7. Kinetic Data for Water/2-Propanol Mixtures

x_{water}	$\ln(k/k_0)$	$\ln(k/k_0) - 2 \ln x_{\text{water}}$
0.995	-0.109	-0.0990
0.990	-0.225	-0.205
0.985	-0.345	-0.315
0.980	-0.480	-0.440
0.975	-0.614	-0.563

Table 8. Kinetic Data for Water/1-Butanol Mixtures

x_{water}	$\ln(k/k_0)$	$\ln(k/k_0) - 2 \ln x_{\text{water}}$
0.997	-0.0786	-0.0735
0.995	-0.171	-0.161
0.993	-0.244	-0.229
0.990	-0.337	-0.317
0.987	-0.442	-0.416

Table 9. Kinetic Data for Water/2-Methyl-2-propanol Mixtures

x_{water}	$\ln(k/k_0)$	$\ln(k/k_0) - 2 \ln x_{\text{water}}$
0.995	-0.120	-0.110
0.990	-0.261	-0.241
0.985	-0.403	-0.373
0.980	-0.587	-0.546
0.976	-0.749	-0.700