

University of Nebraska - Lincoln  
**DigitalCommons@University of Nebraska - Lincoln**

---

Nutrition and Health Sciences -- Faculty  
Publications

Nutrition and Health Sciences, Department of

---

8-25-2015

# Electromyographic Responses from the Vastus Medialis during Isometric Muscle Actions

C. M. Smith

*University of Nebraska-Lincoln*

Terry J. Housh

*University of Nebraska-Lincoln*, [thoush1@unl.edu](mailto:thoush1@unl.edu)

Trent J. Herda

*University of Kansas*, [t.herda@ku.edu](mailto:t.herda@ku.edu)

Jorge M Zuniga


*Creighton University*, [jorgezuniga@creighton.edu](mailto:jorgezuniga@creighton.edu)

Clayton L. Camic

*University of Wisconsin-La Crosse*, [ccamic@uwlax.edu](mailto:ccamic@uwlax.edu)

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/nutritionfacpub>

 Part of the [Human and Clinical Nutrition Commons](#), [Molecular, Genetic, and Biochemical Nutrition Commons](#), and the [Other Nutrition Commons](#)

---

Smith, C. M.; Housh, Terry J.; Herda, Trent J.; Zuniga, Jorge M; Camic, Clayton L.; Bergstrom, Haley; Smith, D.B.; Weir, John R.; Cramer, Joel T.; Hill, E.C.; Cochrane, Kristen C.; Jenkins, N. D. M.; Schmidt, R.; and Johnson, G., "Electromyographic Responses from the Vastus Medialis during Isometric Muscle Actions" (2015). *Nutrition and Health Sciences -- Faculty Publications*. 218.  
<https://digitalcommons.unl.edu/nutritionfacpub/218>

This Article is brought to you for free and open access by the Nutrition and Health Sciences, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nutrition and Health Sciences -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

C. M. Smith, Terry J. Housh, Trent J. Herda, Jorge M Zuniga, Clayton L. Camic, Haley Bergstrom, D.B. Smith, John R. Weir, Joel T. Cramer, E.C. Hill, Kristen C. Cochrane, N. D. M. Jenkins, R. Schmidt, and G. Johnson

# Electromyographic Responses from the Vastus Medialis during Isometric Muscle Actions

## Authors

C. M. Smith<sup>1</sup>, T. J. Housh<sup>1</sup>, T. Herda<sup>2</sup>, J. M. Zuniga<sup>3</sup>, C. L. Camici<sup>4</sup>, H. C. Bergstrom<sup>1</sup>, D. B. Smith<sup>5</sup>, J. P. Weir<sup>2</sup>, J. T. Cramer<sup>6</sup>, E. C. Hill<sup>7</sup>, K. C. Cochrane<sup>1</sup>, N. D. M. Jenkins<sup>1</sup>, R. J. Schmidt<sup>1</sup>, G. Johnson<sup>1</sup>

## Affiliations

Affiliation addresses are listed at the end of the article

## Key words

- VM
- innervation zone
- MVIC
- EMG
- IZ
- electrode placement

## Abstract

This study examined the electromyographic (EMG) responses from the vastus medialis (VM) for electrodes placed over and away from the innervation zone (IZ) during a maximal voluntary isometric contraction (MVIC) and sustained, submaximal isometric muscle action. A linear electrode array was placed on the VM to identify the IZ and muscle fiber pennation angle during an MVIC and sustained isometric muscle action at 50% MVIC. EMG amplitude and frequency parameters were determined from 7 bipolar channels of the electrode array, including over the IZ, as well as 10 mm, 20 mm and 30 mm prox-

imal and distal to the IZ. There were no differences between the channels for the patterns of responses for EMG amplitude or mean power frequency during the sustained, submaximal isometric muscle action; however, there were differences between channels during the MVIC. The results of the present study supported the need to standardize the placement of electrodes on the VM for the assessment of EMG amplitude and mean power frequency. Based on the current findings, it is recommended that electrode placements be distal to the IZ and aligned with the muscle fiber pennation angle during MVICs, as well as sustained, submaximal isometric muscle actions.

## Introduction

The development of the linear array of electrodes has allowed for surface electromyographic (EMG) electrode placement guidelines that consider the location of the innervation zone (IZ: connection between nerve terminals and muscle fibers) and muscle fiber pennation angle. Original guidelines often suggested that electrodes be located over the largest portion of the muscle belly, because this location resulted in the greatest EMG amplitude values [13]. Electrode placements over the muscle belly can, however, result in large variations in EMG signal intensities due to interference from the IZ, which is usually near the muscle belly [1,24,29,30]. Typically, electrode placements over the IZ result in lower EMG amplitude (i.e., root mean square: RMS) and greater EMG mean power frequency (MPF) values than when the electrodes are placed away from the IZ [29]. The lower EMG RMS values recorded over the IZ result from signal cancellation associated with bipolar electrode placements [21,30]. That is, motor unit action potentials (MUAP) propagate in opposite directions from the IZ and result in

MUAPs reaching each electrode simultaneously causing greater signal cancellation [29,30]. The greater EMG MPF values recorded over the IZ are due to non-propagating MUAPs that occur at the IZ. Recent electrode placement guidelines [14,24,29,30] have generally suggested avoiding the IZ due to the large variations in EMG signal intensities; however, not all guidelines [13,14,33] have considered the pennation angle of the muscle fibers. Misalignment of electrodes relative to the muscle fiber pennation angle can lead to over- and under-estimates of the frequency content of the EMG signal [7].

A number of studies have used the vastus medialis (VM) to examine neuromuscular function during isometric muscle actions [3,5,9,10,22,28,29,31], but little is known about the effects of the IZ and muscle fiber pennation angle on EMG parameter responses during a maximal voluntary isometric contraction (MVIC) or sustained isometric muscle actions. Beck et al. [3] reported differences in absolute EMG amplitude and MPF values as a result of step incremental muscle actions at 10–90% MVIC recorded over the IZ compared to away from the IZ, but the differ-

accepted after revision  
August 24, 2015

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0035-1564174>  
Published online:  
May 13, 2016  
Int J Sports Med 2016; 37:  
647–652 © Georg Thieme  
Verlag KG Stuttgart · New York  
ISSN 0172-4622

## Correspondence

**Cory Michael Smith**  
Nutrition and Health Sciences  
University of Nebraska-Lincoln  
110 Ruth Levertton Hall  
Lincoln  
United States 68583  
Tel.: +1/402/472 2690  
Fax: +1/402/472 0589  
csmith@unl.edu

ences disappeared when normalized to values at MVIC. Hedayatpour et al. [12] found differences between proximal, middle and distal electrode placements on the VM aligned with the muscle fiber pennation angle for the fatigue-related patterns of responses of EMG amplitude as a result of a sustained isometric muscle action at 80% MVIC. Yeung et al. [32], however, placed bipolar EMG electrodes "...at the most palpable part of the muscle belly of the VM" (p. 381) and reported a 15% decrease in MVIC, but no changes in EMG amplitude or MPF after 30 repeated MVICs. Gallina et al. [9] found that the EMG amplitude and MPF responses from the VM for electrode placements over the IZ and aligned with the muscle fiber pennation angle were less sensitive to variations in knee flexion angle than when the electrodes were placed away from the IZ. Furthermore, Nozic et al. [25] showed that there were differences in the pennation angles of the proximal (12°) vs. distal (52°) portions of the VM [25], and it has been reported that EMG parameters from electrode placements over the proximal portion of the VM were less reliable than those from placements over the distal portion. These studies [3,9,12,25,32] indicated that electrode placements on the VM can affect reliability and measured EMG parameter responses during MVIC and step incremental muscle actions, as well as the fatigue-related patterns of responses during submaximal sustained isometric muscle actions. Thus, it is important that there are standardized electrode placement guidelines for the VM to reduce the influences of the IZ and muscle fiber pennation angle on the EMG signal, as well as to allow valid comparisons of the results of previous studies. Therefore, the purpose of the present study was to: 1) examine the responses of the EMG time and frequency domain parameters from the VM when electrode placements are located over and away from the IZ during an MVIC and sustained submaximal isometric muscle action of the leg extensors; and 2) propose electrode placement guidelines for the VM that consider the location of the IZ and muscle fiber pennation angle.

## Material and Methods



### Subjects

9 healthy adults (7 men and 2 women, mean±SD age=22.6±2.4 years; body mass=82.2±14.8 kg; height=181.3±10.5 cm) volunteered to participate in the study. All subjects were free of any knee, hip, or ankle pain. The subjects regularly participated in physical activities such as running, bicycling, and resistance training. The study was approved by the University Institutional Review Board for Human Subjects, and all subjects completed a health history questionnaire and signed an informed consent document prior to testing. In addition, the present study was in compliance with the Ethical Standards described by Harris et al. [11].

### Orientation session

The orientation session was used to determine the location of the IZ and the pennation angle of the muscle fibers of the VM. The skin was dry shaven and cleaned with a damp cloth prior to applying a probe with 8 silver bar electrodes (5 mm×1 mm, 10 mm interelectrode distance, Ottino Bioelectronica, Torino, Italy) over the VM on a reference line between the anterior spina iliaca superior and the anterior border of the medial ligament [29]. The length of this reference line was recorded (● Table 1). The reference electrodes were placed around the subject's wrist according to the procedures described by the EMG16 User Man-

**Table 1** The location of the innervation zone and muscle fiber pennation angle identified by a linear array. The SENIAM [12] recommendation suggests placing the electrodes at 20% the distance between the anterior spina iliaca superior and the anterior border of the medial ligament. The SENIAM and Actual IZ were measured from the anterior border of the medial ligament on the reference line.

Subject	Gender	Reference Line (cm)	SENIAM (cm)	Actual IZ (cm)	Pennation Angle (degrees)
1	Male	53	10.6	12	61
2	Female	57	11.4	15	54
3	Male	51	10.2	12	43
4	Male	51	10.2	11.5	50
5	Female	51	10.2	10	62
6	Male	48	9.7	11	50
7	Male	49	9.8	9	50
8	Male	52	10.4	13.5	50
9	Male	46	9.2	10	55
Mean		50.89	10.19	11.56	52.78
SD		3.14	0.62	1.86	5.97

ual [18]. The subjects were instructed to perform a submaximal isometric contraction of their leg extensors to identify their IZ. The location of the IZ was identified by the EMG channel with the minimal amplitude and phase reversal [23]. The distance of the IZ from the anterior border of the medial ligament was recorded (● Table 1). The probe was then moved along the muscle until the IZ was in the center of the electrode array. The pennation angle of the muscle fibers of the VM was determined by rotating the probe around the IZ until the slopes of the 2 lines connecting the EMG waveforms from the channels above and below the IZ had symmetrical propagation of MUAPs [18]. The muscle fibers pennation angle was then recorded for each subject, and an outline of the probe was traced with a non-washable marker before removing the probe so the same electrode position could be obtained in the subsequent session.

### Warm-up

All isometric leg extension testing was performed on the dominant leg based on kicking preference (all subjects were right-leg dominant), using a calibrated Cybex II dynamometer at a knee joint angle of 120°. Prior to the isometric testing, each subject performed a warm-up of five 6-s submaximal isometric muscle actions, followed by a 2-min rest period. The subjects were instructed to provide an effort corresponding to approximately 50% of their maximum during each muscle action.

### Maximum voluntary isometric contraction

After completing the warm-up, each subject performed 2, 6-s MVIC trials. A 2-min rest was given after each trial. Strong verbal encouragement was provided for all trials. The MVIC was calculated for a 2-s time period corresponding to the middle 33% of each 6-s trial. The highest torque value of the 2 MVIC trials was used to calculate each 50% MVIC for the subsequent sustained isometric task.

### Sustained isometric task

Each subject was instructed to perform a sustained isometric muscle action to exhaustion at 50% MVIC. Exhaustion was operationally defined as the subject being unable to maintain a torque value within ±5% of their 50% MVIC. The subjects tracked their torque production on a computer monitor placed in front

of them that displayed the real-time, digitalized torque signal overlaid onto a programmed template identifying their target torque value. The isometric template and real-time torque signal overlay were programmed using LabVIEW (LabVIEW version 7.1 National Instruments, Austin, TX).


**Electromyographic measurements and signal processing**

Surface EMG signals were recorded from the dominant VM muscle with an 8-channel linear electrode array and EMG 16 data acquisition system (EMG 16, LISiN-Prima Biomedical & Sport, Treviso, Italy). The skin over the VM was carefully abraded and cleaned with rubbing alcohol prior to securing the 8-channel linear electrode array to the VM using a double-sided adhesive strip. The adhesive strip had small holes that were cut for each silver bar electrode, and each hole was filled with 30 µl of conductive gel with a gel dispenser (AG22331, Eppendorf, Hamburg, Germany). The raw EMG signals from each electrode of the probe were recorded in a monopolar signal acquisition mode (gain×500) and analog filtered (fourth-order Bessel, bandwidth = 10–500 Hz) with the surface EMG 16 data acquisition system. Electrode 1 corresponded to the most proximal and electrode 8 to the most distal electrode on the linear array. The monopolar EMG signals were converted to a digital form with a 12-bit analog-to-digital converter at a sampling frequency of 2048 Hz and stored on a personal computer for subsequent analyses.

7 bipolar electrode configurations (10 mm interelectrode distance) were obtained by differentiating the digitized monopolar signals forming (● Fig. 1): channel 1 from electrodes 1 and 2 (30 mm proximal to the IZ), channel 2 from electrodes 2 and 3 (20 mm proximal to the IZ), channel 3 from electrodes 3 and 4 (10 mm proximal to the IZ), channel 4 from electrodes 4 and 5 (over the IZ), channel 5 from electrodes 5 and 6 (10 mm distal to the IZ), channel 6 from electrodes 6 and 7 (20 mm distal to the IZ), and channel 7 from electrodes 7 and 8 (30 mm distal to the IZ). The EMG signals were zero-meaned, and bandpass filtered (fourth-order Butterworth) at 10–500 Hz. The EMG RMS and MPF of the middle 33% (2-s epochs) of the MVIC were calculated for each channel. The EMG RMS and MPF values for the sustained isometric muscle action were calculated using 0.5-s epochs extracted every 5%, beginning at 10%, of the normalized time to exhaustion. All signal processing was performed using custom programs written with LabVIEW programming software (Version 13.0, National Instruments, Austin TX).

**Statistical analysis**

Separate one-way (1×7) repeated-measures ANOVAs were performed on the absolute EMG RMS and MPF from the MVIC to determine if there were mean differences among the channels. Significant main effects for time were followed by Tukey LSD corrected dependent sampled *t*-tests on the simple main comparisons. The EMG RMS and MPF from the sustained isometric muscle action were normalized to the MVIC at each electrode location. Linear regression analyses were then used to determine the EMG RMS and MPF vs. normalized time to exhaustion relationships during the sustained isometric muscle action for each channel. The slope coefficients for the EMG RMS and MPF vs. normalized time to exhaustion relationships during the sustained isometric muscle action for all 7 channels for each subject and the composite of all subjects were compared statistically using the F-test procedures of Pedhazur [26]. An alpha of  $p \leq 0.05$  was considered statistically significant for all comparisons.



**Fig. 1** Linear array placement depicting the distance and electrodes that create channels 1 through 7.

**Table 2** Absolute electromyographic mean power frequency (EMG MPF) and root mean square (EMG RMS) values for the maximal voluntary isometric contraction (MVIC) of the vastus medialis.

	EMG MPF (Hz±SD)	EMG RMS (µV±SD)
Channel 1	69.04±9.71	132.09±17.90
Channel 2	71.38±9.67	155.83±19.65
Channel 3	87.11±16.51	106.03±34.70
Channel 4	111.24±19.98	65.58±23.33
Channel 5	105.86±23.64	84.23±32.14
Channel 6	69.81±10.92	119.37±20.25
Channel 7	69.00±10.95	97.79±27.09


Note: Refer to ● Fig. 1 for channel locations

\* $p \leq 0.05$  (EMG RMS: Channel 1>4, 5, 6, and 7; Channel 2>1, 3, 4, 5, 6, and 7; Channel 4 < 3, 6, and 7; Channel 6>5 and 7; EMG MPF: Channel 1<3, 4, and 5; Channel 2>3, 4, and 5; Channel 3>6 and 7; Channel 4>3, 6, and 7; Channel 5>6 and 7)


**Results**

**Maximal voluntary isometric contraction electromyographic measurements**

● Table 2 shows the absolute EMG RMS and MPF values from the 7 channels during the MVIC muscle actions. For EMG RMS, there were significant differences among channels for 15 of the 21 comparisons (● Table 2). For EMG MPF, there were significant differences among channels for 12 of the 21 comparisons (● Table 2).



**Fig. 2** The positive relationship between electromyographic root mean square (EMG RMS) from the vastus medialis and percent time to exhaustion during the sustained isometric muscle action at 50% MVIC. There were no significant differences ( $p > 0.05$ ) in slope coefficients between any of the 7 channels.



**Fig. 3** The negative relationship between electromyographic mean power frequency (EMG MPF) from the vastus medialis and percent time to exhaustion during the sustained isometric muscle action at 50% MVIC. There were no significant differences ( $p > 0.05$ ) in slope coefficients between any of the 7 channels.

### Electromyographic root mean square and mean power frequency patterns of responses during sustained isometric muscle action

There were significant positive relationships for normalized EMG RMS vs. normalized time to exhaustion for the composite of all subjects for all channels (◉ Fig. 2). Furthermore, there were no significant differences among slope coefficients for any of the channels (◉ Fig. 2).

There were significant negative relationships for normalized EMG MPF vs. normalized time to exhaustion for the composite of all subjects for all channels (◉ Fig. 3). Furthermore, there were no significant differences among slope coefficients for any of the channels (◉ Fig. 3).

### Location of the innervation zone and pennation angle for the vastus medialis

The reference line between the anterior border of the medial ligament and the anterior spina iliaca superior was  $50.89 \pm 3.14$  cm. The IZ was  $11.56 \pm 1.86$  cm proximal to the anterior border of the medial ligament (◉ Table 1). The mean muscle fiber pennation angle was  $52.78 \pm 5.97^\circ$  (◉ Table 1).

### Discussion

#### Electromyographic root mean square and mean power frequency during maximal voluntary isometric contraction

Electrode placement recommendations for the VM suggest that bipolar arrangements should be located distal to the IZ to avoid confounding influences that may affect absolute and normalized EMG RMS and MPF values [13, 14, 29, 30]. Electrode placements located over the IZ typically result in lower EMG RMS and higher EMG MPF values compared to those away from the IZ [29, 30]. The lower EMG RMS values are a result of MUAPs that travel in opposite directions from the IZ, simultaneously reach each of the bipolar electrodes, and are cancelled by differential amplification [2]. The greater EMG MPF values over the IZ, compared to away from the IZ, are the result of non-propagating MUAPs that are only detectable over the IZ [2]. Rainoldi et al. [29] showed that for the VM, a change in electrode placement of 10 mm proximal or distal to the IZ can result in EMG amplitude changes in excess of 200%. In addition, Beck et al. [3] showed a 200% decrease in EMG amplitude and 20% increase in EMG MPF when recorded over the IZ compared to 30 mm distal to the IZ of the VM. The current findings indicated that during MVIC muscle actions, electrode placements over the IZ resulted in up to 58% lower EMG RMS and 37% greater EMG MPF values than those recorded proximal or distal to the IZ. Specifically, there were differences in the EMG RMS and MPF values for channel 4 (over the IZ) vs. channels 1, 2, 3, 6, and 7 (◉ Table 2). Furthermore, there were differences among electrode placements proximal to the IZ (channels 1, 2, and 3), as well as those distal to the IZ (channels 5, 6, and 7). Thus, the present findings indicated that the IZ should be avoided during MVIC measurements due to large variations in EMG RMS and MPF values compared to electrode placements away from the IZ (◉ Table 2).

The findings of the present study, in conjunction with previous studies [9, 13, 14, 24, 29, 30], suggested that electrode placement recommendations should be developed on a muscle by muscle basis, because different muscles are characterized by a variety of factors that can affect the validity and reliability of EMG parameters measured during an MVIC including the [1, 9, 16, 20]: 1) number of IZs; 2) location of the IZ; 3) amount of movement or shift of a muscle relative to the skin surface during a muscle action; 4) amount of crosstalk from nearby muscles; 5) location of electrode placements relative to the tendinous region; and 6) muscle fiber pennation angle. The VM has only one IZ that is located at approximately the midpoint of the muscle where it begins to taper proximally, near the location where the sartorius muscle overlaps the proximal portion of the VM [19]. The proximal tapering of the VM, location of the IZ, and overlapping of the sartorius muscle result in a larger EMG recording area at the distal end of the VM than at the proximal end. The larger recording area makes distal electrode placements less likely to be over a tendinous region and affected by crosstalk from the surrounding

muscles, including the sartorius. In addition, it has been suggested [9, 20, 30] that the VM shifts proximally during a muscle action which can cause proximal electrode placements to move over the IZ and result in lower EMG RMS and greater EMG MPF values than electrode placements distal to the IZ [9, 20, 30]. Gallina et al. [9] reported that when performing an MVIC, a small proximal shift in the IZ of the VM resulted in 50–75% lower EMG RMS and 25% greater EMG MPF when electrode placements were located 10 mm proximal to the IZ compared to those 10 mm distal to the IZ. It has also been suggested that the muscle fibers of the VM have different pennation angles distal and proximal to the IZ, with the distal portion being angled at approximately 50° and the proximal at approximately 35–40° [9, 16, 20, 24, 29, 31]. In the present study, the average pennation angle for the VM was 53° with a range of 43–62°, which was in agreement with previous studies that used MRI, dissection, muscle mapping and linear array to measure the VM pennation angle [1, 15, 27, 29]. It is thus recommended that when recording EMG parameters from the VM during an MVIC without a linear array, electrode placements should be located between 10 and 30 mm distal to the midpoint of the muscle to avoid the IZ, and orientated at 53° to approximate the pennation angle of the muscle fibers.

### Electromyographic root mean square and mean power frequency patterns of responses during sustained submaximal isometric muscle actions

Fatiguing submaximal isometric muscle actions are typically characterized by increases in EMG RMS and decreases in EMG MPF [23]. These fatigue-related neuromuscular responses have been attributed to decreases in muscle pH caused by the buildup of metabolic byproducts (hydrogen ions, inorganic phosphate, potassium, and ammonia), which alter muscle contractility resulting in greater muscle activation to maintain a constant force or torque [4, 17]. In addition to a decrease in muscle pH [4], fatigue-induced decreases in EMG MPF have also been associated with increases in extracellular potassium, which results in a progressive loss of membrane excitability [8]. In the present study, increases in EMG RMS and decreases in EMG MPF occurred for all electrode placements (channels 1, 2, 3, 4, 5, 6, and 7) and there were no differences among the slope coefficients for the normalized EMG RMS and EMG MPF vs. time relationships for the composite of all subjects (● Fig. 2,3). These findings were in agreement with those of Silva et al. [31], who reported similar increases in EMG RMS and decreases in EMG MPF from the VM during a sustained isometric muscle action at 50% MVIC of the leg extensors. Thus in the present study, the EMG RMS and EMG MPF vs. time relationships during submaximal isometric muscle actions of the leg extensors for the composite of all subjects suggested that the buildup of metabolic byproducts affected the EMG RMS and EMG MPF from the VM to the same degree for all electrode placements. Proximal electrode placements on the VM can be affected by the shifting of the muscle during an MVIC such that the electrodes become located over the IZ. It has been suggested, however, that the distance of the shift is positively related to torque production. In the present study, any proximal shift of the VM during the sustained isometric muscle action at 50% MVIC had no effect on the patterns or magnitude of the normalized EMG responses. These findings supported those of De Freitas et al. [6] and Beck et al. [3] who reported that normalization to MVIC minimized the effects of electrode movement rela-

tive to the muscle on EMG time and frequency domain responses during submaximal isometric muscle actions.

### Summary

Based on the findings of the present study, it is recommended that during an MVIC muscle action, bipolar electrode placements should be located 10–30 mm distal to the midpoint of the VM to avoid the IZ and oriented at 53° to approximate the muscle fiber pennation angle. Although the EMG RMS and EMG MPF vs. time relationships were not affected by electrode placement, this placement is also recommended during sustained submaximal muscle actions so that MVICs can be measured to normalize the data. The current findings indicated that single EMG RMS and EMG MPF measurements from MVICs were affected by electrode placements relative to the IZ, while the EMG RMS and EMG MPF vs. time relationships during the sustained isometric muscle action were not.

**Conflict of interest:** The authors have conflict of interest to declare.

### Affiliations

- <sup>1</sup> Nutrition and Health Sciences, University of Nebraska-Lincoln, Lincoln, United States
- <sup>2</sup> Health, Sport and Exercise Science, University of Kansas, Lawrence, United States
- <sup>3</sup> Exercise Science, Creighton University, Omaha, United States
- <sup>4</sup> Exercise & Sport Science, University of Wisconsin – La Crosse, La Crosse, United States
- <sup>5</sup> Health and Human Performance, Oklahoma State University, Stillwater, United States
- <sup>6</sup> Department of Nutrition & Health Sciences, University of Nebraska-Lincoln, Lincoln, United States
- <sup>7</sup> Nutrition and Health Sciences, Ethan Hill, Lincoln, United States

### References

- 1 Barbero M, Merletti R, Rainoldi A. Atlas of muscle innervation zones: understanding surface electromyography and its applications. Springer Science & Business Media 2012
- 2 Basmajian JV, De Luca C. Muscles alive: their functions revealed by electromyography. William & Wilkins 1985; 278: 126
- 3 Beck TW, Housh TJ, Cramer JT, Mielke M, Hendrix R. The influence of electrode shift over the innervation zone and normalization on the electromyographic amplitude and mean power frequency versus isometric torque relationships for the vastus medialis muscle. *J Neurosci Methods* 2008; 169: 100–108
- 4 Brody LR, Pollock MT, Roy SH, De Luca CJ, Celli B. pH-induced effects on median frequency and conduction velocity of the myoelectric signal. *J Appl Physiol* 1991; 71: 1878–1885
- 5 Chan AY, Lee FL, Wong PK, Wong CY, Yeung SS. Effects of knee joint angles and fatigue on the neuromuscular control of vastus medialis oblique and vastus lateralis muscle in humans. *Eur J Appl Physiol* 2001; 84: 36–41
- 6 DeFreitas JM, Costa PB, Ryan ED, Herda TJ, Cramer JT, Beck TW. An examination of innervation zone movement with increases in isometric torque production. *Clin Neurophysiol* 2008; 119: 2795–2799
- 7 Farina D, Cescon C, Merletti R. Influence of anatomical, physical, and detection-system parameters on surface EMG. *Biol Cybern* 2002; 86: 445–456
- 8 Fortune E, Lowery MM. The effect of extracellular potassium concentration on muscle fiber conduction velocity examined using model simulation. Conference proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society 2007; 2726–2729
- 9 Gallina A, Merletti R, Gazzoni M. Innervation zone of the vastus medialis muscle: position and effect on surface EMG variables. *Physiol Meas* 2013; 34: 1411–1422
- 10 Grabiner MD, Koh TJ, Miller GF. Fatigue rates of vastus medialis oblique and vastus lateralis during static and dynamic knee extension. *J Orthop Res* 1991; 9: 391–397

- 11 *Harriss DJ, Atkinson G.* Ethical standards in sport and exercise science research: 2016 update. *Int J Sports Med* 2015; 36: 1121–1124
- 12 *Hedayatpour N, Arendt-Nielsen L, Farina D.* Non-uniform electromyographic activity during fatigue and recovery of the vastus medialis and lateralis muscles. *Journal of electromyography and kinesiology: J Electromyogr Kinesiol* 2008; 18: 390–396
- 13 *Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G.* Development of recommendations for SEMG sensors and sensor placement procedures. *Journal of electromyography and kinesiology: J Electromyogr Kinesiol* 2000; 10: 361–374
- 14 *Hermens HJ, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, Disselhorst-Klug C, Hägg G.* European recommendations for surface electromyography. *Roessingh Res Develop* 1999; 8: 13–54
- 15 *Hubbard JK, Sampson HW, Elledge JR.* Prevalence and morphology of the vastus medialis oblique muscle in human cadavers. *Anat Rec* 1997; 249: 135–142
- 16 *Kollmitzer J, Ebenbichler GR, Kopf A.* Reliability of surface electromyographic measurements. *Clin Neurophysiol* 1999; 110: 725–734
- 17 *Lindstrom L, Magnusson R, Petersen I.* Muscular fatigue and action potential conduction velocity changes studied with frequency analysis of EMG signals. *Electromyography* 1970; 10: 341–356
- 18 *Manual EUs.* EMG16 User Manual. 16-channels surface electromyographic signal amplifier. LISiN Bioengineering Center Polytechnic of Turin, Department of Electronics Turin, Italy 2006
- 19 *Martini FH, Timmons MJ, Robert B.* Human Anatomy San Francisco, PA Pearson and Benjamin Cummings 2005
- 20 *Mathur S, Eng JJ, MacIntyre DL.* Reliability of surface EMG during sustained contractions of the quadriceps. *J Electromyogr Kinesiol* 2005; 15: 102–110
- 21 *Merletti R, Farina D, Gazzoni M.* The linear electrode array: a useful tool with many applications. *J Electromyogr Kinesiol* 2003; 13: 37–47
- 22 *Merletti R, Fiorito A, Lo Conte LR, Cisari C.* Repeatability of electrically evoked EMG signals in the human vastus medialis muscle. *Muscle Nerve* 1998; 21: 184–193
- 23 *Merletti R, Knafitz M, De Luca CJ.* Myoelectric manifestations of fatigue in voluntary and electrically elicited contractions. *J Appl Physiol* (Bethesda, Md: 1985) 1990; 69: 1810–1820
- 24 *Mesin L, Merletti R, Rainoldi A.* Surface EMG: the issue of electrode location. *Journal of electromyography and kinesiology: J Electromyogr Kinesiol* 2009; 19: 719–726
- 25 *Nozic M, Mitchell J, de Klerk D.* A comparison of the proximal and distal parts of the vastus medialis muscle. *Aust J Physiother* 1997; 43: 277–281
- 26 *Pedhazur E.* Multiple Regression in Behavioral Research. USA: Thomson Learning Inc; 1982
- 27 *Peeler J, Cooper J, Porter MM, Thliveris JA, Anderson JE.* Structural parameters of the vastus medialis muscle. *Clin Anat* 2005; 18: 281–289
- 28 *Rainoldi A, Bullock-Saxton JE, Cavarretta F, Hogan N.* Repeatability of maximal voluntary force and of surface EMG variables during voluntary isometric contraction of quadriceps muscles in healthy subjects. *J Electromyogr Kinesiol* 2001; 11: 425–438
- 29 *Rainoldi A, Melchiorri G, Caruso I.* A method for positioning electrodes during surface EMG recordings in lower limb muscles. *J Neurosci Methods* 2004; 134: 37–43
- 30 *Rainoldi A, Nazzaro M, Merletti R, Farina D, Caruso I, Gaudenti S.* Geometrical factors in surface EMG of the vastus medialis and lateralis muscles. *J Electromyogr Kinesiol* 2000; 10: 327–336
- 31 *Silva CRd, Silva DdO, Ferrari D, Negrão Filho RdF, Alves N, Azevedo FMd.* Exploratory study of electromyographic behavior of the vastus medialis and vastus lateralis at neuromuscular fatigue onset. *Motriz* 2014; 20: 213–220
- 32 *Yeung SS, Au AL, Chow CC.* Effects of fatigue on the temporal neuromuscular control of vastus medialis muscle in humans. *Eur J Appl Physiol* 1999; 80: 379–385
- 33 *Zipp P.* Recommendations for the standardization of lead positions in surface electromyography. *Eur J Appl Physiol* 1982; 50: 41–54