

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Papers in Veterinary and Biomedical Science

Veterinary and Biomedical Sciences,
Department of

2-9-1999

NUCLEOTIDE SEQUENCES FOR DETECTION OF SERPULINA HYODYSENTERIAE

Gerald E. Duhamel

University of Nebraska- Lincoln

Robert Elder

University of Nebraska- Lincoln

Follow this and additional works at: <https://digitalcommons.unl.edu/vetscipapers>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#), [Cell and Developmental Biology Commons](#), [Immunology and Infectious Disease Commons](#), [Medical Sciences Commons](#), [Veterinary Microbiology and Immunobiology Commons](#), and the [Veterinary Pathology and Pathobiology Commons](#)

Duhamel, Gerald E. and Elder, Robert, "NUCLEOTIDE SEQUENCES FOR DETECTION OF SERPULINA HYODYSENTERIAE" (1999). *Papers in Veterinary and Biomedical Science*. 342.

<https://digitalcommons.unl.edu/vetscipapers/342>

This Article is brought to you for free and open access by the Veterinary and Biomedical Sciences, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Veterinary and Biomedical Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



US005869630A

United States Patent [19]

[11] Patent Number: **5,869,630**

Duhamel et al.

[45] Date of Patent: **Feb. 9, 1999**

[54] **NUCLEOTIDE SEQUENCES FOR DETECTION OF SERPULINA HYODYSENTERIAE**

[75] Inventors: **Gerald E. Duhamel; Robert Elder,** both of Lincoln, Nebr.

[73] Assignee: **Board of Regents, University of Nebraska Lincoln,** Lincoln, Nebr.

[21] Appl. No.: **727,126**

[22] Filed: **Oct. 8, 1996**

Related U.S. Application Data

[62] Division of Ser. No. 252,492, Jun. 1, 1994.

[51] Int. Cl.⁶ **C07H 21/02; C07H 21/04; C12Q 1/68; C12P 19/34**

[52] U.S. Cl. **536/22.1; 536/231; 536/243; 536/24.31; 536/24.32; 536/24.33; 435/6; 435/91.2**

[58] Field of Search **435/6, 91.1, 91.2, 435/91.21; 536/22.1, 23.1, 24.3, 24.31, 24.32, 24.33**

[56] **References Cited**

U.S. PATENT DOCUMENTS

4,469,672	9/1984	Harris	424/23
4,683,202	7/1987	Mullins .	
4,748,019	5/1988	Lysons .	
4,764,370	8/1988	Fields et al. .	
4,818,680	4/1989	Collins et al. .	
4,851,331	7/1989	Vary et al. .	
4,999,191	3/1991	Glisson et al. .	
5,034,315	7/1991	Jensen et al.	435/6
5,298,392	3/1994	Atlas et al. .	
5,306,616	4/1994	Lupski et al. .	

FOREIGN PATENT DOCUMENTS

282965	9/1988	European Pat. Off. .	
350715	1/1990	European Pat. Off. .	
WO 88/04778	6/1988	WIPO .	
WO 90/02565	3/1990	WIPO .	
WO 91/04036	4/1991	WIPO .	

OTHER PUBLICATIONS

Achacha et al., *J. Vet. Diag. Invest.*, 3:211-214 (1991).
 Adachi et al., *Zentralbl. Bakteriologie*, 245:527 (1979).
 Baum et al., *Infect. Immun.*, 25:792-796 (1979).
 Bej et al., "Amplification of Nucleic Acids by Polymerase Chain Reaction (PCR) and Other Methods and Their Applications", *Critic. Rev. Biochem. Mol. Biol.*, 26(3/4):301-334 (1991).
 Belanger et al., *J. Clin. Microbiol.*, 29:1727-1729 (1991).
Bio/Technique, 6:693 (1988).
 Boyden et al., "Cloning and Characterization of *Treponema hyodysenteriae* Antigens and Protection in a CF-1 Mouse Model by Immunization with a Cloned Endoflagellar Antigen," *Infect. Immun.*, 57:3808-3815 (1989).
 Boeckman, "New Swine Dysentery Vaccines May be Just Around the Corner," *Swine Practitioner* at pages 4-6, Jan., 1994.
 Burrows et al., *Vet. Rec.*, 108:187-189 (1981).

Caputa et al., *J. Clin. Microbiol.*, 29:2418-2413 (1991).
 Chatfield et al., "Identification of the Major Antigens of *Treponema hyodysenteriae* and Comparison with Those of *Treponema innocens*," *Inf. & Immun.*, 56:1070-1075 (1988).
 Chenngappa et al., "Laboratory Procedures for the Diagnosis of Swine Dysentery," *Report of the Committee on Swine Dysentery*, American Association of Veterinary Laboratory Diagnosticians, Committee on Swine Dysentery, Aug. (1989).
 Chien et al., *J. Bacteriol.*, 127:1550 (1976).
 Combs et al., *Res. Vet. Sci.*, 50:286-289 (1991).
 Curtiss III et al., *Infection and Immunity*, 55:3035 (1987).
 Cwyk et al., *Arch. Microbiol.*, 122:231-239 (1979).
 Dettori et al., "Molecular Cloning and Characterization of DNA from Human Intestinal Spirochetes," *Europ. J. Epidemiol.*, 8(2):198-205 (1992).
 Dougan et al., *J. Infect. Dis.*, 158:1329 (1988).
 Duhamel et al., "Serodiagnosis of Swine Dysentery Using Recombinant Proteins from *Treponema Hyodysenteriae*," *Nebraska Veterinary Science Newsletter*, 19:3-4, Jun. 1990.
 Duhamel et al., "Update on Prevention and Control of Swine Dysentery," *31st George A. Young Swine Conference and Annual Nebraska SPF Swine Conference*, Lincoln, NE, at pp. 7-20, Aug. 13-14 (1990).
 Duhamel et al., "Freedom From Dysentery", *Swine Practitioner*, at pp. 4-5, 8, and 10, Sep. 1990.
 Duhamel et al., "Application to Restriction Length Fragment Length Polymorphism Typing to Epidemiological Tracing of *Serpulina (Treponema) hyodysenteriae*," *Proc. 12th Int. Cong. Pig. Vet. Soc.*, The Hague, Netherlands, vol. 1, p. 276 (1992).
 Duhamel et al., *J. Vet. Diagn. Invest.*, 4:285-292 (1992).
 Duhamel et al., "Herd Health: Swine Dysentery Probe, Promising Cleanup Tool," *National Hog Farmer*, pp. 48-50, Dec. 15, 1993.
 Edwards et al., *J. Bacteriol.*, 170:3991 (1988).
 Elder et al., "Rapid Identification of *Serpulina hyodysenteriae* By Polymerase Chain Reaction," 53rd Annual North Central Branch Meeting, American Society for Microbiology, Oct. 28-30, at page 21 (1993).

(List continued on next page.)

Primary Examiner—Stephanie W. Zitomer
 Assistant Examiner—Jeffrey Fredman
 Attorney, Agent, or Firm—Suiter & Associates PC

[57] **ABSTRACT**

The invention provides a method for detecting the presence of *Serpulina hyodysenteriae* in a biological sample, an oligonucleotide primer and an *S. hyodysenteriae*-specific oligonucleotide probe useful in that method, and an article of manufacture that contains the primers and/or probe. Also provided are an about 2.3-kb DNA fragment derived from genomic DNA of *S. hyodysenteriae* and encoding for an about 56 kDa polypeptide, a recombinant expression vector containing the DNA fragment, the 56 kDa polypeptide and a monoclonal antibody reactive with the peptide, and a method of assaying for antibodies reactive with the 56 kDa peptide.

17 Claims, 6 Drawing Sheets

OTHER PUBLICATIONS

- Elder et al., "Rapid Detection of *Serpulina hyodysenteriae* in Diagnostic Specimens By PCR," *J. Clin. Microbiol.*, 32:1497 (1994).
- Fisher et al., *Am. J. Vet. Res.*, 46:450-455 (1981).
- Fisher et al., "Development and Evaluation of a Whole Cell ELISA for Detection of Serum Antibodies to *Treponema hyodysenteriae* in Swine," Abstract No. 70, 71st Conference of Research Workers in Animal Disease, Nov. 5-6 (1990).
- Flores et al., "Identification of Human Rotavirus Serotype by Hybridization to Polymerase Chain Reaction-Generated Probes Derived from a Hyperdivergent Region of the Gene Encoding Outer Capsid Protein VP7," *J. Virol.*, 64:4021-4024 (1990).
- Gabe et al., "Molecular Analysis of the vspA-H Genes Which Comprise a Multigene Family Encoding a 39-kDa Variable Surface Protein from *Serpulina hyodysenteriae*," Abstract D156, 94th ASM General Meeting, May 23-27, 1994.
- Hampson et al., *Epidemiol. Infect.*, 103:275 (1989).
- Harris et al., *Diseases of Swine*, 7th ed., Iowa State University Press (Ames, IA), at pp. 599-616 (1992).
- Hopp et al., *Mol. Immunol.*, 20:483 (1983).
- Hornich et al., in *Vet. Med.*, 24:29 (1979).
- Hugo et al., *J. Clin. Microbiol.*, 25:26-30 (1987).
- Hunter et al., *Vet. Rec.*, 104:383-384 (1979).
- Islam et al., "Detection of *Shigella dysenteriae* Type 1 and *Shigella flexneri* Feces by Immunomagnetic Isolation and Polymerase Chain Reaction," *J. Clin. Micro.*, 30:2801-2806 (1992).
- Jenkins, *Am. J. Vet. Res.*, 41:338 (1980).
- Jenkins et al., *Vet. Immunological Immunopathology*, 2:19 (1981).
- Jenkins et al., *Brit. Vet. Journal*, 145:565 (1989).
- Jensen et al., "Detection and Identification of *Treponema hyodysenteriae* by Using Oligodeoxynucleotide Probes Complementary to 16S rRNA," *J. Clin. Microbiol.*, 28:2717-2721 (1990).
- Joens et al., *Vet. Records*, 106:245 (1980).
- Joens et al., "Molecular Characterization of Proteins from Porcine Spirochetes," *Infect. Imm.*, 54:893-896 (1986).
- Jones et al., *J. Clin. Microbiol.*, 24:1071-1074 (1986).
- Judd et al., *Appl. Env. Microbiol.*, 59:1702 (1993).
- Katz et al., "Colorimetric Diagnosis of Prolonged Blue-tongue Viremia in Sheep, Using an Enzyme-Linked Oligonucleotide Sorbent Assay of Amplified Viral Nucleic Acids," *Am. J. Vet. Res.*, 54:2021-2026. (1993).
- Kent et al., *J. Gen. Micro.*, 135:1625 (1989).
- Kinyon et al., *Infect. Immun.*, 15:638-646 (1977).
- Kinyon et al., *Int. J. Syst. Bact.*, 29:102-109 (1979).
- Kohler et al., *Nature*, 256:496-97 (1975).
- Koopman et al., "Cloning and DNA Sequence Analysis of a *Serpulina* (*Treponema*) *hyodysenteriae* Gene Encoding a Periplasmic Flagellar Sheath Protein," *Infect. Imm.*, 60:2920-2925 (1992).
- Kunkle et al., *J. Clin. Microbiol.*, 24:669-671 (1986).
- Kunkle et al., *J. Clin. Microbiol.* 26:2357-2360 (1988).
- Kunkle et al., "Recombinant DNA Technology in Controlling Swine Dysentery," 1992 *Nebraska Swine Report* at pages 3-4 (1992).
- Lee et al., *Vet. Microbiol.*, 34:35-46 (1993).
- Lemcke et al., *Vet. Rec.*, 14:548-551 (1979).
- Lemcke et al., *J. Hygiene*, 86:173 (1981).
- Li et al., *J. Clin. Microbiol.*, 29:2794-2797 (1991).
- Lory, *J. Bacteriol.*, 174:3423-3428 (1992).
- Luneberg et al., "Detection of *Mycoplasma pneumoniae* by Polymerase Chain Reaction and Nonradioactive Hybridization in Microtiter Plates," *J. Clin. Micro.* 31:1035-1094 (1993).
- Lymbery et al., *Vet. Microbiol.*, 22:89-99 (1990).
- Lysons, *Vet. Rec.*, 129:314-315 (1991).
- Mapother et al., *J. Clin. Micro.*, 22:161-164 (1985).
- Mowat et al., *Immunology Today*, 12:383 (1991).
- Muir et al., "Cloning and Expression of a *Serpula* (*Treponema*) *hyodysenteriae* Hemolysin Gene," *Infect. Immun.*, 60:529-535 (1992).
- Mullis et al., *Methods Enzymol.*, 155:335 (1987).
- Norby, "Veterinary Scientists Hope to Break Swine Dysentery Transmission Cycle," *Research Nebraska!* at pages 16-17, Sep., 1993.
- Owen, *Iowa State J. Res.*, 62:293-311 (1987).
- Picard et al., *Can. J. Microbiol.*, 26:985-991 (1980).
- Quaife, "A New Weapon Against Swine Dysentery," *Swine Practitioner* at pages 10-11, Oct., 1990.
- Ramanathan et al. "Identification and Partial Characterization of a Group of Weakly β -hemolytic Intestinal Spirochetes of Swine Distinct from *Serpulina innocens* Isolate," *Vet. Microbiol.*, 37:53-64 (1993).
- Rychlik et al., *Nucleic Acid Res.*, 17:8543-8551 (1989).
- Saiki et al., *Science*, 239:487 (1988).
- Schafer et al., "Rapid Identification of *Serpulina hyodysenteriae* By Polymerase Chain Reaction Amplification of a DNA Sequence Unique to *Serpulina hyodysenteriae*," 32nd North Central Conference Veterinary Laboratory Diagnosticians, Madison, WI, Jun. 6-9 (1993).
- Schlink et al., *Can. J. Comp. Med.*, 47:320 (1983).
- Sellwood et al., "Outer Envelope and Axial Filament Polypeptides of *T. hyodysenteriae*, Characterization and Molecular Cloning," *Proc. Inter. Pig. Vet. Soc.*, at page 119 (1990).
- Sellwood, *Proc. 12th Cong. Int. Pig Vet. Soc.*, The Hague, The Netherlands, at page 264 (1992).
- Smith et al., *Vet. Microbiol.*, 24:29-41 (1990).
- Sommer et al., "Minimal Homology Requirements for PCR Primers," *Nucleic Acids Res.*, 17(16):6749 (1989).
- Songer et al., *Am. J. Vet. Res.*, 39:913-916 (1978).
- Sotiropoulos et al., *J. Clin. Microbiol.*, 31:1746-1752 (1993).
- Sotiropoulos et al., *J. Clin. Microbiol.*, 32:1397-1401 (1994).
- Southern et al., *J. Mol. Biol.*, 98:503-517 (1975).
- Stanton et al., *Vet. Microbiol.*, 18:177-190 (1988).
- Stanton et al., "Reclassification of *Treponema hyodysenteriae* and *Treponema innocens* in a New Genus, *Serpulina* gen. nov., as *Serpula hyodysenteriae* comb. nov. and *Serpula innocens* comb. nov.," *Int. J. Syst. Bacteriol.*, 41:50-58 (1991).
- Stanton et al., "Proposal to Change the Genus Designation *Serpula* to *Serpulina* gen. nov. Containing the Species *Serpulina hyodysenteriae* comb. nov. and *Serpulina innocens* comb. nov.," *Int. J. Syst. Bacteriol.*, 42:189-190 (1992).
- "Swine Dysentery Products" in *Annual Guide to Herd Health: Pork 94* at pages 54-55 (1994).
- Taylor et al., *Proc. 12th Congr. Int. Pig Vet. Soc.*, The Hague, The Netherlands, at page 280 (1992).
- ter Huurne, "Swine Dysentery Pathogenesis and Vaccine Development," Ph.D. Dissertation, University of Ulbrecht, Nov. 1993.
- Thomas et al., *J. Med. Microbiol.*, 37:214-220 (1992).

Torp et al., *Proc. 12th Cong. Int. Pig Vet. Soc.*, The Hague, The Netherlands, at page 270 (1992).

Wannemuehler et al., *Infec. Immun.*, 56:3032–3039 (1988).

Welsh et al., “Fingerprinting Genomes Using POR with Arbitrary Primers”, *Nucleic Acids Res.*, 18:7213–7218 (1990).

Williams et al., *Nucleic Acid. Res.*, 18:6531 (1990).

Wright et al., *J. Clin. Micro.*, 27:411–416 (1989).

Matthews, et al., “Analytical Strategies for the Use of DNA Probes”, *Anal Biochem.* 169:1–25, (1989).

CGGCCAGTGC CAAGCTTTAC CAGTTGAGGG CGACTATTAT TCTGATAAAA AAATGTTAAG	60
AAGATTAGAC CCTTTTATTA ATTTTGAAT ATATGCCGCT CATCATGCAT TTAAGCAGGC	120
TGGTATAGAA CCGAAAACAG GCTTTGATCC TTTAAGAGCC GGTTGTGTTC TTGGTAGCGG	180
TATTGGCGGT ATGACTACTC TTTTATCTAA CCATCAAGTT TTACTTAATG ACGGACCTGG	240
CAGAGTATCA CCTTTCTTTG TACCTATGCA AATAATCAAT ATGACACCTG GTTTAATATC	300
TATGGAATAT GGTATGAACG GACCTAACTA CAGTACAGTT ACTGCATGTG CTTCTTCAAA	360
CCACTCTATA GGTTTAGGTT ATAAACATAT TAAAGATAAT GAAGCTGATA TT ATG	415
	Met
	1
GTA GTT GGA GGT TCT GAA GCT ACT ATA AAT CCT CTT ACT ATA GCT GGT	463
Val Val Gly Gly Ser Glu Ala Thr Ile Asn Pro Leu Thr Ile Ala Gly	
	5
	10
	15
TTC AAT AAT GCT AGA GCT TTA TCT ACT AGA AAT GAT GAT CCT GCT AAA	511
Phe Asn Asn Ala Arg Ala Leu Ser Thr Arg Asn Asp Asp Pro Ala Lys	
	20
	25
	30
GCA TCA AGA CCT TTT GAT AAA GGA AGA GAC GGA CTT GCT ATA GCC AGA	559
Ala Ser Arg Pro Phe Asp Lys Gly Arg Asp Gly Leu Ala Ile Ala Arg	
	35
	40
	45
TAT TTA ATA AAA AAT GGC TAT GAT GTA AAA ATA TAT ATC ACA GGA AAT	607
Tyr Leu Ile Lys Asn Gly Tyr Asp Val Lys Ile Tyr Ile Thr Gly Asn	
	50
	55
	60
	65
CTT GAC AGA GTT AAT AAA GAT ACC TAC TCT AAC TTT AAT ATA TTA AAA	655
Leu Asp Arg Val Asn Lys Asp Thr Tyr Ser Asn Phe Asn Ile Leu Lys	
	70
	75
	80
TCT ATG AAT ATA GAT ATT AAT TAT TTA GGA AGC GAA GAA GAT GCC ATA	703
Ser Met Asn Ile Asp Ile Asn Tyr Leu Gly Ser Glu Glu Asp Ala Ile	
	85
	90
	95
TCA GCT GCA GAA AAT ATA GAA AGA AAA TCA ATA GTA TTA GAT TCA TTA	751
Ser Ala Ala Glu Asn Ile Glu Arg Lys Ser Ile Val Leu Asp Ser Leu	
	100
	105
	110
TTT GGT ACA GGC GGA AAC AGA CCT TTA GAA GGA ATA CAA AAA GCT CTT	799
Phe Gly Thr Gly Gly Asn Arg Pro Leu Glu Gly Ile Gln Lys Ala Leu	
	115
	120
	125
ATA GAT AGT TTG AAT AAA TTA GAT GTT CTT AGA ATA GCA ATA GAT ATA	847
Ile Asp Ser Leu Asn Lys Leu Asp Val Leu Arg Ile Ala Ile Asp Ile	
	130
	135
	140
	145

FIG. 1A

CCT	TCA	GGA	TTA	GCT	TCA	AAA	ATA	AAT	GAT	AAT	GAC	AAT	GTA	TAT	ACT	895
Pro	Ser	Gly	Leu	Ala	Ser	Lys	Ile	Asn	Asp	Asn	Asp	Asn	Val	Tyr	Thr	
				150					150					160		
TGT	TTT	AAA	GCA	CAT	GAA	ACA	TAT	ACT	ATA	TGC	TTC	GCT	AAA	GAT	ATA	943
Cys	Phe	Lys	Ala	His	Glu	Thr	Tyr	Thr	Ile	Cys	Phe	Ala	Lys	Asp	Ile	
			165					170					175			
TTC	TTT	TTA	TAC	AGA	ACA	AGA	GAA	TAT	ATA	GGA	AAA	TTA	TTC	ATA	ATA	991
Phe	Phe	Leu	Tyr	Arg	Thr	Arg	Glu	Tyr	Ile	Gly	Lys	Leu	Phe	Ile	Ile	
		180					185					190				
AAA	TCA	ATA	TTC	CCA	GAT	GAA	ATA	TTA	GAT	AAT	TGG	GGA	TAT	AAA	GCT	1039
Lys	Ser	Ile	Phe	Pro	Asp	Glu	Ile	Leu	Asp	Asn	Trp	Gly	Tyr	Lys	Ala	
	195					200					205					
AAA	CTT	ATA	GAT	TAT	AAT	GAA	AAA	ATA	AAT	ATA	AAT	AGA	AAC	TCT	CTA	1087
Lys	Leu	Ile	Asp	Tyr	Asn	Glu	Lys	Ile	Asn	Ile	Asn	Arg	Asn	Ser	Leu	
210					215				220					225		
TAC	AGC	AAA	AGA	GAA	CAA	GGA	ATG	CTT	GCT	ATA	GTA	GCA	GGA	AGT	GAT	1135
Tyr	Ser	Lys	Arg	Glu	Gln	Gly	Met	Leu	Ala	Ile	Val	Ala	Gly	Ser	Asp	
				230					235					240		
AAT	TAT	ATA	GGG	GCT	GCT	GTT	CTA	GCT	GTA	AAT	GCT	GCT	TAT	AGA	TTG	1183
Asn	Tyr	Ile	Gly	Ala	Ala	Val	Leu	Ala	Val	Asn	Ala	Ala	Tyr	Arg	Leu	
			245				250						255			
GGT	GTA	GGA	TAC	ATA	AGA	TTA	TAT	GTA	CCT	AAA	GGC	ATA	ATA	AAA	AAT	1231
Gly	Val	Gly	Tyr	Ile	Arg	Leu	Tyr	Val	Pro	Lys	Gly	Ile	Ile	Lys	Asn	
		260				265						270				
ATA	AGA	GAT	GCC	ATA	ATG	CCT	TCT	ATG	CCT	GAA	ATT	GTT	ATT	ATA	GGA	1279
Ile	Arg	Asp	Ala	Ile	Met	Pro	Ser	Met	Pro	Glu	Ile	Val	Ile	Ile	Gly	
	275					280					285					
GTT	GGA	GAA	GAA	AAT	CAA	AAA	TTC	TTC	ACA	GAA	AAT	GAC	ATT	GAA	ATA	1327
Val	Gly	Glu	Glu	Asn	Gln	Lys	Phe	Phe	Thr	Glu	Asn	Asp	Ile	Glu	Ile	
290				295						300				305		
GTA	AAT	GAT	ATT	AAT	AAA	AGC	GAT	GCT	TGT	ATA	ATA	GGT	TCT	GGT	ATA	1375
Val	Asn	Asp	Ile	Asn	Lys	Ser	Asp	Ala	Cys	Ile	Ile	Gly	Ser	Gly	Ile	
				310					315					320		
GGC	AGA	GAT	TTG	TCT	ACA	GAA	ATT	TTT	GTA	AAT	ACT	ATA	TTA	AAG	CAA	1423
Gly	Arg	Asp	Leu	Ser	Thr	Glu	Ile	Phe	Val	Asn	Thr	Ile	Leu	Lys	Gln	
			325					330					335			
ATA	AAT	ATA	CCT	ACT	GTT	ATT	GAT	GCT	GAT	GCT	TTA	TAT	TTA	ATG	TTT	1471
Ile	Asn	Ile	Pro	Thr	Val	Ile	Asp	Ala	Asp	Ala	Leu	Tyr	Leu	Met	Phe	
		340					345					350				

FIG. 1B

GAA AGC ACT CTT AAT GAA CTT AAT AAT AAT TTT ATA ATC ACT CCT CAT Glu Ser Thr Leu Asn Glu Leu Asn Asn Asn Phe Ile Ile Thr Pro His 355 360 365	1519
ATA TAT GAA TTT GAA AAA CTT ACA CAG ATA AAT CAT ATA GAG GTT TTA Ile Tyr Glu Phe Glu Lys Leu Thr Gln Ile Asn His Ile Glu Val Leu 370 375 380 385	1567
GAA AAT CCT TAT CAG GCA TTA TTA ATA TAC AGA GAA AAA ACT AAT GCC Glu Asn Pro Tyr Gln Ala Leu Leu Ile Tyr Arg Glu Lys Thr Asn Ala 390 395 400	1615
TCA ATA GTA TTA AAA GAT GCT GTA AGT TTC CTA ATG CAT GAA AAT GAT Ser Ile Val Leu Lys Asp Ala Val Ser Phe Leu Met His Glu Asn Asp 405 410 415	1663
ATA TAT ATA AAT TAT AAC CCT AGA GAA TCT ATG GGG AAA GCA GGT ATG Ile Tyr Ile Asn Tyr Asn Pro Arg Glu Ser Met Gly Lys Ala Gly Met 420 425 430	1711
GGT GAT GTT TTT GCT GGA TTT ATA GGT GCT TTG CTC GCT AGA AAA CTA Gly Asp Val Phe Ala Gly Phe Ile Gly Ala Leu Leu Ala Arg Lys Leu 435 440 445	1759
AAT ATA TTA GAT GCT TCA AAA CTA GCA TTG ATA ATA CAG GCT AAA TCT Asn Ile Leu Asp Ala Ser Lys Leu Ala Leu Ile Ile Gln Ala Lys Ser 450 455 460 465	1807
TTT AAT ATA TTA TCA AAA AAA TTC GGA AAT GAT TAT ATT CAG CCT AAA Phe Asn Ile Leu Ser Lys Lys Phe Gly Asn Asp Tyr Ile Gln Pro Lys 470 475 480	1855
GAT TTG GCA AAT ATT TCA TAT AAA ATA CTA AAA GGA TAT AAA TTT GCC Asp Leu Ala Asn Ile Ser Tyr Lys Ile Leu Lys Gly Tyr Lys Phe Ala 485 490 495	1903
TAGAGAAGTT TACGACCCTA AACAAAAAGA ATTAGAATTC TACGCTAAAA GAGAGGTAAA	1963
GCCCCCTGCT CCTAAAAGAG AGGTAAGCAT ATTTGCTAGA AGATGGTTTA TGTTTTTATA	2023
CGGAACCTTC CTCACATTAG TTGTAATTGG TATGCTTTTA TATAAAAAAG GATTCTTTAA	2083
TAATATACCA TTATTTGAAG CTTTAAAGCC TAAAACAGAT GTTATAGTAA AAATTAATAA	2143
TGCTGAATTC GTTAATGATG CAGTAATTAC AACTATAGAA CTCGAAAATT CAAATTATAC	2203
TAATTCTEAA AGTATAGAAA CACTAAGAAG TTATTTTCA TTGTACAAAA ATAGAAAATT	2263
AATATTTACA GGCAATCGTT CTTTTAATAA TATAAGATTC CCAGTAGGTC AGAGAATAGG	2323
ATTCAATTT	2332

FIG. 1C

FIGURE 2




FIGURE 3




FIGURE 4 A





FIGURE 4 B



**NUCLEOTIDE SEQUENCES FOR
DETECTION OF SERPULINA
HYODYSENTERIAE**

This is a divisional of copending application Ser. No. 08/252,492 filed on Jun. 1, 1994.

**STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH**

The invention described herein was made with the assistance of funds provided by the U.S. Department of Agriculture/CSRS, Animal Health Project No. Neb. 14-048, Immunobiology of Enteric Diseases of Swine and Cattle; U.S. Department of Agriculture, Regional Research Project NC-62, Prevention of Enteric Diseases of Swine; U.S. Department of Agriculture/NC-IPM, No. Neb. 14-062, Integrated Management Practices for Control of Swine Dysentery and Salmonellosis. The Government has certain rights in the invention.

BACKGROUND OF THE INVENTION

Swine dysentery is a highly contagious disease of growing and finishing swine which has a significant economic impact on the United States swine industry. Although the causative agent has been known for nearly 20 years, no effective measures, other than medication of animals and sanitation of premises, are available to prevent the occurrence of the disease or reduce its severity once introduced into a susceptible herd. Recently published data from Iowa State University indicated a projected statewide cost for prevention and control of swine dysentery of approximately \$2.4 million per month. Since the Iowa swine population represents one-fourth of the nation's industry, the annual losses due to swine dysentery for the United States may represent as much as \$115.2 million. Owen, *Iowa State J. Res.*, 62:293-311 (1987).

Serpulina hyodysenteriae is the primary etiologic agent of swine dysentery. Harris and Lysons, *Diseases of Swine*, 7th ed., Iowa State University Press (Ames, IA), at pp. 599-616 (1992); and Stanton et al., *Int. J. Syst. Bacteriol.*, 41:50-58 (1991). Nine serotypes of *S. hyodysenteriae* have been recognized worldwide, with serotypes 1 and 2 being the most prevalent in the United States. Baum and Joens, *Infect. Immun.*, 25:792-796 (1979); Mapother and Joens, *J. Clin. Micro.*, 22:161-164 (1985); and Li et al., *J. Clin. Microbiol.*, 29:2794-2797 (1991). The diagnosis of swine dysentery is based on herd history, clinical signs, observation of characteristic intestinal lesions, and isolation of *S. hyodysenteriae* from feces or intestine using selective agar medium incubated anaerobically for 2 to 4 days. Chengappa et al., *Report of the Committee on Swine Dysentery*, American Association of Veterinary Laboratory Diagnosticians, Inc. (Columbia, Mo.) (1989). Laboratory confirmation of *S. hyodysenteriae* by culture is based upon colony morphology, pattern and intensity of hemolysis, and other growth characteristics, all of which are very similar for the non-pathogenic *Serpulina innocens*, a common inhabitant of the colon of swine. Kinyon and Joens, (1979). As a result, a definitive diagnosis of swine dysentery can be very challenging particularly when the disease occurs on premises where weakly β -hemolytic intestinal spirochetes (WBHIS) are present in the swine population.

An important aspect of swine dysentery is the occurrence of prolonged shedding of *S. hyodysenteriae* in the feces of some animals following recovery from diarrhea. Songer and Harris, *Am. J. Vet. Res.*, 39:913-916 (1978); and Fisher and

Olander, *Am. J. Vet. Res.*, 46:450-455 (1981). Asymptomatic carrier-shedder swine are important reservoirs for maintenance of *S. hyodysenteriae* on infected premises and transmission of the organism to uninfected premises. The solution to swine dysentery prevention lies in being able to quickly and accurately identify carrier-shedder swine and avoid their entry into uninfected herds. However, identification of asymptomatic carrier-shedders of *S. hyodysenteriae*, is difficult due to the detection limits of currently available laboratory isolation procedures.

Direct culture of diagnostic specimens is the only method available for laboratory identification of *S. hyodysenteriae*. However, it is well known that the sensitivity of the direct culture method depends upon the number of organisms present in the sample, which in turn depends on the stage of infection of the animal at the time of collection. Kunkle and Kinyon reported that the numbers of *S. hyodysenteriae* in porcine colonic contents at the onset of swine dysentery ranged between 2×10^6 and 2×10^{10} CFU/g when cultured using the selective BJ medium. Kunkle et al., *J. Clin. Microbiol.*, 26:2357-2360 (1988). In contrast, subclinically affected animals may shed recoverable numbers of spirochetes only sporadically and in much lower numbers than animals with clinical swine dysentery often resulting in false negative culture results. Field cases of swine dysentery also may contain drug residues that adversely affect recovery of viable *S. hyodysenteriae* by culture.

Identification of *S. hyodysenteriae* by culture is highly subjective and can lead to false results, particularly when results of cultures are interpreted by inexperienced laboratory workers. For this reason, several biochemical tests have been proposed for rapid differentiation of enteropathogenic and non-pathogenic intestinal spirochetes of swine. Achacha et al., *J. Vet. Diag. Invest.*, 3:211-214 (1991); Belanger et al., *J. Clin. Microbiol.*, 29:1727-1729 (1991); Hunter et al., *Vet. Rec.*, 104:383-384 (1979); and Smith et al., *Vet. Microbiol.*, 24:29-41 (1990). Although these biochemical characteristics are highly conserved among field isolates of *S. hyodysenteriae*, WBHIS have been shown to yield highly variable results making a conclusive identification of *S. hyodysenteriae* based on biochemical tests alone practically impossible. Achacha et al., cited supra; Belanger et al., cited supra; Burrows et al., *Vet. Rec.*, 108:187-189 (1981); Kinyon et al., *Infect. Immun.*, 15:638-646 (1977); Lymbery et al., *Vet. Microbiol.*, 22:89-99 (1990); Picard et al., *Can. J. Microbiol.*, 26:985-991 (1980); Ramanathan et al., *Vet. Microbiol.*, 37:53-64 (1993); and Torp and Thorensen, *Proc. 12th Congr. Int. Pig Vet. Soc.*, The Hague, The Netherlands, at page 270. (1992). In addition, the biochemical tests require growth of the organism for 2 to 4 days.

Other methods of differentiating *S. hyodysenteriae* from WBHIS include growth inhibition by discs soaked in anti-serum (Lemcke and Burrows, *Vet. Rec.*, 104:548-551 (1979)) and rapid slide agglutination (Burrows and Lemcke, *Vet. Rec.*, 108:187-189 (1981)). In addition to problems of non-specific clumping of spirochetes in the saline control in the slide agglutination test, these tests require large numbers of pure culture of spirochetes which can take up to 3 weeks to grow. Lysons, *Vet. Rec.*, 129:314-315 (1991). Although pre-absorption of reference polyclonal antisera with WBHIS increases the specificity of the serological tests, occasional *S. hyodysenteriae* isolates continue to be falsely classified as non-pathogenic in these tests. An alternative method using microscopic agglutination under phase contrast or dark field microscopy was recently proposed. However, some isolates of *S. hyodysenteriae* gave weaker reactions in that assay than with the slide agglutination test (Lysons, cited supra).

Mouse monoclonal antibodies capable of differentiating *S. hyodysenteriae* from porcine WBHIS have also been proposed as potential diagnostic reagents. Sellwood et al., *Proc. 12th Congr. Int. Pig Vet. Soc.*, The Hague, The Netherlands, at page 264 (1992); and Thomas and Sellwood, *J. Med. Microbiol.*, 37:214–220 (1992). However, other studies, indicate that spirochetes other than *S. hyodysenteriae* can express antigenic determinants recognized by these reagents and cause false positive results. Taylor et al., *Proc. 12th Congr. Int. Pig Vet. Soc.*, The Hague, The Netherlands, at page 280 (1992). The fact that no serological reagents are available commercially also limits the applicability of serological techniques to routine diagnosis of swine dysentery.

Certain genes encoding *S. hyodysenteriae* antigens and capable of eliciting protection against infection in mice have been cloned and expressed in *Escherichia coli* using a phage expression system. Boyden et al., *Infect. Immun.*, 57:3808–3815 (1989). However, none of these reagents have been examined for potential application as diagnostic tools. One of the most recent diagnostic applications of recombinant DNA technology to swine dysentery control used oligodeoxynucleotide probes to 16S rRNA of *S. hyodysenteriae*. Jensen et al., *J. Clin. Microbiol.*, 28:2717–2721 (1990). However, the sensitivity of this probe method for detection of spirochetes in feces was equivalent to routine bacteriological culture (10^5 organisms/g of feces), and further studies question the specificity of the 16S rRNA probe to *S. hyodysenteriae* (Torp and Thoresen, cited supra). Dot blot hybridization with whole-chromosomal probes and DNA probes for identification of *S. hyodysenteriae* have been reported. Combs and Hampson, *Res. Vet. Sci.*, 50:286–289 (1991); Sotiropoulos et al., *J. Clin. Microbiol.*, 31:1746–1752 (1993); and Sotiropoulos et al., *J. Clin. Microbiol.*, 32:1397–1401 (1994). Although the sensitivity of the whole-chromosomal probes was not reported, colony dot blot hybridization with DNA probes was shown to be only slightly better than culture (10^4 organisms/g of feces). These tests are labor intensive, require specialized equipment, and have turn-around times that are incompatible with routine laboratory diagnosis.

A solution to prevention of disease caused by *S. hyodysenteriae* lies in being able to quickly and accurately identify carrier-shedder animals and avoid their entry into uninfected herds. Therefore, there is a need to develop a method and reagents for detecting *S. hyodysenteriae* in low numbers specifically, rapidly and directly from diagnostic and environmental samples. There is also a need to develop a sensitive and specific method for rapid detection of *S. hyodysenteriae* in a biological sample to diagnose and monitor infection in acutely- or subclinically-infected animals before, during and after treatment and in their environment. There is also a need to develop a method for rapid detection of *S. hyodysenteriae* for monitoring disinfection of the environment in contact with infected animals.

SUMMARY OF THE INVENTION

These and other objects are achieved by the present invention which is directed to a method for detecting the presence of at least one serotype of *Serpulina hyodysenteriae* in a biological sample, an oligonucleotide primer, and a *S. hyodysenteriae*-specific oligonucleotide probe useful in that method, and an article of manufacture (i.e., kit) containing the primers and/or probe. Also provided are an about 2.3-kb DNA fragment derived from chromosomal DNA of *S. hyodysenteriae* B204 serotype 2 that encodes for an about 56 kDa polypeptide, a recombinant expression vector containing the DNA fragment, the about 56 kDa polypeptide and a

monoclonal antibody reactive with the peptide. The invention also provides for a method of immunizing animals and vaccine preparations for protecting animals against disease caused by *S. hyodysenteriae*. The methods and compositions of the invention are useful to identify at least one serotype of *S. hyodysenteriae*, to diagnose *S. hyodysenteriae* infection, to detect carrier-shedder animals, to monitor efficacy of treatment for disease caused by *S. hyodysenteriae*, to monitor disinfection of fomites, and to protect animals from infection with *S. hyodysenteriae*.

According to the invention, a biological sample of an animal such as feces, intestinal contents, mucosal scrapings, rectal swabs, and environmental samples among others, is tested for the presence of at least one serotype of *S. hyodysenteriae* by measuring the presence or absence of DNA amplification products from a primer that hybridizes to a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* B204 serotype 2. The 2.3 kb HindIII restriction fragment was obtained from a partial digest with HindIII and preferably has a nucleotide sequence as shown in FIG. 1 (SEQ ID NO:1). The DNA amplification products specific for *S. hyodysenteriae* can be detected by (a) extracting DNA from a biological sample; (b) amplifying a target sequence of the extracted DNA to provide DNA amplification products carrying a selected target DNA sequence; and (c) detecting the presence of *S. hyodysenteriae* by detecting the presence of the DNA amplification products. Preferably, the amplification of the DNA sequence is by polymerase chain reaction (PCR) by amplifying the gene sequence with DNA polymerase and at least one oligonucleotide primer. The primers preferably have a sequence of positive-sense 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3), or negative-sense 5'-TCCTATTCTCTGACCTACTG (SEQ ID NO:4), or complements thereof or mixtures thereof.

The amplification products can be detected, for example, by dot blot or Southern blot analysis including by reacting the DNA amplification products with a labeled oligonucleotide probe that can hybridize to the about 2.3 kb HindIII DNA fragment of *S. hyodysenteriae* B204 serotype 2 shown in FIG. 1. Prior to detection, the DNA amplification products can optionally be separated by electrophoresis. Alternatively, the PCR products can be detected by immobilization to a bead or a multiwell plate by a probe or primer labeled with biotin, followed by hybridization with a detectably labeled probe. The oligonucleotide probe preferably hybridizes to all or a portion of a 2.3 kb HindIII DNA fragment of *S. hyodysenteriae* B204 serotype 2 having the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1). The detection of amplified products preferably allows for the detection of all serotypes of *S. hyodysenteriae* at low levels of bacteria in the sample (i.e., about 1 to 10 microorganisms/0.1 gm sample).

The invention provides an isolated HindIII DNA fragment of about 2.3-kb that is derived from a partial digest of chromosomal DNA of *S. hyodysenteriae* B204 serotype 2 with HindIII, which encodes for an about 56 kDa polypeptide. The DNA fragment preferably has the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1).

The DNA restriction fragment may be incorporated into an expression vector. The expression vectors include the restriction fragment operably linked to transcriptional and translational control regions in the vector. The expression vectors are useful to form transformed cells. The transformed cells can be used to screen monoclonal antibodies specific for *S. hyodysenteriae* to produce a 56 kDa polypeptide of *S. hyodysenteriae*, to prepare mutant sequences of the 2.3 kb HindIII restriction fragments, and as an intermediate

to prepare the 2.3 kb HindIII restriction fragment for DNA sequencing. A pUC18 plasmid carrying the 2.3 kb HindIII restriction fragment from *S. hyodysenteriae* B204 and designated pRED3C6, deposited with the American Type Culture Collection, Rockville, Md., has been given Accession No. 75826.

Also provided are oligonucleotide primers that hybridize to all or a portion of a DNA sequence of the 2.3 kb HindIII fragment of *S. hyodysenteriae* B204 serotype 2 (SEQ ID NO:1). The DNA sequence of an oligonucleotide primer is preferably positive-sense 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3) and negative-sense 5'-TCCTATTCTCTGACCTACTG (SEQ ID NO:4), or complements thereof or mixtures thereof. The primers can be complementary to the 3' or 5' end of the target sequences. The primers can also have mismatches with the target sequence or be degenerate.

The target DNA sequence is also useful in providing an oligonucleotide probe for detecting a target sequence of at least one serotype of *S. hyodysenteriae*. The probe is preferably hybridizable with a DNA sequence located on the 2.3 kb HindIII fragment of the chromosomal DNA of *S. hyodysenteriae*, preferably a 1.55 kb DNA sequence within the 2.3 kb fragment. The DNA sequence of the oligonucleotide probe is preferably hybridizable to all or a portion of the 2.3 kb HindIII fragment of *S. hyodysenteriae* B204, as shown in FIG. 1, and can distinguish *S. hyodysenteriae* from other microorganisms and cells in a biological sample.

The about 56 kDa polypeptide encoded by the about 2.3 kb DNA fragment is useful in eliciting antibodies that are specifically reactive with *S. hyodysenteriae*. The 56 kDa polypeptide and nucleotide sequence encoding the polypeptide may also be useful in a vaccine preparation to provide a protective effect for an infection caused by *S. hyodysenteriae*. The invention further provides an in vitro assay for detecting *S. hyodysenteriae*-specific antibodies in a sample. In that method, a sample to be tested is contacted with a composition containing the about 56 kDa polypeptide, which is preferably labelled, to form a conjugate which is then detected.

An article of manufacture (i.e., kit) is also provided for use in the detection of at least one serotype of *S. hyodysenteriae* in a biological sample. The kit is composed of one or more reagents contained within a packaging material, the reagents useful for detection of the spirochete including at least one oligonucleotide primer that hybridizes to a DNA sequence of the 2.3 kb fragment of *S. hyodysenteriae* B204 as shown in FIG. 1, and/or an oligonucleotide probe which is hybridizable to all or a portion of a DNA gene sequence included in the 2.3 kb gene fragment, preferably an about 1.55 kb gene fragment. The kit may optionally include instruction means with information regarding the use of the reagent and/or other component of the kit such as how to conduct an assay, and the like. The instruction means may be a label or tag attached to the packaging, a printed package insert, and the like. Two or more reagents and/or other components may be combined together to form the kit, preferably packaged together within containing means such as a box or plastic bag, and the like.

The kits and methods of the invention can be used to identify at least one serotype of *S. hyodysenteriae* to diagnose *S. hyodysenteriae* infection, detect carrier-shedder animals, monitor the efficacy of treatment for diseases caused by *S. hyodysenteriae*, or monitor disinfection of the environment. A method for monitoring the efficacy of treatment is useful to ensure that subclinically infected or carrier-

shedder animals are not formed. The method involves monitoring the presence of *S. hyodysenteriae* throughout the course of treatment of the animal. The presence of *S. hyodysenteriae* in a biological sample from the treated animal is detected by detecting DNA amplification products of a target sequence of *S. hyodysenteriae*, wherein the target sequence is a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae*.

The invention also provides for vaccines and methods of immunizing an animal to produce a protective immune response against *S. hyodysenteriae* infection. The vaccines include an amount of a 56 kDa polypeptide encoded on a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* B204 effective to induce a protective response to *S. hyodysenteriae* infection. The vaccine preparations are in admixture with a physiologically acceptable carrier and can be administered to animals using standard methods. The vaccines and methods of the invention are useful to prevent the adverse effects of disease caused by *S. hyodysenteriae*.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the nucleotide sequence of the recombinant 2.3-kb HindIII DNA fragment from *Serpulina hyodysenteriae* (SEQ ID NO:1) which indicates a single open reading frame encoding a protein with a predicted molecular weight of approximately 56 kDa (SEQ ID NO:2).

FIG. 2 is an image of a Southern blot hybridization using [α - 32 P]dCTP labelled 2.3 kb fragment of clone pRED3C6 to purified chromosomal DNA. Purified chromosomal DNA was digested with HindIII, electrophoresed on a 0.8% agarose gel, and transferred to a nylon membrane. Lanes 1 to 8, *Serpulina hyodysenteriae* isolate B78, B234, B204, B169, A1, B8044, B6933, and AcK 300/8, respectively; lanes 9 to 12, WBHIS B256, B359, B1555a, and 4/71, respectively; lane 13, *Treponema succinifaciens*.

FIG. 3 is an image of an ethidium bromide stained agarose gel and Southern blot hybridization of PCR using purified chromosomal DNA obtained from reference isolates of *S. hyodysenteriae* serotypes 1 through 9. The amplified products were electrophoresed on a 0.8% agarose gel and stained with ethidium bromide. The insert at the bottom of the photograph shows hybridization of transferred DNA from the same gel with an internal *S. hyodysenteriae*-specific oligonucleotide probe 5'-end labelled with [γ - 32 P]ATP. Lanes: 1, molecular weight standard (1-kb DNA ladder; GIBCO-BRL); 2, B78; 3, B204; 4, B169; 5, A1; 6, B8044; 7, B6933; 8, AcK 300/8; 9, FM-88-90; 10, FMV 89-3323.

FIG. 4 is an image of an ethidium bromide stained agarose gel (panel A) and Southern blot hybridization (panel B) of PCR using purified DNA obtained from porcine feces inoculated with serial ten fold dilutions of *Serpulina hyodysenteriae* isolate B204 cells. FIG. 4A shows the amplified products electrophoresed on a 0.8% agarose gel and stained with ethidium bromide. FIG. 4B shows hybridization of transferred DNA from the same gel with an internal *S. hyodysenteriae*-specific oligonucleotide probe 5'-end labelled with [γ - 32 P]ATP. Lanes: 1, molecular weight standard (1-kb DNA ladder; GIBCO-BRL); lanes 2 to 9, feces containing 10^5 , 10^4 , 10^3 , 10^2 , 10^1 , 10^0 , 10^{-1} , 10^{-2} *S. hyodysenteriae* cells/0.1 g, respectively.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides for methods, kits and compositions useful for diagnosis and monitoring of infection of animals with at least one serotype of *S. hyodysenteriae*. The

compositions include probes and primers that can hybridize to a target sequence of at least one serotype of *S. hyodysenteriae*. The probes and primers can preferably hybridize to all serotypes of *S. hyodysenteriae* and not other closely related microorganisms. The target sequence is preferably about a 2.3 kb fragment of *S. hyodysenteriae* chromosomal DNA having the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1). The primers and probes can be used in methods and kits for detecting *S. hyodysenteriae* in a biological sample, preferably by detecting amplification products using primers that hybridize to the target sequence.

The invention also provides a 56 kDa polypeptide encoded by a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* B204, as shown in FIG. 1. The polypeptide and nucleotide sequences are useful to elicit antibodies and in vaccine formulations. The vaccine can be administered to animals to induce a protective immune response against infection with *S. hyodysenteriae*.

1. Primers and Probes that Hybridize to a Target Sequence of *S. hyodysenteriae*

Primers and probes of the invention are useful for identification of at least one serotype of *S. hyodysenteriae* in a biological sample. The primers and probes are preferably those that hybridize to a target sequence of about a 2.3 kb fragment of *S. hyodysenteriae* B204 having the nucleotide sequence of FIG. 1 (SEQ ID NO:1). Hybridization of the primers and/or probes to the target sequence preferably can provide for identification of all serotypes of *S. hyodysenteriae* and distinguish *S. hyodysenteriae* from other cells and closely related microorganisms. Primers can be useful in a diagnostic assay to identify at least one serotype of *S. hyodysenteriae* and to distinguish *S. hyodysenteriae* from at least one other microorganism, and to form probes or to form deletion mutants of the target sequence. Probes are useful to identify at least one serotype of *S. hyodysenteriae* and/or to distinguish *S. hyodysenteriae* from at least one other microorganism, to identify amplification products of a target sequence, and to identify other target sequences.

Primers and probes in accord with the invention are selected so that they can specifically identify at least one serotype of *S. hyodysenteriae*, preferably all serotypes. Those primers or probes that can specifically identify *S. hyodysenteriae* are those that hybridize to a target sequence preferably found in all serotypes and not found in other closely related microorganisms. One such target sequence identified in all serotypes of *S. hyodysenteriae* is about a 2.3 kb fragment of *S. hyodysenteriae* B204 shown in FIG. 1. The 2.3 kb fragment was obtained by partial digestion of chromosomal DNA with HindIII.

The target sequence specific for *S. hyodysenteriae* can be identified from a DNA or cDNA library of *S. hyodysenteriae* chromosomal DNA or mRNA, respectively. A DNA or cDNA library can be generated by standard methods using a restriction enzyme such as HindIII. Suitable host cells such as *E. coli* DH5 α and the like are transformed with the library. Transformed cells can be screened by a variety of standard methods including by reactivity with antibodies that react with antigens of *S. hyodysenteriae*.

Once a clone that reacts with the antibodies that react with antigens of *S. hyodysenteriae* is identified, it can be amplified and sequenced. An example of a target sequence is a 2.3 kb fragment having a nucleotide sequence shown in FIG. 1 (SEQ ID NO 1). This sequence encodes a 56 kDa polypeptide having a predicted amino acid sequence as shown in FIG. 1 (SEQ ID NO:2). Transformed cells including this target sequence were immunoreactive with monoclonal antibody 10G6/G10 (available from Dr. Duhamel at University

of Nebraska, Lincoln, Nebr.) raised against cell-free supernatant antigens of *S. hyodysenteriae* B204.

A target sequence can be isolated and labeled with a detectable label such as a radioactive nucleotide. The target sequence can also serve as a probe and can be screened for hybridization to all serotypes of *S. hyodysenteriae* and for lack of hybridization to other microorganisms such as *S. innocens*, WBHIS strains, *Treponema* spp., *E. coli*, *Salmonella* spp., *Campylobacter* spp., *Bacterioidis vulgatus*, *Spirocheta aurantia*, *Borrelia burgdorferi*, and Leptospiroaceae. Hybridization conditions are preferably low stringency conditions. The target sequence, as well as probes and primers derived from the target sequence, can be used to confirm the identity of a pure culture of *S. hyodysenteriae* isolated by conventional microbiological methods or isolated by immunoaffinity methods, to distinguish at least one serotype of *S. hyodysenteriae* from other cells in a mixed biological sample including other microorganisms and eukaryotic cells. Preferably the probe derived from a target sequence can hybridize to all serotypes of *S. hyodysenteriae* and not to closely related *S. innocens*, WBHIS strains, *Treponema* spp. Once a target sequence is identified, screened for specificity for at least one serotype to *S. hyodysenteriae* and sequenced, it can be used to design primers and/or probes.

Once the sequence of a target sequence from one serotype of *S. hyodysenteriae* is known, primers and probes that hybridize to the known target sequence can be used to identify other closely related target sequence from other serotypes that will hybridize to the same primers and/or probes. For example, other target sequences from other serotypes of *S. hyodysenteriae* can have some DNA sequence differences from the 2.3 kb HindIII restriction fragment from B204 serotype 2, shown in FIG. 1, and still be able to hybridize to primers and probes that hybridize to the 2.3 kb sequence as shown in FIG. 1 under low, medium or high stringency conditions. These other target sequences, once identified, can be sequenced and used to provide a template for design of primers and/or probes. Once selected, these target sequences are further preferably screened for the ability to hybridize to sequences in all serotypes of *S. hyodysenteriae* and not to related microorganisms such as *S. innocens*, *Treponema* spp., and WBHIS spp., and the like.


At least one primer is designed to be useful to amplify the target DNA sequence preferably using standard or hot start polymerase chain reaction. A primer can preferably be about 16 to 30 nucleotides long and more preferably about 20 to 21 nucleotides long. Primers can hybridize to sequences flanking the desired target sequence which is preferably all or a portion of a 2.3 kb fragment of *S. hyodysenteriae* B204 having the nucleotide sequence shown in FIG. 1. Primers can hybridize to sequences at the 5' and/or 3' ends of the target sequence. Primers can hybridize to the DNA strand with the coding sequence of a target sequence and are designated sense primers. Primers can hybridize to the DNA strand that is the complement of the coding sequence of a target sequence and are designated anti-sense primers. Primers that hybridize to each strand of DNA in the same location or to one another are known as complements of one another. Primers can be designed to hybridize to a mRNA sequence complementary to a target DNA sequence and are useful in reverse transcriptase PCR.

Hybridization conditions utilized are those preferred for polymerase chain reaction modified as required for the degree of sequence complementarity of the primers to the target sequence. Hybridization conditions for a primer of about 16 to 30 nucleotides long having no mismatches with the target sequence are those described by Elders et al., *J.*

Clin. Micro., 32:1497 (1994). Briefly, a PCR mixture including 4 mM MgCl₂, 0.2 mM dNTPs and a DNA polymerase were mixed with the DNA extracted from the biological sample. Initial denaturing is at 95° C. for 60 seconds followed by 30 cycles (65° C. for 60 seconds and 72° C. for 120 seconds). The conditions selected are those described for GenAmp 480 (Perkin-Elmer, Norwalk, Conn.).

Hybridization conditions for a primer having about 16 to 30 nucleotides and about up to 30% mismatch are modified as described in Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y. (1989). The melting temperature (T_M) for hybridization is decreased about 1° to 1.5° C. for each 1% of mismatch. Primers less than 20 nucleotides preferably only have about one to three mismatches with the target sequence located at either the 5' or 3' end of the primer. PCR methods using mismatched primers or degenerate primers have been described.

Primers can be designed as overlapping sequences or a nested set as long as all or a portion of the target sequence can be amplified. Primers can include at least about 16 nucleotides starting from the flanking sequence immediately adjacent to the 5' end of the target sequence and overlapping primers can be designed to move from the 5' end to the 3' end of the target sequence as shown below:



Likewise, primers can be designed to include about 16 nucleotides starting from the flanking sequence immediately adjacent to the 3' end and then overlapping the sequence until the 5' end of the target sequence. These primers can vary in size from about 16 to 100 nucleotides. A primer that hybridizes to a flanking region preferably hybridizes to about 16 to 30 nucleotides immediately upstream or downstream from the target sequence. These overlapping primers provide for amplification of all or a portion of the target sequence preferably at least about 16 nucleotides long and more preferably about 20 to 2,300 nucleotides long.

Primers can preferably hybridize under standard conditions for polymerase chain reaction to a portion of the target nucleotide sequence or to the flanking sequences immediately adjacent to the target sequence and provide for amplification of all or a portion of a target sequence including preferably a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204. Primers preferably hybridize to the target sequence with no mismatches. However, primers can have additional nucleotide sequence at the 5' or 3' ends, as for example, to provide restriction enzyme recognition sequences. Restriction enzyme recognition sequences are known to those with skill in the art and can provide an amplified product with restriction endonuclease recognition sequence at one or both ends of the amplified products. Other sequences can be added at one or both ends of the primers to act as markers of the amplified products as, for example, a universal primer sequence.

Primers can also be designed that are not perfectly complementary and still can hybridize to a portion of the target sequence or flanking sequence and thereby provide for amplification of all or a portion of a target sequence. Primers of about 20 nucleotides or less preferably have about one to three mismatches located at the 5' and/or 3' ends. Primers of

about 20 to 30 nucleotides have up to 30% mismatches and can still hybridize to a target sequence. Hybridization conditions for primers with mismatch can be determined by the method described in Maniatis at pages 11.55 to 11.57 which is hereby incorporated by reference or by reference to known methods. The ability of the primer to hybridize to the 2.3 kb HindIII sequence of FIG. 1 under varying conditions can be determined using this method. Because a target sequence is known, the effect of mismatches can be determined by methods known to those of skill in the art. Hybridization conditions for polymerase chain reaction using degenerate or mismatched primers are known to those of skill in the art.

In a preferred version, primer pairs are designed to flank a desired portion of the target sequence. A primer pair has a first primer that hybridizes to a sense strand at the 5' end of the target sequence and a second primer that hybridizes to the anti-sense strand at the 3' end of the target sequence. Examples of the preferred primers include sense 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3); anti-sense 5'-TCCTATTCTCTGACCTACTG (SEQ ID NO:4). The preferred primers can hybridize to and amplify all or a portion of a 2.3 kb HindIII restriction fragment sequence of *S. hyodysenteriae* B204 serotype 2, as shown in FIG. 1.

Once the primers are designed as described above, the primers can be prepared by automated DNA synthesis in accord with standard methods.

Once the target sequence has been identified, the target sequence can serve as a probe or can be used to design other probes. A probe can hybridize to all or a portion of a target sequence, preferably under all stringency conditions. The preferred probes hybridize to all or a portion of a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 as shown in FIG. 1. The probes are useful to detect amplified products of the target sequence and to detect and identify at least one serotype of *S. hyodysenteriae*.

A probe can have about 16 to about 2300 nucleotides, more preferably about 25 to 100 nucleotides. A probe can hybridize to all or a portion of a target sequence. If a probe is designed to detect amplified products of the target sequence, it is preferably designed not to hybridize to the sequence of primers used for amplification. A probe designed to detect amplification products can preferably hybridize to sequences between the primer sequences. A probe can hybridize to the DNA strand with coding sequence and is designated a sense probe. A probe can hybridize to the DNA strand that is complementary to the strand with the coding sequence and is designated an anti-sense probe. A probe can be designed to hybridize to mRNA complementary to the target sequence.

Hybridization conditions for forming hybrids for a probe of about 16 to 100 nucleotides or more without any mismatches are preferably those described in Flores et al., *J. of Vir.*, 64:4021 (1990). Briefly, immobilized target sequences are incubated at 53° C. for 16 to 32 hours in 4×SSC (1×SSC=0.15M NaCl plus 0.015M sodium citrate), 0.02% sodium dodecyl sulfate, 50% formamide, 50 mM potassium phosphate buffer, 10% Dextran sulfate, 0.02% Ficoll (Pharmacia, Inc., Piscataway, N.J.), 0.02% bovine serum albumin, 0.02% polyvinylpyrrolidone, 20 μg sheared salmon DNA per ml, and the probe. Prehybridizations are carried out in the same buffer minus dextran sulfate. After incubation, the hybrids are washed four times for 5 minutes at room temperature with 0.1% sodium dodecyl sulfate in 2.5×SSC, and then twice for 15 minutes at 53° C. with 0.05% sodium dodecyl sulfate in 1.25×SSC. These conditions can be modified, if necessary, depending on the method

of detection employed, such as dot blot hybridization, Southern blot hybridization, or multiwell solution hybridization. Modification of hybridization conditions depending on the method employed are known.

A probe of the invention can be perfectly complementary to the target sequence or can have some mismatches with the target sequence. A probe of about 16 to 20 nucleotides preferably has about 1 to 3 mismatches localized near the 5' or 3' ends of the probe (i.e., within 5 base pairs of either end). Probes of about 20 to 2300 nucleotides can have up to about 30% mismatches and still hybridize to the target sequence. Mismatched probes can still hybridize to the target sequence if conditions of hybridization are modified to account for the mismatch, as, for example, by decreasing melting temperature by about 1.0 to 1.5° C. for every 1% of mismatch. Because a target DNA sequence has been cloned and identified, the effect of mismatches in the probe on T_M (melting temperature) can be determined using a standard method such as described by Maniatis at pages 11.47 to 11.57, which is hereby incorporated by reference.

Probes can also be detectably labeled using a variety of standard methods. The probes can be detectably labeled by incorporating one or more labeled nucleotides into the probe. Nucleotides can be labeled with biotin, with a radiolabel, or with a fluorescent moiety or luminescent moiety and the like.

A preferred probe of the invention can hybridize to all or a portion of a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* B204 (SEQ ID NO:1) or to amplification products of a 2.3 kb HindIII restriction fragment from serotype B204 and other serotypes. A preferred probe has a sequence as follows:

sense-5'-TAGGGGCTGCTGTTCTAGCTGTAAATGC
(SEQ ID NO:5)

and can hybridize to about a 1.5 kb amplification product of *S. hyodysenteriae* B204 and other serotypes and to the 2.3 kb HindIII fragment under low stringency conditions.

Once designed, a probe can be prepared by automated DNA synthesis or by polymerase chain reaction using appropriate primers. The probe can also be prepared by digesting the 2.3 kb insert in pRED3C6 with any one or more of the restriction enzymes shown in Table I.

A probe according to the invention can be used in a method for identifying at least one serotype of *S. hyodysenteriae* or distinguishing *S. hyodysenteriae* from at least one other microorganism. The probes are used in methods such as restriction enzyme, PCR amplification, slot blot, dot blot, and Southern blot analysis of DNA taken from biological samples suspected of containing at least one serotype of *S. hyodysenteriae*. The steps of the method include isolating DNA from a biological sample suspected of containing at least one serotype of *S. hyodysenteriae*, digesting the extracted DNA with at least one restriction enzyme, and detecting any sequence specific to *S. hyodysenteriae* by hybridization to a probe specific for at least one serotype of *S. hyodysenteriae* chromosomal DNA.

The extraction of DNA from a biological sample can be accomplished by standard methods. Elder et al., cited supra. Biological samples can include pure cultures of bacteria isolated by standard microbiological methods or a biological sample suspected of having at least one serotype of *S. hyodysenteriae* as well as other microorganisms. Examples of biological samples include feces, intestinal contents, intestinal mucosal scrapings, fecal swabs and the like. Environmental samples are also biological samples such as manure, manure-contaminated soil, fomites, pits, lagoon water, or effluent from a premises suspected of containing *S. hyodysenteriae*.

The extract is then digested with at least one restriction enzyme. The choice of restriction enzyme is based upon the sequence of the DNA target sequence specific for *S. hyodysenteriae* and the recognition sequence of the restriction enzyme. Recognition sequences for restriction enzymes are known to those of skill in the art. If all or a portion of the DNA sequence of the target sequence for *S. hyodysenteriae* is known, then restriction enzymes can be selected based on that sequence. For example, the sequence of a 2.3 kb HindIII target sequence, shown in FIG. 1, indicates that there is one site where HindIII can cut the sequence. Other restriction enzymes can be selected that can cut the sequence at one or more locations, preferably at about 1 to 3 locations, as shown in Table I.

TABLE I

AccI	AcII	AluI	AlwI	AlwNI	ApoI
AvaII	BbvI	BccI	BcgI	BfaI	BseRI
BslI	BsmI	BsmAI	BsmBI	BsoFI	BspMI
BsrI	BsrBI	BsrFI	BsrGI	Cac8I	CjeI
CjePI	CviJI	CviRI	DdeI	DpnI	DraI
EaeI	EarI	EcII	Eco57I	EcoNI	EcoRI
EcoRII	GdiII	HaeIII	HindIII	HinfI	HphI
MaeIII	MboII	MmeI	MnlI	MseI	MslI
MspI	MspAII	MwoI	NlaIII	NsiI	NspI
PstI	PvuII	RsaI	Sau96I	Sau3AI	SexAI
SfaNI	SfcI	SspI	TaqI	TaqII	TfiI
TseI	Tsp5091	TthIII		VspI	

The digested DNA is then contacted with a probe that can hybridize to a target sequence specific for at least one serotype of *S. hyodysenteriae* under conditions of hybridization for a probe of that size and/or sequence complementarity as described previously. The hybridization conditions can be modified as necessary depending on the method of detection of hybrid formation employed including slot blot, dot blot, and/or Southern blot hybridization. The probes are preferably labeled for ease of detection of hybrid formation.

In a preferred version, DNA is extracted from a biological sample such as from an animal suspected of being infected with *S. hyodysenteriae* or from an environmental sample suspected of containing *S. hyodysenteriae* and the extracted DNA is digested with HindIII. The digested DNA extract is optionally separated and contacted with a probe that can hybridize to a target sequence specific for *S. hyodysenteriae* such as a 2.3 kb HindIII restriction fragment of FIG. 1. The preferred probe has a sequence as follows:

sense-5'-TAGGGGCTGCTGTTCTAGCTGTAAATGC
(SEQ ID NO:5)

and is detectably labeled. The presence of *S. hyodysenteriae* is detected by the detection of hybridization of the probe to digested fragments from the biological sample. Methods of detection of hybrids can be utilized depending on the labeled moiety that is attached to the probe, and are standard methods. This method can also be used to distinguish *S. hyodysenteriae* from at least one other microorganism.

2. Recombinant Expression Vectors and Transformed Cells

A target nucleotide sequence that allows for identification of at least one serotype of *S. hyodysenteriae* can be cloned into an expression vector and introduced into suitable host cells to form transformed cells. The transformed cells carrying an expression vector are useful for amplification of all or a portion of a target sequence to provide a probe, to provide any gene products encoded by the target sequence, and/or as vaccine formulations.

A target sequence, such as all or a portion of a 2.3 kb HindIII partial digest fragment from *S. hyodysenteriae* B204 (SEQ ID NO:1), can be cloned into a suitable expression

vector such as pUC18, pKC30, pBR322, pKK177-3, pET-3, and the like by standard methods. Commercially available expression vectors provide for cloning for a target sequence into a site of the vector such that the target sequence is operably linked to transcriptional and translational control regions. It is preferred, but not required, that a target sequence is operably linked to an inducible promoter such as the λ PL promoter, the lac promoter, the tac promoter, or the T₇ promoter, and the like.

The expression vectors can then be introduced into suitable host cells using methods such as calcium phosphate precipitation, liposome mediated transformation, protoplast transformation, electroporation, and the like. Suitable host cells include *E. coli* strains such as *E. coli* DH5 α , and avirulent isogenic Salmonella spp. such as *S. typhimurium* deletion mutants lacking adenylate cyclase and cAMP receptor protein, Salmonella mutants in aro genes, and other Salmonella vaccine strains as described in *Bio/Tech*, 6:693 (1988), and the like.

Transformed cells can be screened by a variety of methods including colony hybridization or reactivity with antibodies specific for *S. hyodysenteriae* B204. A transformed cell is an *E. coli* DH5 α cell carrying a pUC18 plasmid with a 2.3 kb HindIII restriction fragment insert from *S. hyodysenteriae* B204. A pUC18 plasmid carrying a 2.3 kb HindIII insert from *S. hyodysenteriae* B204 designated pRED3C6, deposited with the American Type Culture Collection in Rockville, Md. has Accession No. 75826.

3. Method for Detection of *S. hyodysenteriae* in a Biological Sample

According to the invention, a biological sample may be analyzed for the presence of *S. hyodysenteriae* by detecting the presence of DNA amplification products from primers that hybridize to a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 serotype 2. Compared to other detection methods presently known and used, the invention advantageously provides a method that is highly sensitive in detecting at least one serotype of *S. hyodysenteriae* in a biological sample when it is present in very low concentration, for example, about 1 to 10 organisms per 0.1 gm sample. It is preferred that an about 1.55 kb DNA sequence that lies between two regions of the about 2.3 kb HindIII sequence is amplified. That sequence is unique to all serotypes of *S. hyodysenteriae*, and provides for the specific detection of the spirochete from other closely-related microorganisms including other members of the order Spirochaetales. Thus, this method is also useful to distinguish *S. hyodysenteriae* from at least one other microorganism.

In brief, the DNA amplification products can be detected by (a) extracting DNA from a biological sample; (b) amplifying a target sequence of the extracted DNA to provide DNA amplification products carrying a selected target DNA sequence; and (c) detecting the presence of *S. hyodysenteriae* by detecting the presence of the DNA amplification products.

The biological sample may be, for example, feces, mucosal secretion, mucosal scrapings, mucosal cells, rectal swabs, intestinal wall, intestinal contents, local lymph nodes, and the like. The biological sample may be derived, for example, from an animal infected with *S. hyodysenteriae*, an animal suspected of being a carrier of *S. hyodysenteriae*, an animal being treated for an infection caused by *S. hyodysenteriae*. The biological sample can also be an environmental sample such as manure, manure-contaminated soil, fomites, pits, lagoon water, or effluent from a premises suspected of containing *S. hyodysenteriae*. Animals susceptible to infection with *S. hyodysenteriae*

include swine, ratics (such as rheas), rodents such as rats and mice, dogs, birds, poultry, and other wildlife.

The detection method can also be optionally combined with methods for isolation of microorganisms to provide for confirmation of infection with *S. hyodysenteriae* and/or to increase the sensitivity of the assay. For example, *S. hyodysenteriae* present in a biological sample could be separated from other biological material using an antibody attached to a solid support such as a monoclonal antibody attached to immunomagnetic particle, as described by Islam et al., *J. Clin. Micro.*, 30:2801 (1992). The isolated *S. hyodysenteriae* can then be detected using the polymerase chain reaction as described herein.

In a preferred method, the amplification of the DNA sequence is by polymerase chain reaction (PCR), as described in U.S. Pat. No. 4,683,202 to Mullis; Mullis et al., *Cold Spring Harbor Symp. Quanti. Biol.* 51:263 (1986); Mullis and Faloona, *Methods Enzymol.* 155:335 (1987); Saiki et al., *Science* 239:487 (1988b); and Chien et al., *J. Bacteriol.* 127:1550 (1976). In brief, the DNA sequence is amplified by reaction with at least one oligonucleotide primer or pair of oligonucleotide primers that hybridize to the target sequence or a flanking sequence of the target sequence and a DNA polymerase to extend the primer(s) to amplify the target sequence. The amplification cycle is repeated to increase the concentration of the target DNA sequence.

The biological sample is first treated to extract a nucleic acid sequence specific to *S. hyodysenteriae*, which may be either mRNA or DNA. The nucleic acid fragments may be extracted from the biological sample by standard methods as described in Elders et al., cited supra. Either purified chromosomal DNA or total DNA or total mRNA is extracted from the biological sample for amplification and detection.

In addition, the polymerase chain reaction may be used to amplify cDNA that has been synthesized in vitro by reverse transcriptase of an mRNA template, according to standard methods. The mRNA that is extracted from the biological sample for synthesis of cDNA sequence is suspected of including a messenger RNA that is complementary to a target sequence such as a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204. Primers useful to generate cDNA can be designed as described previously.

Other methods of polymerase chain reaction using various combination of primers including a single primer to about 3 primers are known to those of skill in the art and are described in Maniatis, cited supra. Those methods include asymmetric PCR, PCR using mismatched or degenerate primers, reverse transcriptase PCR, arbitrarily primed PCR (Welsh et al., *Nucleic Acids Res.*, 18:7213 (1990)), or RAPD PCR, IMS-PCR (as described by Islam et al., *J. Clin. Micro.*, 30:2801 (1992)), multiwell PCR (ELOSAs) (as described by Luneberg et al., *J. Clin. Micro.* 31:1088 (1993) and Katz et al., *Am. J. Vet. Res.*, 54:2021 (1993)). The methods also include amplification using a single primer as described by Judd et al., *Appl. Env. Microbiol.*, 59:1702 (1993).

The nucleotide sequences are recovered from the biological sample so as to be substantially free of substances that may interfere with the enzymatic amplification procedure, as for example, enzymes, low molecular weight substances such as peptides, proteins, lipids, carbohydrates, and the like. Such methods are known and used in the art.

In a preferred version, an oligonucleotide primer pair that can hybridize to the 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* B204 shown in FIG. 1 are mixed with deoxynucleotides, Taq polymerase, and the extracted DNA

or CDNA. Initial denaturing is at 95° C. for 60 seconds followed by 30 cycles with 60 seconds at 65° C. and 120 seconds at 72° C. Amplified products are optionally separated by methods such as agarose gel electrophoresis. The amplified products can be detected by either staining with ethidium bromide silver stain or by hybridization to a probe as described previously. The preferred oligonucleotide primer pairs amplify a portion of the target sequence of *S. hyodysenteriae* B204 of FIG. 1 to form an about 1.5 kb amplification product.

In an alternative embodiment, at least one probe that hybridizes to the amplified products is labeled with a biotin moiety and/or at least one probe labeled with fluorescently labeled probe. The hybrids are then bound to a solid support such as a bead, multiwell plate, dipstick or the like that is coated with streptavidin. The presence of bound hybrids can be detected using an antibody to the fluorescent tag conjugated to horseradish peroxidase. The enzymatic activity of horseradish peroxidase can be detected with a colored, luminescent or fluorimetric substrate. Conversion of the substrate to product can be used to detect and/or measure the presence of *S. hyodysenteriae* PCR products.

An oligonucleotide primer preferably has a gene sequence that hybridizes to a sequence flanking one end of the DNA sequence to be amplified. The DNA sequence to be amplified is located adjacent the attachment of the single primer, or between the attachment of the two primers. In the use of a pair of oligonucleotide primers, each of the primers has a different DNA sequence and hybridizes to sequences that flank either end of the target sequence to be amplified. Design of primers and their characteristics have been described previously. The preferred DNA sequence of the oligonucleotide primer is positive-sense 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3), or negative-sense 5'-TCCTATTCTCTGACCTACTG (SEQ ID NO:4), or a complement thereof, or a mixture thereof. The primer may also be a degenerate primer that hybridizes to the target gene sequence under hybridization conditions for a primer of that size and sequence complementarity.

For the binding and amplification, the sample DNA is provided in an aqueous buffer formulated with an effective amount of a divalent cation which is preferably MgCl₂, preferably at a concentration of about 0.05–5mM; an effective amount of DNA polymerase with Taq DNA polymerase being preferred in the form of native purified enzyme or a synthesized form such as Ampli-Taq (available commercially from Perkin-Elmer Cetus Corp., San Francisco, Calif.) an effective amount of dNTPs as a nucleotide source, including, dATP, dCTP, dGTP and dTTP, preferably in a saturating concentration, preferably about 200 uM per dNTP; and an effective amount of one or a pair of oligonucleotide primers. The reaction mixture containing the annealed primer(s) is reacted with a DNA polymerase at about 72° C for about 1–10 minutes, preferably about 3–5 minutes, to extend the primers to make a complementary strand of the target gene sequence. The cycle is then repeated by denaturing the DNA strands with heat, annealing and extending, preferably for about 25–40 cycles, preferably about 30 cycles.

The major resulting product is preferably an about 1.55 kb gene sequence which termini are defined by the oligonucleotide primer(s), and whose length is defined by the distance between the two primers or the length of time of the amplification reaction. The gene sequence then serves as a template for the next amplification cycle.

The amplified DNA products are optionally separated from the reaction mixture and then analyzed. The amplified

gene sequences may be visualized, for example, by separating the gene sequences from undesirable side-products and unreacted reagents by electrophoresis in an agarose or polyacrylamide gel, by HPLC separation in an ion exchange column or size exclusion column, or by the ELOSA technique, and other like techniques known and used in the art.

The amplified gene sequence may be directly or indirectly labelled by incorporation of an appropriate visualizing label, as for example, a radioactive, calorimetric, fluorometric or luminescent signal, or the like. In addition, the gel may be stained before or after electrophoresis with a visualizing dye such as ethidium bromide or silver stain wherein the resulting bands may be visualized under ultraviolet light.

The amplified DNA products may also be detected by Southern blot assay, dot blot assay, slot blot assay, or other like analysis, in which the amplified products are separated by electrophoresis on a polyacrylamide or agarose gel, transferred to a membrane such as a nitrocellulose or nylon membrane, reacted with an oligonucleotide probe, and detected. The amplified products may also be detected by reverse blotting hybridization (dot blot) in which an oligonucleotide probe specific to the gene sequence is adhered to a nitrocellulose or polyvinylchloride (PVC) support such as a multi-well plate, and then the sample containing labelled amplified product is added, reacted, washed to remove unbound substance, and a labelled amplified product attached to the probe or the gene sequence imaged by standard methods. For slot blot analysis, the DNA products may be applied to a nylon membrane using a microfiltration apparatus, and hybridized with a labelled oligonucleotide probe, as described by standard methods.

The PCR amplification products can be detected in a multiwell plate by hybridization to a probe specific for *S. hyodysenteriae* such as described in ELOSA methods by Katz et al., cited supra. For analysis of nucleic acids extracted from more than one biological sample, amplified products of each extract may be placed separately into a different well of a multi-well plate, such as a PVC plate, and coated with a capture molecule such as streptavidin. The amplified products specific for *S. hyodysenteriae* in each well can be preferably detected by hybridization to a probe labelled with a moiety that provides for color change such as a fluorescent-labelled probe. A multi-well plate format allows the screening of many different biological samples in a single plate and can be conducted because of the extreme sensitivity of the PCR assay for detection of 1 to 10 organisms per 0.1 gram of feces or other biological sample. A multi-well format likewise allows for the screening and quantitation of the target sequence of a single sample in a series of dilutions.

In those assays, the amplified products are hybridized with an oligonucleotide probe that hybridizes to the target gene sequence. It is preferred that the oligonucleotide probe is hybridizable to an about 1.5 kb gene sequence located within the about 2.3 kb HindIII partial digest restriction fragment of the chromosomal DNA of *S. hyodysenteriae* B204. The probe is hybridizable to all or a portion of the gene sequence under conditions suitable for a probe of that size and sequence complementarity. The probe can distinguish from other microorganisms and cells in a biological sample. Preferably, the oligonucleotide probe, specific to *S. hyodysenteriae* has the sequence: positive-sense 5'-TAGGGGCTGCTGTTCTAGCTGIAAATGC (SEQ ID NO:5). Other probes including detectably labelled probes may be prepared as described previously.

Excess probe is removed from the reaction vessel or support, for example, by washing with a suitable solution.

The presence or absence of the DNA gene product is then determined by visualization of the label on the membrane or the vessel with an imaging system corresponding to the label that is used, including, for example, autoradiography, radiation counting, X-ray, colorimetric, fluorometric or luminescent signal, and the like.

The detection of amplified gene product in the sample indicates the presence of *S. hyodysenteriae* in the biological sample and in the animal. The method is useful in diagnosing an *S. hyodysenteriae* infection in animals, and for detecting animals that are carrier-shedders of *S. hyodysenteriae* in that they have no outward signs of disease but carry the spirochete internally and shed the organism in feces and other body materials and for detecting *S. hyodysenteriae* in environmental samples contaminated with body fluids from infected animals.

The amplification method is also useful for monitoring the efficacy of treatment for an infection caused by *S. hyodysenteriae* to ensure that subclinically infected or carrier-shedder animals are not formed. For example, the assay may monitor the effectiveness of treatment of an animal with an antimicrobial agent such as carbadox, tiamulin, lincomycin, arsanilic acid, chlortetracycline, oxytetracycline, bacitracin, pyrantel tartrate, fenbendazole, gentamicin, neomycin, roxarsone, tylosin, sulfamethazine, virginiamycin, and the like, or a disinfection of the environment using agents such as chlorhexidine, formaldehyde, cresols, phenols and quaternary ammonium compounds among other treatments. In that method, biological samples such as feces, rectal swabs, mucosal scrapings, environmental samples, and the like, are obtained from an animal under treatment for an infection caused by *S. hyodysenteriae* or the environment of the animal, and DNA amplification products of a target sequence of *S. hyodysenteriae* from those samples are analyzed for the presence of the *S. hyodysenteriae*. Samples are obtained from the animal from time to time on a routine basis over the course of treatment, or afterwards, preferably about every day following treatment. The method can be used to monitor the efficacy of disinfection of equipment, fomites and the environment surrounding an animal infected with *S. hyodysenteriae*.

4. Vaccine.

A 56 kDa polypeptide and nucleotide sequence of a 2.3 kb fragment of *S. hyodysenteriae* B204 shown in FIG. 1 are useful in formulating vaccines for immunizing animals against infection by *S. hyodysenteriae*. The polypeptide is also useful in stimulating antibodies specifically reactive with all the serotypes of *S. hyodysenteriae* and not with other closely related non-pathogenic intestinal spirochetes. The vaccine contains an amount of the 56 kDa polypeptide effective to elicit a protective immune response against *S. hyodysenteriae* in the animal, and achieving clinical efficacy, by stimulating the production of antibodies specifically reactive with *S. hyodysenteriae*, exemplified by monoclonal antibody 10G6/G10 (available from Dr. Duhamel at University of Nebraska, Lincoln, Nebr.). The effectiveness of the vaccine is due, at least in part, to the conservative nature of the nucleotide sequence encoding the 56 kDa polypeptide between different serotypes of *S. hyodysenteriae* and to its uniqueness to *S. hyodysenteriae*.

The vaccine is composed of a substantially pure, 56 kDa polypeptide encoded on an about 2.3 kb HindIII restriction fragment of the chromosomal DNA of *S. hyodysenteriae*, exemplified by the chromosomal DNA of *S. hyodysenteriae* isolate B204 shown in FIG. 1. The 56 kDa protein has a predicted amino acid sequence shown in FIG. 1 (SEQ ID NO:2). As used herein, the term "substantially pure" means

that the polypeptide has been extracted and isolated from its natural association with other proteins, lipids, and other like substances from an appropriate host system. Preferably, the DNA fragment has the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1).

The polypeptide may be produced, for example by incorporating the DNA restriction fragment into an expression vector, which includes the restriction fragment operably linked to transcriptional and translational control regions in the vector. The expression vector may then be used to form a transformed cell that includes the DNA fragment which can be used to produce the 56 kDa polypeptide, as described previously. The 56 kDa polypeptide has a predicted amino acid sequence as shown in FIG. 1 (SEQ ID NO:2). A plasmid designated pRED3C6 and carrying the 2.3 kb HindIII insert from *S. hyodysenteriae* B204 has been deposited with the American Type Culture Collection, Rockville, Md., and given Accession No. 75826.

The polypeptide is administered in combination with a physiologically-acceptable, non-toxic, liquid carrier, compatible with the polypeptide and the animal. Suitable pharmacological carriers include, for example, physiological saline (0.85%), phosphate-buffered saline (PBS), Tris (hydroxymethyl aminomethane (TRIS), Tris-buffered saline, and the like.

The vaccine may further include an adjuvant to enhance the immune response in the animal. Such adjuvants include, for example, aluminum hydroxide, aluminum phosphate, Freund's Incomplete Adjuvant (FCA), liposomes, ISCOMs (Mowat et al., *Immunology Today*, 12:383 (1991)), EMULSIGEN PLUS, and the like. The vaccine may also include additives such as buffers and preservatives to maintain isotonicity, physiological pH, stability, and sterility. Parenteral and intravenous formulations of the vaccine may include an emulsifying and/or suspending agent, together with pharmaceutically-acceptable diluents to control the delivery and the dose amount of the vaccine. Other additives may be included as desired, as for example, preservatives, buffering agents, and the like.

The vaccine may be used for alleviating or minimizing the symptoms of an infection caused by *S. hyodysenteriae*. The vaccine may be delivered to the animal, for example, by parenteral delivery, injection (subcutaneous or intramuscular), oral, intrarectal, and the like, by known techniques in the art. For prophylactic and anti-infectious therapeutic use in vivo, the vaccine contains an amount of the polypeptide to stimulate a level of active immunity in the animal to inhibit and/or eliminate *S. hyodysenteriae* pathogenesis.

Factors bearing on the vaccine dosage include, for example, the age and weight of the animal. The range of a given dose is about 1 to 1000 μg of the purified polypeptide per ml, preferably about 50–200 $\mu\text{g}/\text{ml}$ preferably given in about 1–5 ml doses. The vaccine should be administered to the animal in an amount effective to ensure that the animal will develop an immunity to protect against infection by *S. hyodysenteriae*. For example, for an about 20–30 lb swine, a single dose of a vaccine made with Freund's incomplete adjuvant for subcutaneous or intramuscular injection, would contain about 50–200 μg of the purified polypeptide per ml. Preferably, the vaccine is given in an about 1–5 ml dose before or at the time of weaning (2–4 weeks old). The immunizing dose would then be followed by a booster injection given at about 14–28 days after the first injection.

The invention also provides for a vaccine formulation of a live avirulent microorganism transformed with a vector having a 2.3 kb DNA fragment insert from *S. hyodysenteriae*

B204 having a nucleotide sequence as shown in FIG. 1. The fragment can be inserted in to a suitable expression vector such as pYA292, as described previously. Plasmid pYA292 carries a Salmonella origin of replication, an *asd* gene as the only selectable marker gene, P_{trc} regulator/promoter, a ribosome binding site, and an ATG start codon followed by a multiple cloning site and transcriptional terminators. Plasmid pYA292 is designed to express a recombinant antigen as a non-fusion protein.

An expression vector carrying an insert can be transformed into an avirulent immunogenic host cell such as *Salmonella* spp. having attenuating mutations in genes encoding enzymes involved in the synthesis of vital metabolites such as *aro*, *cya*, *crp*, and *asd* as described by Curtis et al., *Infection and Immunity*, 55:3035 (1987); Dugan et al., *J. Infect. Dis.*, 158:1329 (1988); and Edwards et al., *J. Bacteriol.*, 170:3991 (1988). These microorganisms have the capacity to elicit long-lasting humoral and cell mediated immunity at high levels of safety.

The *Salmonella* spp. transformed with a vector carrying a 2.3 kb insert from *S. hyodysenteriae* can be screened for reactivity with antibodies to *S. hyodysenteriae* or by hybridization of DNA or mRNA to a probe specific for the 2.3 kb insert. Transformed *Salmonella* spp. producing a polypeptide encoded by the 2.3 kb insert (SEQ ID NO:1) can be further selected for avirulence and for generating a protective immune response against *S. hyodysenteriae* in an animal. Animals include swine, poultry, ratites, rodents, birds, rheas, and the like. The transformed microorganism that is avirulent and elicits a protective immune response against *S. hyodysenteriae* infection in an animal is the preferred microorganism for the vaccine formulation.

The microorganisms can be combined with carriers or adjuvants as described previously. The vaccine can be administered parenterally, e.g. subcutaneously, intraperitoneally or intramuscularly, orally or intra-rectally. Preferably, the vaccine is administered orally at least one and preferably at least two or more times at intervals of about 14 to 28 days.

The amount of microorganism included in the vaccine formulation is an amount that is effective to generate a protective immune response against infection with *S. hyodysenteriae* as determined by detecting a decrease in the mortality and/or symptoms of disease caused by *S. hyodysenteriae*. The amount of microorganisms will depend, in part, on the route of administration and/or the animal to be immunized. Preferably, an amount of microorganisms is about 10^5 to 10^{10} CFU/ml and more preferably about 10^6 to 10^9 CFU/ml.

Passive Immunization

The polypeptide may also be utilized to raise polyclonal antibody sera and monoclonal antibodies for use in passive immunization therapies. Polyclonal antibodies may be raised to the polypeptide by hyperimmunizing an animal with an inoculum containing the isolated 56 kDa polypeptide. The blood serum may be removed and contacted with immobilized 56 kDa polypeptide reactive with the protein-specific antibodies. The semi-purified serum may be further treated by chromatographic methods to purify IgG and IgM immunoglobulins to provide a purified polyclonal antibody sera for commercial use.

Monoclonal antibodies reactive with the polypeptide may be raised by hybridoma techniques known and used in the art. In brief, a mouse, rat, rabbit or other appropriate species may be immunized with the 56 kDa polypeptide. The spleen of the animal is then removed and processed as a whole cell preparation. Following the method of Kohler and Milstein (*Nature* 256:496-97 (1975)), the immune cells from the

spleen cell preparation can be fused with myeloma cells to produce hybridomas. The hybridomas may then be cultured and the culture fluid tested for antibodies specific for the 56 kDa polypeptide using, for example, an ELISA in which the 56 kDa polypeptide is immobilized onto a solid surface and act as capture antigens. The hybridoma may then be introduced into the peritoneum of the host species to produce a peritoneal growth of the hybridoma, and ascites fluids containing the monoclonal antibody specific to the spirochete may be collected.

The monoclonal antibodies may be used in diagnostic and therapeutic compositions and methods, including passive immunization. Immunoglobulins specific towards the 56 kDa polypeptide may be used to provide passive immunity against an infection caused by *S. hyodysenteriae*. Animals may be treated by administering immunoglobulins intramuscularly at about 100/mg/kg body weight, about every 3-7 days.

Diagnostic Method

Antibodies to the 56 kDa polypeptide may be used in an *in vitro* method of diagnosing an infection of *S. hyodysenteriae* in an animal. The diagnostic method includes contacting a body material potentially containing *S. hyodysenteriae* such as feces, mucosal scraping, or other like tissue sample or body material with a labelled antibody raised to the 56 kDa polypeptide encoded on an about 2.3 kb HindIII restriction fragment of *S. hyodysenteriae*, and detecting the label in the complex formed between the polypeptide in the body material and the labelled antibody. The method may also be performed by combining the body sample with the antibody to the polypeptide, and then contacting the sample with a labelled anti-species antibody reactive with the polypeptide-specific antibody, and then detecting the label.

In addition, the 56 kDa polypeptide may be used as a capture antigen in a method of monitoring and profiling an infection caused by *S. hyodysenteriae*. For example, the polypeptide may be used in an ELISA technique by immobilizing the polypeptide on a solid support such as a polyvinylchloride plate, and contacting the immobilized peptide with a sample material to react with and detect antibodies present in the sample.

The invention further provides an *in vitro* assay for detecting *S. hyodysenteriae*-specific antibodies in a sample. In that method, a sample to be tested is contacted with a composition containing the about 56 kDa polypeptide, which is preferably labelled, to form a conjugate which is then detected. A method for diagnosing an infection by *S. hyodysenteriae* in a biological sample may be carried out with the polyclonal antibody sera or monoclonal antibodies described hereinabove, in an enzyme-linked immunosorbant assay (ELISA), radioimmunoassay (RIA), immunofluorescent assay (IFA), a Northern, Western or Southern blot assay, and the like. In brief, the antibody or biological sample (i.e., tissue sample, body fluid) may be immobilized, for example, by contact with a polymeric material such as polystyrene, polyvinylchloride, a nitrocellulose paper, or other like support means for immobilizing the antibody or sample. The other antibody or biological sample is then added, incubated, and the non-immobilized material is removed by washing or other means. A labeled species-specific antibody reactive with the later is added. The serum antibody or *S. hyodysenteriae* bacteria in the biological sample, is then added and the presence and quantity of label is determined to indicate the presence and amount of *S. hyodysenteriae* bacteria in the biological sample.

5. Mutants of *S. hyodysenteriae* and Vaccine Composition

Once a target sequence specific for *s. hyodysenteriae* and that can distinguish at least one serotype of *S. hyodysente-*

riae from other closely related microorganisms is identified, the target sequence can be altered or mutated to form mutants of *S. hyodysenteriae*. An example of a target sequence is found in all serotypes of *S. hyodysenteriae* and not in other closely-related microorganisms is a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* serotype B204 having the nucleotide sequence shown in FIG. 1. *S. hyodysenteriae* B204 mutants with alterations in the sequence shown in FIG. 1 can be generated using standard methods. Mutants with alterations or deletions in this region can be selected for reduced virulence and/or the ability to stimulate an immune response, preferably a protective immune response to infection with *S. hyodysenteriae* serotypes. These mutants can be useful in vaccine formulations and to elicit antibodies in animals without causing mortality. The antibodies elicited could be used for passive immunization.

Mutants of *S. hyodysenteriae* B204 with alterations in the 2.3 kb HindIII restriction fragment can be generated using standard methods such as chemical mutagenesis (as described in U.S. Pat. No. 4,999,191 to Glisson et al.); transposon mediated mutagenesis (as described in U.S. Pat. No. 4,764,370 to Fields et al.); ultraviolet irradiation; and methods of site-specific mutagenesis (as described in Maniatis et al., cited supra).

Mutant microorganisms can be screened for alteration to the target sequence in a 2.3 kb HindIII restriction fragment using a variety of methods. The mutant can be screened preferably for lack of production of a functional gene product encoded by a 2.3 kb HindIII restriction fragment by lack of reactivity with an antibody specific for all or a portion of the polypeptide encoded by the fragment as exemplified by monoclonal antibody 10G6/G10 or polyclonal antibodies to the peptides, as described in Example VI. The mutants can also be screened for an inability to hybridize to a 2.3 kb HindIII restriction fragment probe or a change in the restriction enzyme fragments that hybridize to the probe. Other screening methods, such as using PCR and sequencing of the mutated fragment, can also be employed.

Alterations or mutations to the target sequence include nucleotide substitutions, deletions, additions (i.e., especially insertion of a transposon) in the 2.3 kb HindIII restriction fragment of serotype B204. Once mutants with alterations to the sequence are identified, they can be further selected for reduced virulence in animals and for the ability to elicit an immune response, preferably a protective immune response, using standard methods. The especially preferred mutants have a deletion of a portion of a 2.3 kb HindIII restriction fragment, have reduced virulence for animals, and elicit a protective immune response that inhibits wild-type *S. hyodysenteriae* infection with at least one serotype of *S. hyodysenteriae*.

The mutants of the invention are useful to elicit antibodies in animals without causing mortality. These antibodies can be useful in methods of passive immunization as described previously.

The mutants of the invention can also be useful in vaccine formulation. A vaccine formulation includes an amount of a mutant of *S. hyodysenteriae* having reduced virulence for animals effective to inhibit *S. hyodysenteriae* infection in animals, wherein the mutation is an alteration of a 2.3 kb HindIII restriction fragment of *S. hyodysenteriae* B204 in admixture with a physiologically acceptable carrier. The mutant microorganism is administered in combination with a physiologically acceptable, non-toxic liquid carrier compatible with the microorganism and the animal. Suitable

pharmacological carriers include, for example, physiological saline (0.85%), phosphate buffered saline, Tris (hydroxymethylamino methane), Tris buffered saline, and the like.

The vaccine may further include an adjuvant to enhance the immune response in the animal. Such adjuvants include, for example, aluminum hydroxide, aluminum phosphate, Freund's incomplete adjuvant, liposome, ISCOMs, EMULSIGEN, and the like. The vaccine may also include additives such as buffers and preservatives to maintain isotonicity physiological pH instability. Parental and intravenous formulations of the vaccine may include emulsifying and/or suspending agent together with pharmaceutically acceptable diluents to control the delivery and dose amount of the vaccine.

The vaccine may be used for alleviating or minimizing the symptoms of disease caused by *S. hyodysenteriae*. The vaccine may be delivered to the animal, for example, by parental delivery, injection (subcutaneous or intramuscular), or oral delivery by techniques known in the art. For prophylactic and anti-infectious therapeutic use in vivo, the vaccine contains an amount of the microorganism effective to stimulate a level of active immunity in the animal to inhibit and/or eliminate *S. hyodysenteriae* pathogenesis.

Factors bearing on the vaccine dosage include, for example, the age and weight of the animal. The range of a given dose is about 10^5 to 10^{10} CFU of the microorganism per ml, preferably about 10^6 to 10^9 CFU/ml, preferably given in about 1 to 5 ml doses. The vaccine can be administered to the animal as a single dose but is preferably administered as 2 or 3 doses over an 8–10 week period.

The invention will be further described by reference to the following detailed examples, wherein the methodologies are as described below. These examples are not meant to limit the scope of the invention that has been set forth in the foregoing description. Variation within the concepts of the invention are apparent to those skilled in the art. The disclosures of the cited references throughout the specification are incorporated by reference herein.

EXAMPLE I

Bacterial Culture and Growth Conditions

Several genera of bacteria were cultured and grown to provide nucleic acid samples for analysis by polymerase chain reaction and Southern blot hybridization with the 2.3 kb HindIII DNA fragment of clone pRED3C6.

1. Bacterial Strains.

Serpulina hyodysenteriae strain B78 serotype 1 (ATCC# 27164; Harris et al., *Int. J. Syst. Bact.*, 29:102-109 (1979); and Kinyon et al., *Infect. Immun.*, 15:638-646 (1977)), strain B204 serotype 2, strain B169 serotype 3, and strain A1 serotype 4, were obtained from J. M. Kinyon, College of Veterinary Medicine, Iowa State University, Ames, Iowa. Reference *S. hyodysenteriae* strains B234 serotype 1, B8044 serotype 5, B6933 serotype 6, AcK 300/8 serotype 7 were provided by L. A. Joens, Department of Veterinary Science, University of Arizona, Tucson, Ariz. Reference strains FM-88-90 serotype 8, FMV 89-3323 serotype 9 were provided by M. Jacques, Faculté de Médecine Vétérinaire, Université de Montréal, Saint-Hyacinthe, Québec, Canada. Li et al., *J. Clin. Microbiol.*, 29:2794-2797 (1991). The reference isolates for WBHIS, *Serpulina innocens*, isolates B256 (ATCC# 29796; and Harris et al., cited supra) and 4/71 were obtained from the American Type Tissue Culture Collection, Rockville, Md., and T. B. Stanton, National Pig Disease Center, Ames, Iowa, respectively.

A total of 13 field isolates representing three genotypic groups of WBHIS distinct from *S. innocens* were obtained from porcine feces, porcine rectal swabs, and porcine colonic mucosal scrapings submitted either to the Veterinary Diagnostic Center, University of Nebraska-Lincoln or Agriculture Canada, Saint-Hyacinthe, Québec. Lee et al., *Vet. Microbiol.*, 34:35–46 (1993); and Ramanathan et al., *Vet. Microbiol.*, 37:53–64 (1993). The WBHIS isolates B359 and B1555a were obtained from J. M. Kinyon, and isolate D9201243A was provided by R. L. Walker, California Veterinary Diagnostic Laboratory System, University of California, Davis, Calif. The WBHIS isolate 16 (ATCC# 49776; Jones et al., *J. Clin. Microbiol.*, 24:1071–1074 (1986)) obtained from an HIV-positive homosexual male with diarrhea was provided by R. M. Smibert, Virginia Polytechnic Institute, Blacksburg, Va. In addition, species of other genera of bacteria were obtained and cultured accordingly to standard methods as described below. In some instances, chromosomal DNA was obtained.

Treponema succinifaciens, isolate 6091 (ATCC# 33096; Cwyk et al., *Arch. Microbiol.*, 122:231–239 (1979)) and *Bacteroides vulgatus* (ATCC# 31376) were obtained from the American Type Tissue Culture Collection. Chromosomal DNA from *Spirochaeta aurantia* was provided by E. P. Greenberg, University of Iowa, Iowa City, Iowa. The *Treponema pallidum* chromosomal DNA was provided by M. V. Norgard, The University of Texas Health Science Center, Houston, Tex. Chromosomal DNA from representative stains of each of the ten genetic groups in the family Leptospiraceae including *Leptospira biflexa* serovars patoc, semaranga, and codice, *Leptospira interrogans* serovars icterohaemorrhagiae, fortbragg, ballum, celledoni, lyme, and borincana, and *Leptonema illini* serovar illini were provided by R. L. Zuerner, National Pig Disease Center, Ames, Iowa. Chromosomal DNA from *Borrelia burgdorferi*, *Campylobacter coli* and *C. hyointestinalis* were provided by M. P. Murtaugh, University of Minnesota, St. Paul, Minn. Isolates of *Salmonella choleraesuis* and *S. typhimurium* were provided by P. J. Fedorka-Cray, National Pig Disease Center, Ames, Iowa. The *Escherichia coli* DH5 α was purchased from a commercial source (GIBCO-BRL, Gaithersburg, Md.).

2. Medium and growth conditions.

For isolation of DNA, cultures of *Serpulina* spp., WBHIS, and *T. succinifaciens* were propagated in pre-reduced anaerobically-sterilized (PRAS) trypticase soy broth supplemented with 0.5% (wt/vol) glucose (Sigma Chemical Co., St. Louis, Mo.), 0.05% (wt/vol) cysteine hydrochloride monohydrate (Sigma), 1.0% (wt/vol) yeast extract (BBL Microbiology Systems, Becton Dickinson and Co., Cockeysville, Md.), 2.0% (v/v) bovine fetal serum (HyClone Laboratories, Inc., Logan, Utah), 0.2% (wt/vol) sodium bicarbonate and 5.0% (v/v) sterile porcine fecal extract as described by Kunkle et al., *J. Clin. Microbiol.*, 24:669–671 (1986), except that 1% (vol/vol) of room air was injected at the time of inoculation (Stanton et al., *Vet. Microbiol.*, 18:177–190 (1988)). Broth cultures were grown to late logarithmic phase in 5 ml volumes in Hungate tubes or in 250 ml volumes in serum bottles. Cultures were stirred constantly using a magnetic stirrer at 37° C. under a 10% hydrogen, 10% carbon dioxide and 80% nitrogen atmosphere for 48 to 72 hours. Cultures of *Salmonella* spp., *Bacteroides vulgatus* and *Escherichia coli* were grown at 37° C. with shaking in Luria-Bertani broth to late logarithmic phase.

Chromosomal DNA was purified as previously described by Ramanathan et al., *Vet. Microbiol.*, 37:53–64 (1993), except that the final pellet was resuspended in sterile H₂O to a final concentration of 12.5 ng/ μ l.

EXAMPLE II

Library Construction and Recombinant Screening

A library of *S. hyodysenteriae* isolate B204 in *E. coli* DH5 α was constructed.

1. Preparation of the DNA Library.

Serpulina hyodysenteriae isolate B204 and weakly β -hemolytic intestinal spirochetes of swine; isolates B359 and B1555a (courtesy of J. M. Kinyon, College of Veterinary Medicine, Iowa State University, Ames, Iowa); *Serpulina innocens* isolate B256 (American Type Culture Collection, Rockville, Md.); and *E. coli* strain DH5 α (Bethesda Research Laboratories, Inc., Gaithersburg, Md.) were used. *Serpulina* spp. and other spirochetes were propagated in FS medium (Kunkle et al., *J. Clin. Microbiol.*, 24:669–671 (1986)) to a density of 10⁸ to 10⁹ cells per ml at 37° C.

DNA was isolated, by a modification of a previously reported method (Caputa et al., *J. Clin. Microbiol.*, 29:2418–2413 (1991)), from a 500 ml culture of *S. hyodysenteriae*, isolate B204. Briefly, spirochetes were centrifuged, washed twice in 100 ml of TE buffer (10 mM Tris-HCl [pH 8.0], 1 mM Na₂EDTA [pH 8.0]), and resuspended in 25 ml of 50 mM Tris-HCl (pH 8.0), 50 mM Na₂EDTA (pH 8.0). N-lauroylsarcosine was added to a final concentration of 2% (wt/vol), followed by the addition of 100 μ g/ml of proteinase K. The mixture was incubated at 56° C. for 16 hours. Phenylmethylsulfonyl fluoride was added to a final concentration of 1 mM and the mixture was incubated at room temperature for 10 minutes, then the solution was mixed with 0.1 volume of 7.5M ammonium acetate, and the DNA was precipitated with 2 volumes of ethanol. The DNA precipitate was recovered on a glass rod, washed with 70% ethanol, resuspended in TE buffer to a concentration of 0.5 to 1 μ g/ μ l and stored at –20° C. until needed.

Standard cloning protocols were used for DNA manipulations. Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989). Purified chromosomal DNA from *S. hyodysenteriae*, isolate B204 was incompletely digested with HindIII (Stratagene, LaJolla, Calif.) and 4 kb to 9 kb fragments were obtained by centrifugation on a 5% incremental sucrose gradient with a range of 10 to 40%. After dialysis against TE buffer, the DNA fragments were ligated with T4 DNA ligase (Stratagene, LaJolla, Calif.) to plasmid vector pUC18 (Bethesda Research Laboratories, Inc., Gaithersburg, Md.) dephosphorylated with calf intestinal alkaline phosphatase (Bethesda Research Laboratories, Inc., Gaithersburg, Md.). *E. coli*, strain DH5 α cells were transformed with the ligation mix and recombinant clones were selected by growth on Luria-Bertani (LB) agar containing 100 μ g/ml ampicillin (Bethesda Research Laboratories, Inc., Gaithersburg, Md.), 12 μ g/ml isopropyl- β -D-thiogalactosidase (IPTG) (Bethesda Research Laboratories, Inc., Gaithersburg, Md.), and 40 μ g/ml 5-bromo-4-chloro-3-indoyl- β -D-galactoside (X-gal) (Bethesda Research Laboratories, Inc., Gaithersburg, Md.). Replica plating was executed by transferring colonies to nitrocellulose membranes (HAFT, 0.45 μ m pore size) (Millipore Corp., Bedford, Mass.).

Transformed *E. coli* DH5a cells were screened with monoclonal antibodies, such as 10G6/G10, by colony immunoblotting. These monoclonal antibodies can be prepared as described in Example VI or are available from Dr. Duhamel, University of Nebraska, Lincoln, Nebr. A monoclonal antibody used for screening has been designated

10G6/G10 and is an IgM antibody that is specific for cell free supernatants derived from *S. hyodysenteriae*. Briefly, replica membranes (Millipore cat. #HATF 137 50, Millipore Corp., Bedford, Mass.) were lysed in chloroform vapor followed by overnight incubation in lysis/blocking solution (5% nonfat dry milk, 0.5 M MgCl₂, 40 mg/ml lysozyme, 100 mg/ml chloramphenicol, 2 mg/ml DNase) at room temperature. Membranes then were incubated sequentially at room temperature with ascites fluid for 2 hours, followed by biotin-labeled goat anti-mouse [IgA+IgG+IgM(H+L)] antibody (Kirkegaard and Perry Laboratories Inc., Gaithersburg, Md.) for 1 hour, peroxidase-labeled streptavidin (Kirkegaard and Perry) for 45 minutes, and 4-Chloro-1-Naphthol (Kirkegaard and Perry) for 5 minutes. Five 5-minute washes with wash buffer (1M Tris-base, 2M NaCl, 5% nonfat dry milk and 0.05% NP-40 (pH 7.5)) were performed between each incubation step. One immunopositive clone, designated pRED3C6, was identified based on development of a dark purple precipitate.

EXAMPLE III

DNA Sequencing and Primer Selection

The immunopositive clone was amplified and the plasmid DNA with insert was isolated and sequenced. The insert was identified as about a 2.3 kb HindIII partial digest fragment. The nucleotide sequence of the 2.3 kb insert fragment was used to design primers.

1. Plasmid and insert DNA isolation and sequencing

The recombinant plasmids were isolated (Magic™ Minipreps, Promega Corp., Madison, Wis.), digested with the restriction enzyme HindIII (GIBCO-BRL), and the DNA fragments separated by electrophoresis in a 0.8% agarose gel using TAE running buffer (16 mM Tris-base, 8 mM sodium acetate, and 1 mM EDTA, pH 7.5) containing 0.66 µg of ethidium bromide per ml. The resulting bands were visualized and photographed under ultraviolet light with a Polaroid MP 4 land camera. Recombinant DNA bands were excised from the gels with a razor blade, isolated (GeneClean II, BioRad, Richmond, Calif.), and subjected to the restriction enzymes AccI, AluI, EcoRI, DraI, HaeIII, SspI and XbaI (GIBCO-BRL). Resulting fragments were purified and subcloned into the vector pUC18 for sequencing using a Model

4,000 DNA sequencer (Li-Cor, Inc. Lincoln, Nebr.). Sequencing data were analyzed and assembled using the Program manual for the GCG Package Version 7, Apr. 1991 from the Genetics Computer Group (1991). Results are shown in FIG. 1.

Presence of -10 and -35 sequences along with a probable ribosomal binding site upstream from the transcription start codon of the gene suggested a single protein transcript. Analysis of the deduced amino acid sequence of the ORF for prediction of a membrane translocation signal failed to identify any significant homology with other known N-terminal export signal sequences. However, the C-terminal sequence of the gene contained a highly charged amino acid followed by a Ser Thr-rich amphophilic region suggestive of an extracellularly secreted protein. Lory, *J. Bacteriol.*, 174:3423-3428 (1992). The sequence encodes a polypeptide with a predicted molecular weight of a 56 kDa protein. The nucleotide sequence may also encode related polypeptides of smaller molecular weight that could be post-translationally processed.

Furthermore, comparisons of the DNA sequence of the recombinant 2.3-kb DNA fragment of pRED3C6 with the sequences of the tly A gene (Muir et al., *Infect. Immun.*, 60:529-535 (1992)), tly B, and tly C gene of *S. hyodysenteriae* (A. Agnes H. M. ter Huurne, *Ph.D. Dissertation*, University of Utrecht, Utrecht, Netherlands, (1993)), and the fla A gene of Koopman et al., *Inf. and Imm.*, 60:2920 (1992) indicated that these sequences had less than 45% nucleotide sequence identity and less than 25% amino acid identity. The nucleotide and amino acid sequences were also compared with nucleotide sequences encoding a 39 kDa protein from *S. hyodysenteriae*, shown in ML Technology Ventures' PCT Application No. WO91/04036 published Apr. 4, 1991, and had less than 47% DNA sequence identity and less than 25% amino acid identity. A comparison to sequences published in EP0350715 also showed less than 45% DNA sequence identity and less than 20% amino acid identity. The nucleotide sequence comparison also revealed in most cases that there were little or no regions of contiguous sequence identity of greater than 10 base pairs. The sequence comparisons were conducted using GCG Package, version 7.3, Jun. 1993 (Genetics Computer Group, Madison Wis.). See Table II.

TABLE II

Sequence analysis comparisons between
the 2.3-kb fragment of clone pRED3C6 and sequences currently available.†

Sequence	Gene	Amino acid		Nucleotide	
		% Similarity	% Identity	% Identity	GAPs
Muir et al., 1992	tly A	55.7	24.4	44.5	8
Ter Huurne et al., 1993	tly B	45.4	19.9	42.2	16
Ter Huurne et al., 1993	tly C	42.0	19.5	44.4	20
Koopman et al., 1993	fla A	42.6	14.8	44.1	11
ML Tech. Vent., 1991	Copy #1	42.9	18.2	44.6	
	Copy #2	47.1	19.9	43.2	
	Copy #3	44.7	19.8	44.4	
	Copy #4	43.3	17.2	43.6	
	Copy #5	42.9	19.0	43.1	
	Copy #6	42.8	18.8	41.2	
	Copy #7	42.9	21.6	45.0	
	Copy #8	43.7	19.8	46.5	
ML Tech. Vent., 1990	38 kDa	46.0	17.0	42.0	
	60 kDa	43.6	18.4	41.7	

†Gap, In; Genetics Computer Group. 1991. Program Manual for the GCG Package, Version 7.3, June 1993, 575 Science Drive, Madison, Wisconsin 53711, p. 5-27 to 5-53.

Additionally, an exhaustive search of the EMBL database failed to identify any sequence with significant DNA or amino acid homology with the recombinant 2.3 kb HindIII DNA fragment and its deduced amino acid sequence.

While not meant to limit the invention in any way, there is evidence that the 56 kDa polypeptide encoded by the recombinant 2.3 kb DNA fragment of pRED3C6 represents a putative *S. hyodysenteriae* hemolysin distinct from those described previously. This evidence indicates that transformation of a non-hemolytic *E. coli* host with pRED3C6 conferred hemolytic activity to the *E. coli* host. Cleavage and subcloning of the ORF of the recombinant 2.3 kb DNA fragment resulted in complete loss of hemolytic activity of the *E. coli* host. The fact that the 2.3 kb DNA sequence reacts specifically with all the serotypes of *S. hyodysenteriae* and none of the non-pathogenic intestinal spirochetes further indicates that it may be associated with a virulence determinant of *S. hyodysenteriae* such as hemolysin.

2. Southern blotting

Approximately 2 μ g of chromosomal DNA from *S. hyodysenteriae* serotypes 1 to 7, *S. innocens* isolates B256 and 4/71, WBHIS isolates B359 and B1555a and *T. succinifaciens* were digested with HindIII, electrophoretically separated on a 0.8% agarose gel, and transferred by capillary diffusion as described by Southern et al., *J. Mol. Biol.*, 98:503-517 (1975), to nylon membranes (HybondTM-N, Amersham, Arlington Heights, Ill.). Prehybridization, hybridization, and washing steps with a recombinant DNA fragment (2.3 kb) obtained from the immunopositive clone pRED3C6 labelled with [α -³²P] dCTP using an oligolabeling kit (Pharmacia LKB Biotechnology, Piscataway, N.J.) were performed as described by Ramanathan et al., *Vet. Microbiol.*, 37:53-64 (1993). For slot blot analysis purified genomic DNAs from cultivable reference isolates of the order Spirochaetales including *S. hyodysenteriae* serotypes 1 through 9, *S. innocens* isolates B256 and 4/71, 16 isolates of WBHIS belonging to 3 genotypic groups distinct from *S. innocens*, *Spirochaeta aurantia*, *Treponema* spp., *Borrelia burgdorferi*, and representatives of each of the 10 genetic groups of the family Leptospiraceae, as well as enteric bacteria including *Escherichia coli*, *Salmonella* spp., *Campylobacter* spp., and *Bacteroides vulgatus* were applied to nylon membranes (Zeta-probeTM, Bio-Rad) using a microfiltration apparatus (Bio-Dot SF[®], Bio-Rad). Prehybridization, hybridization and washing steps with a recombinant DNA fragment obtained from the immunopositive clone pRED3C6 labelled with [α -³²P] dCTP were carried out as described Ramanathan et al., *Vet. Microbiol.*, 37:53-64 (1993). The results are shown in FIG. 2.

Southern blot hybridization of [α -³²P]dCTP labelled 2.3-kb fragment from clone pRED3C6 yielded a strong hybridization signal with chromosomal DNA from reference isolates of *S. hyodysenteriae* serotypes 1 through 7, but not with *S. innocens* isolates B256 and 4/71, WBHIS isolates B359 and B1555a, and *T. succinifaciens* (FIG. 2). When the same probe was reacted with chromosomal DNAs obtained from other cultivable reference isolates of the order Spirochaetales as well as enteric bacteria including *Escherichia coli*, *Salmonella* spp., *Campylobacter* spp., and *Bacteroides vulgatus* in a slot blot hybridization assay, a specific signal was observed only with chromosomal DNA obtained from reference isolates of *S. hyodysenteriae* serotypes 1 through 9.

3. Primers and internal probe for PCR and Southern blot analysis

An oligonucleotide primer pair (positive-sense 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3) and negative-sense 5'-TCCTATTCTCTGACCTACTG

(SEQ ID NO:4)) and an internal *S. hyodysenteriae*-specific oligonucleotide probe (positive-sense 5'-TAGGGGCTGCTGTTCTAGCTGTAAATGC (SEQ ID NO:5)) were designed and synthesized (Integrated DNA Technologies, INC. Coralville, Iowa) based on results of DNA sequence analysis of the recombinant DNA fragment of the immunopositive clone pRED3C6. The primers were used for amplification either of purified chromosomal DNAs or total DNA extracted by the method described in Example I and Example V either from normal porcine feces inoculated with *S. hyodysenteriae* cells, or from porcine feces, porcine rectal swabs, and porcine colonic mucosal scrapings obtained from swine with clinical an infection caused by *S. hyodysenteriae*.

Primers or probes can be designed based on the sequence of 2.3 kb HindIII fragment shown in FIG. 1. Primers can be designed using primer search algorithms such as Primer Detective (Clontech Laboratories, Inc., Palo Alto, Calif.). Probes can be designed using OLIGO Computer Program (Rychlik and Rhoades, "A Computer Program for Choosing Optimal-Oligonucleotides for Filter Hybridization, Sequencing and In Vitro Amplification of DNA", *Nucleic Acid Res.*, 17:8543-8551 (1989) or other commercially-available computer software with similar applications.

EXAMPLE IV

Analysis of PCR Products from Several Strains of Bacteria

The primers designed as described in Example III were used to amplify either purified chromosomal DNA or total DNA extracted from normal porcine feces inoculated with *S. hyodysenteriae* cells or porcine feces, porcine rectal swabs, and porcine colonic mucosal scrapings obtained from swine with an infection caused by *S. hyodysenteriae*. The PCR products were analyzed by hybridization to an internal probe as described in Example III.

The DNA was amplified using a hot start PCR as described by the manufacturer (GeneAmpTM PCR System 480, Perkin Elmer, Norwalk, Conn.) in a total volume of 75 μ l containing 4 mM MgCl₂; 1 \times of PCR buffer; 0.2 mM of each dATP, dCTP, dGTP, dTTP (Perkin-Elmer Cetus); 75 pmol of primers; and 1.5 U of Taq DNA polymerase (Perkin-Elmer Cetus) in sterile filtered autoclaved water. Initial denaturing was for 60 s at 95 $^{\circ}$ C., followed by 30 cycles (60 s at 65 $^{\circ}$ C. and 120 s at 72 $^{\circ}$ C.). The amplified products were visualized in 1.25% agarose gels ran at 3 V/cm and stained with ethidium bromide. Southern blots were prehybridized and hybridized each for 1 hour at T_m-10 $^{\circ}$ C. with an internal *S. hyodysenteriae*-specific oligonucleotide probe 5'-end labelled with [γ -³²P]ATP using T4 polynucleotide kinase (Pharmacia) as described by Maniatis et al., *Molecular Cloning: Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1982). Washes consisted of 1 \times SSPE (3.6M NaCl, 0.2M NaH₂PO₄, 0.02M EDTA, pH 7.7) with 0.1% SDS (3 times for 5 min at room temperature and once for 5 min at T_m-10 $^{\circ}$ C.). The membranes were exposed to X-OMAT AR Cronex radiograph film (Eastman Kodak Company, Rochester, N.Y.) in a cassette with lightning plus intensifying screens (DuPont, Wilmington, Del.) at -70 $^{\circ}$ C.

With purified chromosomal DNA from each of the 9 serotypes of *S. hyodysenteriae* as template, PCR assay resulted in 1.55 kb products (FIG. 3). The specificity of the 1.55 kb products for *S. hyodysenteriae* was confirmed based on production of a restriction endonuclease pattern of the

PCR products identical to the predicted restriction map analysis of pRED3C6 (data not shown) and positive hybridization signal with the *S. hyodysenteriae*-specific internal oligonucleotide probe (FIG. 3). The specificity of the reaction for *S. hyodysenteriae* was further confirmed by the absence of products and hybridization signal, respectively after gel electrophoresis and Southern blot hybridization with internal *S. hyodysenteriae*-specific oligonucleotide probe of PCR amplified chromosomal DNA obtained from other cultivable reference isolates of the order Spirochaetales, including *S. innocens* isolates B256 and 4/71, other genotypic groups of WBHIS distinct from *S. innocens*, as well as enteric bacteria including *Escherichia coli*, *Salmonella* spp., *Campylobacter* spp., and *Bacteroides vulgatus*.

EXAMPLE V

PCR Detection of *S. hyodysenteriae* from Diagnostic Samples

Samples from uninfected and infected swine were evaluated by the PCR method for diagnosis of the presence of *S. hyodysenteriae*.

1. PCR detection of *S. hyodysenteriae* in porcine feces

The sensitivity of the PCR detection of *S. hyodysenteriae* in porcine feces was determined by two separate methods. In a first method, 10-fold serial dilutions of spirochete broth cultures were added to constant volumes of undiluted normal porcine feces in two separate experiments. The sensitivity of the PCR assay was estimated based on the numbers of spirochete cells in the original culture as determined by a Petroff-Houser cell counting chamber. Briefly, sterile tubes containing 0.1 g of normal feces were inoculated with 1 ml containing either sterile PBS (negative control) or serial ten-fold dilutions of *S. hyodysenteriae* isolate B204 cells in sterile PBS, from 10^5 to 10^{-2} and vortexed for 5 minutes. The samples were allowed to stand for 10 minutes, then the supernatant (approximately 0.9 ml) was drawn off and processed for total DNA extraction as described by the manufacturer (Nucleon DNA extraction kit, Scotlab, Shelton, Conn.) except that the samples were heated at 100° C. for 15 min prior to the cell lysis step and 5M sodium perchlorate deproteinization was replaced by 100 μ g of proteinase K per ml. Total DNA from each tube was used for PCR amplification followed by agarose gel electrophoresis and Southern blot analysis using the *S. hyodysenteriae*-specific oligonucleotide probe.

In a second method, feces collected from two untreated swine at the onset of an infection caused by *S. hyodysenteriae* were serially diluted ten-fold (10^{-1} to 10^{-12}) in 2 ml volumes of sterile PBS. One-ml fractions from each dilution then were processed for determination of the total numbers of viable *S. hyodysenteriae* by a plate counting method, and detection of *S. hyodysenteriae*-specific products by the PCR assay, respectively. For the plate counting method, a total of 10 drops of 10 μ l each were placed onto freshly made BJ selective medium, and the number of colony forming units (CFU) per 0.1 ml was determined after incubation at 42° C. in the Gas Pak Anaerobic System (BBL) for 9 days. One ml fractions from each ten-fold dilution were processed for total DNA extraction, and PCR detection of *S. hyodysenteriae*, as described above. The specificity of the PCR assay for detection of *S. hyodysenteriae* in diagnostic specimens was compared with conventional bacteriological culture on BJ medium incubated anaerobically at 42° C. for 10 days. Porcine feces (n=3), porcine rectal swabs (n=2), and porcine colonic mucosal scrapings (n=4) obtained from six different

premises where clinical signs of an infection caused by *S. hyodysenteriae* were reported by the referring veterinarians (Duhamel et al., *J. Vet. Diagn. Invest.*, 4:285–292 (1992)), were processed for PCR assays and cultures. For PCR assays, total DNA was extracted from 100 μ l of supernatants from either dysenteric porcine feces or porcine colonic mucosal scrapings, as described above. Rectal swabs were mixed with 1 ml of sterile PBS for 2 minutes, and the total DNA was extracted from the supernatants.

Negative controls were included in all PCR assays for detection of *S. hyodysenteriae* in porcine feces. In the spiked feces experiments, unspiked feces and feces spiked with 10^{-2} dilution of broth culture (100-fold dilution beyond the numbers of spirochetes estimated by the Petroff-Houser cell counting chamber) were used as negative controls. In the experiments using fecal samples from swine at the onset of an infection caused by *S. hyodysenteriae*, dilutions beyond 10^{-10} were considered as negative controls (according to Kunkle et al., *J. Clin. Microbiol.*, 26:2357–2360 (1988), dysenteric feces contain between 10^6 and 10^{10} CFU/g). In the experiments testing diagnostic specimens, the fecal sample containing a WBHIS was used as the negative control.

The sensitivity of the PCR assay for detection of *S. hyodysenteriae* in serial ten-fold dilutions of spirochete broth cultures added to normal porcine feces was 1 organism per 0.1 g of feces in the first experiment (data not shown), and 10 organisms per 0.1 g of feces in the second experiment (FIG. 3). The number of spirochetes in dysenteric feces from 2 untreated swine were comparable to those reported previously for the BJ selective culture medium; 1×10^5 and 2×10^5 CFU/0.1 ml, respectively. Kunkle et al., *J. Clin. Microbiol.*, 26:2357–2360 (1988). Presence of *S. hyodysenteriae*-specific products at dilutions up to 10^{-9} in both fecal specimens by PCR assay indicated a 1,000 fold increase in sensitivity compared with conventional culture. The 10^{-1} to 10^{-12} dilutions yielded negative results by both methods.

Examination of porcine feces, porcine rectal swabs, and porcine colonic mucosal scrapings obtained from nine swine on six different premises by PCR assays yielded 1.55-kb products in all samples where *S. hyodysenteriae* was identified by conventional bacteriological culture method, as shown in Table III below. The one sample which was negative by PCR assay yielded WBHIS by culture. Table III shows a comparison of conventional bacteriological culture method on selective BJ agar medium and polymerase chain reaction (PCR) assay for detection of *Serpulina hyodysenteriae* in diagnostic specimens.

TABLE III

Premises	Sample		Result		
	Number	Type	Culture	PCR	
A	1	Feces	<i>S. hyodysenteriae</i>	+	
	2	RS†	<i>S. hyodysenteriae</i>	+	
B	1	MS	<i>S. hyodysenteriae</i>	+	
C	1	MS	<i>S. hyodysenteriae</i>	+	
D	1	MS	<i>S. hyodysenteriae</i>	+	
E	2	Feces	<i>S. hyodysenteriae</i>	+	
F	1	MS	WBHIS	-	
Total:	6	9			

†RS = Porcine rectal swab.

MS = Porcine colonic mucosal scraping.

EXAMPLE VI

Preparation of Monoclonal Antibodies to *S. hyodysenteriae* Antigens

Monoclonal antibodies were raised against cell-free supernatant antigens from *S. hyodysenteriae* produced by a previously described method (Dupont et al., in press) according to standard methods (Hugo et al., *J. Clin. Microbiol.*, 25:26–30 (1987)). Polyclonal antibodies were raised against two different synthetic peptides selected from the predicted amino acid sequence of the 2.3 kb insert of clone pRED3C6.

1. Monoclonal antibody production

Eight- to ten-week old BALB/c mice were immunized intraperitoneally with 100 μ g of cell-free supernatant antigens from *S. hyodysenteriae*, isolate B204, concentrated 10 times using a YM5 Diaflow ultrafilter (Amicon, Beverly, Mass.) and mixed with equal volumes of Freund's complete adjuvant. Dupont et al., *Vet. Microbiol.*, in press (VETMIC 723). Identical booster injections containing 50 μ g of cell-free supernatant antigens in Freund's incomplete adjuvant were given 14, 28, and 42 days later. Four days after the booster injection, spleen cells were harvested and fused with SP 2/0 cells using 50% polyethylene glycol. Hybridomas producing antibodies that reacted with cell-free supernatant antigens from *S. hyodysenteriae*, isolate B204, by ELISA were cloned by limiting dilution and stabilized before injection into mice for ascites production. Hugo et al., *J. Clin. Microbiol.*, 25:26–30 (1987). Monoclonal antibodies 467, F11, 1D8/E11, 3E1D/F1, 6C1D/F8 and 10G6/G10 were identified and are available from Dr. Duhamel, University of Nebraska, Lincoln, Nebr. Hybridoma 10G6/G10 producing an IgM monoclonal antibody that reacted with cell-free supernatant antigens of *S. hyodysenteriae* by ELISA was cloned by limiting dilution and stabilized before injection into mice for ascites production (Hugo et al., cited supra) and is available from Dr. Duhamel, University of Nebraska, Lincoln, Nebr.

2. Polyclonal antibodies to synthetic peptides

Information on the predicted amino acid sequence encoded by the 2.3 kb fragment of clone pRED3C6 provides a basis for identification of antigenic domains. Using Hopp-Woods hydrophobicity plots as an indicator of surface orientation and potential antigenicity, two peptides were synthesized and used for production of hyperimmune sera in guinea pigs. Hopp et al., *Mol. Immunol.*, 20:483 (1983).

Polyclonal antibodies were produced in adult Hartley Albino guinea pigs against synthetic peptides A (DPAKASRPFD) and B (IPLFEALKPKT) derived from the predicted amino acid sequence of nucleotides 500–529 (peptide A) and 2093–2126 (peptide B) of the 2.3 kb fragment of clone pRED3C6. The initial injection consisted of 50 μ g of each peptide diluted in 100 μ l of sterile water and mixed with 100 μ l of Freund's complete adjuvant administered subcutaneously. Booster injections containing 100 μ g of each peptide in 100 μ l of sterile water mixed with equal volumes of Freund's incomplete adjuvant were given 14 and 28 days after the initial injection. Final bleeding was completed 7 days after the last booster injection.

EXAMPLE VII

Identification of the Site of *S. hyodysenteriae* Persistence in Carrier Swine

A PCR test may be used for screening replacement stock and during herd elimination programs by analyzing fecal

shedding patterns by subclinically-infected swine including those on medication as described below.

1. Define the pattern and identify the site of *S. hyodysenteriae* persistence in carrier swine

About 20 specific pathogen-free (SPF) swine will be inoculated with *S. hyodysenteriae*, isolate B204, as previously described. Elder et al., cited supra. Each swine will be monitored daily for clinical signs of an infection caused by *S. hyodysenteriae* (usually within 14 days post-inoculation). At least about 30 percent of the swine most likely will die or need to be euthanatized because of severe clinical signs. The remaining naturally-recovered swine will be randomly allocated to two experimental groups and housed in separate isolation rooms of the Pig Research Facility (ARF) of the Department of Veterinary and Biomedical Sciences.

The pattern of *S. hyodysenteriae* shedding in naturally-recovered swine (continuous versus sporadic) will be assessed using the PCR assay described in Example V, and bacterial culture of fecal specimens collected every other day for 30 days from the day of cessation of bloody diarrhea (usually 14 to 21 days after the onset of clinical signs). At the end of 30 days, the effect of stress on fecal shedding of the spirochetes will be determined. Swine from one experimental group will be taken for a 40-mile truck-ride, placed into a disinfected room of the ARF, and the pattern of *S. hyodysenteriae* fecal shedding will be monitored for an additional 7 days.

At the end of the observation period, the swine in each group will be euthanatized and the distribution of *S. hyodysenteriae* in 15 predetermined sites along the wall of the distal ileum, cecum, spiral colon, descending colon, and rectum of each swine will be determined by PCR and culture. Duhamel et al., *J. Vet. Diag. Invest.*, 4:285 (1992) and Elder et al., cited supra. DNA extracted from fecal samples (live swine) and mucosal scrapings (necropsy specimens) obtained from each swine will be subjected to PCR amplification followed by agarose gel electrophoresis and Southern blot analysis using the *S. hyodysenteriae*-specific oligonucleotide probe, as described in Examples IV and V. Results of PCR assay will be compared with conventional bacterial anaerobic culture and will provide an indication of the site of *S. hyodysenteriae* levels in feces and tissues of the swine over time post-infection.

2. Analysis of the pattern of *S. hyodysenteriae* shedding in feces of medicated swine

A total of 30 SPF swine will be infected with *S. hyodysenteriae*, isolate B204. At the onset of clinical signs, swine will be randomly allocated to five experimental groups of 5 swine each and medicated with the following antimicrobials: Group 1. Carbadox: 50 g/t in feed continuous; Group 2. Tiamulin: 3.5 mg/lb/day in water for 5 days; Group 3.35 g/t in feed continuous; Group 4. Lincomycin: 3.8 mg/lb/day in water for 6 days; Group 5. 100 g/t in feed continuous. The pattern of *S. hyodysenteriae* fecal shedding will be monitored using PCR and bacterial culture every other day for 3 weeks after the initiation of medication. At the end of the observation period, the swine will be euthanatized and the presence of *S. hyodysenteriae* in the intestinal tract of each swine will be determined by PCR and culture. Results will show the efficacy of the treatment regimens to decrease infection with *S. hyodysenteriae* even to very low levels of infection that can be detected by PCR methods.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i i i) NUMBER OF SEQUENCES: 5

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2332 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 413..1903

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

CGGCCAGTGC CAAGCTTTAC CAGTTGAGGG CGACTATTAT TCTGATAAAA AAATGTTAAG      60
AAGATTAGAC CCTTTTATTA ATTTTGGAAAT ATATGCCGCT CATCATGCAT TTAAGCAGGC      120
TGGTATAGAA CCGAAAACAG GCTTTGATCC TTTAAGAGCC GGTGTGTGTTT TTGGTAGCGG      180
TATTGGCGGT ATGACTACTC TTTTATCTAA CCATCAAGTT TTAAGTAAATG ACGGACCTGG      240
CAGAGTATCA CCTTTCTTTG TACCTATGCA AATAATCAAT ATGACACCTG GTTTAATATC      300
TATGGAATAT GGTATGAACG GACCTAACTA CAGTACAGTT ACTGCATGTG CTCTTTCAAA      360
CCTCTCTATA GGTTTAGGTT ATAAACATAT TAAAGATAAT GAAGCTGATA TT ATG      415
                               Met
                               1

GTA GTT GGA GGT TCT GAA GCT ACT ATA AAT CCT CTT ACT ATA GCT GGT      463
Val Val Gly Gly Ser Glu Ala Thr Ile Asn Pro Leu Thr Ile Ala Gly
                               5
                               10
                               15

TTC AAT AAT GCT AGA GCT TTA TCT ACT AGA AAT GAT GAT CCT GCT AAA      511
Phe Asn Asn Ala Arg Ala Leu Ser Thr Arg Asn Asp Asp Pro Ala Lys
                               20
                               25
                               30

GCA TCA AGA CCT TTT GAT AAA GGA AGA GAC GGA CTT GCT ATA GCC AGA      559
Ala Ser Arg Pro Phe Asp Lys Gly Arg Asp Gly Leu Ala Ile Ala Arg
                               35
                               40
                               45

TAT TTA ATA AAA AAT GGC TAT GAT GTA AAA ATA TAT ATC ACA GGA AAT      607
Tyr Leu Ile Lys Asn Gly Tyr Asp Val Lys Ile Tyr Ile Thr Gly Asn
                               50
                               55
                               60
                               65

CTT GAC AGA GTT AAT AAA GAT ACC TAC TCT AAC TTT AAT ATA TTA AAA      655
Leu Asp Arg Val Asn Lys Asp Thr Tyr Ser Asn Phe Asn Ile Leu Lys
                               70
                               75
                               80

TCT ATG AAT ATA GAT ATT AAT TAT TTA GGA AGC GAA GAA GAT GCC ATA      703
Ser Met Asn Ile Asp Ile Asn Tyr Leu Gly Ser Glu Glu Asp Ala Ile
                               85
                               90
                               95

TCA GCT GCA GAA AAT ATA GAA AGA AAA TCA ATA GTA TTA GAT TCA TTA      751
Ser Ala Ala Glu Asn Ile Glu Arg Lys Ser Ile Val Leu Asp Ser Leu
                               100
                               105
                               110

TTT GGT ACA GGC GGA AAC AGA CCT TTA GAA GGA ATA CAA AAA GCT CTT      799
Phe Gly Thr Gly Gly Asn Arg Pro Leu Glu Gly Ile Gln Lys Ala Leu
                               115
                               120
                               125

ATA GAT AGT TTG AAT AAA TTA GAT GTT CTT AGA ATA GCA ATA GAT ATA      847
Ile Asp Ser Leu Asn Lys Leu Asp Val Leu Arg Ile Ala Ile Asp Ile
                               130
                               135
                               140
                               145

CCT TCA GGA TTA GCT TCA AAA ATA AAT GAT AAT GAC AAT GTA TAT ACT      895
Pro Ser Gly Leu Ala Ser Lys Ile Asn Asp Asn Asp Asn Val Tyr Thr
                               150
                               155
                               160

```

-continued

TGT Cys	TTT Phe	AAA Lys	GCA Ala 165	CAT His	GAA Glu	ACA Thr	TAT Tyr	ACT Thr 170	ATA Ile	TGC Cys	TTC Phe	GCT Ala	AAA Lys 175	GAT Asp	ATA Ile	943
TTC Phe	TTT Phe	TTA Leu 180	TAC Tyr	AGA Arg	ACA Thr	AGA Arg	GAA Glu 185	TAT Tyr	ATA Ile	GGA Gly	AAA Lys	TTA Leu 190	TTC Phe	ATA Ile	ATA Ile	991
AAA Lys	TCA Ser 195	ATA Ile	TTT Phe	CCA Pro	GAT Asp	GAA Glu 200	ATA Ile	TTA Leu	GAT Asp	AAT Asn	TGG Trp 205	GGA Gly	TAT Tyr	AAA Lys	GCT Ala	1039
AAA Lys 210	CTT Leu	ATA Ile	GAT Asp	TAT Tyr	AAT Asn 215	GAA Glu	AAA Lys	ATA Ile	AAT Asn	ATA Ile 220	AAT Asn	AGA Arg	AAC Asn	TCT Ser	CTA Leu 225	1087
TAC Tyr	AGC Ser	AAA Lys	AGA Arg	GAA Glu 230	CAA Gln	GGA Gly	ATG Met	CTT Leu	GCT Ala 235	ATA Ile	GTA Val	GCA Ala	GGA Gly	AGT Ser	GAT Asp 240	1135
AAT Asn	TAT Tyr	ATA Ile	GGG Gly 245	GCT Ala	GCT Ala	GTT Val	CTA Leu	GCT Ala 250	GTA Val	AAT Asn	GCT Ala	GCT Ala	TAT Tyr 255	AGA Arg	TTG Leu	1183
GGT Gly	GTA Val	GGA Gly 260	TAC Tyr	ATA Ile	AGA Arg	TTA Leu	TAT Tyr 265	GTA Val	CCT Pro	AAA Lys	GGC Gly	ATA Ile 270	ATA Ile	AAA Lys	AAT Asn	1231
ATA Ile	AGA Arg 275	GAT Asp	GCC Ala	ATA Ile	ATG Met	CCT Pro 280	TCT Ser	ATG Met	CCT Pro	GAA Glu	ATT Ile 285	GTT Val	ATT Ile	ATA Ile	GGA Gly	1279
GTT Val 290	GGA Gly	GAA Glu	GAA Glu	AAT Asn	CAA Gln 295	AAA Lys	TTC Phe	TTC Phe	ACA Thr	GAA Glu 300	AAT Asn	GAC Asp	ATT Ile	GAA Glu	ATA Ile 305	1327
GTA Val	AAT Asn	GAT Asp	ATT Ile	AAT Asn 310	AAA Lys	AGC Ser	GAT Asp	GCT Ala	TGT Cys 315	ATA Ile	ATA Ile	GGT Gly	TCT Ser	GGT Gly 320	ATA Ile	1375
GGC Gly	AGA Arg	GAT Asp	TTG Leu 325	TCT Ser	ACA Thr	GAA Glu	ATT Ile	TTT Phe 330	GTA Val	AAT Asn	ACT Thr	ATA Ile	TTA Leu 335	AAG Lys	CAA Gln	1423
ATA Ile	AAT Asn	ATA Ile 340	CCT Pro	ACT Thr	GTT Val	ATT Ile	GAT Asp 345	GCT Ala	GAT Asp	GCT Ala	TTA Leu	TAT Tyr 350	TTA Leu	ATG Met	TTT Phe	1471
GAA Glu	AGC Ser 355	ACT Thr	CTT Leu	AAT Asn	GAA Glu	CTT Leu 360	AAT Asn	AAT Asn	AAT Asn	TTT Phe	ATA Ile 365	ATC Ile	ACT Thr	CCT Pro	CAT His	1519
ATA Ile 370	TAT Tyr	GAA Glu	TTT Phe	GAA Glu	AAA Lys 375	CTT Leu	ACA Thr	CAG Gln	ATA Ile	AAT Asn 380	CAT His	ATA Ile	GAG Glu	GTT Val	TTA Leu 385	1567
GAA Glu	AAT Asn	CCT Pro	TAT Tyr	CAG Gln 390	GCA Ala	TTA Leu	TTA Leu	ATA Ile	TAC Tyr 395	AGA Arg	GAA Glu	AAA Lys	ACT Thr	AAT Asn 400	GCC Ala	1615
TCA Ser	ATA Ile	GTA Val	TTA Leu 405	AAA Lys	GAT Asp	GCT Ala	GTA Val	AGT Ser 410	TTC Phe	CTA Leu	ATG Met	CAT His	GAA Glu 415	AAT Asn	GAT Asp	1663
ATA Ile	TAT Tyr	ATA Ile 420	AAT Asn	TAT Tyr	AAC Asn	CCT Pro	AGA Arg 425	GAA Glu	TCT Ser	ATG Met	GGG Gly	AAA Lys 430	GCA Ala	GGT Gly	ATG Met	1711
GGT Gly	GAT Asp 435	GTT Val	TTT Phe	GCT Ala	GGA Gly	TTT Phe 440	ATA Ile	GGT Gly	GCT Ala	TTG Leu	CTC Leu 445	GCT Ala	AGA Arg	AAA Lys	CTA Leu	1759
AAT Asn 450	ATA Ile	TTA Leu	GAT Asp	GCT Ala	TCA Ser 455	AAA Lys	CTA Leu	GCA Ala	TTG Leu 460	ATA Ile	ATA Ile	CAG Gln	GCT Ala	AAA Lys	TCT Ser 465	1807
TTT Phe	AAT Asn	ATA Ile	TTA Leu	TCA Ser 470	AAA Lys	AAA Lys	TTC Phe	GGA Gly	AAT Asn 475	GAT Asp	TAT Tyr	ATT Ile	CAG Gln	CCT Pro	AAA Lys 480	1855

-continued

GAT	TTG	GCA	AAT	ATT	TCA	TAT	AAA	ATA	CTA	AAA	GGA	TAT	AAA	TTT	GCC	1903
Asp	Leu	Ala	Asn	Ile	Ser	Tyr	Lys	Ile	Leu	Lys	Gly	Tyr	Lys	Phe	Ala	
			485					490					495			
TAGAGAAGTT	TACGACCCTA	AACAAAAAGA	ATTAGAATTC	TACGCTAAAA	GAGAGGTAAA	1963										
GCCCCCTGCT	CCTAAAAGAG	AGGTAAGCAT	ATTTGCTAGA	AGATGGTTTA	TGTTTTTATA	2023										
CGGAACTTTC	CTCACATTAG	TTGTAATTGG	TATGCTTTTA	TATAAAAAAG	GATTCTTTAA	2083										
TAATATACCA	TTATTTGAAG	CTTTAAAGCC	TAAAACAGAT	GTTATAGTAA	AAATTAATAA	2143										
TGCTGAATTC	GTAAATGATG	CAGTAATTAC	AACTATAGAA	CTCGAAAATT	CAAATTATAC	2203										
TAATTCTGAA	AGTATAGAAA	CACTAAGAAG	TTATTTTTCA	TTGTACAAAA	ATAGAAAATT	2263										
AATATTTACA	GGCAATCGTT	CTTTAATAA	TATAAGATTC	CCAGTAGGTC	AGAGAATAGG	2323										
ATTC AATTT						2332										

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 497 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Val	Val	Gly	Gly	Ser	Glu	Ala	Thr	Ile	Asn	Pro	Leu	Thr	Ile	Ala	
1				5					10					15		
Gly	Phe	Asn	Asn	Ala	Arg	Ala	Leu	Ser	Thr	Arg	Asn	Asp	Asp	Pro	Ala	
			20					25					30			
Lys	Ala	Ser	Arg	Pro	Phe	Asp	Lys	Gly	Arg	Asp	Gly	Leu	Ala	Ile	Ala	
			35				40					45				
Arg	Tyr	Leu	Ile	Lys	Asn	Gly	Tyr	Asp	Val	Lys	Ile	Tyr	Ile	Thr	Gly	
			50			55					60					
Asn	Leu	Asp	Arg	Val	Asn	Lys	Asp	Thr	Tyr	Ser	Asn	Phe	Asn	Ile	Leu	
					70					75					80	
Lys	Ser	Met	Asn	Ile	Asp	Ile	Asn	Tyr	Leu	Gly	Ser	Glu	Glu	Asp	Ala	
				85					90					95		
Ile	Ser	Ala	Ala	Glu	Asn	Ile	Glu	Arg	Lys	Ser	Ile	Val	Leu	Asp	Ser	
			100					105					110			
Leu	Phe	Gly	Thr	Gly	Gly	Asn	Arg	Pro	Leu	Glu	Gly	Ile	Gln	Lys	Ala	
		115					120					125				
Leu	Ile	Asp	Ser	Leu	Asn	Lys	Leu	Asp	Val	Leu	Arg	Ile	Ala	Ile	Asp	
		130				135					140					
Ile	Pro	Ser	Gly	Leu	Ala	Ser	Lys	Ile	Asn	Asp	Asn	Asp	Asn	Val	Tyr	
					150					155					160	
Thr	Cys	Phe	Lys	Ala	His	Glu	Thr	Tyr	Thr	Ile	Cys	Phe	Ala	Lys	Asp	
				165					170					175		
Ile	Phe	Phe	Leu	Tyr	Arg	Thr	Arg	Glu	Tyr	Ile	Gly	Lys	Leu	Phe	Ile	
			180					185					190			
Ile	Lys	Ser	Ile	Phe	Pro	Asp	Glu	Ile	Leu	Asp	Asn	Trp	Gly	Tyr	Lys	
		195					200					205				
Ala	Lys	Leu	Ile	Asp	Tyr	Asn	Glu	Lys	Ile	Asn	Ile	Asn	Arg	Asn	Ser	
		210				215					220					
Leu	Tyr	Ser	Lys	Arg	Glu	Gln	Gly	Met	Leu	Ala	Ile	Val	Ala	Gly	Ser	
		225			230					235					240	
Asp	Asn	Tyr	Ile	Gly	Ala	Ala	Val	Leu	Ala	Val	Asn	Ala	Ala	Tyr	Arg	
				245					250					255		

-continued

Leu	Gly	Val	Gly 260	Tyr	Ile	Arg	Leu	Tyr 265	Val	Pro	Lys	Gly	Ile 270	Ile	Lys
Asn	Ile	Arg 275	Asp	Ala	Ile	Met	Pro 280	Ser	Met	Pro	Glu	Ile 285	Val	Ile	Ile
Gly	Val 290	Gly	Glu	Glu	Asn	Gln 295	Lys	Phe	Phe	Thr	Glu 300	Asn	Asp	Ile	Glu
Ile 305	Val	Asn	Asp	Ile	Asn 310	Lys	Ser	Asp	Ala	Cys 315	Ile	Ile	Gly	Ser	Gly 320
Ile	Gly	Arg	Asp	Leu 325	Ser	Thr	Glu	Ile	Phe 330	Val	Asn	Thr	Ile	Leu 335	Lys
Gln	Ile	Asn	Ile 340	Pro	Thr	Val	Ile	Asp 345	Ala	Asp	Ala	Leu	Tyr 350	Leu	Met
Phe	Glu	Ser 355	Thr	Leu	Asn	Glu	Leu 360	Asn	Asn	Asn	Phe	Ile 365	Ile	Thr	Pro
His	Ile 370	Tyr	Glu	Phe	Glu	Lys 375	Leu	Thr	Gln	Ile	Asn 380	His	Ile	Glu	Val
Leu 385	Glu	Asn	Pro	Tyr	Gln 390	Ala	Leu	Leu	Ile	Tyr 395	Arg	Glu	Lys	Thr	Asn 400
Ala	Ser	Ile	Val	Leu 405	Lys	Asp	Ala	Val	Ser 410	Phe	Leu	Met	His	Glu 415	Asn
Asp	Ile	Tyr	Ile 420	Asn	Tyr	Asn	Pro	Arg 425	Glu	Ser	Met	Gly	Lys 430	Ala	Gly
Met	Gly	Asp 435	Val	Phe	Ala	Gly	Phe 440	Ile	Gly	Ala	Leu	Leu 445	Ala	Arg	Lys
Leu	Asn 450	Ile	Leu	Asp	Ala	Ser 455	Lys	Leu	Ala	Leu	Ile 460	Ile	Gln	Ala	Lys
Ser 465	Phe	Asn	Ile	Leu	Ser 470	Lys	Lys	Phe	Gly	Asn 475	Asp	Tyr	Ile	Gln	Pro 480
Lys	Asp	Leu	Ala	Asn 485	Ile	Ser	Tyr	Lys	Ile 490	Leu	Lys	Gly	Tyr	Lys 495	Phe

Ala

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGTACAGGCG GAAACAGACC TT

22

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TCCTATTCTC TGACCTACTG

20

(2) INFORMATION FOR SEQ ID NO:5:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TAGGGGCTGC TGTTCTAGCT GTAAATGC

2 8

What is claimed is:

1. An isolated HindIII DNA fragment of about 2.3 kb derived from chromosomal DNA of *S. hyodysenteriae* B204 comprising the nucleotide sequence of SEQ ID NO:1.
2. An oligonucleotide probe that specifically hybridizes under stringent conditions to a target DNA sequence of *S. hyodysenteriae*, wherein the target sequence is about a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 which target DNA sequence comprises SEQ ID NO:1 and wherein the probe does not hybridize to at least one other microorganism.
3. The oligonucleotide probe according to claim 2, labeled with a detectable moiety selected from the group consisting of a biotin labeled nucleotide, radiolabeled nucleotide and a fluorescent tagged nucleotide.
4. The oligonucleotide probe according to claim 2, wherein the probe is hybridizable to DNA amplification products of primers that specifically hybridize under stringent conditions to the about 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 which restriction fragment comprises SEQ ID NO:1 and which primers do not hybridize to at least one other microorganism.
5. The oligonucleotide probe according to claim 2, wherein the probe is about 20 to 2300 nucleotides long.
6. A recombinant DNA expression vector comprising a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 which restriction fragment comprises SEQ ID NO:1 operably linked to transcriptional and translational control regions of the expression vector.
7. An expression vector according to claim 6 having the characteristics of a plasmid pRED3C6, ATCC No. 75826.
8. A transformed cell carrying a recombinant expression vector of claim 6.
9. An oligonucleotide primer for amplifying a target sequence of *S. hyodysenteriae*, wherein the primer specifically hybridizes under stringent conditions to a target sequence, wherein the target sequence is a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 which restriction fragment comprises SEQ ID NO: 1 and which primer does not hybridize to at least one other microorganism.
10. The oligonucleotide primer according to claim 9, wherein the oligonucleotide has a DNA sequence comprising positive sense
 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3), or negative sense
 5'-TCCTATTCTCTGACCTACTG (SEQ ID NO:4), or complements thereof.
11. A kit for use in the detection of *S. hyodysenteriae* in a biological sample, comprising:
 - (a) a first reagent comprising at least one oligonucleotide primer that specifically hybridizes under stringent conditions to a target sequence of a 2.3 kb HindIII partial digest restriction fragment of *S. hyodysenteriae* B204 which restriction fragment comprises SEQ ID NO: 1 and which primers do not hybridize to at least one other microorganism.
 12. A kit according to claim 11, further comprising in combination: a second reagent comprising an oligonucleotide probe which specifically hybridizes under stringent conditions to either a 2.3 kb HindIII restriction fragment comprising SEQ ID NO: 1 or to an amplified product produced using a primer of claim 9 and which probe does not hybridize to at least one other microorganism; and the reagents packaged within containing means.
 13. The kit according to claim 11, wherein the containing means are selected from the group consisting of a vial, jar, tubes and multiwell plate.
 14. A kit according to claim 11, further comprising reagents for extracting nucleic acids from a biological sample.
 15. The kit according to claim 11, wherein the oligonucleotide primer has a DNA sequence comprising positive sense
 5'-GGTACAGGCGGAAACAGACCTT (SEQ ID NO:3), or negative sense
 5'-TCCTATTCTCTGACCTACTG (SEQ ID NO:4), complements thereof, or mixtures thereof.
 16. A kit according to claim 11, further comprising reagents for conducting the polymerase chain reaction.
 17. An isolated nucleic acid consisting of SEQ ID NO: 1.

* * * * *