

University of Groningen

## End-stage renal disease in Slovak children

Kolvek, G.; Reijneveld, S.A.; Podracka, L.; Rosenberger, J.; Nagyova, I.; Stewart, R.E.; van Dijk, J.P.

*Published in:*  
European Journal of Pediatrics

*DOI:*  
[10.1007/s00431-011-1462-1](https://doi.org/10.1007/s00431-011-1462-1)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2011

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Kolvek, G., Reijneveld, S. A., Podracka, L., Rosenberger, J., Nagyova, I., Stewart, R. E., & van Dijk, J. P. (2011). End-stage renal disease in Slovak children: epidemiology from a European perspective. *European Journal of Pediatrics*, 170(11), 1445-1451. <https://doi.org/10.1007/s00431-011-1462-1>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# End-stage renal disease in Slovak children: epidemiology from a European perspective

Gabriel Kolvek · Sijmen A. Reijneveld ·  
Ludmila Podracka · Jaroslav Rosenberger ·  
Iveta Nagyova · Roy E. Stewart · Jitse P. van Dijk

Received: 21 February 2011 / Accepted: 22 March 2011 / Published online: 8 April 2011  
© Springer-Verlag 2011

**Abstract** The aim of this study was to examine the occurrence of end-stage renal disease (ESRD) in Slovak children, to compare it with earlier Slovak data and with data from other European countries, and to explore etiology. Over the years 2003–2009, data on the incidence and prevalence of all cases of ESRD in children from all four Slovak tertiary pediatric centers were collected. The data were compared with two earlier Slovak studies and with European data from the European Society of Paediatric Nephrology. The median annual incidence rate of ESRD in Slovak children under 15 years of age was 6.6 per million age-related population (pmarp). The prevalence rate on 31 December 2009 was 24.1 pmarp. Compared with the last study (18.6 pmarp), the differences were not statistically significant. The comparison with neighboring countries and with the European average shows no significant difference in incidence, while prevalence is significantly lower compared to neighboring Austria

and some other (mostly western) European countries as well as the European average. In conclusion, during the past decade, the incidence and prevalence rates of ESRD in Slovak children have remained stable. Compared to the European average, the prevalence in Slovak children is significantly lower.

**Keywords** End-stage renal disease · Incidence · Prevalence · Etiology · Renal replacement therapy

## Introduction

The pediatric population suffering from end-stage renal disease (ESRD) in Europe is growing. Children with formerly lethal diseases nowadays survive until adulthood and beyond [11]. The reported number of children on renal

---

A list of the participating centers is provided in the Acknowledgements.

---

G. Kolvek (✉) · L. Podracka  
1st Paediatric Department, Faculty of Medicine,  
P.J. Safarik University,  
Tr. SNP 1,  
040 11 Kosice, Slovak Republic  
e-mail: gabriel.kolvek@upjs.sk

G. Kolvek · L. Podracka · J. Rosenberger · I. Nagyova ·  
J. P. van Dijk  
Graduate School Kosice Institute for Society and Health, P.J.  
Safarik University,  
Kosice, Slovak Republic

S. A. Reijneveld · R. E. Stewart · J. P. van Dijk  
Department of Social Medicine, University Medical Center  
Groningen, University of Groningen,  
Groningen, The Netherlands

J. Rosenberger  
Nephrology and Dialysis Center Fresenius,  
Kosice, Slovak Republic

J. Rosenberger  
Transplantation Department, Faculty Hospital L. Pasteur,  
Kosice, Slovak Republic

J. Rosenberger  
1st Internal Department, Faculty of Medicine, P.J. Safarik  
University,  
Kosice, Slovak Republic

I. Nagyova  
Institute of Public Health–Department of Social Medicine,  
Faculty of Medicine, P.J. Safarik University,  
Kosice, Slovak Republic

replacement therapy (RRT) has almost tripled during a relatively short period of 15 years [8, 13, 14].

Nevertheless, reliable epidemiological information on ESRD and RRT in European children is not easy to obtain. One source of information is the annual reports of the European Renal Association–European Dialysis Transplantation Association (ERA-EDTA), but these reports are somewhat limited since data are often lacking, especially from Central and Eastern European countries. This is probably a consequence of the absence of official national or regional registries in most of these countries, including Slovakia [15]. In 2007, the European Society of Paediatric Nephrology (ESPN) began collecting data on pediatric renal replacement therapy (RRT) patients exclusively on the basis of national or large regional registries in an attempt to solve the problem of insufficient information.

In Slovakia, only two pediatric epidemiologic studies on this topic have been published so far. The first such study was by Zvara et al. in the years 1975–1977 and included children under the age of 15 years with chronic kidney disease (CKD) [17]. As a result of unavailability of dialysis and transplantation, the prevalence of RRT at that time was only 1.5 per million age-related population (pmarp) under 15 years of age (0.4 per million total population—pmp). The next study on Slovak children was done in the year 2002 by Podracka et al., but this study was done on CKD children in a different age category (under 18 years of age) revealing a prevalence rate of 29.3 pmarp for this age group (7.4 children pmp) [10].

Up to now, a national register with the potential of monitoring health care of pediatric end-stage renal disease patients has not been established in Slovakia. Therefore, the main aims of the present study are:

1. To evaluate the actual incidence and prevalence rates of ESRD and RRT in Slovak children;
2. To compare the results regarding 2003–2009 with those from earlier epidemiological surveys carried out in Slovakia;
3. To compare the incidence and prevalence rates of ESRD and RRT in Slovak children with those of children from other European countries;
4. To compare the etiology, i.e., primary renal diseases and treatment modes, with earlier studies on Slovak children and on children from other European countries.

## Materials and methods

### Patients

According to a new ERA-EDTA policy, this epidemiologic study evaluated data from all four tertiary centers for

pediatric nephrology in Slovakia (Bratislava, Košice, Banská Bystrica, and Martin) treating all pediatric RRT patients [2]. The total sample consisted of 83 patients below the age of 19 years (32 girls, 51 boys) treated for ESRD in Slovakia in the period 1 January 2003 to 31 December 2009. Population data were obtained from the Statistical Office of the Slovak Republic.

### Procedure and measures

Data were collected using questionnaires sent to the heads of all four pediatric dialysis and transplantation centers in Slovakia. For each patient, *age*, *gender*, and *clinical variables* were retrieved. Clinical variables were derived from the medical files by one of the authors (GK) regarding the primary renal disease, the date and mode of the first RRT (peritoneal dialysis (PD), hemodialysis (HD), or preemptive transplantation), and changes in treatment including date and type of each transplantation (preemptive or non-preemptive, from a living or deceased donor). The primary renal diseases were recorded as diagnosed by all four centers in Slovakia and divided into the categories according to ERA-EDTA [4].

For comparison of incidence and prevalence rates of RRT with other European countries on the basis of ESPN datasets, the upper age limit was set to 14 years. Patients were divided into 5-year categories (0–4, 5–9, 10–14). For complete analysis of the Slovak situation, we added an extra category of 15–18-year-old children, as these patients are considered pediatric (under 19 years old) in Slovakia and other countries.

Incidence and prevalence rates were calculated as pmarp and pmp in the different age categories (0–4, 5–9, 10–14, 15–18). Prevalence rates were expressed as the number of ESRD patients alive and on RRT on 31st December of each year. Data on RRT from other European countries were derived from ESPN/ERA-EDTA annual reports [5, 6].

### Statistical analyses

Descriptive data regarding incidence and prevalence rates of RRT from 2003 to 2009 were presented. Poisson regression was used to find significant differences between the last study from 2002 and actual data from 2009. Differences between Slovakia and neighboring countries were compared over the years 2007–2008 using test of proportions. The differences were considered statistically significant at a *p* level of 0.05. SAS 9.2 was used for the statistical analyses.

## Results

Between 1 January 2003 and 31 December 2009, 52 new patients reached ESRD before the age of 19 years and

initiated RRT in Slovakia. The median age of the patients at the start of RRT was 12 years, and the range was 0–18 years. The male/female ratio of the patients was 1.58.

#### Age distribution and incidence and prevalence rates

The largest portion of the patients ( $n=20$ ) was aged 10 to 14 years when they initiated RRT; 18 patients were 15 to 18 years old; and six were younger than age of 5 years at the time of RRT initiation (Table 1). The median annual incidence rate of RRT (during the study period 2003–2009) was 6.6 pmarp under age 15 years (1.5 pmtp). The current prevalence rate of RRT in Slovak children under 15 years old (measured on 31 December 2009) was found to be 24.1 pmarp (3.7 pmtp). Table 1 shows the incidence and prevalence rates in detail during the whole study period. The average mortality of patients under the age of 15 years on RRT during the whole period was 1.3%.

#### Comparison with earlier Slovak epidemiological surveys

In the first Slovak study, done in 1977 by Zvara et al., only the prevalence rate under the age of 15 years was explicitly

described (1.5 pmarp, 0.4 pmtp) [17]. The study performed in 2002 presented data regarding children under the age of 18 years [10]. To allow for a direct comparison, we recalculated the rates for children under the age of 15 years. The incidence rate was then 4.1 pmarp under 15 years (0.7 pmtp), and the prevalence rate was 18.6 pmarp under 15 years (3.2 pmtp). A slightly positive trend both for incidence and prevalence is currently present, although the differences between 2002 and 2009 are not statistically significant.

#### Slovakia compared to other European countries

The comparison of incidence and prevalence of RRT among Slovak children with children from other European countries using data published by ESPN in 2009–2010 for the years 2007–2008 is presented in Table 2. The comparison with Slovakia's neighboring countries and with the European average shows no significant difference in incidence. The Slovak prevalence does not differ than that of most of its neighbors, but it is significantly lower compared with neighboring Austria. The same holds true in the comparison of Slovak prevalence with some other

**Table 1** Incidence and prevalence rates of RRT in Slovak children according to age and gender over the years 2003–2009

|                   | Total                    | Gender                    |                          | Age groups              |                         |                           |                           |
|-------------------|--------------------------|---------------------------|--------------------------|-------------------------|-------------------------|---------------------------|---------------------------|
|                   | 0–18<br><i>n</i> [pmarp] | Girls<br><i>n</i> [pmarp] | Boys<br><i>n</i> [pmarp] | 0–4<br><i>n</i> [pmarp] | 5–9<br><i>n</i> [pmarp] | 10–14<br><i>n</i> [pmarp] | 15–18<br><i>n</i> [pmarp] |
| <b>Incidence</b>  |                          |                           |                          |                         |                         |                           |                           |
| 2002              | n.a.                     | n.a.                      | n.a.                     | 0 [0.0]                 | 1 [3.2]                 | 3 [7.7]                   | n.a.                      |
| 2003              | 6 [4.6]                  | 2 [3.2]                   | 4 [6.1]                  | 0 [0.0]                 | 1 [3.3]                 | 2 [5.2]                   | 3 [8.9]                   |
| 2004              | 9 [7.2]                  | 4 [6.6]                   | 5 [8.2]                  | 2 [7.7]                 | 1 [3.4]                 | 4 [10.9]                  | 2 [6.1]                   |
| 2005              | 8 [6.6]                  | 2 [3.7]                   | 6 [9.6]                  | 2 [7.7]                 | 1 [3.5]                 | 5 [14.3]                  | 0 [0.0]                   |
| 2006              | 6 [5.1]                  | 5 [9.3]                   | 1 [1.6]                  | 0 [0.0]                 | 2 [7.8]                 | 2 [6.0]                   | 2 [6.3]                   |
| 2007              | 9 [8.3]                  | 4 [7.6]                   | 5 [9.0]                  | 1 [3.8]                 | 0 [0.0]                 | 3 [9.5]                   | 5 [16.2]                  |
| 2008              | 4 [3.8]                  | 3 [5.8]                   | 1 [1.8]                  | 1 [3.7]                 | 1 [3.8]                 | 1 [3.3]                   | 1 [3.3]                   |
| 2009              | 10 [8.9]                 | 1 [1.8]                   | 9 [15.7]                 | 0 [0.0]                 | 2 [7.7]                 | 3 [10.3]                  | 5 [17.3]                  |
| Median 2003–2009  | 8 [6.6]                  | 3 [5.8]                   | 5 [8.2]                  | 1 [3.7]                 | 1 [3.5]                 | 3 [9.5]                   | 2 [6.3]                   |
| <b>Prevalence</b> |                          |                           |                          |                         |                         |                           |                           |
| 2002              | n.a.                     | n.a.                      | n.a.                     | 1 [3.8]                 | 6 [19.1]                | 11 [28.4]                 | n.a.                      |
| 2003              | 35 [27.3]                | 13 [20.7]                 | 22 [35.1]                | 1 [3.8]                 | 3 [10.0]                | 9 [23.6]                  | 22 [64.9]                 |
| 2004              | 41 [32.8]                | 14 [23.0]                 | 27 [44.3]                | 1 [3.9]                 | 6 [20.6]                | 9 [24.5]                  | 25 [75.9]                 |
| 2005              | 43 [35.3]                | 14 [23.6]                 | 29 [48.8]                | 2 [7.7]                 | 6 [21.1]                | 12 [34.3]                 | 23 [71.4]                 |
| 2006              | 42 [35.4]                | 18 [31.1]                 | 24 [41.4]                | 1 [3.8]                 | 8 [29.1]                | 14 [42.1]                 | 19 [59.9]                 |
| 2007              | 44 [37.9]                | 18 [31.8]                 | 26 [46.0]                | 2 [7.5]                 | 6 [22.5]                | 15 [47.3]                 | 21 [68.0]                 |
| 2008              | 39 [34.3]                | 20 [36.0]                 | 19 [34.2]                | 3 [11.0]                | 5 [19.1]                | 12 [39.8]                 | 19 [62.9]                 |
| 2009              | 40 [35.7]                | 18 [32.9]                 | 22 [40.3]                | 2 [7.1]                 | 4 [15.4]                | 14 [48.1]                 | 20 [69.1]                 |
| Median 2003–2009  | 41 [35.3]                | 18 [31.1]                 | 24 [41.4]                | 2 [7.1]                 | 6 [20.6]                | 12 [39.8]                 | 21 [68.0]                 |

*n.a.* not available

**Table 2** Incidence and prevalence rates of RRT in European children under 15 years from 2007–2008 with Slovak data derived from Table 1

|                              | Incidence rates |           |                     |                       | Prevalence rates |             |                     |                       | Significance |
|------------------------------|-----------------|-----------|---------------------|-----------------------|------------------|-------------|---------------------|-----------------------|--------------|
|                              | 2007            | 2008      | Average (2007–2008) | Contrast vs. Slovakia | 2007             | 2008        | Average (2007–2008) | Contrast vs. Slovakia |              |
| Slovakia                     | 4.6 (4.7)       | 3.6 (3.6) | 4.1 (4.2)           | 1                     | 26.7 (27.0)      | 26.1 (23.9) | 26.4 (25.5)         | 1                     | –            |
| Neighboring countries        |                 |           |                     |                       |                  |             |                     |                       |              |
| Austria                      | 7.7             | 4.4       | 6.1                 | 1.49                  | 40.1             | 39.5        | 39.8                | 1.51                  | 0.02         |
| Czech republic               | 6.1             | 3.4       | 4.8                 | 1.17                  | 22.3             | 21.0        | 21.7                | 0.82                  | n.s.         |
| Hungary                      | 2.3             | 2.0       | 2.2                 | 0.54                  | 28.3             | 20.7        | 24.5                | 0.93                  | n.s.         |
| Poland                       | 6.9             | 6.6       | 6.8                 | 1.66                  | 34.4             | 34.8        | 34.6                | 1.31                  | n.s.         |
| Slavic countries             |                 |           |                     |                       |                  |             |                     |                       |              |
| Belarus                      | n.a.            | 5.6       | n.c.                | n.c.                  | n.a.             | 18.3        | n.c.                | n.c.                  | n.c.         |
| Croatia                      | 2.9             | 8.8       | 5.9                 | 1.44                  | 27.7             | 29.4        | 28.6                | 1.08                  | n.s.         |
| FYR of Macedonia             | 7.9             | 0.0       | 4.0                 | 0.98                  | 18.3             | 16.0        | 17.2                | 0.65                  | n.s.         |
| Montenegro                   | 0.0             | 24.5      | 12.3                | 3.00                  | 8.1              | 32.7        | 20.4                | 0.77                  | n.s.         |
| Serbia                       | 4.4             | 11.5      | 8.0                 | 1.95                  | 24.5             | 33.7        | 29.1                | 1.10                  | n.s.         |
| Russia                       | 7.8             | 3.4       | 5.6                 | 1.37                  | 16.1             | 9.8         | 13.0                | 0.49                  | <0.0001      |
| Slovenia                     | 7.1             | 14.2      | 10.7                | 2.61                  | 21.3             | 35.4        | 28.4                | 1.08                  | n.s.         |
| Ukraine                      | n.a.            | 1.7       | n.c.                | n.c.                  | n.a.             | 2.3         | n.c.                | n.c.                  | n.c.         |
| Remaining European countries |                 |           |                     |                       |                  |             |                     |                       |              |
| Belgium                      | n.a.            | 3.3       | n.c.                | n.c.                  | n.a.             | 57.2        | n.c.                | n.c.                  | n.c.         |
| Denmark                      | 4.9             | 6.8       | 5.9                 | 1.44                  | 37.0             | 40.1        | 38.6                | 1.46                  | 0.04         |
| Estonia                      | 0.0             | 5.0       | 2.5                 | 0.61                  | 10.0             | 15.0        | 12.5                | 0.47                  | n.s.         |
| Finland                      | 12.3            | 3.4       | 7.9                 | 1.93                  | 92.4             | 79.5        | 86.0                | 3.26                  | <0.0001      |
| France                       | 6.5             | 5.4       | 6.0                 | 1.46                  | 34.2             | 30.4        | 32.3                | 1.22                  | n.s.         |
| Greece                       | 8.1             | 7.5       | 7.8                 | 1.90                  | 28.8             | 30.5        | 29.7                | 1.13                  | n.s.         |
| Iceland                      | 15.2            | 0.0       | 7.6                 | 1.85                  | 30.5             | 30.2        | 30.4                | 1.15                  | n.s.         |
| Italy                        | 4.2             | 4.3       | 4.3                 | 1.05                  | 29.7             | 31.6        | 30.7                | 1.16                  | n.s.         |
| Latvia                       | 3.2             | 0.0       | 1.6                 | 0.39                  | 3.2              | 3.2         | 3.2                 | 0.12                  | 0.003        |
| Lithuania                    | 5.7             | 2.0       | 3.9                 | 0.95                  | 32.2             | 33.3        | 32.8                | 1.24                  | n.s.         |
| Norway                       | 3.3             | 6.6       | 5.0                 | 1.22                  | 45.2             | 42.9        | 44.1                | 1.67                  | 0.006        |
| Portugal                     | 9.8             | 8.0       | 8.9                 | 2.17                  | 39.2             | 42.4        | 40.8                | 1.55                  | 0.01         |
| Spain                        | n.a.            | 7.4       | n.c.                | n.c.                  | n.a.             | 39.5        | n.c.                | n.c.                  | n.c.         |
| Sweden                       | 7.1             | 6.5       | 6.8                 | 1.66                  | 48.5             | 44.1        | 46.3                | 1.75                  | 0.001        |
| Switzerland                  | n.a.            | 5.1       | n.c.                | n.c.                  | n.a.             | 44.1        | n.c.                | n.c.                  | n.c.         |
| The Netherlands              | 7.5             | 5.1       | 6.3                 | 1.54                  | 44.4             | 45.7        | 45.1                | 1.71                  | 0.0009       |
| United Kingdom               | 7.7             | 7.6       | 7.7                 | 1.88                  | 51.6             | 51.5        | 51.6                | 1.95                  | <0.0001      |
| Average (ESPN)               | 6.5             | 5.2       | 5.9                 | 1.44                  | 33.6             | 28.6        | 31.1                | 1.19                  | 0.01         |

Source: ESPN and ERA-EDTA datasets 2007, 2008; our recalculations in brackets  
*n.a.* not available, *n.c.* not calculated, *Slovakia* reference category

(mostly western) European countries as well as the European average.

### Etiology

The distribution of primary diseases leading to ESRD in Slovak children between 1 January 2003 and 31 December 2009 is shown in Table 3. Congenital anomalies of the kidneys and urinary tract (CAKUT) with kidney (hypo) dysplasia comprised the major cause. Cystic diseases were the next in importance. The predominance of congenital and hereditary disorders as a leading cause of ESRD compared with acquired nephropathies was evident.

The distribution of primary diseases could not be compared directly with previous studies because the etiology in those studies was only described in regard to all CKD patients but not regarding those in ESRD separately. Regarding all CKD patients, the reported distribution of primary diseases was: CAKUT, 52.4%; hereditary disorders, 14.3%; glomerular diseases, 28.6%; and interstitial nephritides, 4.8% in 1977; and CAKUT, 37.5%, hereditary disorders, 26.5%, glomerular diseases, 11.5%; and interstitial nephritides, 18.5% in 2002.

**Table 3** Etiology of ESRD in new patients (cohort 2003–2009)

| Etiology                  | Total      | Gender        |                |
|---------------------------|------------|---------------|----------------|
|                           |            | 0–18<br>n (%) | Girls<br>n (%) |
| 1. CAKUT                  | 18 (34.6)  | 5 (23.9)      | 13 (42.0)      |
| Renal hypo-/dysplasia     | 8 (15.3)   | 1 (4.8)       | 7 (22.7)       |
| Vesico-ureteric reflux    | 3 (5.8)    | 2 (9.5)       | 1 (3.2)        |
| Obstructive uropathy      | 7 (13.4)   | 2 (9.5)       | 5 (16.1)       |
| Urethral valve            | 0 (0.0)    | 0 (0.0)       | 0 (0.0)        |
| Obstruction without valve | 6 (11.5)   | 1 (4.8)       | 5 (16.1)       |
| Neurogenic bladder        | 1 (1.9)    | 1 (4.8)       | 0 (0.0)        |
| 2. Cystic disease         | 10 (19.2)  | 3 (14.4)      | 7 (22.7)       |
| Polycystic kidney disease | 4 (7.7)    | 1 (4.8)       | 3 (9.7)        |
| Juvenile nephronophthisis | 4 (7.7)    | 2 (9.5)       | 2 (6.5)        |
| 3. Hereditary nephropathy | 5 (9.6)    | 3 (14.4)      | 2 (6.5)        |
| Alport syndrome           | 2 (3.8)    | 2 (9.5)       | 0 (0.0)        |
| 4. Glomerulonephritis     | 8 (15.4)   | 5 (23.9)      | 3 (9.7)        |
| FSGS                      | 5 (9.6)    | 3 (14.3)      | 2 (6.5)        |
| 5. HUS                    | 1 (1.9)    | 1 (4.8)       | 0 (0.0)        |
| 6. Vasculitis             | 2 (3.8)    | 0 (0.0)       | 2 (6.4)        |
| 7. Miscellaneous          | 6 (11.5)   | 3 (14.4)      | 3 (9.7)        |
| 8. Unknown                | 2 (3.8)    | 1 (4.8)       | 1 (3.2)        |
| Total                     | 52 (100.0) | 21 (100.0)    | 31 (100.0)     |

*ESRD* end-stage renal disease, *CAKUT* congenital anomalies of kidneys and urinary tract, *HUS* hemolytic-uremic syndrome, *FSGS* focal segmental glomerulosclerosis

### Treatment mode

During the study period, 27 patients (51.9%) (younger than 19 years old) started RRT on PD and 24 patients (46.2%) started their RRT on HD. Only one patient (1.9%) was transplanted preemptively. Out of all the first transplantations, 22.0% were from living-related donors. In previous Slovak surveys, initial treatment modalities were not described [10, 17]. Compared to published data on the overall proportion of preemptive transplantations in Europe (15.1%), the proportion of preemptive transplantations in Slovak children is significantly lower. Regarding the prevalent treatment modalities measured on 31 December 2008, the proportion of HD patients ( $n=8$ ; 20.5%) does not significantly differ from the ERA-EDTA data ( $n=180$ ; 13.1%), and the proportion of PD patients ( $n=12$ ; 30.8% vs  $n=100$ ; 7.3%) is significantly higher and of transplanted patients ( $n=19$ ; 48.7% vs  $n=1097$ ; 79.7%) significantly lower compared to the ERA-EDTA 2010 annual report.

### Discussion

This study presents an overview of the epidemiology and etiology of ESRD and RRT in Slovak children, focusing on changes observed in the period 2003–2009 compared to historical data and data on the European perspective. The median annual incidence rate of RRT in Slovak children under age 15 years in this period was 6.6 pmarp. These rates were somewhat higher than that in the preceding study on Slovak children in 2002 [17], but without statistical significance. Data on the incidence of RRT regarding the first Slovak study from 1978 are not available, but it seems likely that the number of patients initiating RRT was lower as a consequence of the unavailability of dialysis and transplantation in the past. Compared with neighboring countries and with the European average, no significant differences in incidence rates were found. Explanations for these similarities are the similar demographic structure but also the seemingly comparable availability of dialysis. An increase in the incidence rates of RRT can thus be expected, as the European RRT incidence is currently rising [6, 13, 15].

On the basis of the available prevalence data (1.5 pmarp in 1977, 18.6 pmarp in 2002, 24.1 pmarp in 2009), it could be assumed that the initially steep rise of RRT prevalence rates stabilized a couple of years after introducing dialysis and transplantation into the treatment regime of Slovak children. Later variations could easily be due to fluctuation by chance (18.6 pmarp in 2002, 27.0 pmarp in 2007, 24.1 pmarp in 2009). Regarding neighboring countries, prevalence rates were significantly higher in the neighboring Austria but not in other neighboring countries. Moreover,

the average European prevalence rate was higher. The higher prevalence in Austria and in some other mostly western European countries may be due to the combination of improved medical care for patients in earlier stages of CKD in combination with earlier treatment of ESRD, especially in younger ages where a low weight can also play a significant role for establishing dialysis [13, 15].

The predominance of congenital and hereditary reasons as the main causes of ESRD in children has similarly been found in a study on neighboring Poland [16] and in many other previous studies on Europe [1, 3, 7] and other parts of the world [9, 12]. A study of the British Association of Paediatric Nephrology published in December 2009, the most recent European study, reported a higher proportion of CAKUT (50.0%) and congenital nephrosis (9.3%) [7]. A Dutch study on the period 1987–2001 reported a relatively very high proportion of urethral valves (14.3%) as the reason for ESRD, which contrasts with the Slovak results, in which there was zero contribution of urethral valves among RRT patients. Environmental, ethnic, and other differences might explain the different occurrence of primary renal diseases in Europe compared to our findings [15].

#### Strengths and limitations

An important strength of our study is that we obtained data on the entire country using a rigorous data collection. A limitation is the relatively small number of children, which limits the potential for subgroup analyses.

#### Conclusion

A slightly positive trend both for incidence and prevalence rates of pediatric RRT patients has been present in Slovakia during the past decade, although no significant difference compared to the last study (2002) was found. The comparison with the neighboring countries and with the European average shows no significant differences in incidence. The prevalence does not significantly differ from that of most of its neighbors but is significantly lower compared with neighboring Austria and some other (mostly western) European countries as well as the European average.

A central national registry for children on RRT should be established, not only to facilitate epidemiological analyses and longitudinal studies but also as a first step toward monitoring and improving the quality of health care and quality of life of Slovak children.

**Acknowledgments** The authors would like to thank all participating centers for their invaluable contribution to the data collection: Z. Kizekova MD, Bratislava; M. Dluholucky MD, Banska Bystrica; and

M. Antonyova MD, Martin. This work was supported by the Slovak Research and Development Agency under contract no. APVV-20-038305. This work was also supported by the project “Centre for Excellent Research of Atherosclerosis and its Complications—Myocardial Infarction and Stroke,” Operational Program of Research and Development financed by the European Fund for Regional Development.

**Conflict of interest** The authors declare that they have no conflict of interest.

#### References

1. Ardissino G, Dacco V, Testa S, Bonaudo R, Claris-Appiani A, Taioli E, Marra G, Edefonti A, Sereni F, ItalKid Project (2003) Epidemiology of chronic renal failure in children: data from the ItalKid project. *Pediatrics* 111:e382–e387
2. Briggs JD (2000) The ERA-EDTA Registry returns to Amsterdam. *Nephrol Dial Transplant* 15:1326–1327
3. British Association for Paediatric Nephrology (2005) Report from the paediatric renal registry. In: Ansell D, Feest TG (eds) *The UK Renal Registry 8th Annual Report*, pp 269–291
4. ERA-EDTA Registry (2008) ERA-EDTA Registry 2006 annual report. Academic Medical Center, Department of Medical Informatics, Amsterdam
5. ESPN/ERA-EDTA Registry (2009) ESPN/ERA-EDTA Registry 2007 annual report. Academic Medical Center, Department of Medical Informatics, Amsterdam
6. ESPN/ERA-EDTA Registry (2010) ESPN/ERA-EDTA Registry 2008 annual report. Academic Medical Center, Department of Medical Informatics, Amsterdam
7. Lewis MA, Shaw J, Sinha MD, Adalat S, Hussain F, Castledine C, van Schalkwyk D, Inward C (2010) UK Renal Registry 12th annual report (December 2009): chapter 14: demography of the UK paediatric renal replacement therapy population in 2008. *Nephron Clin Pract* 115(Suppl 1):c279–c288
8. Mehls O, Rigden S, Ehrlich JH, Berthoux F, Jones EH, Valderrabano F (1996) Report on management of renal failure in Europe, XXV, 1994. The child-adult interface. The EDTA-ERA Registry. *European Dialysis and Transplant Association-European Renal Association. Nephrol Dial Transplant* 11(Suppl 1):22–36
9. NAPRTCS (2007) North American Pediatric Renal Trials and Collaborative Studies 2007. <http://www.emmes.com/study/ped/annlrept/annlrept2007.pdf>. Accessed 2 Nov 2011
10. Podracka L, Kovacs L (2004) Epidemiologia chronickej renálnej insuficiencie u deti na Slovensku. (Epidemiology of chronic renal insufficiency in children in Slovakia). *Cesk Pediatr* 7:331
11. Reiss U, Wingen AM, Scharer K (1996) Mortality trends in pediatric patients with chronic renal failure. *Pediatr Nephrol* 10:41–45
12. U.S. Renal Data System (2005) USRDS 2005 Annual Data Report: Atlas of End-stage renal disease in the United States. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda
13. van der Heijden BJ, van Dijk PC, Verrier-Jones K, Jager KJ, Briggs JD (2004) Renal replacement therapy in children: data from 12 registries in Europe. *Pediatr Nephrol* 19:213–221
14. van Stralen KJ, Tizard EJ, Verrina E, Schaefer F, Jager KJ, European Society for Paediatric Nephrology/European Renal Association-European Dialysis and Transplant Association (ESPN/

- ERA-EDTA) registry study group (2010) Demographics of paediatric renal replacement therapy in Europe: 2007 annual report of the ESPN/ERA-EDTA registry. *Pediatr Nephrol* 25:1379–1382
15. Warady BA, Chadha V (2007) Chronic kidney disease in children: the global perspective. *Pediatr Nephrol* 22:1999–2009
  16. Zurowska A, Zagodzón I, Balasz I, Boguszewska A, Prokurat C, Pietrzyk J, Drozd D, Szczepanska M, Stefaniak E, Jander A, Roszkowska-Blaim D, Ziolkowska H, Makulska I, Kollataj B, Jarmolinski T, Siten G, Stankiewicz R, Wiercinski R (2006) Congenital and genetic related causes of end-stage renal disease—data from Polish Registry of Renal Replacement Therapy in Children 2000–2004. *Przegl Lek* 63(Suppl 3):57–59
  17. Zvara V, Sasinka M, Pribylincova V, Kaiserova E, Borosova E, Reznicek J (1978) Epidemiology of chronic renal insufficiency in children in Slovakia. *Cesk Pediatr* 33:321–325