



University of Groningen

## Inflammation Is Associated with Voriconazole Trough Concentrations

van Wanrooy, Marjolijn J. P.; Span, Lambert F. R.; Rodgers, Michael G. G.; van den Heuvel, Edwin R.; Uges, Donald R. A.; van der Werf, Tjip S.; Kosterink, Jos G. W.; Alffenaar, Jan-Willem C.

*Published in:*  
Antimicrobial Agents and Chemotherapy

*DOI:*  
[10.1128/AAC.03820-14](https://doi.org/10.1128/AAC.03820-14)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2014

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

van Wanrooy, M. J. P., Span, L. F. R., Rodgers, M. G. G., van den Heuvel, E. R., Uges, D. R. A., van der Werf, T. S., ... Alffenaar, J-W. C. (2014). Inflammation Is Associated with Voriconazole Trough Concentrations. *Antimicrobial Agents and Chemotherapy*, 58(12), 7098-7101. <https://doi.org/10.1128/AAC.03820-14>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Inflammation Is Associated with Voriconazole Trough Concentrations

Marjolijn J. P. van Wanrooy,<sup>a</sup> Lambert F. R. Span,<sup>b</sup> Michael G. G. Rodgers,<sup>c</sup> Edwin R. van den Heuvel,<sup>d</sup> Donald R. A. Uges,<sup>a</sup> Tjip S. van der Werf,<sup>e</sup> Jos G. W. Kosterink,<sup>a,f</sup> Jan-Willem C. Alffenaar<sup>a</sup>

University of Groningen, University Medical Center Groningen, Department of Clinical Pharmacy and Pharmacology, Groningen, The Netherlands<sup>a</sup>; University of Groningen, University Medical Center Groningen, Department of Hematology, Groningen, The Netherlands<sup>b</sup>; University of Groningen, University Medical Center Groningen, Department of Epidemiology, Groningen, The Netherlands<sup>c</sup>; University of Groningen, University Medical Center Groningen, Departments of Internal Medicine and Pulmonary Diseases & Tuberculosis, Groningen, The Netherlands<sup>d</sup>; University of Groningen, Department of Pharmacy, Division of Pharmacotherapy and Pharmaceutical Care, Groningen, The Netherlands<sup>e</sup>; University of Groningen, Department of Pharmacy, Division of Pharmacotherapy and Pharmaceutical Care, Groningen, The Netherlands<sup>f</sup>

**Voriconazole concentrations display a large variability, which cannot completely be explained by known factors. Inflammation may be a contributing factor, as inflammatory stimuli can change the activities and expression levels of cytochrome P450 isoenzymes. We explored the correlation between inflammation, reflected by C-reactive protein (CRP) concentrations, and voriconazole trough concentrations. A retrospective chart review of patients with at least one steady-state voriconazole trough concentration and a CRP concentration measured on the same day was performed. A total of 128 patients were included. A significantly ( $P < 0.001$ ) higher voriconazole trough concentration was observed in patients with severe inflammation (6.2 mg/liter; interquartile range [IQR], 3.4 to 8.7 mg/liter;  $n = 20$ ) than in patients with moderate inflammation (3.4 mg/liter; IQR, 1.6 to 5.4 mg/liter;  $n = 60$ ) and in patients with no to mild inflammation (1.6 mg/liter; IQR, 0.8 to 3.0 mg/liter;  $n = 48$ ). The patients in all three groups received similar voriconazole doses based on mg/kg body weight ( $P = 0.368$ ). Linear regression analyses, both unadjusted and adjusted for covariates of gender, age, dose, route of administration, liver enzymes, and interacting coadministered medications, showed a significant association between voriconazole and CRP concentration ( $P < 0.001$ ). For every 1-mg/liter increase in the CRP concentration, the voriconazole trough concentration increased by 0.015 mg/liter (unadjusted 95% confidence interval [CI], 0.011 to 0.020 mg/liter; adjusted 95% CI, 0.011 to 0.019 mg/liter). Inflammation, reflected by the C-reactive protein concentration, is associated with voriconazole trough concentrations. Further research is necessary to assess if taking the inflammatory status of a patient into account is helpful in therapeutic drug monitoring of voriconazole to maintain concentrations in the therapeutic window, thereby possibly preventing suboptimal treatment or adverse events.**

Voriconazole, a broad-spectrum antifungal agent, is considered a first-line agent for the treatment of invasive aspergillosis (1). Several studies showed a relation between the efficacy and safety of voriconazole and voriconazole trough concentrations (2–7). Voriconazole trough concentrations of  $>1.5$  mg/liter are associated with a favorable response to treatment. High voriconazole trough concentrations are associated with an increased incidence of adverse events, such as visual disturbances, hallucinations, and abnormalities in liver enzyme levels. The Infectious Diseases Society of America guidelines recommend determination of voriconazole concentrations in conjunction with other measures of clinical assessment to evaluate potential toxicity or to document adequate voriconazole exposure (1).

In daily practice, a large variability in voriconazole concentrations is observed, not only between patients (8) but also within individual patients over time (9). Polymorphisms of cytochrome P450 isoenzymes (10), impaired liver function (11), and drug-drug interactions (12) influence voriconazole pharmacokinetics but do not completely explain the observed variability. Therefore, other factors may contribute to the variability of voriconazole pharmacokinetics.

We expect that inflammation may be one of these factors. Infections or inflammatory stimuli can change the activities and expression levels of various forms of cytochrome P450 isoenzymes (13). The downregulation of cytochrome P450 isoenzymes during inflammation decreases the hepatic clearance of drugs that are metabolized by these enzymes. As voriconazole is extensively metabolized by cytochrome P450 isoenzymes 2C19, 3A4, and 2C9 (14), inflammation may contribute to the variability of the pharmacokinetics of voriconazole.

The aim of this retrospective study was to investigate whether inflammation, reflected by C-reactive protein (CRP) concentrations, is associated with voriconazole trough concentrations.

## MATERIALS AND METHODS

A retrospective chart review was performed for all patients aged  $\geq 18$  years who had at least one steady-state voriconazole trough concentration and a CRP concentration measured on the same day. The patients were treated at the University Medical Center Groningen, The Netherlands, between January 2006 and December 2010. Patients who were concomitantly using a strong inhibitor or inducer of cytochrome P450 isoenzymes were excluded. Steady state was considered to be achieved after 6 doses when 2 loading doses were administered (15), after 10 doses when  $<2$  loading doses were administered (16), and at 6 doses after a dosage adjustment (17). We measured voriconazole serum concentrations using a validated method that involved liquid chromatography coupled with tandem-mass spectrometry (18). This study was evaluated by the local ethics committee (IRB 2013-491) and was in accordance with Dutch law because of its retrospective nature.

For each eligible patient, data that included demographic details, medical history, and laboratory parameters were collected from the medical chart. In addition to the voriconazole trough and CRP concentrations,

Received 8 July 2014 Returned for modification 10 August 2014

Accepted 7 September 2014

Published ahead of print 15 September 2014

Address correspondence to Jan-Willem C. Alffenaar, j.w.c.alffenaar@umcg.nl.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.03820-14

TABLE 1 Patient characteristics<sup>a</sup>

Characteristic	No. (%) of patients or median (IQR)
<b>Demographic</b>	
Gender (female)	59 (46)
Age (yr)	55 (42–62)
Weight (kg)	75 (61–85)
Height (m)	1.75 (1.67–1.81)
BMI <sup>b</sup> (kg/m <sup>2</sup> )	24.2 (21.6–26.6)
<b>Underlying disease</b>	
Hematologic malignancy	67 (52)
Solid organ transplant	34 (27)
Other <sup>c</sup>	27 (21)
<b>Voriconazole treatment</b>	
Twice-daily dose (mg/kg)	3.3 (2.6–3.9)
Intravenous administration	64 (50)
<b>Laboratory parameter</b>	
Voriconazole (mg/liter)	2.7 (1.3–4.8)
CRP (mg/liter)	71 (15–152)
Alkaline phosphatase (U/liter)	114 (71–192)
ALT (U/liter)	36 (17–69)
AST (U/liter)	31 (19–61)
γ-Glutamyl transferase (U/liter)	105 (55–232)
Total bilirubin (μmol/liter)	9 (6–16)
<b>Coadministered medication</b>	
(Es)omeprazole	81 (63)

<sup>a</sup> *n* = 128.<sup>b</sup> BMI, body mass index.<sup>c</sup> Other diagnosed diseases included chronic pulmonary obstructive disease, cystic fibrosis, and granulomatosis with polyangiitis.

alkaline phosphatase, alanine aminotransferase (ALT), aspartate aminotransferase (AST), γ-glutamyl transferase (γGT), and total bilirubin concentrations were collected. Information on voriconazole treatment was also gathered. In addition, we evaluated medication used concomitantly that may have influenced the voriconazole concentrations by means of interactions with cytochrome P450 isoenzymes 2C19 and 3A4.

Numerical variables are summarized as medians and interquartile ranges (IQRs), and categorical variables are summarized as frequencies and percentages.

To explore the possible association of inflammation with voriconazole trough concentrations, voriconazole trough concentrations of patients with no to mild inflammation (CRP, ≤40 mg/liter), those with moderate inflammation (CRP, 41 to 200 mg/liter), and those with severe inflammation (CRP, >200 mg/liter) (19, 20) were compared using a Kruskal-Wallis test. A linear regression analysis was performed to assess the contribution of inflammation, reflected by CRP concentrations, to the variability in voriconazole trough concentrations, both unadjusted and adjusted for the gender, age, dose, route of administration, alkaline phosphatase, ALT, AST, γGT, total bilirubin, and interacting comedication covariates. We corrected for these variables as much as possible to determine the direct association of inflammation (reflected by CRP concentrations) with voriconazole trough concentrations. For each patient, only the first steady-state voriconazole trough concentration with a CRP concentration measured on the same day was used for analysis. These analyses were performed with IBM SPSS 22.0 (IBM Corp., Armonk, NY, USA).

## RESULTS

Based on data from the hospital information system, we selected patients who had a steady-state voriconazole trough concentra-

tion and a CRP concentration measured on the same day. Of these 133 patients, 128 patients were included in our analysis. Five patients were excluded because they concomitantly used phenytoin (*n* = 2), rifampin (*n* = 2), or HIV medication (*n* = 1).

The patient characteristics are shown in Table 1. The median values of most laboratory parameters were inside their respective normal ranges. CRP and γGT concentrations were higher than the upper limit of normal. Regarding medication used concomitantly, which potentially influenced voriconazole concentrations, only omeprazole and esomeprazole were used simultaneously with voriconazole.

A significantly (*P* < 0.001) higher voriconazole trough concentration was observed in patients with severe inflammation (6.2 [IQR, 3.4 to 8.7] mg/liter; *n* = 20) than in patients with moderate inflammation (3.4 [1.6 to 5.4] mg/liter; *n* = 60) and no to mild inflammation (1.6 [0.8 to 3.0] mg/liter; *n* = 48). These differences are shown in Fig. 1. The patients in all three groups received similar doses of voriconazole (*P* = 0.368) twice daily. The median (IQR) voriconazole doses were 3.0 (2.5 to 3.8) mg/liter for patients with no to mild inflammation, 3.6 (2.7 to 3.9) mg/liter for patients with moderate inflammation, and 3.5 (2.9 to 4.0) mg/liter for patients with severe inflammation. Of the 48 patients with a CRP concentration of ≤40 mg/liter, only one patient (2.1%) had a voriconazole trough concentration of >5 mg/liter (5.8 mg/liter). Of the 20 patients with a CRP concentration of >200 mg/liter, none had a voriconazole trough concentration of <1.5 mg/liter.

The association between CRP concentrations and voriconazole trough concentrations was significant in both the unadjusted (*P* < 0.001) and adjusted (*P* < 0.001) linear regression analyses. For every 1-mg/liter increase in the CRP concentration, the voriconazole trough concentration increased by 0.015 mg/liter in each of the two analyses (unadjusted 95% confidence interval [CI], 0.011 to 0.020 mg/liter; adjusted 95% CI, 0.011 to 0.019 mg/liter). The adjusted linear regression analysis explained 46% of the variance in the voriconazole trough concentrations. Figure 2 shows the voriconazole and CRP concentrations of a patient during treatment with voriconazole.




FIG 1 Box (median and 25th to 75th percentiles) and whisker (5th and 95th percentiles) plots of voriconazole trough concentrations for patients with no to mild inflammation (*n* = 48), moderate inflammation (*n* = 60), and severe inflammation (*n* = 20). The filled circles represent outliers. The median (interquartile range [IQR]) voriconazole dose (in mg/kg of body weight) for each group is displayed above each box-and-whisker plot. The dotted lines represent the therapeutic window of voriconazole trough concentrations.




FIG 2 Voriconazole trough concentrations (solid curve) and CRP concentrations (dashed curve) of a 46-year-old male patient with cystic fibrosis who was admitted to the intensive care unit with respiratory insufficiency and was treated with voriconazole after *Aspergillus fumigatus* was isolated from his respiratory secretions. On day 24, the patient received a bilateral lung transplant. The dotted lines represent the therapeutic window of voriconazole.

## DISCUSSION

This is the first study to show that inflammation, reflected by C-reactive protein concentrations, is associated with voriconazole trough concentrations. The multiple linear regression analysis showed that, for every 1-mg/liter increase in the CRP concentration, the voriconazole trough concentration was 0.015 mg/liter higher. These results provide a possible explanation for some of the variability in voriconazole concentrations.

Inflammation is associated with the variability observed between and within patients at different time points, regardless of polymorphisms of cytochrome P450 isoenzymes (10), impaired liver function (11), and drug-drug interactions (12), which are already known to influence voriconazole pharmacokinetics. The inflammatory status of patients is different and can change over time during voriconazole treatment, which alters inflammatory cytokine concentrations. The inflammatory cytokines cause a downregulation of cytochrome P450 isoenzymes at the level of gene transcription, which results in a decrease in the corresponding mRNA, protein, and enzyme activities (13, 21–23). These decreases change the rate of the metabolism of voriconazole, resulting in variable voriconazole concentrations. Triflino et al. hypothesized that the observed variability is caused by multiple factors, including changes in absorption, patient protein status, liver function, and disease-modifying effects, such as graft-versus-host disease (9). Long-term exposure to voriconazole has been suggested to cause autoinduction of cytochrome P450 isoenzymes, resulting in decreasing voriconazole concentrations (24). However, autoinduction has not been seen in healthy human volunteers (25).

A limitation of this study is its retrospective nature, which possibly introduced selection bias and limited the availability of desired data. The impact of selection bias on the results was expected to be limited, given that voriconazole concentrations were measured in most patients in our hospital, not only when efficacy or safety was in question. Information on the genetic polymorphisms of the cytochrome P450 isoenzymes in individual patients was not available, because these data were not determined in routine patient care. Therefore, we could explain only a part of the variability of voriconazole pharmacokinetics. Cytochrome P450 polymorphisms contribute to the variability of voriconazole pharmacokinetics between patients (10). Polymorphisms might explain some of the pharmacokinetic variability of voriconazole, in particular the unexplained variability in our model. It is unlikely that polymorphisms completely explain the observed variability, because the prevalence of poor metabolizers of CYP2C19, who likely contribute most to the variability in voriconazole trough concentrations between patients, is only 3% to 5% in the Caucasian population (26).

CRP concentrations were used as markers for inflammatory status because the concentrations of inflammatory cytokines were not available in this retrospective study. Cytokines stimulate the production of acute-phase proteins; CRP concentrations in particular change rapidly when the inflammatory status of a patient changes (27). As a result, CRP concentrations are used extensively in daily practice as markers for inflammation.

Further research that includes information about cytochrome P450 isoenzyme polymorphisms is necessary. We suggest a prospective study to investigate if a dosing algorithm can be devel-



oped to predict the voriconazole concentration on the basis of patient characteristics and voriconazole dose and route of administration.

In conclusion, we have shown that inflammation, reflected by the CRP concentration, is associated with voriconazole trough concentrations. Further research is necessary to assess if accounting for the inflammatory status of a patient is helpful in therapeutic drug monitoring of voriconazole to maintain concentrations in the therapeutic window, thereby possibly preventing suboptimal treatment or adverse events.

## REFERENCES

- Walsh TJ, Anaissie EJ, Denning DW, Herbrecht R, Kontoyiannis DP, Marr KA, Morrison VA, Segal BH, Steinbach WJ, Stevens DA, van Burik JA, Wingard JR, Patterson TF, Infectious Diseases Society of America. 2008. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin. Infect. Dis.* 46:327–360. <http://dx.doi.org/10.1086/525258>.
- Dolton MJ, Ray JE, Chen SC, Ng K, Pont LG, McLachlan AJ. 2012. Multicenter study of voriconazole pharmacokinetics and therapeutic drug monitoring. *Antimicrob. Agents Chemother.* 56:4793–4799. <http://dx.doi.org/10.1128/AAC.00626-12>.
- Park WB, Kim NH, Kim KH, Lee SH, Nam WS, Yoon SH, Song KH, Choe PG, Kim NJ, Jang IJ, Oh MD, Yu KS. 2012. The effect of therapeutic drug monitoring on safety and efficacy of voriconazole in invasive fungal infections: a randomized controlled trial. *Clin. Infect. Dis.* 55:1080–1087. <http://dx.doi.org/10.1093/cid/cis599>.
- Smith J, Safdar N, Knasinski V, Simmons W, Bhavnani SM, Ambrose PG, Andes D. 2006. Voriconazole therapeutic drug monitoring. *Antimicrob. Agents Chemother.* 50:1570–1572. <http://dx.doi.org/10.1128/AAC.50.4.1570-1572.2006>.
- Miyakis S, van Hal SJ, Ray J, Marriott D. 2010. Voriconazole concentrations and outcome of invasive fungal infections. *Clin. Microbiol. Infect.* 16:927–933. <http://dx.doi.org/10.1111/j.1469-0691.2009.02990.x>.
- Pascual A, Calandra T, Bolay S, Buclin T, Bille J, Marchetti O. 2008. Voriconazole therapeutic drug monitoring in patients with invasive mycoses improves efficacy and safety outcomes. *Clin. Infect. Dis.* 46:201–211. <http://dx.doi.org/10.1086/524669>.
- Pascual A, Csajka C, Buclin T, Bolay S, Bille J, Calandra T, Marchetti O. 2012. Challenging recommended oral and intravenous voriconazole doses for improved efficacy and safety: population pharmacokinetics-based analysis of adult patients with invasive fungal infections. *Clin. Infect. Dis.* 55:381–390. <http://dx.doi.org/10.1093/cid/cis437>.
- Trifilio S, Ortiz R, Pennick G, Verma A, Pi J, Stosor V, Zembower T, Mehta J. 2005. Voriconazole therapeutic drug monitoring in allogeneic hematopoietic stem cell transplant recipients. *Bone Marrow Transplant.* 35:509–513. <http://dx.doi.org/10.1038/sj.bmt.1704828>.
- Trifilio SM, Yarnold PR, Scheetz MH, Pi J, Pennick G, Mehta J. 2009. Serial plasma voriconazole concentrations after allogeneic hematopoietic stem cell transplantation. *Antimicrob. Agents Chemother.* 53:1793–1796. <http://dx.doi.org/10.1128/AAC.01316-08>.
- Weiss J, Ten Hoevel MM, Burhenne J, Walter-Sack I, Hoffmann MM, Rengelshausen J, Haefeli WE, Mikus G. 2009. CYP2C19 genotype is a major factor contributing to the highly variable pharmacokinetics of voriconazole. *J. Clin. Pharmacol.* 49:196–204. <http://dx.doi.org/10.1177/0091270008327537>.
- Jeu L, Piacenti FJ, Lyakhovetskiy AG, Fung HB. 2003. Voriconazole. *Clin. Ther.* 25:1321–1381. [http://dx.doi.org/10.1016/S0149-2918\(03\)80126-1](http://dx.doi.org/10.1016/S0149-2918(03)80126-1).
- Brüggemann RJ, Alffenaar JW, Blijlevens NM, Billaud EM, Kosterink JG, Verweij PE, Burger DM. 2009. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with other coadministered agents. *Clin. Infect. Dis.* 48:1441–1458. <http://dx.doi.org/10.1086/598327>.
- Morgan ET. 1997. Regulation of cytochromes P450 during inflammation and infection. *Drug Metab. Rev.* 29:1129–1188. <http://dx.doi.org/10.3109/03602539709002246>.
- Theuretzbacher U, Ihle F, Derendorf H. 2006. Pharmacokinetic/pharmacodynamic profile of voriconazole. *Clin. Pharmacokinet.* 45:649–663. <http://dx.doi.org/10.2165/00003088-200645070-00002>.
- Purkins L, Wood N, Greenhalgh K, Eve MD, Oliver SD, Nichols D. 2003. The pharmacokinetics and safety of intravenous voriconazole—a novel wide-spectrum antifungal agent. *Br. J. Clin. Pharmacol.* 56(Suppl 1):S2–S9. <http://dx.doi.org/10.1046/j.1365-2125.2003.01992.x>.
- Purkins L, Wood N, Greenhalgh K, Allen MJ, Oliver SD. 2003. Voriconazole, a novel wide-spectrum triazole: oral pharmacokinetics and safety. *Br. J. Clin. Pharmacol.* 56(Suppl 1):S10–S16. <http://dx.doi.org/10.1046/j.1365-2125.2003.01993.x>.
- Purkins L, Wood N, Ghahramani P, Greenhalgh K, Allen MJ, Kleinermans D. 2002. Pharmacokinetics and safety of voriconazole following intravenous-to oral-dose escalation regimens. *Antimicrob. Agents Chemother.* 46:2546–2553. <http://dx.doi.org/10.1128/AAC.46.8.2546-2553.2002>.
- Alffenaar JW, Wessels AM, van Hateren K, Greijdanus B, Kosterink JG, Uges DR. 2010. Method for therapeutic drug monitoring of azole antifungal drugs in human serum using LC/MS/MS. *J. Chromatogr. B Analyt. Technol. Biomed. Life. Sci.* 878:39–44. <http://dx.doi.org/10.1016/j.jchromb.2009.11.017>.
- Manian FA. 1995. A prospective study of daily measurement of C-reactive protein in serum of adults with neutropenia. *Clin. Infect. Dis.* 21:114–121. <http://dx.doi.org/10.1093/clinids/21.1.114>.
- Clyne B, Olshaker JS. 1999. The C-reactive protein. *J. Emerg. Med.* 17:1019–1025. [http://dx.doi.org/10.1016/S0736-4679\(99\)00135-3](http://dx.doi.org/10.1016/S0736-4679(99)00135-3).
- Aitken AE, Morgan ET. 2007. Gene-specific effects of inflammatory cytokines on cytochrome P450 2C, 2B6 and 3A4 mRNA levels in human hepatocytes. *Drug Metab. Dispos.* 35:1687–1693. <http://dx.doi.org/10.1124/dmd.107.015511>.
- Aitken AE, Richardson TA, Morgan ET. 2006. Regulation of drug-metabolizing enzymes and transporters in inflammation. *Annu. Rev. Pharmacol. Toxicol.* 46:123–149. <http://dx.doi.org/10.1146/annurev.pharmtox.46.120604.141059>.
- Renton KW. 2004. Cytochrome P450 regulation and drug biotransformation during inflammation and infection. *Curr. Drug Metab.* 5:235–243. <http://dx.doi.org/10.2174/1389200043335559>.
- Moriyama B, Elinoff J, Danner RL, Gea-Banacloche J, Pennick G, Rinaldi MG, Walsh TJ. 2009. Accelerated metabolism of voriconazole and its partial reversal by cimetidine. *Antimicrob. Agents Chemother.* 53:1712–1714. <http://dx.doi.org/10.1128/AAC.01221-08>.
- Roffey SJ, Cole S, Comby P, Gibson D, Jezequel SG, Nedderman AN, Smith DA, Walker DK, Wood N. 2003. The disposition of voriconazole in mouse, rat, rabbit, guinea pig, dog, and human. *Drug Metab. Dispos.* 31:731–741. <http://dx.doi.org/10.1124/dmd.31.6.731>.
- Tamminga WJ, Wemer J, Oosterhuis B, Weiling J, Wilffert B, de Leij LF, de Zeeuw RA, Jonkman JH. 1999. CYP2D6 and CYP2C19 activity in a large population of Dutch healthy volunteers: indications for oral contraceptive-related gender differences. *Eur. J. Clin. Pharmacol.* 55:177–184. <http://dx.doi.org/10.1007/s002280050615>.
- Gabay C, Kushner I. 1999. Acute-phase proteins and other systemic responses to inflammation. *N. Engl. J. Med.* 340:448–454. <http://dx.doi.org/10.1056/NEJM199902113400607>.