



University of Groningen

Iron status and heart failure

Klip, IJsbrand Thomas

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Klip, IJ. T. (2016). Iron status and heart failure: From prediction to prognosis. [Groningen]: Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER

**SERUM FERRITIN AND
RISK FOR NEW ONSET
HEART FAILURE AND
CARDIOVASCULAR EVENTS
IN THE COMMUNITY**

IJsbrand T. Klip
Adriaan A. Voors
Dorine W. Swinkels
Stephan J.L. Bakker
Jenny E. Kootstra-Ros
Carolyn S. Lam
Pim van der Harst
Dirk J. van Veldhuisen
Peter van der Meer

Accepted Eur J Heart Fail 2016

5

ABSTRACT

Aims:

Heart failure (HF) is a common manifestation of patients with primary and secondary causes of iron overload, whereas in patients with established HF iron deficiency impairs outcome. Whether body iron, either depleted or in overload, amplifies the risk for new onset HF among healthy individuals is unknown. The present study aimed to assess whether markers of iron status, or iron-regulatory hormone hepcidin are associated with new onset HF or cardiovascular (CV) events in the general population.

Methods and results:

In 6386 subjects from the Prevention of REnal and Vascular ENd-stage Disease (PREVEND), a prospective, community-based, cohort study, markers of iron status and iron-regulatory hormone hepcidin were measured. Mean age was 53.1 ± 12.0 years and 50.7% of the cohort was female. During a mean follow-up of 8.3 years, 199 subjects (3.1%) were newly diagnosed with HF, 456 (7.1%) experienced a CV event and 356 died from all causes. A higher annual HF incidence per ferritin quartile was observed in women ($P < 0.001$), but not in men (P for interaction 0.032). Multivariable analyses demonstrated ferritin levels to remain independently predictive for new onset HF in women only ($P = 0.024$). This association persisted within strata defined by markers of the metabolic syndrome, markers of inflammation or other markers of iron homeostasis, including hepcidin. No association between ferritin or hepcidin and incident CV events or all-cause mortality was observed in either sex.

Conclusions:

Increased serum ferritin levels independently amplify the risk for new onset HF in women in the community. Studies need to elucidate the pathophysiological mechanisms through which ferritin levels associate with specific outcomes.

ABBREVIATIONS

BMI	=	body mass index
CV	=	cardiovascular
HF	=	heart failure
HFpEF	=	heart failure with preserved ejection fraction
HFrfEF	=	heart failure with reduced ejection fraction
hs-CRP	=	high-sensitive C-reactive protein
LBBB	=	left bundle branch block
LVH	=	left ventricular hypertrophy
MI	=	myocardial infarction
PREVEND	=	Prevention of REnal and Vascular ENd-stage Disease
ROS	=	Reactive oxygen species
TSAT	=	Transferrin saturation
UAE	=	urinary albumin excretion

INTRODUCTION

Iron is an essential micronutrient for a wide range of metabolic and biological processes. Systemic iron homeostasis is orchestrated by hepcidin, a liver-derived peptide that has emerged as the key circulating regulator of iron absorption and tissue distribution.¹ In response to low body iron stores, decreased erythropoiesis and hypoxia, hepcidin levels decrease allowing efficient iron mobilization. In contrast, inflammation and increased body iron stores cause excessive hepcidin production, which subsequently inhibits intestinal iron absorption and iron release from reticuloendothelial macrophages.¹

In patients with primary and secondary causes of iron overload, cardiomyopathy is a common manifestation.² In contrast, severe iron deficiency has also been associated with abnormalities in systolic and diastolic cardiac function, suggesting an U-shaped relationship between body iron stores and the process leading to heart failure.^{3,4} Prior epidemiological studies have proposed an association between increasing body iron stores (mostly expressed as serum ferritin levels) and risk for cardiovascular (CV) disease in the general population, whereas others have found conflicting results.⁵ Likewise, the role of hepcidin as a risk modifier in cardiovascular disease is also controversial.^{6,7}

Whether markers of iron status, either depleted or in overload, or regulatory hormone hepcidin predict the risk of new onset heart failure (HF) among healthy individuals is unknown. Therefore, we aimed to determine whether markers of iron homeostasis antedate the development of HF and cardiovascular outcome. For these purposes, we used data obtained in the Prevention of RENal and Vascular ENd-stage Disease (PREVEND) study, a large prospective, well-characterized, contemporary observational cohort with a long-term follow-up period.

METHODS

This study was performed using data of subjects participating in the PREVEND study. Details of the study protocol have been published elsewhere (www.prevend.org).⁸ In brief, from 1997 to 1998, all inhabitants of the city of Groningen, The Netherlands, aged 28-75 years, were sent a questionnaire on demographics, disease history, smoking habits, use of medication, and a vial to collect an early morning urinary sample (n = 85,421). Of these subjects, 40,856 responded (47.8%). After exclusion of subjects with type 1 diabetes

mellitus (defined as the use of insulin), and pregnant women, subjects with an urinary albumin excretion (UAE) ≥ 10 mg/L (n=6000) and a randomly selected control group with an UAE < 10 mg/L (n=2592) completed the protocol and formed the baseline PREVEND cohort (n=8592). For the current analyses we used data from the second survey, which took place between 2001 and 2003 (n=6894), as hepcidin and iron measurements were only available from this time period. We excluded 508 subjects due to missing ferritin or hepcidin values, and missing follow-up data, leaving 6386 subjects for the current analysis. The PREVEND study was approved by the institutional medical ethics committee and conducted in accordance with the Declaration of Helsinki. All subjects provided written informed consent.

Definitions & calculations

Blood pressure was calculated as the mean of the measurements of the two study visits of the second survey. Hypertension was defined as systolic blood pressure > 140 mmHg, diastolic blood pressure > 90 mmHg, or self-reported use of antihypertensive medication. Body mass index (BMI) was calculated as the ratio of weight and height squared (kg/m^2), and obesity was defined as a BMI > 30 kg/m^2 . Hypercholesterolemia was defined as total serum cholesterol > 6.5 mmol/L (251 mg/dL), a serum cholesterol ≥ 5.0 mmol/L (193mg/dL) when a history of myocardial infarction (MI) was present or when lipid-lowering medication was used. Type 2 diabetes mellitus was defined as a fasting glucose level of ≥ 7.0 mmol/L (126 mg/dL), a non-fasting glucose level of ≥ 11.1 mmol/L (200 mg/dL), or the use of anti-diabetic drugs. Smoking was defined as current smoking or smoking cessation within the previous year. Urinary albumin excretion was calculated as the average UAE in the two consecutive 24-h urine collections. An estimate of the glomerular filtration rate was calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation.⁹ History of myocardial infarction or stroke was defined as a self-reported condition requiring hospitalization for at least 3 days. Anemia was defined according to the World Health Organization (WHO) criteria (hemoglobin < 13 g/dL for men and < 12 g/dL for women).¹⁰ Standard 12-lead electrocardiograms were recorded using the computer Modular ECG Analysis System.¹¹ The presence of a left bundle branch block (LBBB) was defined as a QRS duration on ECG > 120 msec. Left ventricular hypertrophy (LVH) was defined using the Cornell criteria: $\text{RaVL} + \text{SV}_3$ (with 6mm added in women) multiplied by the QRS duration. A threshold of 2440 mm*ms was used to identify LVH.¹²

Analytical methods

Fasting blood samples were obtained in the morning from all participants from 2001-2003. Aliquots of these samples were stored immediately at -80°C until further analysis. Hemoglobin (g/dL) was measured using a Coulter Counter STKS sum (Coulter Corporation, Miami, Florida, USA). Serum iron (umol/L), ferritin (ug/L) and transferrin (g/L) were measured using a colorimetric assay, immunoassay and immunoturbidimetric assay, respectively (Roche Diagnostics, Mannheim, Germany). The total iron binding capacity was calculated by multiplying transferrin x 25.2. Subsequently, transferrin saturation (TSAT [%]) was computed by dividing serum iron by TIBC and multiplied by 100. Serum hepcidin concentrations were measured with a competitive enzyme-linked immunosorbent assay, as described elsewhere.¹³ The between-plates and inter-assay coefficient of variation (CV) were 8.6% and 16.2%, respectively. Hepcidin concentrations are expressed in nanomoles per liter (nmol/L). Concentrations of total cholesterol, plasma glucose and serum creatinine were measured using standard methods. Urinary albumin concentration was determined by nephelometry, with a threshold of 2.3 mg/L and intra- and interassay CV of 2.2 and 2.6%, respectively (BN II, Dade Behring Diagnostica, Marburg, Germany). High-sensitive C-reactive protein (hs-CRP) was also determined using nephelometry with a threshold of 0.175 mg/L and intra- and interassay coefficients of less than 4.4 and 5.7% respectively.

Definition of cardiovascular events and new onset heart failure

Follow-up for the present study was defined as the time between the first follow-up visit to the outpatient department and the date of a CV event, new onset HF, death, or January 1st 2011. Subjects were censored on the date they moved to an unknown destination or at the last date of follow-up (January 1st 2011), whichever date came first. Information on dates and causes of death for every participant was obtained from Statistics Netherlands and coded according to the 10th revision of the International Classification of Diseases.¹⁴

The combined incidence of CV morbidity and mortality after the second survey was used. Information on hospitalization for CV morbidity was obtained from PRISMANT, the Dutch national registry of hospital discharge diagnoses, which has proven validity with 84% of the primary diagnosis and 87% of the secondary diagnosis matching those recorded in the patients' charts.¹⁵ In the present study, CV events were defined as the following: acute MI (ICD-code 410), acute and subacute ischaemic heart disease (411), subarachnoid haemorrhage (430), intra-cerebral haemorrhage (431), other intra-cranial haemorrhage (432), occlusion or stenosis of the pre-cerebral

(433) or cerebral arteries (434), coronary artery bypass grafting or percutaneous transluminal coronary angioplasty, and other vascular interventions as percutaneous transluminal angioplasty or bypass grafting of aorta and peripheral vessels.

Participants with a new diagnosis of HF were identified using criteria described in the Heart Failure Guidelines of the European Society of Cardiology and an endpoint adjudication committee ascertained the diagnosis of HF, as described elsewhere.¹⁶ Additionally, HF was classified as HF with a reduced (HFrEF) or preserved ejection fraction (HFpEF) based on the left ventricular ejection fraction at the time of diagnosis. To acknowledge the most recent trends in cut-off for HFrEF and HFpEF in accordance with the most recent HF guidelines, we set the cut-off for HFpEF at $\geq 50\%$.¹⁷ In all subjects developing new onset heart failure in the present study data on LVEF was available.

Statistical analyses

Baseline continuous variables are expressed as means with standard deviation (SD) when normally distributed, as medians with interquartile range (IQR) when distribution is skewed, or as numbers and percentages when categorical. Inter-group differences were tested using trend-analyses. For further analyses, skewed variables were transformed to a 2-log scale to achieve a normal distribution.

By design, the PREVEND study over-selected subjects with an elevated UAE ($> 10\text{mg/L}$). To overcome this over-selection of subjects with UAE, a statistical weighted method was applied in our regression analyses, allowing our conclusions to be generalized to the general population.^{18,19} The association of ferritin with baseline variables was examined by bootstrapping the linear regression model 1000 times. Variables selected >700 times were assumed to be accurate and included in the multivariate model.

Overall and sex-stratified adjusted annual incidence rates of new onset HF, CV events, and all-cause mortality were assessed assuming a Poisson distribution. The test for trend was calculated using the Mantel-Haenszel method. To preserve ferritin (and hepcidin) as continuous variables, multi-variable fractional polynomials were employed to determine the best fitting functional form of both markers for all outcome parameters. To account for death as a competing risk for new onset HF, a subject's follow-up time was censored at the time of death whenever death occurred before the onset of HF. Hazard ratios are then interpreted as the covariate effects on the cause-specific (sub)hazard function of the transition from heart failure-free to new onset HF in a competing risks model with new onset HF and all-cause

mortality as the two absorbing states.²⁰ Crude analyses were consecutively adjusted for 1) age and sex; 2) a multivariate model consisting of the following established HF and CV risk factors: age, sex, BMI, current smoking, diabetes, hypertension, hypercholesterolemia, history of MI or stroke, LVH, renal function, and levels of hemoglobin, erythropoietin, UAE, and presence of inflammation (hs-CRP>5 mg/L); and 3) time-varying development of CV events. Similar analyses were done for incident CV events and all-cause mortality. All reported probability values are two-tailed and a value of $P < 0.05$ was used as the nominal level of statistical significance. Models and analyses were performed using STATA software version 13.0 (StataCorp LP, College station, Texas, USA).

RESULTS

Baseline characteristics of the 6386 subjects, stratified per ferritin quartiles, are displayed in *Table 1*. Mean age was 53.1 ± 12.0 years and 50.7% of the cohort was female. Median ferritin concentration was 97 (47 - 173) ug/L and was higher in men (142 ug/L) compared to women (61 ug/L; $P < 0.001$). Iron deficiency, defined by ferritin levels < 30 ug/L, was present in 928 subjects (14.5%) and more frequently observed in women (24.6% vs. 4.2% in men; $P < 0.001$). Although postmenopausal women had significantly higher ferritin levels in relation to their premenopausal counterparts (94 ug/L vs. 36 ug/L; $P < 0.001$), iron deficiency was still more frequently seen when compared to men (8.7% vs. 4.2% in men; $P < 0.001$).

Per increasing ferritin quartile, individuals were generally older, more often male, and more likely to have cardiovascular risk factors (e.g. diabetes, hypertension). Additionally, higher concentrations of hemoglobin, hs-CRP, TSAT and hepcidin were observed, whereas renal function worsened and EPO concentrations declined. Baseline ferritin quartiles were additionally stratified by sex (*Table S1A&1B*). Similar results were detected for both sexes, with the exception of prior MI and the presence of either LBBB or LVH, which were more frequently observed in women.

Clinical associates of ferritin levels

In both sexes, serum ferritin remained positively correlated with hemoglobin and MCV, other markers of iron homeostasis and glucose concentrations. Whereas ferritin was strongly related to cardiovascular risk factors (BMI, cholesterol) in men, inflammation and decreasing erythropoietin levels were correlated with ferritin levels in women (*Table 2*). In a bootstrapped

model for ferritin (per doubling), these variables were highly selected. The presence of LVH (a strong HF risk factor) or LBBB was associated with ferritin levels in women (but not in men) in univariable regression only (Table S2). The multivariable model in women had a better model fit compared to men (Table 2). Similar cardiovascular risk factors related to ferritin levels in men and women without prior CV disease and adjusted R² was comparable (Table S3).

Markers of iron homeostasis and new onset heart failure.

After a median follow-up of 8.3 (7.8-8.9), years, 199 individuals (3.1%) were diagnosed with new onset HF, 456 (7.1%) experienced a CV event and 356 (5.6%) died from all-causes. Compared to other markers of iron status, ferritin had the best accuracy in predicting incident HF and other events (Figure S7). The adjusted annual incidence of new onset HF rose per increasing fer-




Figure 1. Total annual incidence of new onset heart failure according to ferritin (1A) and hepcidin (1B) quartiles and stratified by sex.



Table 1. Baseline characteristics according to ferritin quartiles.

Variables	Total	Q1	Q2	Q3	Q4	P - value for trend
<i>n</i>	6386	1636	1579	1576	1595	NA
Ferritin (ug/L), min-max	97 (1-1925)	27 (1-48)	71 (49-97)	130 (98-172)	257 (173-1925)	NA
Clinical signs						
Age (years)	53.1 ± 12.0	48.3 ± 10.9	53.3 ± 12.0	54.6 ± 11.7	56.4 ± 11.7	< 0.001
Female sex (%)	50.7	82.0	56.6	40.5	23.1	< 0.001
BMI (kg/m ²)	26.7 ± 4.3	25.5 ± 4.3	26.2 ± 4.3	26.9 ± 4.2	28.1 ± 4.1	< 0.001
> 30 (%)	18.7	12.7	15.5	19.1	27.6	< 0.001
Systolic BP (mmHg)	126.3 ± 18.8	119.5 ± 17.1	125.0 ± 18.7	128.5 ± 18.1	132 ± 18.6	< 0.001
Heart rate (b.p.m.)	68.5 ± 10.0	68.9 ± 9.4	68.4 ± 10.3	68.4 ± 10.0	68.3 ± 10.3	0.022
LVH according to Cornell (%)	2.2	1.0	2.3	2.2	3.2	0.001
LBBB (%)	5.8	3.4	4.8	7.1	8.1	< 0.001
Medical history (%)						
Smoking or quit < 1 year	30.4	29.7	32.8	31.0	28.0	0.19
MI	2.9	2.0	2.5	3.1	4.2	< 0.001
Stroke	0.9	0.9	1.1	0.6	0.9	0.68
Type 2 diabetes	7.8	5.0	6.3	7.8	12.1	< 0.001
Hypercholesterolemia	24.9	15.8	24.1	26.6	33.1	< 0.001
Hypertension	31.2	19.1	27.8	34.4	43.7	< 0.001
Anemia	9.3	20.9	7.2	5.0	4.1	< 0.001

Table 1. Baseline characteristics according to ferritin quartiles. (continued)

Variables	Total	Q1	Q2	Q3	Q4	P - value for trend
<i>Laboratory measurements</i>						
Hemoglobin (g/dL)	13.7 ± 1.2	12.9 ± 1.2	13.7 ± 1.1	14.0 ± 1.1	14.3 ± 1.1	< 0.001
MCV (fL)	90.4 ± 4.6	88.9 ± 5.3	90.7 ± 4.1	90.8 ± 4.2	91.3 ± 4.4	< 0.001
Hepcidin (nmol/L)	3.1 (1.7-4.9)	1.1 (0.6-1.7)	2.6 (1.8-3.5)	3.8 (2.8-5.1)	5.6 (4.2-8.0)	< 0.001
Serum iron (umol/L)	15.9 ± 5.6	14.1 ± 6.3	16.0 ± 5.2	16.1 ± 5.1	17.2 ± 5.5	< 0.001
TSAT (%)	24.8 ± 9.4	20.6 ± 9.5	25.0 ± 8.3	25.8 ± 8.4	28.2 ± 9.6	< 0.001
Erythropoietin (IU/L)	7.8 (5.9-10.3)	8.6 (6.5-12.0)	7.7 (5.9-10.1)	7.5 (5.7-9.7)	7.3 (5.5-9.7)	< 0.001
hs-CRP (mg/L)	1.4 (0.6-3.0)	1.0 (0.4-2.3)	1.3 (0.6-2.9)	1.4 (0.7-3.2)	1.8 (0.8-3.6)	< 0.001
Cholesterol (mmol/L)	5.4 ± 1.0	5.1 ± 1.0	5.4 ± 1.0	5.5 ± 1.0	5.6 ± 1.1	< 0.001
Glucose (mmol/L)	4.8 (4.4-5.3)	4.6 (4.3-5.1)	4.7 (4.4-5.2)	5.0 (4.5-5.4)	5.1 (4.6-5.6)	< 0.001
Creatinine (umol/L)	84.8 ± 19.1	78.9 ± 14.1	83.4 ± 16.9	86.5 ± 19.0	90.4 ± 23.6	< 0.001
eGFR (ml/min/1.73m ²)	89.6 ± 17.7	94.1 ± 17.1	89.9 ± 17.6	88.2 ± 17.4	86.2 ± 17.7	< 0.001
< 60 (%)	5.7	3.2	5.5	6.7	7.7	< 0.001
UAE (mg/24h)	8.5 (6.0-15.1)	7.8 (5.7-12.4)	7.9 (5.8-13.9)	8.5 (6.1-16.0)	10.0 (6.7-20.3)	< 0.001

Continuous variables are presented as mean ± standard deviation or as median (inter-quartile range) when non-normally distributed. Binary categorical variables are presented as percentages. Abbreviations: BMI = Body mass index, BP = Blood pressure, eGFR = Estimated glomerular filtration rate, hs-CRP = high-sensitive C-reactive protein, LBBB = Left bundle branch block, LVH = Left ventricular hypertrophy, MCV = Mean corpuscular volume, MI = Myocardial infarction, TSAT = Transferrin saturation, UAE = Urinary albumin excretion.

Table 2. Multivariable regression beta-coefficients between cardiovascular risk factors and serum ferritin in men and women (selected after bootstrapping).

Variables (selected after bootstrapping)	Men (n = 3145)			Women (n = 3241)		
	Standardized β	T	P - value	Standardized β	T	P - value
Menstruation (yes vs. no)	-	-	-	-0.129	-7.4	< 0.001
BMI (per kg/m ²)	0.090	5.4	< 0.001	-	-	-
Smoking (yes vs. no)	-	-	-	-0.035	-2.6	0.011
Hemoglobin (per g/dL)	0.108	6.5	< 0.001	0.071	4.1	< 0.001
MCV (per fL)	0.099	6.0	< 0.001	0.124	7.6	< 0.001
Log 2 hepcidin (nmol/L)	0.680	38.6	< 0.001	0.625	30.4	< 0.001
TSAT (per 5%)	0.094	5.1	< 0.001	0.131	5.7	< 0.001
Log 2 erythropoietin (IU/L)	-	-	-	-0.031	-2.0	0.042
Log 2 hs-CRP (mg/L))	-	-	-	0.059	3.5	< 0.001
Cholesterol (mmol/L)	0.044	2.5	0.011	-	-	-
Log 2 glucose (mmol/L)	0.043	3.6	< 0.001	0.052	4.4	< 0.001
	R² total model			R² total model		
			0.60			0.74

For abbreviations, see Table 1.

ritin quartile (*Figure 1*). This trend only appeared to be significant in women ($P < 0.001$). Crude and multivariable fractional polynomials revealed the association between ferritin levels and the development of new onset HF to be best described by a linear function. In subsequent Cox regression analyses (with all-cause mortality as a competing risk) increasing ferritin concentrations were strongly associated with the development of new onset HF ($P < 0.001$; *Table 3*) However, this association was lost in multivariable analyses. Interaction analyses showed a significant interaction between sex, ferritin levels and new onset HF (P_{int} 0.032). When stratified by sex, this association persisted only in women ($P < 0.001$) and remained significant in all multivariable models. Adjusting for menstrual status did not alter the results. No interaction between menstrual status, ferritin levels and development of new onset HF was found in women ($P = 0.59$). Additional subgroup analyses in women demonstrated increasing ferritin levels to amplify risk for incident HF in almost every subgroup, but were significantly more predictive in women without anemia, without microalbuminuria and with an eGFR >60 ml/min/1.73m². (*Figure S2*).

Since body iron stores are firmly regulated at the systemic level by hepcidin, we additionally explored the role of hepcidin and risk of new onset HF. Similar observations regarding adjusted annual HF incidence were made for hepcidin, both overall and in women (*Figure 1*). Likewise, a strong associa-

Table 3. Levels of ferritin and hepcidin and risk for incident heart failure with all-cause mortality as a competing risk.

Ferritin (per doubling)	sHR	95%CI	Harrell's C	P-value
Total population (n = 6386)¹				
Univariable	1.48	1.30-1.69	0.64	< 0.001
Model 1	1.24	1.05-1.46	0.83	0.011
Model 2	1.06	0.89 - 1.26	0.88	0.45
Men (n = 3145)				
Univariable	1.00	0.81-1.23	0.52	0.98
Model 1	1.06	0.87-1.29	0.82	0.54
Model 2	0.92	0.75-1.14	0.87	0.45
Women (n = 3241)				
Univariable	2.07	1.63-2.61	0.74	< 0.001
Model 1	1.57	1.20-2.05	0.83	0.001
Model 1A	1.69	1.26-2.26	0.79	< 0.001
Model 2	1.35	1.02-1.79	0.87	0.038
Model 3	1.39	1.05-1.86	0.87	0.024
Hepcidin (per doubling)	HR	95%CI	Harrell's C	P-value
Total population (n = 6386)²				
Univariable	1.45	1.23-1.72	0.61	< 0.001
Model 1	1.20	0.98-1.45	0.83	0.07
Model 2	1.09	0.92-1.30	0.88	0.39
Men (n = 3145)				
Univariable	0.96	0.79-1.16	0.54	0.76
Model 1	1.00	0.81-1.25	0.82	0.94
Model 2	0.95	0.74-1.21	0.87	0.49
Women (n = 3241)				
Univariable	2.10	1.63-2.70	0.75	< 0.001
Model 1	1.59	1.17-2.17	0.84	0.003
Model 1A	1.68	1.19-2.37	0.79	0.003
Model 2	1.44	1.03-2.05	0.87	0.039
Model 3	1.60	1.09-2.37	0.88	0.018

Abbreviations: CI = Confidence interval, sHR = Sub-hazard ratio.

1P for interaction ferritin, sex and new onset heart failure = 0.032.

2P for interaction hepcidin, sex and new onset heart failure = 0.045.

Model 1 is adjusted for age (per 10 years) and sex (in total population).

Model 1A is adjusted for menstrual state instead of age.

Model 2 is adjusted for model 1 + cardiovascular and heart failure risk factors: BMI, current smoking, the presence of diabetes, hypertension, hypercholesterolemia, history of MI or stroke, LVH, renal function, and levels of hemoglobin, erythropoietin, hs-CRP and UAE.

Model 3 is adjusted for model 2 + time-varying incident cardiovascular events.




Figure 2. Sex stratified risk for new onset heart failure according to the joint classification of ferritin levels and tertiles of TSAT (A), hepcidin (B), glucose (C), BMI (D), glucose (E) or systolic blood pressure (F).

tion between levels of hepcidin and the development of new onset heart failure was observed in women only (P_{int} 0.045). Removing potential outliers (highest and lowest 1%) did not change the results. In subjects without prior myocardial infarction and/or stroke, both ferritin and hepcidin levels remained strongly associated with the development of HF in women (*Table S4*). Sensitivity analyses in postmenopausal women showed ferritin - and not hepcidin - to remain predictive for new-onset HF (*Table S5*).

To assess whether the association between body iron stores and risk of new onset HF were modified by other markers of iron homeostasis (TSAT, hepcidin), inflammation (hs-CRP) or the metabolic syndrome (BMI, glucose, systolic blood pressure), the joint association of ferritin levels with either one of these 6 variables and the development of HF was examined. In these joint analyses, ferritin levels remained independently associated with new onset HF in women in almost all stratified analyses (*Figure 2*).

Finally, 79 of the 199 subjects newly diagnosed with HF were women. Of these women, 37 developed HFrEF and 42 HFpEF. Unadjusted cause-specific hazard analyses demonstrated serum ferritin to be associated with either HFrEF or HFpEF, with no significant interaction between both entities (*Table S6*). Multivariable analyses showed that levels of ferritin remained strongly associated with an increased risk for HFpEF only. However, again no significant interaction was found.

Markers of iron homeostasis and risk for cardiovascular disease and all-cause mortality.

A higher total annual incidence of new CV events and all-cause mortality was observed per increasing ferritin and hepcidin quartile (*Figure S3*). When stratified by sex, these observations only persisted in women. Multivariable fractional polynomials again showed the best fitting functional form between levels of ferritin or hepcidin and new CV events to be linear. Crude regression analyses showed increasing ferritin and hepcidin concentrations to be both strongly predictive for new CV events and all-cause mortality in the overall population and in women (both $P < 0.001$). However, age- and multivariable analyses were non-significant. No interaction between variables was demonstrated.

DISCUSSION

In this contemporary, prospective, population-based cohort, increasing ferritin levels amplify the risk for new onset HF in women, but not in men.

This relationship is independent of established conventional CV and HF risk factors and the development of CV events over time. These observations provide evidence that increased ferritin levels are an independent risk factor for the development of HF in women and might play a direct or indirect role in the pathogenic mechanism leading to HF.

How do elevated iron stores relate to incident heart failure?

Whereas iron deficiency has detrimental effects on outcome in established heart failure,^{17,21,22} the role of iron markers or markers of erythropoiesis in the development of HF has only been investigated to some extent.^{23,24} Previous literature suggested a U-shaped relationship between iron stores and new onset HF.^{3,4} In the present study, we observed a linear relationship between increasing ferritin levels and the development of HF. Excess body iron is known to accumulate in organs such as the heart and lead to cardiomyopathy. The generation of reactive oxygen species (ROS) in mitochondria is believed to be the most important pathogenic pathway determining cardiomyocyte damage. Production of ROS causes lipid peroxidation and DNA damage, leading to cell death, fibrosis, and eventually cardiac dysfunction. However, iron overload is only observed when the binding capacity of transferrin is fully saturated and non-transferrin bound iron is formed. Interestingly, joint analyses showed an association between serum ferritin and new onset HF in all TSAT tertiles, suggesting that another pathway linking increased ferritin levels with cardiomyocyte damage may also be involved.

Iron-mediated cell damage does not only occur under conditions of iron overload. It has been proposed that iron maldistribution among organs, tissues and cellular compartments can also attenuate cell integrity and cell life. Dysmetabolic hyperferritinemia, meaning elevated ferritin levels with normal TSAT, is commonly observed in clinical practice and has been associated with insulin resistance, obesity, hypertension, and other manifestations of the metabolic syndrome.²⁵ During circumstances of subclinical systemic inflammation, iron efflux in organs may be halted due to increased hepcidin production and subsequent internalization and degradation of its receptor, iron exporter ferroportin.²⁶ Therefore, elevated ferritin levels (and hepcidin) might be a reflection to a low-grade inflammatory response of another pathophysiological process (e.g. metabolic syndrome), causally responsible for the development of HF.²⁷ Supporting this hypothesis is the fact that we observed a positive trend in blood pressure, BMI, and levels of glucose, cholesterol and hs-CRP per increasing ferritin quartile. Additionally, strong associations between ferritin and levels of hepcidin, hs-CRP, and glucose in women were observed. To minimize for potential confounding, we adjusted

for the presence of obesity, diabetes, hypertension, hypercholesterolemia and inflammation in our multivariable analyses. Moreover, joint analyses showed increasing ferritin levels to remain predictive for incident HF in almost all tertiles. Nevertheless, to which extent elevated ferritin levels truly reflect iron stores (inflammation and oxidative stress may complicate the picture) is unclear. Moreover, the exact mechanism linking increased ferritin levels, iron-regulatory hormone hepcidin and the development of HF is still not completely understood.

To our surprise, the association between increasing ferritin levels and new onset HF was only present in women. It is known that some traditional cardiovascular risk factors may have a different impact on males compared to females.²⁸ Additional literature also suggests that the link between ferritin, hepcidin and dysmetabolic features may be particularly relevant in women.^{29,30} The role of sex hormones in the regulation of ferritin and hepcidin expression may also be responsible to a certain extent for the sex-related differences observed in the present study and has recently been studied in animal models.³¹ Yet, although we adjusted for menopausal status and other established cardiovascular/metabolic risk factors, levels of ferritin remained associated with incident HF in women.

Finally, a stronger association between levels of ferritin and the development of HFpEF was observed in women. To date there is a limited understanding of the underlying pathophysiology of HFpEF. A novel concept in the pathogenesis of HFpEF was proposed involving the role of comorbidities (e.g. hypertension, diabetes) to induce a subclinical inflammatory state, eventually leading to microvascular endothelial inflammation and subsequent stiffening and diastolic dysfunction.³² Given the aforementioned relationship between dysmetabolic hyperferritinemia and diabetes or metabolic syndrome, iron misdistribution and consecutive oxidative stress might play a direct or indirect role in the development of HFpEF. However, no significant interaction between both HF entities was observed and the number of HFpEF cases in the present study is low. Therefore, these results need cautious interpretation and merit further investigation.

Markers of iron status and risk of cardiovascular events

In the present study, we did not observe an association between ferritin or hepcidin concentrations and development of CV events or all-cause mortality in multivariable analyses. It has been hypothesized that both elevated and decreased markers of iron status enhance the risk for CV disease^{33,34}. However, conflicting results have also been reported.³⁵ Furthermore, studies in hemodialysis patients or women from the general population have

suggested a role for elevated hepcidin levels in the pathogenesis of atherosclerosis.³⁶⁻³⁸ In contrast, Kautz and colleagues did not detect any increase in hepatic hepcidin expression during progression of atherosclerosis or atherosclerotic plaque size in mice with elevated macrophage iron.⁶ A recently published population-based study also found no relationship between hepcidin or hepcidin-to-ferritin ratio and prevalent carotid atherosclerosis or incident CV events.⁷ Future population-based studies are needed to finally establish whether an abnormal iron homeostasis is associated with the development of CV events in the general population.

Implications for clinical practice

Although the present associations do not prove causality between iron stores and incident HF, the present study provides evidence that increased ferritin levels may directly or indirectly be associated with the pathogenesis of new onset HF in women. Identification of subjects at risk for HF using biomarkers alongside clinical characteristics has grown interest over the years. Markers of iron homeostasis may provide additional information to the clinician regarding etiology, clinical risk (in our case heart failure) and disease severity. Secondly, ferritin is a widely available routine marker and a relatively inexpensive measurement. Finally, elimination of iron may confer a beneficial effect. Iron removal, by means of phlebotomy, improved coronary vascular function in patients with type 2 diabetes and endothelial function in patients with known coronary artery disease.^{39,40} Still, the findings of this study need to be validated in other population-based studies.

Strengths and limitations

The large size of this contemporary, prospective community-based cohort, long follow-up period, and thorough validation of outcome parameters, are strengths of this study. Furthermore, all blood samples were taken in the morning, minimizing the influence of circadian rhythm on markers of iron status. The present study is limited by the fact that the subjects from the PREVEND study are predominantly Caucasian and our results can therefore not be extrapolated to subjects from other ethnicities. Second, our analyses rely on one baseline measurement of iron markers and hepcidin, which is used as a proxy for the concentration in the years before and after this measurement. Therefore, we can't comment on the effects of changes in markers over time. More population-based studies with serial measurements over time are warranted. Third, long-term storage at -80° Celsius might be associated, to some extent, with less precise measurements of hepcidin. A recent paper by Laarakkers and coworkers investigated the stability of hepcidin during long-

term storage.⁴¹ The authors concluded that hepcidin results are not changed during 2 years of storage at -80° Celsius. However, after 2 years storage at -80° Celsius, hepcidin results may become less precise. Although we believe that average results for a population may not be affected, perhaps existing differences in hepcidin levels between groups, or correlations of hepcidin with other parameters might be more difficult to assess. Finally, the PREVEND cohort is enriched for increased UAE. For this reason, we corrected for study design by conducting a design-based analysis. Furthermore, compared with the Framingham cohort, UAE was not higher in PREVEND.⁴²

CONCLUSIONS

In conclusion, increased ferritin levels independently enhance the risk for new onset heart failure in apparently healthy women. These findings provide evidence that elevated ferritin concentrations may be directly or indirectly involved in the pathogenesis of new onset HF in women in the community. Studies need to elucidate the pathophysiological mechanisms through which ferritin levels associate with specific outcomes.

REFERENCES

1. Ganz T, Nemeth E. Heparin and iron homeostasis. *Biochim Biophys Acta* 2012;1823:1434-1443.
2. Gujja P, Rosing DR, Tripodi DJ, Shizukuda Y. Iron overload cardiomyopathy: better understanding of an increasing disorder. *J Am Coll Cardiol* 2010;56:1001-1012.
3. Dong F, Zhang X, Culver B, Chew HG, Jr, Kelley RO, Ren J. Dietary iron deficiency induces ventricular dilation, mitochondrial ultrastructural aberrations and cytochrome c release: involvement of nitric oxide synthase and protein tyrosine nitration. *Clin Sci (Lond)* 2005;109:277-286.
4. Walter PB, Knutson MD, Paler-Martinez A, Lee S, Xu Y, Viteri FE, Ames BN. Iron deficiency and iron excess damage mitochondria and mitochondrial DNA in rats. *Proc Natl Acad Sci U S A* 2002;99:2264-2269.
5. Munoz-Bravo C, Gutierrez-Bedmar M, Gomez-Aracena J, Garcia-Rodriguez A, Navajas JF. Iron: protector or risk factor for cardiovascular disease? Still controversial. *Nutrients* 2013;5:2384-2404.
6. Kautz L, Gabayan V, Wang X, Wu J, Onwuzurike J, Jung G, Qiao B, Lusis AJ, Ganz T, Nemeth E. Testing the iron hypothesis in a mouse model of atherosclerosis. *Cell Rep* 2013;5:1436-1442.
7. Pechlaner R, Kiechl S, Mayr M, Santer P, Weger S, Haschka D, Bansal SS, Willeit J, Weiss G. Correlates of serum hepcidin levels and its association with cardiovascular disease in an elderly general population. *Clin Chem Lab Med* 2015;
8. Hillege HL, Fidler V, Diercks GF, van Gilst WH, de Zeeuw D, van Veldhuisen DJ, Gans RO, Janssen WM, Grobbee DE, de Jong PE, Prevention of Renal and Vascular End Stage Disease (PREVEND) Study Group. Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population. *Circulation* 2002;106:1777-1782.
9. Matsushita K, Mahmoodi BK, Woodward M, Emberson JR, Jafar TH, Jee SH, Polkinghorne KR, Shankar A, Smith DH, Tonelli M, Warnock DG, Wen CP, Coresh J, Gansevoort RT, Hemmelgarn BR, Levey AS, Chronic Kidney Disease Prognosis Consortium. Comparison of risk prediction using the CKD-EPI equation and the MDRD study equation for estimated glomerular filtration rate. *JAMA* 2012;307:1941-1951.
10. Nutritional anaemias. Report of a WHO scientific group. *World Health Organ Tech Rep Ser* 1968;405:5-37.
11. van Bommel JH, Kors JA, van Herpen G. Methodology of the modular ECG analysis system MEANS. *Methods Inf Med* 1990;29:346-353.
12. Smilde TD, Asselbergs FW, Hillege HL, Voors AA, Kors JA, Gansevoort RT, van Gilst WH, de Jong PE, Van Veldhuisen DJ. Mild renal dysfunction is associated with electrocardiographic left ventricular hypertrophy. *Am J Hypertens* 2005;18:342-347.
13. Kroot JJ, Laarakkers CM, Geurts-Moespot AJ, Grebenchtchikov N, Pickkers P, van Ede AE, Peters HP, van Dongen-Lases E, Wetzels JF, Sweep FC, Tjalsma H, Swinkels DW. Immunochemical and mass-spectrometry-based serum hepcidin assays for iron metabolism disorders. *Clin Chem* 2010;56:1570-1579.
14. Statistics Netherlands. Available at <http://www.cbs.nl/en-GB/menu/home/default.htm?Languageswitch=on> [data specifically requested] (20 February 2013).
15. Stricker BH, Herings RM. Plea for the retention of the Dutch National Medical Registration (LMR) to provide reliable information regarding public health

- and healthcare. *Ned Tijdschr Geneeskd* 2006;150:1916-1917.
16. Brouwers FP, de Boer RA, van der Harst P, Voors AA, Gansevoort RT, Bakker SJ, Hillege HL, van Veldhuisen DJ, van Gilst WH. Incidence and epidemiology of new onset heart failure with preserved vs. reduced ejection fraction in a community-based cohort: 11-year follow-up of PREVEND. *Eur Heart J* 2013;
 17. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P, Authors/Task Force Members, Document Reviewers. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016;
 18. Linssen GC, Bakker SJ, Voors AA, Gansevoort RT, Hillege HL, de Jong PE, van Veldhuisen DJ, Gans RO, de Zeeuw D. N-terminal pro-B-type natriuretic peptide is an independent predictor of cardiovascular morbidity and mortality in the general population. *Eur Heart J* 2010;31:120-127.
 19. Skinner CJ, Holt D, Smith TM. In: Editors: *Analyses of Complex Surveys, 1989*, Chichester, John Wiley and Sons, pp 195-198, ed.
 20. Putter H, Fiocco M, Geskus RB. Tutorial in biostatistics: competing risks and multi-state models. *Stat Med* 2007;26:2389-2430.
 21. Klip IT, Jankowska EA, Enjuanes C, Voors AA, Banasiak W, Bruguera J, Rozentryt P, Polonski L, van Veldhuisen DJ, Ponikowski P, Comin-Colet J, van der Meer P. The additive burden of iron deficiency in the cardiorenal-anaemia axis: scope of a problem and its consequences. *Eur J Heart Fail* 2014;16:655-662.
 22. Nunez J, Comin-Colet J, Minana G, Nunez E, Santas E, Mollar A, Valero E, Garcia-Blas S, Cardells I, Bodi V, Chorro FJ, Sanchis J. Iron deficiency and risk of early readmission following a hospitalization for acute heart failure. *Eur J Heart Fail* 2016;
 23. Klip IT, Postmus D, Voors AA, Brouwers FP, Gansevoort RT, Bakker SJ, Hillege HL, de Boer RA, van der Harst P, van Gilst WH, van Veldhuisen DJ, van der Meer P. Hemoglobin levels and new-onset heart failure in the community. *Am Heart J* 2015;169:94-101.e2.
 24. Grote Beverborg N, van der Wal HH, Klip IT, Voors AA, de Boer RA, van Gilst WH, van Veldhuisen DJ, Gansevoort RT, Hillege HL, van der Harst P, Bakker SJ, van der Meer P. High serum erythropoietin levels are related to heart failure development in subjects from the general population with albuminuria: data from PREVEND. *Eur J Heart Fail* 2016;
 25. Adams PC, Barton JC. A diagnostic approach to hyperferritinemia with a non-elevated transferrin saturation. *J Hepatol* 2011;55:453-458.
 26. Hentze MW, Muckenthaler MU, Andrews NC. Balancing acts: molecular control of mammalian iron metabolism. *Cell* 2004;117:285-297.
 27. Wang J, Sarnola K, Ruotsalainen S, Moilanen L, Lepisto P, Laakso M, Kuusisto J. The metabolic syndrome predicts incident congestive heart failure: a 20-year follow-up study of elderly Finns. *Atherosclerosis* 2010;210:237-242.
 28. He J, Ogden LG, Bazzano LA, Vupputuri S, Loria C, Whelton PK. Risk factors for congestive heart failure in US men and women: NHANES I epidemiologic follow-up study. *Arch Intern Med* 2001;161:996-1002.

29. Jiang R, Manson JE, Meigs JB, Ma J, Rifai N, Hu FB. Body iron stores in relation to risk of type 2 diabetes in apparently healthy women. *JAMA* 2004;291:711-717.
30. Sheu WH, Chen YT, Lee WJ, Wang CW, Lin LY. A relationship between serum ferritin and the insulin resistance syndrome is present in non-diabetic women but not in non-diabetic men. *Clin Endocrinol (Oxf)* 2003;58:380-385.
31. Brewer CJ, Wood RI, Wood JC. mRNA regulation of cardiac iron transporters and ferritin subunits in a mouse model of iron overload. *Exp Hematol* 2014;42:1059-1067.
32. Paulus WJ, Tschope C. A novel paradigm for heart failure with preserved ejection fraction: comorbidities drive myocardial dysfunction and remodeling through coronary microvascular endothelial inflammation. *J Am Coll Cardiol* 2013;62:263-271.
33. Hsu HS, Li CI, Liu CS, Lin CC, Huang KC, Li TC, Huang HY, Lin WY. Iron deficiency is associated with increased risk for cardiovascular disease and all-cause mortality in the elderly living in long-term care facilities. *Nutrition* 2013;
34. Salonen JT, Nyyssonen K, Korpela H, Tuomilehto J, Seppanen R, Salonen R. High stored iron levels are associated with excess risk of myocardial infarction in eastern Finnish men. *Circulation* 1992;86:803-811.
35. Lapice E, Masulli M, Vaccaro O. Iron deficiency and cardiovascular disease: an updated review of the evidence. *Curr Atheroscler Rep* 2013;15:358-013-0358-0.
36. Saeed O, Otsuka F, Polavarapu R, Karmali V, Weiss D, Davis T, Rostad B, Pachura K, Adams L, Elliott J, Taylor WR, Narula J, Kolodgie F, Virmani R, Hong CC, Finn AV. Pharmacological suppression of hepcidin increases macrophage cholesterol efflux and reduces foam cell formation and atherosclerosis. *Arterioscler Thromb Vasc Biol* 2012;32:299-307.
37. van der Weerd NC, Grooteman MP, Bots ML, van den Dorpel MA, den Hoedt CH, Mazairac AH, Nube MJ, Penne EL, Wetzels JF, Wiegerinck ET, Swinkels DW, Blankestijn PJ, Ter Wee PM, CONTRAST Investigators. Hepcidin-25 is related to cardiovascular events in chronic haemodialysis patients. *Nephrol Dial Transplant* 2013;28:3062-3071.
38. Galesloot TE, Holewijn S, Kiemeneij LA, de Graaf J, Vermeulen SH, Swinkels DW. Serum hepcidin is associated with presence of plaque in postmenopausal women of a general population. *Arterioscler Thromb Vasc Biol* 2014;34:446-456.
39. Nitenberg A, Ledoux S, Valensi P, Sachs R, Antony I. Coronary microvascular adaptation to myocardial metabolic demand can be restored by inhibition of iron-catalyzed formation of oxygen free radicals in type 2 diabetic patients. *Diabetes* 2002;51:813-818.
40. Duffy SJ, Biegelsen ES, Holbrook M, Russell JD, Gokce N, Keaney JF, Jr, Vita JA. Iron chelation improves endothelial function in patients with coronary artery disease. *Circulation* 2001;103:2799-2804.
41. Laarakkers CM, Wiegerinck ET, Klaver S, Kolodziejczyk M, Gille H, Hohlbaum AM, Tjalsma H, Swinkels DW. Improved mass spectrometry assay for plasma hepcidin: detection and characterization of a novel hepcidin isoform. *PLoS One* 2013;8:e75518.
42. Velagaleti RS, Gona P, Larson MG, Wang TJ, Levy D, Benjamin EJ, Selhub J, Jacques PF, Meigs JB, Tofler GH, Vasan RS. Multimarker approach for the prediction of heart failure incidence in the community. *Circulation* 2010;122:1700-1706.

SUPPLEMENTAL FILES

Table S1A. Baseline characteristics according to ferritin quartiles in men.

Variables	Total	Q1	Q2	Q3	Q4	P - value for trend
<i>n</i>	3145	785	794	783	783	NA
Ferritin (ug/L), min-max	142 (3 - 1925)	56 (3 - 84)	111 (85 - 142)	179 (143 - 229)	319 (230 - 1925)	NA
Age (years)	54.0 ± 12.4	54.7 ± 13.0	52.6 ± 12.2	53.0 ± 12.5	55.7 ± 11.6	< 0.001
BMI (kg/m ²)	26.8 ± 3.7	25.7 ± 3.5	26.3 ± 3.5	27.0 ± 3.7	28.1 ± 3.8	< 0.001
>30 (%)	17.3	11.8	12.4	17.6	27.4	< 0.001
Systolic BP (mmHg)	130.6 ± 17.6	128.9 ± 17.7	129.5 ± 17.7	129.8 ± 16.6	134.1 ± 18.1	< 0.001
Heart rate (b.p.m.)	67.2 ± 10.4	66.6 ± 10.8	67.0 ± 9.8	67.0 ± 10.0	68.1 ± 10.7	0.011
LVH according to Cornell (%)	3.0	3.4	2.6	2.1	4.0	0.72
LBBB (%)	9.4	9.8	9.7	8.3	9.7	0.62
Medical history (%)						
Smoking or quit < 1 year	30.3	31.4	31.9	31.6	26.4	0.040
MI	4.7	5.6	4.7	4.2	4.2	0.19
Stroke	1.0	1.8	0.4	0.8	1.2	0.33
Type 2 diabetes	8.7	8.2	6.9	8.0	11.7	0.010
Hypercholesterolemia	25.2	20.9	22.6	23.9	33.4	< 0.001
Hypertension	35.1	30.4	31.7	33.6	44.8	< 0.001
Anemia	5.1	9.8	3.7	4.2	2.9	< 0.001
Laboratory measurements						
Hemoglobin (g/dl)	14.5 ± 1.0	14.2 ± 1.1	14.5 ± 0.9	14.5 ± 0.9	14.6 ± 0.9	< 0.001
MCV (fl)	90.8 ± 4.4	90.0 ± 4.7	90.7 ± 4.1	90.8 ± 4.1	91.6 ± 4.5	< 0.001
Hepcidin (nM/L)	3.7 (2.3 - 5.6)	1.8 (1.1 - 2.6)	3.2 (2.3 - 4.3)	4.4 (3.3 - 5.9)	6.1 (4.5 - 8.5)	< 0.001
Serum iron (umol/L)	16.7 ± 5.5	15.7 ± 5.8	16.4 ± 5.3	16.8 ± 5.2	17.8 ± 5.5	< 0.001
Transferrin saturation (%)	26.6 ± 9.2	24.0 ± 9.1	26.2 ± 8.8	27.2 ± 8.8	29.0 ± 9.5	< 0.001
Erythropoietin (IU/L)	7.6 (5.8 - 9.9)	8.1 (6.2 - 10.9)	7.5 (5.9 - 9.9)	7.2 (5.5 - 9.2)	7.5 (5.8 - 9.7)	< 0.001
hs-CRP (mg/L)	1.3 (0.6 - 2.8)	1.0 (0.5 - 2.3)	1.1 (0.5 - 2.6)	1.3 (0.7 - 3.1)	1.6 (0.8 - 3.3)	< 0.001
Cholesterol (mmol/L)	5.4 ± 1.0	5.2 ± 1.0	5.4 ± 1.0	5.4 ± 1.0	5.6 ± 1.1	< 0.001
Glucose (mmol/L)	4.8 (4.4 - 5.3)	4.8 (4.4 - 5.3)	4.8 (4.4 - 5.3)	5.0 (4.5 - 5.4)	5.1 (4.6 - 5.6)	< 0.001
Creatinine (umol/L)	93.0 ± 18.9	92.7 ± 17.6	91.8 ± 14.3	93.0 ± 15.7	94.5 ± 25.8	0.001
eGFR (ml/min/1.73m ²)	88.4 ± 17.9	88.5 ± 18.6	90.2 ± 17.8	88.8 ± 17.0	86.1 ± 17.9	0.001
<60 (%)	6.8	7.0	6.6	5.5	8.3	0.46
UAE (mg/24h)*	9.5 (6.6 - 19.2)	9.2 (6.3 - 18.3)	8.7 (6.3 - 18.4)	9.5 (6.7 - 17.7)	10.8 (7.1 - 22.9)	< 0.001

For abbreviations, see Table 1.

Table S1B. Baseline characteristics according to ferritin quartiles in women.

Variables	Total	Q1	Q2	Q3	Q4	P - value for trend
<i>n</i>	3241	837	811	785	808	NA
Ferritin ($\mu\text{g/L}$) min-max	61 (3 - 1414)	17 (3 - 30)	45 (31 - 61)	83 (62 - 114)	166 (114 - 1414)	NA
Age (years)	52.2 \pm 11.6	45.5 \pm 9.0	49.6 \pm 10.7	54.2 \pm 11.2	59.9 \pm 10.0	<0.001
Menstruation (%)	47.4	80.8	57.3	35.3	14.3	<0.001
BMI (kg/m^2)	26.6 \pm 4.9	25.4 \pm 4.6	25.8 \pm 4.5	26.7 \pm 4.8	28.4 \pm 4.9	<0.001
>30 (%)	20.0	12.6	14.9	20.1	32.7	<0.001
Systolic BP (mmHg)	122.1 \pm 18.8	119.1 \pm 15.2	119.1 \pm 17.9	123.1 \pm 19.2	129.6 \pm 20.1	<0.001
Heart rate (b.p.m.)	69.7 \pm 9.5	69.5 \pm 9.1	68.9 \pm 9.4	70.5 \pm 9.8	69.9 \pm 9.7	0.026
LVH according to Cornell (%)	1.3	0.6	1.6	0.8	2.6	0.002
LBBS (%)	2.4	1.4	1.9	2.3	4.0	0.001
Medical history (%)						
Smoking or quit < 1 year	30.4	27.1	33.9	30.6	30.1	0.43
MI	1.3	0.8	0.7	1.2	2.4	0.005
Stroke	0.7	0.7	1.0	0.6	0.6	0.63
Type 2 diabetes	6.9	3.9	4.8	5.5	13.3	<0.001
Hypercholesterolemia	24.5	11.6	21.1	27.5	38.3	<0.001
Hypertension	27.3	14.5	20.0	29.1	46.0	<0.001
Anemia	13.5	30.3	9.6	6.7	6.4	<0.001
Laboratory measurements						
Hemoglobin (g/dL)	13.0 \pm 1.0	12.4 \pm 1.1	13.0 \pm 0.9	13.2 \pm 0.8	13.4 \pm 1.0	<0.001
MCV (fl)	90.1 \pm 4.8	88.0 \pm 5.7	90.7 \pm 4.1	90.8 \pm 4.0	91.2 \pm 4.5	<0.001
Hepcidin (nM/L)	2.4 (1.2 - 4.2)	0.7 (0.3 - 1.2)	2.0 (1.4 - 2.7)	3.2 (2.3 - 4.3)	5.2 (3.6 - 7.1)	<0.001
Serum iron ($\mu\text{mol/L}$)	15.5 \pm 5.7	13.1 \pm 6.3	15.5 \pm 5.7	15.7 \pm 5.0	16.0 \pm 5.1	<0.001
Transferrin saturation (%)	23.1 \pm 9.2	18.3 \pm 9.2	23.9 \pm 8.6	24.6 \pm 8.2	25.9 \pm 8.9	<0.001
Erythropoietin (IU/L)	8.0 (6.0 - 10.7)	9.3 (6.8 - 13.4)	7.8 (5.8 - 10.0)	7.5 (5.9 - 10.0)	7.8 (5.8 - 9.9)	<0.001
hs-CRP (mg/L)	1.4 (0.6 - 3.3)	0.9 (0.4 - 2.2)	1.1 (0.5 - 2.9)	1.6 (0.8 - 3.2)	2.2 (1.0 - 4.3)	<0.001
Cholesterol (mmol/L)	5.4 \pm 1.1	5.0 \pm 1.0	5.4 \pm 1.0	5.6 \pm 1.0	5.8 \pm 1.1	<0.001
Glucose (mmol/L)	4.7 (4.4 - 5.2)	4.6 (4.3 - 5.0)	4.6 (4.3 - 5.0)	4.7 (4.4 - 5.2)	5.0 (4.5 - 5.6)	<0.001
Creatinine ($\mu\text{mol/L}$)	76.8 \pm 15.5	75.9 \pm 10.5	75.9 \pm 11.6	76.8 \pm 11.5	78.5 \pm 24.1	0.007
eGFR (ml/min/1.73m^2)	90.9 \pm 17.5	95.9 \pm 16.3	93.7 \pm 17.0	89.0 \pm 16.9	84.7 \pm 17.7	<0.001
<60 (%)	4.6	1.9	3.1	5.2	8.4	<0.001
UAE (mg/24h)*	7.7 (5.7 - 12.4)	7.6 (5.6 - 11.7)	7.4 (5.7 - 11.3)	7.4 (5.6 - 11.3)	8.7 (5.9 - 15.8)	<0.001

For abbreviations, see Table 1.

Table S2. Univariable regression beta-coefficients between cardiovascular risk factors and ferritin in men and women.

Variables	Men			Women		
	Standardized β	T	P - value	Standardized β	T	P - value
Age (per 10 years)	0.008	0.3	0.74	0.447	21.1	< 0.001
Menstruation (yes vs. no)	-	-	-	-0.508	-24.5	< 0.001
BMI (per kg/m ²)	0.230	9.6	< 0.001	0.229	10.0	< 0.001
Systolic BP (per 5 mmHg)	0.103	4.3	< 0.001	0.214	8.1	< 0.001
Heart rate (per 5 b.p.m.)	0.061	2.5	0.013	0.046	2.0	0.05
LVH (yes vs. no)	-0.023	-0.9	0.38	0.078	4.9	< 0.001
LBBB (yes vs. no)	0.009	0.3	0.74	0.094	4.6	< 0.001
Smoking (yes vs. no)	0.029	1.2	0.22	0.062	2.8	0.006
MI (yes vs. no)	-0.068	-2.4	0.017	0.056	3.6	< 0.001
Stroke (yes vs. no)	-0.018	-0.5	0.98	-0.002	-0.2	0.82
Hemoglobin (per g/dL)	0.223	8.1	< 0.001	0.407	16.5	< 0.001
MCV (per fL)	0.165	6.5	< 0.001	0.241	9.9	< 0.001
TSAT (per 5 %)	0.265	10.0	< 0.001	0.342	13.0	< 0.001
Hepcidin (per doubling)	0.742	41.3	< 0.001	0.822	56.9	< 0.001
EPO (per doubling)	-0.103	-3.8	< 0.001	-0.258	-9.0	< 0.001
hs-CRP (per doubling)	0.172	6.6	< 0.001	0.269	11.8	< 0.001
Cholesterol (per mmol/L)	0.143	5.6	< 0.001	0.254	11.4	< 0.001
Glucose (per doubling)	0.115	4.4	< 0.001	0.163	7.3	< 0.001
eGFR (per 5ml/min/1.73m ²)	-0.039	-1.7	0.10	-0.099	-4.4	< 0.001
UAE (per doubling)	0.034	1.8	0.08	0.043	2.1	0.033

For abbreviations, see Table 1.

Table S3. Multivariable regression beta-coefficients between cardiovascular risk factors and serum ferritin in men and women without prior myocardial infarction and/or stroke (selected after bootstrapping).

Variables	Men (n = 2969)			Women (n = 3178)		
	Standardized β	T	P - value	Standardized β	T	P - value
Menstruation (yes vs. no)	-	-	-	-0.127	-7.2	< 0.001
BMI (per kg/m ²)	0.086	5.0	< 0.001	-	-	-
Smoking	-	-	-	-0.030	-2.2	0.028
Hemoglobin (per g/dL)	0.106	6.1	< 0.001	0.071	4.0	< 0.001
MCV (per fL)	0.097	5.6	< 0.001	0.122	7.4	< 0.001
Log 2 ferritin (ug/L)	0.679	37.4	< 0.001	0.728	30.4	< 0.001
TSAT (per 5%)	0.094	4.8	< 0.001	0.016	5.9	< 0.001
Log 2 erythropoietin (IU/L)	-	-	-	-0.064	-3.9	0.044
Cholesterol (per mmol/L)	0.045	2.51	0.012	-	-	-
Log 2 hs-CRP (mg/L))	-	-	-	0.079	5.4	< 0.001
Log 2 glucose (mmol/L)	0.015	0.9	0.001	0.053	4.8	< 0.001
	R² total model		0.59	R² total model		0.75

For abbreviations, see Table 1.

Table S4. Ferritin and hepcidin levels and incident heart failure with death as competing risk in subjects without prior myocardial infarction and/or stroke.

Ferritin (per doubling)	sHR	95% CI	P - value
Total population (n = 6147)¹			
Unadjusted	1.49	1.28-1.74	< 0.001
Model 1	1.30	1.07-1.56	0.007
Model 2	1.09	0.91-1.32	0.345
Men (n = 2969)			
Unadjusted	1.06	0.81-1.38	0.678
Model 1	1.06	0.83-1.34	0.657
Model 2	0.89	0.68-1.15	0.359
Women (n = 3178)			
Unadjusted	2.04	1.61-2.59	< 0.001
Model 1	1.54	1.19-2.00	0.001
Model 1A	1.71	1.28-2.27	< 0.001
Model 2	1.32	1.03-1.71	0.033
Model 3	1.33	1.05-1.85	0.026
Hepcidin (per doubling)	sHR	95% CI	P - value
Total population (n = 6147)²			
Unadjusted	1.57	1.32-1.87	< 0.001
Model 1	1.29	1.05-1.58	0.012
Model 2	1.11	0.92-1.36	0.259
Men (n = 2969)			
Unadjusted	1.09	0.82-1.42	0.598
Model 1	1.05	0.82-1.35	0.695
Model 2	0.90	0.72-1.11	0.321
Women (n = 3178)			
Unadjusted	2.09	1.64-2.68	< 0.001
Model 1	1.60	1.19-2.15	0.002
Model 1A	1.73	1.23-2.45	0.002
Model 2	1.41	1.04-1.90	0.027
Model 3	1.41	1.07-2.23	0.019

Abbreviations: CI = Confidence interval, sHR = Sub-hazard ratio.

1P for interaction ferritin, sex and new onset heart failure = 0.046.

2P for interaction hepcidin, sex and new onset heart failure = 0.034.

Model 1 is adjusted for age (per 10 years).

Model 1A is adjusted for menstrual state instead of age.

Model 2 is adjusted for model 1 + cardiovascular and heart failure risk factors: BMI, current smoking, the presence of diabetes, hypertension, hypercholesterolemia, LVH, renal function, and levels of hemoglobin, erythropoietin, hs-CRP and UAE.

Model 3 is adjusted for model 2 + time-varying incident cardiovascular events.

57

Table S5. Hepcidin and ferritin levels and incident heart failure with death as competing risk in postmenopausal women.

Postmenopausal women	sHR	95% CI	P - value
Ferritin (per doubling)			
Unadjusted	1.75	1.29-2.36	< 0.001
Model 1	1.57	1.18-2.09	0.002
Model 2	1.34	1.03-1.78	0.040
Model 3	1.30	1.02-1.69	0.043
Hepcidin (per doubling)			
Unadjusted	1.64	1.15-2.35	0.008
Model 1	1.52	1.08-2.12	0.017
Model 2	1.31	0.95-1.83	0.10
Model 3	1.48	0.99-2.21	0.06

Abbreviations: CI = Confidence interval, sHR = Sub-hazard ratio.

Model 1 is adjusted for age (per 10 years).

Model 2 is adjusted for model 1 + cardiovascular and heart failure risk factors: BMI, current smoking, the presence of diabetes, hypertension, hypercholesterolemia, history of MI or stroke, LVH, renal function, and levels of hemoglobin, erythropoietin, hs-CRP and UAE.

Model 3 is adjusted for model 2 + time-varying incident cardiovascular events.

Table S6. Levels of ferritin and cause-specific risk for heart failure with a reduced or preserved ejection fraction in women.

Ferritin (per doubling)	HR	95% CI	P - value	P - int
HFrEF (n = 37)				
Unadjusted	1.96	1.55 - 2.48	< 0.001	0.28
Model 1	1.48	1.14 - 1.90	0.003	0.28
Model 2	1.25	0.97 - 1.62	0.09	0.17
Model 3	1.27	0.98 - 1.64	0.07	0.17
HFpEF (n = 42)				
Unadjusted	2.06	1.64 - 2.59	< 0.001	NA
Model 1	1.55	1.21 - 2.00	0.001	NA
Model 2	1.34	1.04 - 1.71	0.022	NA
Model 3	1.36	1.06 - 1.73	0.013	NA

Abbreviations: CI = Confidence interval, HR = Hazard ratio, NA = Not applicable.

Model 1 is adjusted for age (per 10 years).

Model 2 is adjusted for model 1 + cardiovascular and heart failure risk factors: BMI, current smoking, the presence of diabetes, hypertension, hypercholesterolemia, history of MI or stroke, LVH, renal function, and levels of hemoglobin, erythropoietin, hs-CRP and UAE.

Model 3 is adjusted for model 2 + time-varying incident cardiovascular events.




Figure S1. ROC curves and areas under the curve of serum iron, transferrin saturation and ferritin in predicting new onset heart failure (A), cardiovascular events (B), cardiac events (C) and all-cause mortality (D).




Figure S2. Ferritin levels and risk for new onset heart failure among subgroups in women.




Figure S3. Total annual incidence of cardiovascular events (A + B) and all-cause mortality (C + D) according to ferritin and hepcidin quartiles and stratified by sex. ACM = All-cause mortality; CV = Cardiovascular.

**NOVEL INSIGHTS ON IRON
DEFICIENCY IN HEART
FAILURE**

