



University Medical Center Groningen

University of Groningen

## Somatic monitoring of patients with mood and anxiety disorders

Simoons, Mirjam

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

### *Document Version*

Publisher's PDF, also known as Version of record

### *Publication date:*

2018

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Simoons, M. (2018). Somatic monitoring of patients with mood and anxiety disorders: Problem definition, implementation and further explorations. [Groningen]: Rijksuniversiteit Groningen.

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# 8

## LIMITED EVIDENCE FOR RISK FACTORS FOR PROARRHYTHMIA AND SUDDEN CARDIAC DEATH IN PATIENTS USING ANTIDEPRESSANTS: DUTCH CONSENSUS ON ECG MONITORING

Mirjam Simoons\*  
Adrie Seldenrijk\*  
Hans Mulder  
Tom K. Birkenhäger  
Mascha M. Groothedde-Kuyvenhoven  
Rob M. Kok  
Cornelis Kramers  
Wim Verbeeck  
Mirjam Westra  
Eric N. van Roon  
P. Roberto Bakker#  
Henricus G. Ruhé#

\* These authors share first authorship  
# These authors share senior authorship

## ABSTRACT

Currently, there is a lack of international and national guidelines or consensus documents with specific recommendations for electrocardiogram (ECG) screening and monitoring during antidepressant treatment. To make a proper estimation of the risk of cardiac arrhythmias and sudden (cardiac) death during antidepressant use, both the drug and patient-specific factors should be taken into account; however, solid evidence on how this should be done in clinical practice is lacking. Available recommendations on the management of QT(c)-prolongation (with antidepressant treatment) emphasize that special attention should be given to high-risk patients; however, clinicians are in need of more concrete suggestions about how to select patients for ECG screening and monitoring.

Based on a review of the literature, a Dutch multidisciplinary expert panel aimed to formulate specific guidelines to identify patients at risk for cardiac arrhythmias and sudden death by developing a consensus statement regarding ECG screening before, and monitoring during, antidepressant use.

We first reviewed the literature to identify the relative risks of various risk factors on cardiac arrhythmia and sudden (cardiac) death during antidepressant use. These relative contributions of risk factors could not be determined since no systematic reviews or meta-analyses quantitatively addressed this topic. Because evidence was insufficient, additional expert opinion was used to formulate recommendations. This resulted in readily applicable recommendations for clinical practice for selection of high-risk patients for ECG screening and monitoring. ECG screening and monitoring is recommended before and following the start of QTc-prolonging antidepressants in the presence of vulnerability to QTc-prolongation or two or more risk factors (age >65 years, female sex, concomitant use of a QTc-prolonging drug or concomitant use of a drug that influences the metabolism of a QTc-prolonging drug, cardiac disease, excessive dosing and specific electrolyte disturbances).

## INTRODUCTION

The effect of psychotropic drugs on cardiac repolarization has increasingly gained attention in research and clinical practice over the last two decades. The absolute risk of cardiac arrhythmia, such as Torsade de Pointes (TdP), is generally low (fourteen per 10,000 patients over one year (95% confidence interval 11-17/10,000)) and sudden cardiac death as a consequence of all cardiac arrhythmias in general occurs even more rarely.<sup>1</sup> However, the tragic anecdotes of physically healthy patients encountering cardiac arrest and sudden (cardiac) death after the use of psychotropic drugs have underscored that some of these drugs may increase the risk of arrhythmias. This proarrhythmic effect is often marked by a prolongation of the QT-interval/QTc-interval (QT-interval corrected for heart rate) on an electrocardiogram (ECG).<sup>2</sup>

Of the psychotropic drugs, antipsychotics are well known for their QT(c)-prolonging effects and association with TdP and sudden cardiac death, although the available evidence may not support this reputation per se.<sup>3</sup> The incidence rate of sudden cardiac death in users of antipsychotics was 2.9 per 1000 patient-years – a significantly doubled risk compared with (non-psychiatric) non-users.<sup>3</sup> Some antidepressants have proven to also prolong the QT(c)-interval. Although a recent meta-analysis showed significant QTc-prolongation by tricyclic antidepressants (TCAs; doxepin, nortriptyline and amitriptyline) and some selective serotonin reuptake inhibitors (SSRIs; citalopram, escitalopram and sertraline) relative to placebo<sup>4</sup>, CredibleMeds, the internationally renowned source for evidence-based classification of drugs according to their QTc-prolonging abilities, only classifies citalopram and escitalopram as antidepressant drugs with known risk of TdP and clomipramine, desipramine, imipramine, nortriptyline, mirtazapine, trimipramine and venlafaxine as antidepressants with a possible risk of TdP.<sup>5</sup> Moreover, the United States Food and Drug Administration (FDA) has issued several drug safety communications, including for citalopram in 2011 and 2012. The warnings stated that citalopram use could lead to abnormal heart rhythms and prescription doses should not exceed 40 mg per day in adults and 20 mg in patients >65 years of age.<sup>6,7</sup>

To make a proper estimation of the risk of arrhythmia, the combination of characteristics of the antidepressant therapy and patient-specific factors should be taken into account. A number of risk factors that add to the arrhythmia risk, including non-cardiac risk factors, have been proposed, including female sex, older age, (ischemic) heart disease or a history thereof, electrolyte disturbances (including hypomagnesaemia, hypokalaemia and hypocalcaemia), pharmacokinetic and pharmacodynamic genetic factors, congenital long QTc-syndrome and a range of other medical conditions.<sup>8-11</sup> Several studies aimed to verify and synthesize the available evidence for QTc-prolongation into a risk score for use in non-psychiatric hospitalized patients.<sup>12,13</sup> In addition, it has been shown that the risk for arrhythmia increases with increasing numbers of such risk factors<sup>8-10</sup>; however, the relative risks of the individual risk factors for the outcome arrhythmia are still unclear.

Apart from the ECG monitoring recommendations in the Summary of Product Characteristics (SmPC) for individual QT(c)-prolonging antidepressant drugs, there are no clear and concrete national or international guidelines on if, when, and how often an ECG should be performed before and during treatment with antidepressant drugs. In the Dutch multidisciplinary guideline for depression, only with TCA treatment in elderly patients is an ECG recommended before the start of treatment (although this is to rule out contraindications such as a right bundle branch block to assist in drug choice, which is not the focus of this consensus document) and ECG monitoring with nortriptyline treatment in elderly with cardiac risk.<sup>14</sup> Although some of the available international depression guidelines do not mention ECG monitoring with antidepressant treatment<sup>15,16</sup>, other depression guidelines and some consensus documents for the management of QT(c)-prolongation emphasize that special attention should be given to 'high-risk' patients in order to prevent unfavourable cardiac outcomes.<sup>8,10,17-19</sup> For example, in a consensus document, Dodd and colleagues suggest ECG monitoring during the use of TCAs.<sup>20</sup> They also recommend considering ECG monitoring during SSRI and serotonin noradrenaline reuptake inhibitor (SNRI) treatment in 'high-risk' individuals, although they consider it usually unnecessary.<sup>20</sup> However, how 'high-risk' must be quantified in clinical practice remains an enigma that hampers implementation of such recommendations in clinical practice.

Therefore, we aimed to answer the following important, unanswered questions in order to prevent cardiac arrhythmia and sudden (cardiac) death during antidepressant use: 1) should the ECG be monitored with antidepressant treatment, and 2) if so, for which patients (with which risk factors), when, and how often?

## METHODS

### The multidisciplinary expert panel

In 2015, the Dutch Network for Quality Development in Mental Health Care funded the development of recommendations about the prevention, monitoring, and treatment of side effects of psychotropic drugs. A multidisciplinary expert panel for antidepressant drugs addressed the association between the use of antidepressants and proarrhythmic effects and related ECG monitoring issues. The expert panel for antidepressants consisted of four psychiatrists (two being specialized in the treatment of children/adolescents or elderly patients, one additionally in training as a clinical pharmacologist), a general practitioner, an internist-clinical pharmacologist, a nursing specialist, a patient representative, three (hospital) pharmacists, and a postdoctoral researcher.

### Literature search

In order to retrieve relevant literature as a base for our recommendations, two authors (AS and MS) searched for studies addressing the relative risks of risk factors (such as older age and female sex) for cardiac arrhythmia and sudden (cardiac) death, associated with

antidepressant use. We did not use a restriction in the time period of publication and applied no language restrictions. For practical reasons and because of time restrictions for the project, we limited our search to systematic reviews and meta-analyses.

We conducted a search of Medline using the search strategy “‘Psychotropic Drugs’[Mesh]) AND (((Arrhythmias, Cardiac[MeSH] OR arrhythmi\*[tiab] OR proarrhythmi\*[tiab] OR long QT[tiab] OR (prolong\*[tiab] AND QT[tiab]) OR torsade de pointes[tiab] OR torsades de pointes[tiab] OR Death, Sudden, Cardiac[Mesh] OR cardiac death[tiab] OR cardiac mortality[tiab] OR ‘Cardiovascular Diseases/mortality’[Mesh]) AND (drug-induced[tiab] OR drug effects[sh] OR adverse effects[sh])) AND (Risk factors[Mesh] OR risk factors[tiab] OR prognost\*[tiab] OR predict\*[tiab])) AND (systematic\*[tiab] OR review\*[tiab] OR meta-analysis[tiab] OR Meta-Analysis[ptyp] OR systematic[sb])’”. We used the broader term ‘psychotropic drugs’ instead of antidepressant-specific search terms, because a pilot search showed insufficient studies on risk factors for cardiac arrhythmia and sudden (cardiac) death when we used antidepressants in the search. Bibliographies of retrieved studies were scanned for additional systematic reviews and meta-analyses. We performed the last update of the search on 14 March 2017.

### Selection criteria

The study selection was performed independently by two authors (AS and MS) and discrepancies were resolved through discussion. We selected systematic reviews and meta-analyses of studies in (older) adult patients using psychotropic drugs registered in The Netherlands. Papers limited to overdose, intoxication, supratherapeutic dosage or antidepressants as add-on intervention were excluded. In our selection, we focused on cardiac arrhythmia and sudden (cardiac) death as outcomes and excluded publications solely on QTc-interval as a surrogate measure for cardiac arrhythmia and sudden (cardiac) death. The latter was applied because QTc intervals are of limited value due to the use of different formulas to correct for heart-rate (e.g. Bazett or Fridericia) and because the relationship between drug-induced QTc-prolongation and the likelihood of arrhythmia appears to be, at best, modest and neither linear nor straightforward.<sup>21,22</sup> Eligible studies should report on the risk for cardiac arrhythmia and sudden (cardiac) death in association with risk factors (e.g. age, sex, cardiac disease, electrolyte disturbances, prolonged QT(c)-interval (syndrome), use of QT(c)-prolonging drugs).

### Formulation of recommendations

In order to formulate recommendations, the multidisciplinary expert panel received a summary of extracted data from selected studies. The odds ratios, relative risks and/or prevalences of cardiac arrhythmia and sudden (cardiac) death associated with the investigated risk factors were presented for each review/meta-analysis. This evidence served as input for discussion on the indication and timing for ECG screening and monitoring at treatment initiation and during use of antidepressants. When consensus was reached, each recommendation was graded evidence level 1-4 according to the Dutch

criteria for evidence-based guideline development (EBRO), based on the Appraisal of Guidelines Research and Evaluation (AGREE) Collaboration.<sup>23-25</sup>

## RESULTS OF THE LITERATURE SEARCH

We retrieved 100 publications of potential systematic reviews and/or meta-analyses (see Supplementary Material). Seven titles and abstracts appeared to match our inclusion criteria; all seven studies were in English. Based on their full-text, four publications met our selection criteria.<sup>26-29</sup> The major reasons for exclusion of publications were failure to meet our population criterion, no investigation of risk factors or solely outcomes other than cardiac arrhythmia and sudden (cardiac) death.

In a systematic review of TdP cases, Meyer-Massetti et al. investigated the prevalences of a predefined set of risk factors (age, sex, dose, electrolyte imbalance, cardiac disease, concomitant pro-arrhythmic drugs and other drugs influencing cardiac function and baseline QTc) in 54 patients with intravenous administration of the antipsychotic haloperidol.<sup>27</sup> Vieweg et al. provided prevalences of various risk factors in only a small number of TdP cases (n=4) associated with the antipsychotic risperidone.<sup>28</sup> Because of the low number of cases and the focus on antipsychotics, the results of these studies were not generalizable to patients using antidepressants and these two systematic reviews were therefore excluded.

Zeltser et al. performed a systematic review of TdP cases and described the prevalence of six risk factors for arrhythmia in 249 cases of TdP induced by non-cardiac drugs.<sup>29</sup> In brief, 70 of these cases (28.1%) were caused by psychotropic drugs, mainly, but not exclusively, by antipsychotics (not further specified). Of the psychotropic drug-induced TdP cases, 71.4% were female, 44.7% used additional drugs that caused drug interactions (i.e. impairs the metabolism of QT-prolonging drugs or concomitant use of two or more QT-prolonging drugs), 43.1% had existing cardiac disease, 27% used an excessive dose (leading to drug toxicity but excluding cases of suicidal overdose), 17.9% had hypokalaemia, and 17.1% had a vulnerability to QT-prolongation (familial history of long QT-syndrome, history of drug-induced TdP, prolonged QT-interval before drug administration). Cases using psychotropic drugs had, on average, 2.2 risk factors. No odds ratios for the different risk factors were given.

A study by Åström-Lilja et al. investigated the prevalence of a small set of risk factors for arrhythmia in drug-induced TdP.<sup>26</sup> Since Zeltser et al. did not assess age as a risk factor, the study by Åström-Lilja et al. was considered of substantial added value for the expert panel, despite the fact that it was not a systematic review or meta-analysis. This series of 88 cases was based on data from the Swedish pharmacovigilance database, which was not restricted to psychotropic drugs (8/88 (9%) used antidepressants). The age of the TdP cases ranged from 15 to 90 years, with the median age being 74 years. Seventy-two percent of the TdP cases were over 65 years of age. Existing heart disease, female sex and hypokalaemia were present in 90%, 70% and 12% of cases, respectively, while two or more established risk factors were present in 85% of cases (75/88).

## CONSIDERATIONS OF THE EXPERT PANEL

In order to translate the available evidence to recommendations for daily clinical practice, we addressed several issues.

First, the expert panel concluded that the relative contributions of risk factors to cardiac arrhythmias and sudden (cardiac) death during antidepressant use could not be determined since no systematic reviews or meta-analyses addressed this topic specifically.

Second, the absence of a reference group (e.g. drug-free TdP cases or patients using psychotropic drugs that did not develop TdP) in the included reports hampered us to put the prevalences of risk factors into perspective and/or calculate odds ratios or relative risks. Therefore, it is impossible to draw firm conclusions on the relative importance of certain factors or the definition of a high-risk population for ECG screening and monitoring.

Third, we took the average time to steady-state into account when drafting the advice about the timing of the follow-up ECG after reaching the target dose of treatment with a QTc-prolonging antidepressant. Time to steady state is four to five times the half-life of the drug, which is approximately one week for antidepressant drugs.

Fourth, the expert panel would like to point out that the current literature is inconclusive about the intraindividual circadian variation in the length of the QTc-interval, which may range from less than 10 ms to up to 75-100 ms.<sup>21,30,31</sup> It is therefore preferable to register ECG recordings at fixed time points during the day in order to avoid bias in the change between two subsequent ECGs as a result of circadian variation.

Fifth, because threshold values for a prolonged QTc-interval, such as those of the AHA/ACCF/HRS, are the result of consensus, the expert panel suggested to consider a similar threshold value for women and men (i.e. >450 ms, instead of >450 ms for men and >460 ms for women<sup>32</sup>), which will reduce complexity for clinical practice. This is supported by the consideration, that women are at increased risk for QTc-prolongation and TdP; lowering the threshold for a prolonged QTc-interval to >450 ms in women also, will result in earlier identification than with >460 ms.<sup>33</sup>

Last, the expert panel noted that because of the actions of sex hormones on the QTc-interval, the difference between women and men in the prolonged QTc-interval risk is disappears during menopause.<sup>34,35</sup>

## RECOMMENDATIONS

Based on the limited literature results and the consensus reached by our expert panel, we formulated the following recommendations, all graded 4 – expert opinion (Figure 1).<sup>23-25</sup>

- ECG screening and monitoring is recommended for antidepressants with a known or possible risk of TdP according to CredibleMeds (citalopram, clomipramine, desipramine, escitalopram, imipramine, mirtazapine, nortriptyline, trimipramine and venlafaxine<sup>5</sup>) if the patient
  1. is having a known risk of QTc-prolongation:



- › known prolonged QTc-interval;
  - › history of TdP;
  - › family history of long QTc-syndrome or sudden cardiac death;
- and choosing an antidepressant without QTc-prolonging ability is not possible;  
*Remark: if the risk of QTc-prolongation cannot be determined, it can be considered absent for decision making purposes.*

or

2. has two or more of the risk factors listed below (based on the two reviews from the literature search discussed above<sup>26,29</sup>).

- › age over 65 years;
- › female sex;
- › concomitant use of a QTc-prolonging drug (list of drugs with a known risk of TdP according to CredibleMeds<sup>5</sup>) or concomitant use of a drug that influences the metabolism of the antidepressant with a known or possible risk of TdP (i.e. citalopram, clomipramine, desipramine, escitalopram, imipramine, mirtazapine, nortriptyline, trimipramine and venlafaxine<sup>5</sup>);
- › cardiac disease (myocardial infarction, heart failure, valvulopathy, cardiomyopathy);
- › excessive dosing (higher than highest dose according to SmPC, or standard dose with relevant kidney or liver problem);
- › specific electrolyte disturbances (hypocalcaemia, hypokalaemia, hypomagnesemia).<sup>26,29</sup>

*Additional remark: in case of a strong suspicion of electrolyte disturbances, e.g. with alcoholism, anorexia nervosa, diarrhoea, the use of loop diuretics, etc., the calcium, potassium and magnesium serum level should be quantified.*

- Paroxetine, duloxetine and fluoxetine do not affect the QTc-interval in comparison to placebo, while fluvoxamine shortens the QTc-interval compared to placebo.<sup>4</sup> For other antidepressants, evidence for determining their QTc-prolonging ability is currently insufficient.
  - In general, it is unnecessary to make an ECG prior to starting with any antidepressant if the patient is not at risk of QTc-prolongation and all of the abovementioned risk factors for cardiac arrhythmia or conduction disorders are absent.
  - ECG screening and monitoring should consist of an ECG before treatment initiation and one week after reaching the target dose of a the QTc-prolonging antidepressant (i.e. when steady-state has been reached).
  - In case the ECG shows a QTc-interval of 450 ms or above, it is advisable to consult a cardiologist.
- Remark: although we focus on the QTc-interval in this consensus document, the PQ- and QRS-intervals may also be relevant during psychotropic drug use.*
- If needed, for example in acute situations or highly severe mentally ill cases, the treating physician can deviate from the recommended ECG at treatment initiation.




Figure 1. Decision tree for ECG monitoring with antidepressant treatment. <sup>a</sup> Based on CredibleMeds.org<sup>5</sup>, which represents available and evolving evidence that is constantly re-evaluated when new evidence becomes available; <sup>b</sup> If the risk of QTc-prolongation cannot be determined, it can be considered absent for decision making purposes; <sup>c</sup> In case of a strong suspicion of electrolyte disturbances, e.g. with alcoholism, anorexia nervosa, diarrhoea, use of loop diuretics, etc., the calcium, potassium and magnesium serum level should be quantified. QTc QT-interval corrected for heart rate; ECG electrocardiogram; TdP Torsade de Pointes.

## DISCUSSION

In this Dutch consensus document on ECG screening and monitoring in patients using antidepressants, no need for ECG screening and monitoring is recommended in patients without QTc-prolongation vulnerability and other risk factors for cardiac arrhythmia during antidepressant therapy. ECG screening and monitoring is recommended before and following the start of antidepressants with a known or possible risk of TdP (i.e. citalopram, clomipramine, desipramine, escitalopram, imipramine, mirtazapine, nortriptyline, trimipramine and venlafaxine<sup>5</sup>) in the presence of vulnerability to QTc-prolongation or two or more risk factors (age >65 years, female sex, concomitant use of a QTc-prolonging

drug or concomitant use of a drug that influences the metabolism of the QTc-prolonging antidepressant, cardiac disease, excessive dosing and specific electrolyte disturbances).

Some authors, guidelines and drug labels state that all patients receiving QTc-prolonging psychotropic medication should be monitored, but most experts and authors of previous reviews emphasize that ECG screening and monitoring is necessary only in high risk patients.<sup>20,22,36</sup> Current guidelines do not provide uniform recommendations with respect to ECG screening and monitoring in patients using antidepressants. Unfortunately, there is a lack of evidence that ECG screening and monitoring indeed can prevent cases of arrhythmia or sudden cardiac death. For this, the 'number needed to ECG' (NNE) would be an interesting number to indicate the number of patients that should be monitored (with consecutive ECGs) to prevent one additional death or adverse event due to QTc-prolongation. Given the rising costs of (mental) health, the apparent pressure to increase productivity and high administrative workload, it would also be appropriate to perform a proper health economical cost-benefit assessment before issuing general recommendations on ECG screening and monitoring. An optimized balance between costs and yield of ECG screening and monitoring to detect aberrances would support clinicians in their treatment decisions, while unnecessary ECGs could be eliminated as much as possible.

Because of a relative lack of systematic reviews and meta-analyses that compare risk factors (despite a number of studies on risk scores for QTc-prolongation), the relative risks of risk factors for cardiac arrhythmia and sudden (cardiac) death cannot be established/quantified, which hampers a better determination of a 'high-risk' population. Therefore, we decided to consider the various risk factors that were put forward in two earlier reviews of TdP cases.<sup>26,29</sup> The contribution of these risk factors to cardiac arrhythmia and sudden (cardiac) death have not yet been quantified. However, the results of our literature study are corroborated by a recent systematic review by Vandael et al. of large randomized controlled trials and observational studies that assessed the level of evidence for several factors to increase the risk of QTc-prolongation in a general population.<sup>37</sup> Although strong evidence was found for a few risk factors (including hypokalaemia and use of drugs with known risk of TdP as listed by CredibleMeds<sup>5</sup>), little or no evidence was found for many other risk factors.<sup>37</sup>

Based on the identified risk factors, Vandael et al. subsequently aimed at developing a risk score to identify patients at low/high risk for QTc-prolongation in an observational study in hospital patients with a first prescription for haloperidol or a QTc-prolonging antibiotic or antimicrobial.<sup>38</sup> The RISQ-PATH index, using more and some other risk factors than we mention in this paper, was able to exclude low-risk patients from further ECG follow-up when starting QTc-prolonging drugs (high sensitivity (96.2%)), but also resulted in many false positives (low specificity (32.9%)).<sup>38</sup> Because the aim of a risk score would be to safely exclude patients with low risk from the total population, this tool seems a promising instrument. The RISQ-PATH index is similar to our recommendations for ECG screening and monitoring, except that it additionally contains a weighing of risk factors

based on the level of evidence – not relative risk - found in their systematic review. Instead of a risk score, we chose the dichotomous cut-off of two risk factors as a threshold, based on the evidence of a higher risk of cardiac arrhythmia outcomes with an increasing number of risk factors, the average number of risk factors found in the two reviews from our literature study, the general low absolute risk of our primary outcomes and approximation of the same high sensitivity (i.e. yield of 'high risk' cases for cardiac arrhythmia and/or sudden (cardiac) death vs. costs and logistics of obtaining ECGs in mental health). Further research is warranted to quantify the relative risks of each risk factor and the sensitivity/specificity of a risk score in people prescribed an antidepressant before implementation of such a weighted risk index in clinical practice. To obtain efficacy and (cost-)effectiveness data, ideally a randomized controlled trial is performed to assign patients to be monitored according to their risk score versus treatment as usual. This would also enable to determine an NNE to guide future ECG screening and monitoring guidelines. However, given the low prevalence of QTc-prolongation and outcomes like cardiac arrhythmia and sudden death, performing such an RCT requires large numbers of patients and resources. Therefore a non-randomized approach as used by Vandael et al. might be most feasible.<sup>38</sup>

For implementation, the counting of risk factors to determine the need for ECG screening and monitoring would ideally be incorporated in automated clinical decision support systems that alerts prescribers to obtain ECG screening and monitoring.<sup>39</sup> The combination of information from electronic medical records and an electronic prescribing system would be supportive in preventing undesirable outcomes of QTc-prolonging drugs and increased efficacy of ECG screening and monitoring in specifically high-risk patients.<sup>39</sup>

### Strength and limitations

The strength of this paper are the clear recommendations to select 'high risk patients' for ECG screening and monitoring when prescribing a QTc-prolonging antidepressant, which are easily applicable in clinical practice. However, some critical issues must be addressed.

The first limitation of our work is the restriction of our search to systematic reviews and meta-analyses, which provided a summary of available case-series but does not exclude the possibility that reports such as additional case-series or cohort studies might have been missed. Furthermore, we could only identify papers on patients using psychotropic drugs in general, without a focus on antidepressants. We loosened our restriction to include reports on antidepressants only after our pilot search showed insufficient studies when we used 'antidepressants' as a search-term. Given our restriction of systematic reviews and meta-analyses, we think this was the best compromise to retrieve reviews on patients using antidepressants and risk factors for cardiac arrhythmia. From four studies initially selected, only two studies were informative as they included a larger number of patients also using antidepressants. One of these was not a systematic review, but was still considered relevant in the absence of additional adequate evidence from systematic reviews and meta-analyses. Our recommendations are based on the currently limited

available literature and almost entirely on expert opinion (specific for antidepressants) to make them readily applicable in clinical practice.

Second, risk factors for unfavourable cardiac (arrhythmia) outcomes may be generalizable across populations using psychotropic and other drugs. If so, a broader review of studies in patients using non-psychotropic drugs might have revealed more information, e.g. Vandael et al.<sup>37</sup> We suggest to evaluate and amend our recommendations after new relevant studies have been released, with additional practice-based experience following implementation in clinical practice.

Third, we did not include a recommendation on the need for periodic follow-up (e.g. yearly) after the baseline and follow-up ECG during antidepressant use. Because evidence on the timing of cardiac arrhythmia and sudden (cardiac) death relative to the start of antidepressant therapy is scarce, and existence of risk factors may vary with time, it cannot be expected that a normal ECG after treatment initiation will indefinitely predict a low risk of later cardiac arrhythmia outcomes. Therefore, it might be necessary to repeat ECG screening and monitoring over time, for which additional recommendations must be formulated when more data is available.

Fourth, the evidence for the QTc-prolonging abilities of psychotropic drugs is scarce with inconsistent results. For sertraline for example, results on QTc-prolongation are contradictory.<sup>40,41</sup> The initial classification based on a recent comparative systematic review and meta-analysis<sup>4</sup>, was carefully re-evaluated. Despite this systematic review, and in the absence of more specific evidence for fatal cardiac events for the remaining antidepressants on the list from this review, we decided to follow the CredibleMeds classification for our recommendations. Together with this choice, we would like to emphasize again that our recommendations are not final and may change when additional evidence appears.

Finally, ECG screening and monitoring recommendations should be actively and adequately implemented. Previous research has shown that the introduction of new guidelines, consensus statements, or (national) quality improvement programs alone has been minimally effective in improving screening and monitoring rates.<sup>42-45</sup> Specifically, compliance with these recommendations for the risk management of QTc-prolongation by antidepressants may be poor.<sup>46-48</sup>

## Conclusions

We present specific consensus recommendations for ECG screening and monitoring in patients using antidepressants. Although these recommendations are based on limited evidence in the currently scarcely available literature, the elaboration of the available evidence in combination with clinical expertise and the multidisciplinary consensus process resulted in readily applicable recommendations, which, as a next step, need to be empirically validated. Future research should evaluate these recommendations, ideally in an RCT comparing their implementation with care as usual, combined with a health

technology assessment, to assess the NNE and cost-benefit ratio. The recommendations should thereafter be evaluated and amended if necessary.

## **FUNDING**

This project was funded by the Dutch Network for Quality Development in Mental Health Care (Grants P140019 and P140040 to Henricus G. Ruhé and P. Roberto Bakker).

## REFERENCES

1. Coupland C, Hill T, Morriss R, Moore M, Arthur A, Hippisley-Cox J. Antidepressant use and risk of cardiovascular outcomes in people aged 20 to 64: cohort study using primary care database. *BMJ* 2016 Mar 22;352:i1350.
2. Frommeyer G, Eckardt L. Drug-induced proarrhythmia: risk factors and electrophysiological mechanisms. *Nat Rev Cardiol* 2016 Jan;13(1):36-47.
3. Ray WA, Chung CP, Murray KT, Hall K, Stein CM. Atypical antipsychotic drugs and the risk of sudden cardiac death. *N Engl J Med* 2009 Jan 15;360(3):225-235.
4. Beach SR, Kostis WJ, Celano CM, Januzzi JL, Ruskin JN, Noseworthy PA, et al. Meta-analysis of selective serotonin reuptake inhibitor-associated QTc prolongation. *J Clin Psychiatry* 2014 May;75(5):e441-9.
5. CredibleMeds (AZCERT I). QTDrugs Lists. 2017; Available at: <https://crediblemeds.org/new-drug-list/>. Accessed November 29th, 2017.
6. US Food and Drug Administration. FDA Drug Safety Communication: Revised recommendations for Celexa (citalopram hydrobromide) related to a potential risk of abnormal heart rhythms with high doses. 2012; Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm297391.htm>. Accessed September 28th, 2017.
7. US Food and Drug Administration. FDA Drug Safety Communication: Abnormal heart rhythms associated with high doses of Celexa (citalopram hydrobromide). 2011; Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm269086.htm>. Accessed March 2nd, 2017.
8. Drew BJ, Ackerman MJ, Funk M, Gibling WB, Kligfield P, Menon V, et al. Prevention of torsade de pointes in hospital settings: a scientific statement from the American Heart Association and the American College of Cardiology Foundation. *J Am Coll Cardiol* 2010 Mar 2;55(9):934-947.
9. Roden DM. Drug-induced prolongation of the QT interval. *N Engl J Med* 2004 Mar 4;350(10):1013-1022.
10. Ames D, Camm J, Cook P, Falkai P, Gury C, Hurley R, et al. Minimizing the risks associated with QTc prolongation in people with schizophrenia. A consensus statement by the Cardiac Safety in Schizophrenia Group. *Encephale* 2002 Nov-Dec;28(6 Pt 1):552-562.
11. Niemeijer MN, van den Berg ME, Eijgelsheim M, Rijnbeek PR, Stricker BH. Pharmacogenetics of Drug-Induced QT Interval Prolongation: An Update. *Drug Saf* 2015 Oct;38(10):855-867.
12. Anderson HN, Bos JM, Haugaa KH, Morlan BW, Tarrell RF, Caraballo PJ, et al. Prevalence and Outcome of High-Risk QT Prolongation Recorded in the Emergency Department from an Institution-Wide QT Alert System. *J Emerg Med* 2018;54(1):8-15.
13. Tisdale JE, Jaynes HA, Kingery JR, Mourad NA, Trujillo TN, Overholser BR, et al. Development and validation of a risk score to predict QT interval prolongation in hospitalized patients. *Circ Cardiovasc Qual Outcomes* 2013 Jul;6(4):479-487.
14. Spijker J, Bockting CLH, Meeuwissen JAC, Vliet van IM, Emmelkamp PMG, Hermens MLM, et al. Multidisciplinary Guideline Depression (third revision). 2013; Available at: <https://www.ggzrichtlijnen.nl/depressie>. Accessed September 28th, 2017 - Article in Dutch.
15. National Institute for Health and Care Excellence (NICE). Depression in adults: recognition and management (CG 90). 2009; Available at: <https://www.nice.org.uk/guidance/cg90/resources/depression-in-adults-recognition-and-management-pdf-975742636741>. Accessed September 28th, 2017.
16. Kennedy SH, Lam RW, McIntyre RS, Tourjman SV, Bhat V, Blier P, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical Guidelines for the Management of Adults with Major Depressive Disorder: Section 3. Pharmacological Treatments. *Can J Psychiatry* 2016 Sep;61(9):540-560.
17. Royal College of Psychiatrists. CR190: Consensus statement on high-dose antipsychotic medication. 2014; Available at: <http://www.rcpsych.ac.uk/files/pdfversion/CR190.pdf>. Accessed September 28th, 2017.

18. Cleare A, Pariante CM, Young AH, Anderson IM, Christmas D, Cowen PJ, et al. Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2008 British Association for Psychopharmacology guidelines. *J Psychopharmacol* 2015 May;29(5):459-525.
19. American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder. 2010; Available at: [https://psychiatryonline.org/pb/assets/raw/sitewide/practice\\_guidelines/guidelines/mdd.pdf](https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/mdd.pdf). Accessed November 28th, 2017.
20. Dodd S, Malhi GS, Tiller J, Schweitzer I, Hickie I, Khoo JP, et al. A consensus statement for safety monitoring guidelines of treatments for major depressive disorder. *Aust N Z J Psychiatry* 2011 Sep;45(9):712-725.
21. Beach SR, Celano CM, Noseworthy PA, Januzzi JL, Huffman JC. QTc prolongation, torsades de pointes, and psychotropic medications. *Psychosomatics* 2013 Jan-Feb;54(1):1-13.
22. Shah AA, Aftab A, Coverdale J. QTc prolongation with antipsychotics: is routine ECG monitoring recommended? *J Psychiatr Pract* 2014 May;20(3):196-206.
23. Quality institute for health care, CBO. Evidence-based guideline development (EBRO): Manual for working group members. 2007; Available at: [www.ha-ring.nl/download/literatuur/EBRO\\_handl\\_totaal.pdf](http://www.ha-ring.nl/download/literatuur/EBRO_handl_totaal.pdf). Accessed September 28th, 2017 - Article in Dutch.
24. Burgers JS, van Everdingen JJ. Evidence-based guideline development in the Netherlands: the EBRO platform. *Ned Tijdschr Geneeskd* 2004 Oct 16;148(42):2057-2059 - Article in Dutch.
25. AGREE Collaboration. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: the AGREE project. *Qual Saf Health Care* 2003 Feb;12(1):18-23.
26. Astrom-Lilja C, Odeberg JM, Ekman E, Hagg S. Drug-induced torsades de pointes: a review of the Swedish pharmacovigilance database. *Pharmacoepidemiol Drug Saf* 2008 Jun;17(6):587-592.
27. Meyer-Massetti C, Cheng CM, Sharpe BA, Meier CR, Guglielmo BJ. The FDA extended warning for intravenous haloperidol and torsades de pointes: how should institutions respond? *J Hosp Med* 2010 Apr;5(4):E8-16.
28. Vieweg WV, Hasnain M, Hancox JC, Baranchuk A, Digby GC, Kogut C, et al. Risperidone, QTc interval prolongation, and torsade de pointes: a systematic review of case reports. *Psychopharmacology (Berl)* 2013 Aug;228(4):515-524.
29. Zeltser D, Justo D, Halkin A, Prokhorov V, Heller K, Viskin S. Torsade de pointes due to noncardiac drugs: most patients have easily identifiable risk factors. *Medicine (Baltimore)* 2003 Jul;82(4):282-290.
30. Zhang J, Dang Q, Malik M. Baseline correction in parallel thorough QT studies. *Drug Saf* 2013 Jun;36(6):441-453.
31. Smetana P, Batchvarov V, Hnatkova K, Camm AJ, Malik M. Circadian rhythm of the corrected QT interval: impact of different heart rate correction models. *Pacing Clin Electrophysiol* 2003 Jan;26(1 Pt 2):383-386.
32. Rautaharju PM, Surawicz B, Gettes LS, Bailey JJ, Childers R, Deal BJ, et al. AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram: part IV: the ST segment, T and U waves, and the QT interval: a scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society: endorsed by the International Society for Computerized Electrocardiology. *Circulation* 2009 Mar 17;119(10):e241-50.
33. European Medicines Agency. ICH Topic E 14 The Clinical Evaluation of QT/QTc Interval Prolongation and Proarrhythmic Potential for Non-Antiarrhythmic Drugs. 2005; Available at: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2009/09/WC500002879.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500002879.pdf). Accessed September 28th, 2017.
34. Sedlak T, Shufelt C, Iribarren C, Merz CN. Sex hormones and the QT interval: a review. *J Womens Health (Larchmt)* 2012 Sep;21(9):933-941.



35. Kurokawa J, Furukawa T. Non-genomic action of sex steroid hormones and cardiac repolarization. *Biol Pharm Bull* 2013;36(1):8-12.
36. Warnier MJ, Holtkamp FA, Rutten FH, Hoes AW, de Boer A, Mol PG, et al. Quality of drug label information on QT interval prolongation. *Int J Risk Saf Med* 2014;26(2):89-98.
37. Vandael E, Vandenberg B, Vandenberghe J, Willems R, Foulon V. Risk factors for QTc-prolongation: systematic review of the evidence. *Int J Clin Pharm* 2017 Feb;39(1):16-25.
38. Vandael E, Vandenberg B, Vandenberghe J, Spriet I, Willems R, Foulon V. Development of a risk score for QTc-prolongation: the RISQ-PATH study. *Int J Clin Pharm* 2017;39(2):424-432.
39. Schwartz PJ, Woosley RL. Predicting the Unpredictable: Drug-Induced QT Prolongation and Torsades de Pointes. *J Am Coll Cardiol* 2016 Apr 5;67(13):1639-1650.
40. Castro VM, Clements CC, Murphy SN, Gainer VS, Fava M, Weilburg JB, et al. QT interval and antidepressant use: a cross sectional study of electronic health records. *BMJ* 2013 Jan 29;346:f288.
41. Fisch C, Knoebel SB. Electrographic findings in sertraline depression trials. *Drug Invest* 1992;4(4):305-312.
42. Mitchell AJ, Delaffon V, Vancampfort D, Correll CU, De Hert M. Guideline concordant monitoring of metabolic risk in people treated with antipsychotic medication: systematic review and meta-analysis of screening practices. *Psychol Med* 2012 Jan;42(1):125-147.
43. Dhamane AD, Martin BC, Brixner DI, Hudson TJ, Said Q. Metabolic monitoring of patients prescribed second-generation antipsychotics. *J Psychiatr Pract* 2013 Sep;19(5):360-374.
44. Paton C, Adroer R, Barnes TR. Monitoring lithium therapy: the impact of a quality improvement programme in the UK. *Bipolar Disord* 2013;15(8):865-75.
45. Morrato EH, Druss B, Hartung DM, Valuck RJ, Allen R, Campagna E, et al. Metabolic testing rates in 3 state Medicaid programs after FDA warnings and ADA/APA recommendations for second-generation antipsychotic drugs. *Arch Gen Psychiatry* 2010 Jan;67(1):17-24.
46. Cheung D, Wolfe B, Wald H, Cumbler E. Unsafe use of intravenous haloperidol: evaluation of recommendation-concordant care in hospitalized elderly adults. *J Am Geriatr Soc* 2013 Jan;61(1):160-161.
47. Jardin CG, Putney D, Michaud S. Assessment of drug-induced torsade de pointes risk for hospitalized high-risk patients receiving QT-prolonging agents. *Ann Pharmacother* 2014 Feb;48(2):196-202.
48. Warnier MJ, Rutten FH, Souverein PC, de Boer A, Hoes AW, De Bruin ML. Are ECG monitoring recommendations before prescription of QT-prolonging drugs applied in daily practice? The example of haloperidol. *Pharmacoepidemiol Drug Saf* 2015 Jul;24(7):701-708.

## SUPPLEMENTARY MATERIAL

Full list of all retrieved publications from the literature review. Because evidence was insufficient, additional expert opinion was used to formulate the consensus recommendations.

1. Aichhorn W, Whitworth AB, Weiss EM, Hinterhuber H, Marksteiner J. Differences between men and women in side effects of second-generation antipsychotics. *Nervenarzt* 2007 Jan;78(1):45-52.
2. Al-Abri SA, Woodburn C, Olson KR, Kearney TE. Ventricular dysrhythmias associated with poisoning and drug overdose: a 10-year review of statewide poison control center data from California. *Am J Cardiovasc Drugs* 2015 Feb;15(1):43-50.
3. Alvarez W, Jr, Pickworth KK. Safety of antidepressant drugs in the patient with cardiac disease: a review of the literature. *Pharmacotherapy* 2003 Jun;23(6):754-771.
4. Ames D, Camm J, Cook P, Falkai P, Gury C, Hurley R, et al. Minimizing the risks associated with QTc prolongation in people with schizophrenia. A consensus statement by the Cardiac Safety in Schizophrenia Group. *Encephale* 2002 Nov-Dec;28(6 Pt 1):552-562.
5. Astrom-Lilja C, Odeberg JM, Ekman E, Hagg S. Drug-induced torsades de pointes: a review of the Swedish pharmacovigilance database. *Pharmacoepidemiol Drug Saf* 2008 Jun;17(6):587-592.
6. Auquier P, Lancon C, Rouillon F, Lader M, Holmes C. Mortality in schizophrenia. *Pharmacoepidemiol Drug Saf* 2006 Dec;15(12):873-879.
7. Bailey B, Buckley NA, Amre DK. A meta-analysis of prognostic indicators to predict seizures, arrhythmias or death after tricyclic antidepressant overdose. *J Toxicol Clin Toxicol* 2004;42(6):877-888.
8. Beach SR, Celano CM, Noseworthy PA, Januzzi JL, Huffman JC. QTc prolongation, torsades de pointes, and psychotropic medications. *Psychosomatics* 2013 Jan-Feb;54(1):1-13.
9. Brewer TL, Collins M. A review of clinical manifestations in adolescent and young adults after use of synthetic cannabinoids. *J Spec Pediatr Nurs* 2014 Apr;19(2):119-126.
10. Buckley NA, Faunce TA. 'Atypical' antidepressants in overdose: clinical considerations with respect to safety. *Drug Saf* 2003;26(8):539-551.
11. Bullock R. Treatment of behavioural and psychiatric symptoms in dementia: implications of recent safety warnings. *Curr Med Res Opin* 2005 Jan;21(1):1-10.
12. Camm AJ, Karayal ON, Meltzer H, Kolluri S, O'Gorman C, Miceli J, et al. Ziprasidone and the corrected QT interval: a comprehensive summary of clinical data. *CNS Drugs* 2012 Apr 1;26(4):351-365.
13. Carreiro SV, Martins RR, De Carvalho A. Psychotropic drugs and sudden death. *Acta Med Port* 2006 Mar-Apr;19(2):151-164.

14. Castillo Sanchez M, Fabregas Escurriola M, Berge Baquero D, Goday Arno A, Valles Callol JA. Psychosis, cardiovascular risk and associated mortality: are we on the right track? *Clin Investig Arterioscler* 2014 Jan-Feb;26(1):23-32.
15. Chung AK, Chua SE. Effects on prolongation of Bazett's corrected QT interval of seven second-generation antipsychotics in the treatment of schizophrenia: a meta-analysis. *J Psychopharmacol* 2011 May;25(5):646-666.
16. Ciranni MA, Kearney TE, Olson KR. Comparing acute toxicity of first- and second-generation antipsychotic drugs: a 10-year, retrospective cohort study. *J Clin Psychiatry* 2009 Jan;70(1):122-129.
17. Cubeddu LX. QT prolongation and fatal arrhythmias: a review of clinical implications and effects of drugs. *Am J Ther* 2003 Nov-Dec;10(6):452-457.
18. de Groot MH, van Campen JP, Moek MA, Tulner LR, Beijnen JH, Lamoth CJ. The effects of fall-risk-increasing drugs on postural control: a literature review. *Drugs Aging* 2013 Nov;30(11):901-920.
19. De Hert M, Detraux J, van Winkel R, Yu W, Correll CU. Metabolic and cardiovascular adverse effects associated with antipsychotic drugs. *Nat Rev Endocrinol* 2011 Oct 18;8(2):114-126.
20. De Ponti F, Poluzzi E, Cavalli A, Recanatini M, Montanaro N. Safety of non-antiarrhythmic drugs that prolong the QT interval or induce torsade de pointes: an overview. *Drug Saf* 2002;25(4):263-286.
21. De Ponti F, Poluzzi E, Montanaro N. QT-interval prolongation by non-cardiac drugs: lessons to be learned from recent experience. *Eur J Clin Pharmacol* 2000 Apr;56(1):1-18.
22. Del Rosario ME, Weachter R, Flaker GC. Drug-induced QT prolongation and sudden death. *Mo Med* 2010 Jan-Feb;107(1):53-58.
23. Dorado P, Berecz R, Penas-Lledo EM, Llerena A. Antipsychotic drugs and QTc prolongation: the potential role of CYP2D6 genetic polymorphism. *Expert Opin Drug Metab Toxicol* 2007 Feb;3(1):9-19.
24. Fanoë S, Kristensen D, Fink-Jensen A, Jensen HK, Toft E, Nielsen J, et al. Risk of arrhythmia induced by psychotropic medications: a proposal for clinical management. *Eur Heart J* 2014 May 21;35(20):1306-1315.
25. Fazekas T, Liskai G. Clinical proarrhythmias induced by anti-arrhythmic drugs, non-cardiovascular agents and implantable cardioverter-defibrillators. *Orv Hetil* 2002 Jan 13;143(2):61-69.
26. Fazekas T, Liskai G. Proarrhythmic (torsadogenic) effects of QT-prolonging non-antiarrhythmic drugs. *Orv Hetil* 2005 Mar 6;146(10):451-460.
27. Foulke GE, Albertson TE, Walby WF. Tricyclic antidepressant overdose: emergency department findings as predictors of clinical course. *Am J Emerg Med* 1986 Nov;4(6):496-500.

28. Freeman MP, Freeman SA. Lithium: clinical considerations in internal medicine. *Am J Med* 2006 Jun;119(6):478-481.
29. Frimas V, Roberge C, Perroux D, Dauvillier JM. Cardiological monitoring of antipsychotic-treated patients: evaluation and evolution of a hospital protocol. *Encephale* 2008 Oct;34(5):467-476.
30. Funk KA, Bostwick JR. A comparison of the risk of QT prolongation among SSRIs. *Ann Pharmacother* 2013 Oct;47(10):1330-1341.
31. Girardin FR, Gex-Fabry M, Berney P, Shah D, Gaspoz JM, Dayer P. Drug-induced long QT in adult psychiatric inpatients: the 5-year cross-sectional ECG Screening Outcome in Psychiatry study. *Am J Psychiatry* 2013 Dec;170(12):1468-1476.
32. Glassman AH. Clinical management of cardiovascular risks during treatment with psychotropic drugs. *J Clin Psychiatry* 2002;63 Suppl 9:12-17.
33. Glassman AH, Bigger JT, Jr. Antipsychotic drugs: prolonged QTc interval, torsade de pointes, and sudden death. *Am J Psychiatry* 2001 Nov;158(11):1774-1782.
34. Greco KE, Tune LE, Brown FW, Van Horn WA. A retrospective study of the safety of intramuscular ziprasidone in agitated elderly patients. *J Clin Psychiatry* 2005 Jul;66(7):928-929.
35. Groleau G, Jotte R, Barish R. The electrocardiographic manifestations of cyclic antidepressant therapy and overdose: a review. *J Emerg Med* 1990 Sep-Oct;8(5):597-605.
36. Haas SJ, Hill R, Krum H, Liew D, Tonkin A, Demos L, et al. Clozapine-associated myocarditis: a review of 116 cases of suspected myocarditis associated with the use of clozapine in Australia during 1993-2003. *Drug Saf* 2007;30(1):47-57.
37. Haddad PM, Sharma SG. Adverse effects of atypical antipsychotics : differential risk and clinical implications. *CNS Drugs* 2007;21(11):911-936.
38. Hale AS. A review of the safety and tolerability of sertindole. *Int Clin Psychopharmacol* 1998 Mar;13 Suppl 3:S65-70.
39. Hasnain M, Vieweg WV. QTc interval prolongation and torsade de pointes associated with second-generation antipsychotics and antidepressants: a comprehensive review. *CNS Drugs* 2014 Oct;28(10):887-920.
40. Hasnain M, Vieweg WV, Howland RH, Kogut C, Breden Crouse EL, Koneru JN, et al. Quetiapine and the need for a thorough QT/QTc study. *J Clin Psychopharmacol* 2014 Feb;34(1):3-6.
41. Hays JT. Tobacco dependence treatment in patients with heart and lung disease: implications for intervention and review of pharmacological therapy. *J Cardiopulm Rehabil* 2000 Jul-Aug;20(4):215-223.
42. Herrmann N, Lanctot KL. Atypical antipsychotics for neuropsychiatric symptoms of dementia: malignant or maligned? *Drug Saf* 2006;29(10):833-843.

43. Ho JM, Gomes T, Straus SE, Austin PC, Mamdani M, Juurlink DN. Adverse cardiac events in older patients receiving venlafaxine: a population-based study. *J Clin Psychiatry* 2014 Jun;75(6):e552-8.
44. Howland RH. Psychiatric medications and sudden cardiac death: putting the risk in perspective. *J Psychosoc Nurs Ment Health Serv* 2015 Feb;53(2):23-25.
45. Howland RH. The comparative cardiac effects of haloperidol and quetiapine: parsing a review. *J Psychosoc Nurs Ment Health Serv* 2014 Jun;52(6):23-26.
46. Joy JP, Coulter CV, Duffull SB, Isbister GK. Prediction of torsade de pointes from the QT interval: analysis of a case series of amisulpride overdoses. *Clin Pharmacol Ther* 2011 Aug;90(2):243-245.
47. Kao LW, Kirk MA, Evers SJ, Rosenfeld SH. Droperidol, QT prolongation, and sudden death: what is the evidence? *Ann Emerg Med* 2003 Apr;41(4):546-558.
48. Karamatskos E, Lambert M, Mulert C, Naber D. Drug safety and efficacy evaluation of sertindole for schizophrenia. *Expert Opin Drug Saf* 2012 Nov;11(6):1047-1062.
49. Kelly DL, Love RC. Ziprasidone and the QTc interval: pharmacokinetic and pharmacodynamic considerations. *Psychopharmacol Bull* 2001 Autumn;35(4):66-79.
50. Kireyev D, Arkhipov MV, Zador ST, Paris JA, Boden WE. Clinical utility of aVR-The neglected electrocardiographic lead. *Ann Noninvasive Electrocardiol* 2010 Apr;15(2):175-180.
51. Koegelenberg CF, Joubert ZJ, Irusen EM. Tricyclic antidepressant overdose necessitating ICU admission. *S Afr Med J* 2012 Mar 1;102(5):293-294.
52. Koenig X, Hilber K. The anti-addiction drug ibogaine and the heart: a delicate relation. *Molecules* 2015 Jan 29;20(2):2208-2228.
53. Koponen H, Alaraisanen A, Saari K, Pelkonen O, Huikuri H, Raatikainen MJ, et al. Schizophrenia and sudden cardiac death: a review. *Nord J Psychiatry* 2008;62(5):342-345.
54. Kovacs D, Arora R. Cardiovascular effects of psychotropic drugs. *Am J Ther* 2008 Sep-Oct;15(5):474-483.
55. Kurth J, Maguire G. Pediatric case report of quetiapine overdose and QTc prolongation. *Ann Clin Psychiatry* 2004 Oct-Dec;16(4):229-231.
56. Lange-Asschenfeldt C, Lederbogen F. Antidepressant therapy in coronary artery disease. *Nervenarzt* 2011 May;82(5):657-64; quiz 665-6.
57. Laursen TM, Munk-Olsen T, Vestergaard M. Life expectancy and cardiovascular mortality in persons with schizophrenia. *Curr Opin Psychiatry* 2012 Mar;25(2):83-88.
58. Lindstrom E, Farde L, Eberhard J, Haverkamp W. QTc interval prolongation and antipsychotic drug treatments: focus on sertindole. *Int J Neuropsychopharmacol* 2005 Dec;8(4):615-629.

59. Lopponen P, Tetri S, Juvela S, Huhtakangas J, Saloheimo P, Bode MK, et al. Association between warfarin combined with serotonin-modulating antidepressants and increased case fatality in primary intracerebral hemorrhage: a population-based study. *J Neurosurg* 2014 Jun;120(6):1358-1363.
60. Mackin P. Cardiac side effects of psychiatric drugs. *Hum Psychopharmacol* 2008 Jan;23 Suppl 1:3-14.
61. Maldonado JR. Delirium in the acute care setting: characteristics, diagnosis and treatment. *Crit Care Clin* 2008 Oct;24(4):657-722, vii.
62. Marder SR, Essock SM, Miller AL, Buchanan RW, Casey DE, Davis JM, et al. Physical health monitoring of patients with schizophrenia. *Am J Psychiatry* 2004 Aug;161(8):1334-1349.
63. Menkes DB, Knight JC. Cardiotoxicity and prescription of thioridazine in New Zealand. *Aust N Z J Psychiatry* 2002 Aug;36(4):492-498.
64. Merrill DB, Dec GW, Goff DC. Adverse cardiac effects associated with clozapine. *J Clin Psychopharmacol* 2005 Feb;25(1):32-41.
65. Meyer-Masseti C, Cheng CM, Sharpe BA, Meier CR, Guglielmo BJ. The FDA extended warning for intravenous haloperidol and torsades de pointes: how should institutions respond? *J Hosp Med* 2010 Apr;5(4):E8-16.
66. Meyer-Masseti C, Vaerini S, Ratz Bravo AE, Meier CR, Guglielmo BJ. Comparative safety of antipsychotics in the WHO pharmacovigilance database: the haloperidol case. *Int J Clin Pharm* 2011 Oct;33(5):806-814.
67. Mittal V, Kurup L, Williamson D, Muralee S, Tampi RR. Risk of cerebrovascular adverse events and death in elderly patients with dementia when treated with antipsychotic medications: a literature review of evidence. *Am J Alzheimers Dis Other Demen* 2011 Feb;26(1):10-28.
68. Mohr WK, Petti TA, Mohr BD. Adverse effects associated with physical restraint. *Can J Psychiatry* 2003 Jun;48(5):330-337.
69. Neuhut R, Lindenmayer JP, Silva R. Neuroleptic malignant syndrome in children and adolescents on atypical antipsychotic medication: a review. *J Child Adolesc Psychopharmacol* 2009 Aug;19(4):415-422.
70. Olgun H, Yildirim ZK, Karacan M, Ceviz N. Clinical, electrocardiographic, and laboratory findings in children with amitriptyline intoxication. *Pediatr Emerg Care* 2009 Mar;25(3):170-173.
71. Peter L, Dome P, Rihmer Z, Kovacs G, Faludi G. Cardiovascular disorders and depression: a review of epidemiological and possible etiological data. *Neuropsychopharmacol Hung* 2008 May;10(2):81-90.
72. Pizzi C, Rutjes AW, Costa GM, Fontana F, Mezzetti A, Manzoli L. Meta-analysis of selective serotonin reuptake inhibitors in patients with depression and coronary heart disease. *Am J Cardiol* 2011 Apr 1;107(7):972-979.

73. Postema PG, Wolpert C, Amin AS, Probst V, Borggrefe M, Roden DM, et al. Drugs and Brugada syndrome patients: review of the literature, recommendations, and an up-to-date website ([www.brugadadrugs.org](http://www.brugadadrugs.org)). *Heart Rhythm* 2009 Sep;6(9):1335-1341.
74. Rabkin SW. Impact of Age and Sex on QT Prolongation in Patients Receiving Psychotropics. *Can J Psychiatry* 2015 May;60(5):206-214.
75. Raedler TJ. Cardiovascular aspects of antipsychotics. *Curr Opin Psychiatry* 2010 Nov;23(6):574-581.
76. Reingardiene D, Vilcinskaite J. QTc-prolonging drugs and the risk of sudden death. *Medicina (Kaunas)* 2007;43(4):347-353.
77. Richards JR, Schneir AB. Droperidol in the emergency department: is it safe? *J Emerg Med* 2003 May;24(4):441-447.
78. Sala M, Lazzaretti M, De Vidovich G, Caverzasi E, Barale F, d'Allio G, et al. Electrophysiological changes of cardiac function during antidepressant treatment. *Ther Adv Cardiovasc Dis* 2009 Feb;3(1):29-43.
79. Saravane D, Feve B, Frances Y, Corruble E, Lancon C, Chanson P, et al. Drawing up guidelines for the attendance of physical health of patients with severe mental illness. *Encephale* 2009 Sep;35(4):330-339.
80. Sicouri S, Antzelevitch C. Sudden cardiac death secondary to antidepressant and antipsychotic drugs. *Expert Opin Drug Saf* 2008 Mar;7(2):181-194.
81. Singh N, Singh HK, Khan IA. Serial electrocardiographic changes as a predictor of cardiovascular toxicity in acute tricyclic antidepressant overdose. *Am J Ther* 2002 Jan-Feb;9(1):75-79.
82. Smithburger PL, Seybert AL, Armahizer MJ, Kane-Gill SL. QT prolongation in the intensive care unit: commonly used medications and the impact of drug-drug interactions. *Expert Opin Drug Saf* 2010 Sep;9(5):699-712.
83. Straus SM, Bleumink GS, Dieleman JP, van der Lei J, 't Jong GW, Kingma JH, et al. Antipsychotics and the risk of sudden cardiac death. *Arch Intern Med* 2004 Jun 28;164(12):1293-1297.
84. Tamargo J. Drug-induced torsade de pointes: from molecular biology to bedside. *Jpn J Pharmacol* 2000 May;83(1):1-19.
85. Taylor DM. Antipsychotics and QT prolongation. *Acta Psychiatr Scand* 2003 Feb;107(2):85-95.
86. Titier K, Canal M, Deridet E, Abouelfath A, Gromb S, Molimard M, et al. Determination of myocardium to plasma concentration ratios of five antipsychotic drugs: comparison with their ability to induce arrhythmia and sudden death in clinical practice. *Toxicol Appl Pharmacol* 2004 Aug 15;199(1):52-60.
87. Vieweg WV. Mechanisms and risks of electrocardiographic QT interval prolongation when using antipsychotic drugs. *J Clin Psychiatry* 2002;63 Suppl 9:18-24.

88. Vieweg WV, Hasnain M, Hancox JC, Baranchuk A, Digby GC, Kogut C, et al. Risperidone, QTc interval prolongation, and torsade de pointes: a systematic review of case reports. *Psychopharmacology (Berl)* 2013 Aug;228(4):515-524.
89. Vieweg WV, Hasnain M, Howland RH, Hettema JM, Kogut C, Wood MA, et al. Citalopram, QTc interval prolongation, and torsade de pointes. How should we apply the recent FDA ruling? *Am J Med* 2012 Sep;125(9):859-868.
90. Vieweg WV, Wood MA, Fernandez A, Beatty-Brooks M, Hasnain M, Pandurangi AK. Proarrhythmic risk with antipsychotic and antidepressant drugs: implications in the elderly. *Drugs Aging* 2009;26(12):997-1012.
91. Viskin S, Justo D, Halkin A, Zeltser D. Long QT syndrome caused by noncardiac drugs. *Prog Cardiovasc Dis* 2003 Mar-Apr;45(5):415-427.
92. Waring WS, Good AM, Bateman DN. Lack of significant toxicity after mirtazapine overdose: a five-year review of cases admitted to a regional toxicology unit. *Clin Toxicol (Phila)* 2007;45(1):45-50.
93. Warner B, Hoffmann P. Investigation of the potential of clozapine to cause torsade de pointes. *Adverse Drug React Toxicol Rev* 2002;21(4):189-203.
94. Wilens TE, Biederman J, Baldessarini RJ, Geller B, Schleifer D, Spencer TJ, et al. Cardiovascular effects of therapeutic doses of tricyclic antidepressants in children and adolescents. *J Am Acad Child Adolesc Psychiatry* 1996 Nov;35(11):1491-1501.
95. Wilens TE, Stern TA, O'Gara PT. Adverse cardiac effects of combined neuroleptic ingestion and tricyclic antidepressant overdose. *J Clin Psychopharmacol* 1990 Feb;10(1):51-54.
96. Wobrock T, Schwaab B, Bohm M, Schafers HJ, Wanke K, Supprian T. Pharmacotherapeutical approaches to insomnia patients with cardiac diseases and after heart transplantation. *Z Kardiol* 2001 Oct;90(10):717-728.
97. Xue F, Strombom I, Turnbull B, Zhu S, Seeger J. Treatment with duloxetine in adults and the incidence of cardiovascular events. *J Clin Psychopharmacol* 2012 Feb;32(1):23-30.
98. Zeltser D, Justo D, Halkin A, Prokhorov V, Heller K, Viskin S. Torsade de pointes due to noncardiac drugs: most patients have easily identifiable risk factors. *Medicine (Baltimore)* 2003 Jul;82(4):282-290.
99. Zemrak WR, Kenna GA. Association of antipsychotic and antidepressant drugs with Q-T interval prolongation. *Am J Health Syst Pharm* 2008 Jun 1;65(11):1029-1038.
100. More evidence on risks of antipsychotics in adults. Commentaries provide advice about putting the results in perspective. *Harv Ment Health Lett* 2009 Apr;25(10):6.



