



University of Groningen

The organic ties of iron

Slagter, Hans Arent

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Slagter, H. A. (2018). The organic ties of iron: Or the origin and fate of Fe-binding organic ligands. [Groningen]: Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

- Aagaard, K., Swift, J.H., Carmack, E.C., 1985. Thermohaline circulation in the Arctic Mediterranean Seas. *J. Geophys. Res.* 90, 4833–4846.
- Abualhaija, M.M., van den Berg, C.M.G., 2014. Chemical speciation of iron in seawater using catalytic cathodic stripping voltammetry with ligand competition against salicylaldoxime. *Mar. Chem.* 164, 60–74.
- Abualhaija, M.M., Whitby, H., van den Berg, C.M.G., 2015. Competition between copper and iron for humic ligands in estuarine waters. *Mar. Chem.* 172, 46–56.
- Achterberg, E.P., Steigenberger, S., Marsay, C.M., Lemoigne, F.A.C., Painter, S.C., Baker, A.R., Connelly, D.P., Moore, C.M., Tagliabue, A., Tanhua, T., 2018. Iron Biogeochemistry in the High Latitude North Atlantic Ocean. *Sci. Rep.* 8, 1–15.
- Aguilar-Islas, A.M., Rember, R.D., Mordy, C.W., Wu, J., 2008. Sea ice-derived dissolved iron and its potential influence on the spring algal bloom in the Bering Sea. *Geophys. Res. Lett.* 35, L24601.
- Aguilar-Islas, A.M., Séguret, M.J.M., Rember, R., Buck, K.N., Proctor, P., Mordy, C.W., Kachel, N.B., 2016. Temporal variability of reactive iron over the Gulf of Alaska shelf. *Deep. Res. Part II Top. Stud. Oceanogr.* 132, 90–106.
- Aguilar-islas, A.M., Wu, J., Rember, R., Johansen, A.M., Shank, L.M., 2010. Dissolution of aerosol-derived iron in seawater: Leach solution chemistry, aerosol type, and colloidal iron fraction. *Mar. Chem.* 120, 25–33.
- Ambar, I., Serra, N., Neves, F., Ferreira, T., 2008. Observations of the Mediterranean Undercurrent and eddies in the Gulf of Cadiz during 2001. *J. Mar. Syst.* 71, 195–220.
- Amon, R.M.W., Gereon, B., Benedikt, M., 2003. Dissolved organic carbon distribution and origin in the Nordic Seas: Exchanges with the Arctic Ocean and the North Atlantic. *J. Geophys. Res.* 108, 3221.
- Apte, S.C., Gardner, M.J., Ravenscroft, J.E., 1988. An evaluation of voltammetric titration procedures for the determination of trace metal complexation in natural waters by use of computer simulation. *Anal. Chim. Acta* 212, 1–21.
- Arrigo, K.R., van Dijken, G., Pabi, S., 2008. Impact of a shrinking Arctic ice cover on marine primary production. *Geophys. Res. Lett.* 35, 1–6.
- Arrigo, K.R., Van Dijken, G.L., 2011. Secular trends in Arctic Ocean net primary production. *J. Geophys. Res. Ocean.* 116, 1–15.
- Baars, O., Morel, F.M.M., Perlman, D.H., 2014. ChelomEx: Isotope-assisted discovery of metal chelates in complex media using high-resolution LC-MS. *Anal. Chem.* 86, 11298–11305.
- Baker, A.R., Croot, P.L., 2010. Atmospheric and marine controls on aerosol iron solubility in seawater. *Mar. Chem.* 120, 4–13.
- Baker, A.R., Jickells, T.D., 2006. Mineral particle size as a control on aerosol iron solubility. *Geophys. Res. Lett.* 33, 1–4.
- Baker, E.T., Edmonds, H.N., Michael, P.J., Bach, W., Dick, H.J.B., Snow, J.E., Walker, S.L., Banerjee, N.R., Langmuir, C.H., 2004. Hydrothermal venting in magma deserts: The ultraslow-spreading Gakkel and Southwest Indian Ridges. *Geochemistry, Geophys. Geosystems* 5, 1–29.
- Barbeau, K., Rue, E.L., Bruland, K.W., Butler, a, 2001. Photochemical cycling of iron in the surface ocean mediated by microbial iron(III)-binding ligands. *Nature* 413, 409–413.
- Barbeau, K., Rue, E.L., Trick, C.G., Bruland, K.W., Butler, A., 2003. Photochemical reactivity of siderophores produced by marine heterotrophic bacteria and

- cyanobacteria, based on characteristic Fe(III) binding groups. Limnol. Oceanogr. 48, 1069–1078.
- Bartual, S.G., Otero, J.M., Garcia-Doval, C., Llamas-Saiz, A.L., Kahn, R., Fox, G.C., van Raaij, M.J., 2010. Structure of the bacteriophage T4 long tail fiber receptor-binding tip. Proc. Natl. Acad. Sci. U. S. A. 107, 20287–92.
- Batchelli, S., Muller, F.L.L., Chang, K.C., Lee, C.L., 2010. Evidence for strong but dynamic iron-humic colloidal associations in humic-rich coastal waters. Environ. Sci. Technol. 44, 8485–8490.
- Bauch, D., Cherniavskaya, E., Timokhov, L., 2016. Shelf basin exchange along the Siberian continental margin: Modification of Atlantic Water and Lower Halocline Water. Deep Sea Res. Part I 115, 188–198.
- Bauch, D., van der Loeff, M.R., Andersen, N., Torres-Valdes, S., Bakker, K., Abrahamsen, E.P., 2011. Origin of freshwater and polynya water in the Arctic Ocean halocline in summer 2007. Prog. Oceanogr. 91, 482–495.
- Baudoux, A.-C., Brussaard, C.P.D., 2005. Characterization of different viruses infecting the marine harmful algal bloom species *Phaeocystis globosa*. Virology 341, 80–90.
- Baudoux, A.C., Brussaard, C.P.D., 2005. Characterization of different viruses infecting the marine harmful algal bloom species *Phaeocystis globosa*. Virology 341, 80–90.
- Baudoux, A.C., Noordeloos, A.A.M., Veldhuis, M.J.W., Brussaard, C.P.D., 2006. Virally induced mortality of *Phaeocystis globosa* during two spring blooms in temperate coastal waters. Aquat. Microb. Ecol. 44, 207–217.
- Baudoux, A.C., Veldhuis, M.J.W., Witte, H.J., Brussaard, C.P.D., 2007. Viruses as mortality agents of picophytoplankton in the deep chlorophyll maximum layer during IRONAGES III. Limnol. Oceanogr. 52, 2519–2529.
- Beaulieu, S.E., Baker, E.T., German, C.R., 2015. Where are the undiscovered hydrothermal vents on oceanic spreading ridges? Deep. Res. Part II Top. Stud. Oceanogr. 121, 202–212.
- Behrenfeld, M.J., Bale, A.J., Kolber, Z.S., Aiken, J., Falkowski, P.G., 1996. Confirmation of iron limitation of phytoplankton photosynthesis in the equatorial Pacific Ocean. Nature.
- Behrenfeld, M.J., Worthington, K., Sherrell, R.M., Chavez, F.P., Strutton, P., McPhaden, M., Shea, D.M., 2006. Controls on tropical Pacific Ocean productivity revealed through nutrient stress diagnostics. Nature 442, 1025–8.
- Bennett, S.A., Achterberg, E.P., Connelly, D.P., Statham, P.J., Fones, G.R., German, C.R., 2008. The distribution and stabilisation of dissolved Fe in deep-sea hydrothermal plumes. Earth Planet. Sci. Lett. 270, 157–167.
- Bergeron, M., Tremblay, J.-É., 2014. Shifts in biological productivity inferred from nutrient drawdown in the southern Beaufort Sea (2003–2011) and northern Baffin Bay (1997–2011), Canadian Arctic. Geophys. Res. Lett. 41, 3979–3987.
- Beşiktepe, Ş.T., Sur, H.İ., Özsoy, E., Latif, M.A., Oğuz, T., Ünlüata, Ü., 1994. The circulation and hydrography of the Marmara Sea. Prog. Oceanogr. 34, 285–334.
- Bhatt, U.S., Walker, D.A., Walsh, J.E., Carmack, E.C., Frey, K.E., Meier, W.N., Moore, S.E., Parmentier, F.-J.W., Post, E., Romanovsky, V.E., Simpson, W.R., 2014. Implications of Arctic Sea Ice Decline for the Earth System. Annu. Rev. Environ. Resour. 39, 57–89.
- Birchill, A.J., Milne, A., Woodward, E.M.S., Harris, C., Annett, A., Rusiecka, D., Achterberg, E.P., Gledhill, M., Ussher, S.J., Worsfold, P.J., Geibert, W., Lohan, M.C., 2017. Seasonal iron depletion in temperate shelf seas. Geophys. Res. Lett. 44, 8987–8996.
- Blain, S., Bonnet, S., Guieu, C., 2008. Dissolved iron distribution in the tropical and sub-tropical South Eastern Pacific. Biogeosciences 5, 269–280.

- Bluhm, B.A., Kosobokova, K.N., Carmack, E.C., 2015. A tale of two basins: An integrated physical and biological perspective of the deep Arctic Ocean. *Prog. Oceanogr.* 139, 89–121.
- Boiteau, R.M., Fitzsimmons, J.N., Repeta, D.J., Boyle, E.A., 2013. Detection of iron ligands in seawater and marine cyanobacteria cultures by high-performance liquid chromatography-inductively coupled plasma-mass spectrometry. *Anal Chem* 85, 4357–4362.
- Boiteau, R.M., Mende, D.R., Hawco, N.J., McIlvin, M.R., Fitzsimmons, J.N., Saito, M.A., Sedwick, P.N., DeLong, E.F., Repeta, D.J., 2016. Siderophore-based microbial adaptations to iron scarcity across the eastern Pacific Ocean. *Proc. Natl. Acad. Sci.* 113, 14237–14242.
- Bonini, M., Mazzarini, F., 2010. Mud volcanoes as potential indicators of regional stress and pressurized layer depth. *Tectonophysics* 494, 32–47.
- Bonnain, C., Breitbart, M., Buck, K.N., 2016. The Ferrojan Horse Hypothesis: Iron-Virus Interactions in the Ocean. *Front. Mar. Sci.* 3, 1–11.
- Bonnet, S., Guieu, C., 2006. Atmospheric forcing on the annual iron cycle in the western Mediterranean Sea: A 1-year survey. *J. Geophys. Res.* 111.
- Boyd, P.W., Arrigo, K.R., Strzepek, R., Dijken, G.L. Van, van Dijken, G.L., 2012. Mapping phytoplankton iron utilization: Insights into Southern Ocean supply mechanisms. *J. Geophys. Res.* 117, 1–18.
- Boyd, P.W., Ibsanmi, E., Sander, S.G., Hunter, K.A., Jackson, G.A., 2010. Remineralization of upper ocean particles: Implications for iron biogeochemistry. *Limnol. Oceanogr.* 55, 1271–1288.
- Boyd, P.W., Jickells, T., Law, C.S., Blain, S., Boyle, E. a, Buesseler, K.O., Coale, K.H., Cullen, J.J., de Baar, H.J.W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N.P.J., Pollard, R., Rivkin, R.B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., Watson, A.J., 2007. Mesoscale iron enrichment experiments 1993–2005: synthesis and future directions. *Science* 315, 612–617.
- Boyle, E.A., Edmond, J.M., Sholkovitz, E.R., 1977. The mechanism of iron removal in estuaries. *Geochim. Cosmochim. Acta* 41, 1313–1324.
- Brand, L.E., 1991. Minimum iron requirements of marine phytoplankton and the implications for the biogeochemical control of new production. *Limnol. Oceanogr.* 36, 1756–1771.
- Brand, L.E., 1991. Minimum iron requirements of marine phytoplankton and the implications for the biogeochemical control of new production. *Limnol. Oceanogr.* 36, 1756–1771.
- Bratbak, G., Jacobsen, A., Heldal, M., Nagasaki, K., Thingstad, F., 1998. Virus production in *Phaeocystis pouchetii* and its relation to host cell growth and nutrition. *Aquat. Microb. Ecol.* 16, 1–9.
- Breitbarth, E., Achterberg, E., Ardelan, M. V., Baker, A.R., Bucciarelli, E., Chever, F., Croot, P.L., Duggen, S., Gledhill, M., Hassellöv, M., Hassler, C., Hoffmann, L.J., Hunter, K.A., Hutchins, D.A., Ingri, J., Jickells, T., Lohan, M.C., Nielsdóttir, M.C., Sarthou, G., Schoemann, V., Trapp, J.M., Turner, D.R., Ye, Y., 2010. Iron biogeochemistry across marine systems—progress from the past decade. *Biogeosciences* 7, 1075–1097.
- Bronk, D.A., 2002. Dynamics of DON. In: Hansell, D.A., Carlson, C.A. (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*. Academic Press, London, pp. 153–247.
- Brown, M.T., Lippiatt, S.M., Lohan, M.C., Bruland, K.W., 2012. Trace metal distributions within a Sitka eddy in the northern Gulf of Alaska. *Limnol. Oceanogr.* 57, 503–518.

- Browning, T.J., Achterberg, E.P., Rapp, I., Engel, A., Bertrand, E.M., Tagliabue, A., Moore, C.M., 2017. Nutrient co-limitation at the boundary of an oceanic gyre. *Nature* 551, 242–246.
- Bruland, K.W., Middag, R., Lohan, M.C., 2014. Controls of Trace Metals in Seawater. In: *Treatise on Geochemistry*. Elsevier, pp. 19–51.
- Brussaard, C.P.D., 2004a. Viral control of phytoplankton populations - a review. *J. Eukaryot. Microbiol.* 51, 125–138.
- Brussaard, C.P.D., 2004b. Optimization of Procedures for Counting Viruses by Flow Cytometry. *Appl. Environ. Microbiol.* 70, 1506–1513.
- Brussaard, C.P.D., Mari, X., Van Bleijswijk, J.D.L., Veldhuis, M.J.W., 2005. A mesocosm study of *Phaeocystis globosa* (Prymnesiophyceae) population dynamics: II. Significance for the microbial community. *Harmful Algae* 4, 875–893.
- Brussaard, C.P.D., Martínez, J.M., 2008. Algal Bloom Viruses. *Plant Viruses* 2, 1–10.
- Brussaard, C.P.D., Payet, J.P., Winter, C., Weinbauer, M.G., Wilhelm, D.W., Suttle, C.A., 2010. Quantification of aquatic viruses by flow cytometry. In: *Manual of Aquatic Viral Ecology*. ASLO, pp. 102–109.
- Brussaard, C.P.D., Thyrhaug, R., Marie, D., Bratbak, G., 1999. Flow Cytometric Analyses of Viral Infection in Two Marine Phytoplankton Species, *Micromonas Pusilla* (Prasinophyceae) and *Phaeocystis Pouchetii* (Prymnesiophyceae). *J. Phycol.* 35, 941–948.
- Brussaard, C.P.D., Wilhelm, S.W., Thingstad, T.F., Weinbauer, M.G., Bratbak, G., Heldal, M., Kimmance, S.A., Middelboe, M., Nagasaki, K., Paul, J.H., Schroeder, D.C., Suttle, C.A., Vaqué, D., Wommack, K.E., 2008. Global-scale processes with a nanoscale drive: the role of marine viruses. *Isme J.* 0, 1–4.
- Buck, C.S., Landing, W.M., Resing, J.A., Measures, C.I., 2010. The solubility and deposition of aerosol Fe and other trace elements in the North Atlantic Ocean: Observations from the A16N CLIVAR/CO2 repeat hydrography section. *Mar. Chem.* 120, 57–70.
- Buck, K.N., Gerringa, L.J.A., Rijkenberg, M.J.A., 2016. An Intercomparison of Dissolved Iron Speciation at the Bermuda Atlantic Time-series Study (BATS) Site: Results from GEOTRACES Crossover Station A. *Front. Mar. Sci.* 3, 262.
- Buck, K.N., Lohan, M.C., Berger, C.J.M., Bruland, K.W., 2007. Dissolved iron speciation in two distinct river plumes and an estuary: Implications for riverine iron supply. *Limnol. Oceanogr.* 52, 843–855.
- Buck, K.N., Moffett, J., Barbeau, K.A., Bundy, R.M., Kondo, Y., Wu, J., 2012. The organic complexation of iron and copper: an intercomparison of competitive ligand exchange – adsorptive cathodic stripping voltammetry (CLE-ACSV) techniques. *Limnol. Oceanogr. Methods* 10, 496–515.
- Buck, K.N., Sohst, B., Sedwick, P.N., 2015. The organic complexation of dissolved iron along the U.S. GEOTRACES (GA03) North Atlantic Section. *Deep. Res. Part II Top. Stud. Oceanogr.* 116, 152–165.
- Buffle, J., 1988. *Complexation Reactions in Aquatic Systems; An Analytical Approach*, 1st ed. Ellis Horwood, Chichester, UK.
- Buffle, J., 1990. The analytical challenge posed by fulvic and humic compounds. *Anal. Chim. Acta* 232, 1–2.
- Bullard, J.E., 2017. The distribution and biogeochemical importance of high-latitude dust in the Arctic and Southern Ocean-Antarctic regions. *J. Geophys. Res. Atmos.* 122, 3098–3103.
- Bundy, R.M., Abdulla, H.A.N., Hatcher, P.G., Biller, D. V., Buck, K.N., Barbeau, K.A., 2015. Iron-binding ligands and humic substances in the San Francisco Bay estuary and estuarine-influenced shelf regions of coastal California. *Mar. Chem.* 173, 183–

- Bundy, R.M., Boiteau, R.M., McLean, C., Turk-Kubo, K.A., McIlvin, M.R., Saito, M.A., Van Mooy, B.A.S., Repeta, D.J., Mooy, B.A. Van, Repeta, D.J., Van Mooy, B.A.S., Repeta, D.J., 2018. Distinct Siderophores Contribute to Iron Cycling in the Mesopelagic at Station ALOHA. *Front. Mar. Sci.* 5, 1–15.
- Bundy, R.M., Jiang, M., Carter, M., Barbeau, K.A., 2016. Iron-Binding Ligands in the Southern California Current System: Mechanistic Studies. *Front. Mar. Sci.* 3, 1–17.
- Butler, A., 2005. Marine siderophores and microbial iron mobilization. *Biometals* 18, 369–374.
- Calbet, A., Landry, M.R., 2004. Phytoplankton growth, microzooplankton grazing, and carbon cycling in marine systems. *Limnol. Oceanogr.* 49, 51–57.
- Caprara, S., Buck, K.N., Gerringa, L., Rijkenberg, M., Monticelli, D., 2016. A compilation of iron speciation data for open oceanic waters. *Front. Mar. Sci.* 3, 221.
- Carlson, C.A., 2002. Production and Removal Processes. In: Hansell, D.A., Carlson, C.A. (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*. Academic Press, London, pp. 91–151.
- Carmack, E.C., Yamamoto-Kawai, M., Haine, T.W.N., Bacon, S., Bluhm, B.A., Lique, C., Melling, H., Polyakov, I. V., Straneo, F., Timmermans, M.L., Williams, W.J., 2016. Freshwater and its role in the Arctic Marine System: Sources, disposition, storage, export, and physical and biogeochemical consequences in the Arctic and global oceans. *J. Geophys. Res. G Biogeosciences* 121, 675–717.
- Coble, P.G., 2007. Marine optical biogeochemistry: The chemistry of ocean color. *Chem. Rev.* 107, 402–418.
- Coble, P.G., Del Castillo, C.E., Avril, B., 1998. Distribution and optical properties of CDOM in the Arabian Sea during the 1995 Southwest Monsoon. *Deep Sea Res. Part II* 45, 2195–2223.
- Codispoti, L.A., Kelly, V., Thessen, A., Matrai, P., Suttles, S., Hill, V., Steele, M., Light, B., 2013. Synthesis of primary production in the Arctic Ocean: III. Nitrate and phosphate based estimates of net community production. *Prog. Oceanogr.* 110, 126–150.
- Cooper, L.W., Whitledge, T.E., Grebmeier, J.M., Weingartner, T., 1997. The nutrient, salinity, and stable oxygen isotope composition of Bering and Chukchi Sea waters. *J. Geophys. Res.* 102, 12563–12573.
- Cottrell, M.T., Suttle, C. a., Aransas, P., 1995. Dynamics of lytic virus infecting the photosynthetic marine picoflagellate *Micromonas pusilla*. *Limnol. Oceanogr.* 40, 730–739.
- Croot, P.L., Johansson, M., 2000. Determination of iron speciation by cathodic stripping voltammetry in seawater using the competing ligand 2-(2-thiazolylazo)-p-cresol (TAC). *Electroanalysis* 12, 565–576.
- Croot, P.L., Streu, P., Baker, A.R., 2004. Short residence time for iron in surface seawater impacted by atmospheric dry deposition from Saharan dust events. *Geophys. Res. Lett.* 31.
- Cuttler, G., Andersson, P.S., Codispoti, L.A., Croot, P.L., Francois, R., Lohan, M.C., Obata, H., Rutgers van der Loeff, M., 2010. Sampling and sample-handling protocols for GEOTRACES cruises 1–238.
- Dai, M., Martin, J.M., Cauwet, G., 1995. The Significant Role of Colloids in the Transport and Transformation of Organic-Carbon and Associated Trace-Metals (Cd, Cu and Ni) in the Rhone Delta (France). *Mar. Chem.* 51, 159–175.
- de Baar, H.J.W., Boyd, P.W., Coale, K.H., Landry, M.R., Tsuda, A., Assmy, P., Bakker, D.C.E., Bozec, Y., Barber, R.T., Brzezinski, M.A., Buesseler, K.O., Boye, M., Croot, P.L., Gervais, F., Gorbunov, M.Y., Harrison, P.J., Hiscock, W.T., Laan, P., Lancelot,

- C., Law, C.S., Nojiri, Y., Oijen, T. Van, Riebesell, U., Rijkenberg, M.J.A., Waite, A.M., Wong, C., de Baar, H.J.W., 2005. Synthesis of iron fertilization experiments: From the Iron Age in the Age of Enlightenment. *J. Geophys. Res.* 110, 1–24.
- de Baar, H.J.W., Buma, A.G.J., Nolting, R.F., Cadee, G.C., Jacques, G., Treguer, P.J., 1990. On iron limitation of the Southern Ocean: experimental observations in the Weddell and Scotia Seas. *Mar. Ecol. Prog. Ser.* 65, 105–122.
- de Baar, H.J.W., de Jong, J.T.M., Bakker, D.C.E., Löscher, B.M., Veth, C., Bathmann, U., Smetacek, V., 1995. Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean. *Nature* 373, 412–415.
- de Baar, H.J.W., La Roche, J., 2003. Trace Metals in the Oceans: Evolution, Biology and Global Change. In: Wefer, G., Lamy, F., Mantoura, F. (Eds.), *Marine Science Frontiers for Europe*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 79–105.
- de Baar, H.J.W., Timmermans, K.R., Laan, P., De Porto, H.H., Ober, S., Blom, J.J., Bakker, M.C., Schilling, J., Sarthou, G., Smit, M.G., Klunder, M., Baar, H.J.W. De, Timmermans, K.R., Laan, P., Porto, H.H. De, Ober, S., 2008. Titan: A new facility for ultraclean sampling of trace elements and isotopes in the deep oceans in the international Geotraces program. *Mar. Chem.* 111, 4–21.
- de Baar, H.J.W., van Heuven, S.M.A.C., Middag, R., 2017. Ocean Biochemical Cycling and Trace Elements. In: White, W.M. (Ed.), *Encyclopedia of Geochemistry: A Comprehensive Reference Source on the Chemistry of the Earth*. Springer International Publishing, Cham, pp. 1–21.
- de Jong, J., Schoemann, V., Lannuzel, D., Croot, P., de Baar, H., Tison, J., Jong, J. De, Schoemann, V., Lannuzel, D., Croot, P., Baar, H. De, Tison, J., 2012. Natural iron fertilization of the Atlantic sector of the Southern Ocean by continental shelf sources of the Antarctic Peninsula. *J. Geophys. Res.* 117.
- Dulaquais, G., Waeles, M., Gerringsa, L.J.A., Middag, R., Rijkenberg, M.J.A., Riso, R., 2018. The biogeochemistry of electroactive humic substances and its connection to iron chemistry in the North East Atlantic and the Western Mediterranean Sea. *J. Geophys. Res. Ocean*.
- Edmonds, H.N., Michael, P.J., Baker, E.T., Connelly, D.P., Snow, J.E., Langmuir, C.H., Dick, H.J.B., Mühe, R., German, C.R., Graham, D.W., 2003. Discovery of abundant hydrothermal venting on the ultraslow-spreading Gakkel ridge in the Arctic Ocean. *Nature* 421, 252–256.
- Fernández-Méndez, M., Katlein, C., Rabe, B., Nicolaus, M., Peeken, I., Bakker, K., Flores, H., Boetius, A., 2015. Photosynthetic production in the central Arctic Ocean during the record sea-ice minimum in 2012. *Biogeosciences* 12, 3525–3549.
- Field, C.B., Behrenfeld, M.J., Randerson, J.T., Falkowski, P.G., 1998. Primary Production of the Biosphere: Integrating Terrestrial and Oceanic Components. *Science* (80-.). 281, 237–240.
- Fishwick, M.P., Sedwick, P.N., Lohan, M.C., Worsfold, P.J., Buck, K.N., Church, T.M., Ussher, S.J., 2014. Global Biogeochemical Cycles on the dissolution of aerosol iron.
- Fitzsimmons, J.N., Boyle, E.A., Jenkins, W.J., 2014. Distal transport of dissolved hydrothermal iron in the deep South Pacific Ocean. *Proc. Natl. Acad. Sci.* 111, 16654–16661.
- Fitzsimmons, J.N., Bundy, R.M., Al-Subiai, S.N., Barbeau, K.A., Boyle, E.A., 2015. The composition of dissolved iron in the dusty surface ocean: An exploration using size-fractionated iron-binding ligands. *Mar. Chem.* 173, 125–135.
- Fitzsimmons, J.N., John, S.G., Marsay, C.M., Hoffman, C.L., Nicholas, S.L., Toner, B.M., German, C.R., Sherrell, R.M., 2017. Iron persistence in a distal hydrothermal plume supported by dissolved–particulate exchange. *Nat. Geosci.* 10, 195–201.
- Fortier, M., Fortier, L., Michel, C., Legendre, L., 2002. Climatic and biological forcing of

- the vertical flux of biogenic particles under seasonal Arctic sea ice. Mar. Ecol. Prog. Ser. 225, 1–16.
- Francis, J., Madinaveitia, J., Macturk, H.M., Snow, G.A., 1949. Isolation from Acid-Fast Bacteria of a Growth-Factor for *Mycobacterium johnei* and of a Precursor of Phthiocol. Nature 163, 365–366.
- Frey, K.E., McClelland, J.W., 2009. Impacts of permafrost degradation on arctic river biogeochemistry. Hydrol. Process. 23, 169–182.
- Garg, S., Rose, A.L., Waite, T.D., 2007. Superoxide-mediated reduction of organically complexed iron(III): Impact of pH and competing cations (Ca^{2+}). Geochim. Cosmochim. Acta 71, 5620–5634.
- Gascard, J.-C., 1973. Vertical motions in a region of deep water formation. In: Deep Sea Research and Oceanographic Abstracts. Elsevier, pp. 1011–1027.
- Gauglitz, J.M., Zhou, H., Butler, A., 2012. A suite of citrate-derived siderophores from a marine *Vibrio* species isolated following the Deepwater Horizon oil spill. J. Inorg. Biochem. 107, 90–95.
- Geider, R.J., La Roche, J., 1994. The role of iron in phytoplankton photosynthesis, and the potential for iron-limitation of primary productivity in the sea. Photosynth. Res. 39, 275–301.
- Geider, R.J., La Roche, J., Greene, R.M., Olaizola, M., 1993. Response of the photosynthetic apparatus of *Phaeodactylum Tricornutum* (Bacillariophyceae) to nitrate, phosphate, or iron starvation. J. Phycol. 29, 755–766.
- Genty, B., Briantais, J.-M., Baker, N.R., 1989. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. Biochim. Biophys. Acta 990, 87–92.
- Gerringa, L.J.A., Rijkenberg, M.J.A., Bown, J., Margolin, A.R., Laan, P., de Baar, H.J.W., 2016. Fe-Binding Dissolved Organic Ligands in the Oxic and Suboxic Waters of the Black Sea. Front. Mar. Sci. 3, 1–16.
- Gerringa, L.J.A., Rijkenberg, M.J.A., Schoemann, V., Laan, P., de Baar, H.J.W., 2015. Organic complexation of iron in the West Atlantic Ocean. Mar. Chem. 177, 434–446.
- Gerringa, L.J.A., Rijkenberg, M.J.A., Thuróczy, C.E., Maas, L.R.M., 2014. A critical look at the calculation of the binding characteristics and concentration of iron complexing ligands in seawater with suggested improvements. Environ. Chem. 11, 114–136.
- Gerringa, L.J.A., Rijkenberg, M.J.A., Wolterbeek, H.T., Verburg, T.G., Boye, M., de Baar, H.J.W., 2007. Kinetic study reveals weak Fe-binding ligand, which affects the solubility of Fe in the Scheldt estuary. Mar. Chem. 103, 30–45.
- Gerringa, L.J.A., Slagter, H.A., Bown, J., van Haren, H., Laan, P., de Baar, H.J.W., Rijkenberg, M.J.A., 2017. Dissolved Fe and Fe-binding organic ligands in the Mediterranean Sea - GEOTRACES G04. Mar. Chem. 194, 100–113.
- Gerringa, L.J.A., Veldhuis, M.J.W., Timmermans, K.R., Sarthou, G., de Baar, H.J.W., 2006. Co-variance of dissolved Fe-binding ligands with phytoplankton characteristics in the Canary Basin. Mar. Chem. 102, 276–290.
- Gledhill, M., Buck, K.N., 2012. The organic complexation of iron in the marine environment: a review. Front. Microbiol. 3, 69.
- Gledhill, M., Gerringa, L.J.A., 2017. The Effect of Metal Concentration on the Parameters Derived from Complexometric Titrations of Trace Elements in Seawater—A Model Study. Front. Mar. Sci. 4, 254.
- Gledhill, M., McCormack, P., Ussher, S., Achterberg, E.P., Mantoura, R.F.C., Worsfold, P.J., 2004. Production of siderophore type chelates by mixed bacterioplankton populations in nutrient enriched seawater incubations. Mar. Chem. 88, 75–83.
- Gledhill, M., van den Berg, C.M.G., 1994. Determination of complexation of iron(III) with

- natural organic complexing ligands in seawater using cathodic stripping voltammetry. *Mar. Chem.* 47, 41–54.
- Gobler, C.J., Hutchins, D.A., Fisher, N.S., 1997. Release and bioavailability of C, N, P, Se, and Fe following viral lysis of a marine chrysophyte. *Limnol. Oceanogr.* 42, 1492–1504.
- González, A.G., Santana-Casiano, J.M., González-Dávila, M., Pérez-Almeida, N., Suárez De Tangil, M., 2014. Effect of *Dunaliella tertiolecta* organic exudates on the Fe(II) oxidation kinetics in seawater. *Environ. Sci. Technol.* 48, 7933–7941.
- Gordienko, P.A., Laktionov, A.F., 1969. Circulation and physics of the Arctic basin waters. In: Gordon, A.L., Baker, F.W.G. (Eds.), *Oceanography: Annals of The International Geophysical Year*, Volume 46. Pergamon Press, London, pp. 94–112.
- Grasshoff, K., 1983. Determination of nitrate. In: Grasshoff, K., Erhardt, M., Kremling, K. (Eds.), *Methods of Seawater Analysis*. Verlag Chemie, pp. 143–187.
- Gregor, D.J., Loeng, H., Barrie, L., 1998. The Influence of Physical and Chemical Processes on Contaminant Transport into and within the Arctic, Arctic Monitoring and Assessment Report: Arctic Pollution Issues.
- Guieu, C., Chester, R., Nimmo, M., Martin, J.M., Guerzoni, S., Nicolas, E., Mateu, J., Keyse, S., 1997. Atmospheric input of dissolved and particulate metals to the northwestern Mediterranean. *Deep. Res. Part II Top. Stud. Oceanogr.* 44, 655–674.
- Guieu, C., Dulac, F., Desboeufs, K., Wagener, T., Pulido-Villena, E., Grisoni, J.M., Louis, F., Ridame, C., Blain, S., Brunet, C., Bon Nguyen, E., Tran, S., Labiad, M., Dominici, J.M., 2010a. Large clean mesocosms and simulated dust deposition: a new methodology to investigate responses of marine oligotrophic ecosystems to atmospheric inputs. *Biogeosciences* 7, 2765–2784.
- Guieu, C., Loÿe-Pilot, M.D., Benyahya, L., Dufour, A., 2010b. Spatial variability of atmospheric fluxes of metals (Al, Fe, Cd, Zn and Pb) and phosphorus over the whole Mediterranean from a one-year monitoring experiment: Biogeochemical implications. *Mar. Chem.* 120, 164–178.
- Guieu, C., Martin, J.M., Thomas, A.J., Elbaz-Poulichet, F., 1991. Atmospheric versus river inputs of metals to the Gulf of Lions. Total concentrations, partitioning and fluxes. *Mar. Pollut. Bull.* 22, 176–183.
- Haas, C., Pfaffling, A., Hendricks, S., Rabenstein, L., Etienne, J.L., Rigor, I., 2008. Reduced ice thickness in Arctic Transpolar Drift favors rapid ice retreat. *Geophys. Res. Lett.* 35, 1–5.
- Hassler, C.S., Alasonati, E., Mancuso Nichols, C.A., Slaveykova, V.I., 2011a. Exopolysaccharides produced by bacteria isolated from the pelagic Southern Ocean – Role in Fe binding, chemical reactivity, and bioavailability. *Mar. Chem.* 123, 88–98.
- Hassler, C.S., Schoemann, V., Nichols, C.M., Butler, E.C. V, Boyd, P.W., 2011b. Saccharides enhance iron bioavailability to Southern Ocean phytoplankton. *Proc. Natl. Acad. Sci. U. S. A.* 108, 1076–81.
- Hassler, C.S., van den Berg, C.M.G., Boyd, P.W., 2017. Toward a Regional Classification to Provide a More Inclusive Examination of the Ocean Biogeochemistry of Iron-Binding Ligands. *Front. Mar. Sci.* 4, 19.
- Hatta, M., Measures, C.I., Wu, J., Roshan, S., Fitzsimmons, J.N., Sedwick, P., Morton, P., 2015. An overview of dissolved Fe and Mn distributions during the 2010–2011 U.S. GEOTRACES north Atlantic cruises: GEOTRACES GA03. *Deep. Res. Part II Top. Stud. Oceanogr.* 116, 117–129.
- Hawkes, J.A., Connelly, D.P., Gledhill, M., Achterberg, E.P., 2013. The stabilisation and transportation of dissolved iron from high temperature hydrothermal vent systems. *Earth Planet. Sci. Lett.* 375, 280–290.

- Heimbürger, L.E., Migon, C., Losno, R., Miquel, J.C., Thibodeau, B., Stabholz, M., Dufour, A., Leblond, N., 2014. Vertical export flux of metals in the Mediterranean Sea. Deep. Res. Part I Oceanogr. Res. Pap. 87, 14–23.
- Helms, J.R., Stubbins, A., Ritchie, J.D., Minor, E.C., Kieber, D.J., Mopper, K., 2008. Absorption spectral slopes and slope ratios as indicators of molecular weight, source, and photobleaching of chromophoric dissolved organic matter. Limnology Oceanogr. 53, 955–969.
- Hirst, C., Andersson, P.S., Shaw, S., Burke, I.T., Kutscher, L., Murphy, M.J., Maximov, T., Pokrovsky, O.S., Mört, C.M., Porcelli, D., 2017. Characterisation of Fe-bearing particles and colloids in the Lena River basin, NE Russia. Geochim. Cosmochim. Acta 213, 553–573.
- Hofmann, T., Hanlon, A.R.M., Taylor, J.D., Ball, A.S., Osborn, A.M., Underwood, G.J.C., 2009. Dynamics and compositional changes in extracellular carbohydrates in estuarine sediments during degradation. Mar. Ecol. Prog. Ser. 379, 45–58.
- Hogle, S.L., Barbeau, K.A., Gledhill, M., 2014. Heme in the marine environment: from cells to the iron cycle. Metallomics 6, 1107–1120.
- Hölemann, J.A., Schirmacher, M., Kassens, H., Prange, A., 1999. Geochemistry of surficial and ice-rafted sediments from the Laptev Sea (Siberia). Estuar. Coast. Shelf Sci. 49, 45–59.
- Hopkinson, B.M., Barbeau, K.A., 2007. Organic and redox speciation of iron in the eastern tropical North Pacific suboxic zone. Mar. Chem. 106, 2–17.
- Hopkinson, B.M., Morel, F.M., Franc, B.M.H.Æ., Morel, M.M., 2009. The role of siderophores in iron acquisition by photosynthetic marine microorganisms. Biometals 22, 659–669.
- Hudson, R.J.M., Rue, E.L., Bruland, K.W., 2003. Modeling complexometric titrations of natural water samples. Environ. Sci. Technol. 37, 1553–1562.
- Hutchins, D.A., Wang, W.-X., Fisher, N.S., 1995. Copepod Grazing and the Biogeochemical Fate of Diatom Iron. Limnol. Oceanogr. 40, 989–994.
- Ibisanmi, E., Sander, S.G., Boyd, P.W., Bowie, A.R., Hunter, K.A., 2011. Vertical distributions of iron-(III) complexing ligands in the Southern Ocean. Deep Sea Res. Part II Top. Stud. Oceanogr. 58, 2113–2125.
- IOC, S., 2010. The international thermodynamic equation of seawater—2010: calculation and use of thermodynamic properties. Intergovernmental Oceanographic Commission, Manuals and Guides No. 56. In: Manual and Guides. UNESCO.
- IPCC, 2013. Summary for Policymakers. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1–30.
- IPCC, 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland.
- Jiao, N., Herndl, G.J., Hansell, D.A., Benner, R., Kattner, G., Wilhelm, S.W., Kirchman, D.L., Weinbauer, M.G., Luo, T., Chen, F., Azam, F., 2010. Microbial production of recalcitrant dissolved organic matter: Long-term carbon storage in the global ocean. Nat. Rev. Microbiol. 8, 593–599.
- Johansen, A.M., Siefert, R.L., Hoffmann, M.R., 2000. Chemical composition of aerosols collected over the tropical North Atlantic Ocean. J. Geophys. Res. 105, 15277–15312.
- Johnson, K.S., Boyle, E., Bruland, K., Coale, K., Measures, C., Moffett, J., Aguilar-Islas,

- A., Barbeau, K., Bergquist, B., Bowie, A., Buck, K., Cai, Y., Chase, Z., Cullen, J., Doi, T., Elrod, V., Fitzwater, S., Gordon, M., King, A., Laan, P., Laglera-Baquer, L., Landing, W., Lohan, M., Mendez, J., Milne, A., Obata, H., Ossiander, L., Plant, J., Sarthou, G., Sedwick, P., Smith, G.J., Sohst, B., Tanner, S., Van den Berg, S., Wu, J., 2007. Developing standards for dissolved iron in seawater. *Eos (Washington, DC)*. 88, 131–132.
- Jones, M.E., Beckler, J.S., Taillefert, M., 2011. The flux of soluble organic-iron(III) complexes from sediments represents a source of stable iron(III) to estuarine waters and to the continental shelf. *Limnol. Oceanogr.* 56, 1811–1823.
- Journet, E., Desboeufs, K. V., Caquineau, S., Colin, J.-L., 2008. Mineralogy as a critical factor of dust iron solubility. *Geophys. Res. Lett.* 35, n/a-n/a.
- Kalle, K., 1937. Meereskundliche chemische Untersuchungen mit Hilfe des Zeisschen Pulfrich Photometers. *Ann. Hydrogr.*, Berl. 65, 276.
- Kalle, K., 1949. Fluoreszenz und Gelbstoff im Bottnischen und Finnischen Meerbusen. *Dtsch. Hydrogr. Zeitschrift* 2, 117–124.
- Kessner, D., Chambers, M., Burke, R., Agus, D., Mallick, P., 2008. ProteoWizard: Open source software for rapid proteomics tools development. *Bioinformatics* 24, 2534–2536.
- King, A.L., Buck, K.N., Barbeau, K.A., 2012. Quasi-Lagrangian drifter studies of iron speciation and cycling off Point Conception, California. *Mar. Chem.* 128–129, 1–12.
- Kleint, C., Hawkes, J.A., Sander, S.G., Koschinsky, A., 2016. Voltammetric Investigation of Hydrothermal Iron Speciation. *Front. Mar. Sci.* 3, 1–11.
- Klunder, M.B., Bauch, D., Laan, P., de Baar, H.J.W., van Heuven, S., Ober, S., 2012a. Dissolved iron in the Arctic shelf seas and surface waters of the central Arctic Ocean: Impact of Arctic river water and ice-melt. *J. Geophys. Res.* 117.
- Klunder, M.B., Laan, P., Middag, R., de Baar, H.J.W., Bakker, K., 2012b. Dissolved iron in the Arctic Ocean: Important role of hydrothermal sources, shelf input and scavenging removal. *J. Geophys. Res.* 117, C04014.
- Klunder, M.B., Laan, P., Middag, R., De Baar, H.J.W., van Ooijen, J.C., Baar, H.J.W. De, Ooijen, J.C. Van, 2011. Dissolved iron in the Southern Ocean (Atlantic sector). *Deep Sea Res. Part II* 58, 2678–2694.
- Koron, N., Ogrinc, N., Metzger, E., Riedel, B., Faganelli, J., 2015. The impact of induced redox transitions on nutrient diagenesis in coastal marine sediments (Gulf of Trieste, northern Adriatic Sea). *J. Soils Sediments* 15, 2443–2452.
- Korte, L.F., Brummer, G.J.A., Van Der Does, M., Guerreiro, C. V., Hennekam, R., Van Hateren, J.A., Jong, Di., Munday, C.I., Schouten, S., Stuut, J.B.W., 2017. Downward particle fluxes of biogenic matter and Saharan dust across the equatorial North Atlantic. *Atmos. Chem. Phys.* 17, 6023–6040.
- Kustka, A.B., Shaked, Y., Milligan, A.J., King, D.W., Morel, F.M.M., 2005. Extracellular production of superoxide by marine diatoms: Contrasting effects on iron redox chemistry and bioavailability. *Limnol. Oceanogr.* 50, 1172–1180.
- Kwok, R., Cunningham, G.F., Wensnahan, M., Rigor, I., Zwally, H.J., Yi, D., 2009. Thinning and volume loss of the Arctic Ocean sea ice cover: 2003–2008. *J. Geophys. Res. Ocean.* 114, 2003–2008.
- Laglera, L.M., Battaglia, G., van den Berg, C.M.G., 2007. Determination of humic substances in natural waters by cathodic stripping voltammetry of their complexes with iron. *Anal. Chim. Acta* 599, 58–66.
- Laglera, L.M., Battaglia, G., van den Berg, C.M.G., 2011. Effect of humic substances on the iron speciation in natural waters by CLE/CSV. *Mar. Chem.* 127, 134–143.
- Laglera, L.M., Downes, J., Santos-Echeandia, J., Santos-Echeandía, J., 2013. Comparison and combined use of linear and non-linear fitting for the estimation of complexing

- parameters from metal titrations of estuarine samples by CLE/AdCSV. *Mar. Chem.* 155, 102–112.
- Laglera, L.M., Filella, M., 2015. The relevance of ligand exchange kinetics in the measurement of iron speciation by CLE–AdCSV in seawater. *Mar. Chem.* 173, 100–113.
- Laglera, L.M., Santos-Echeandía, J., Caprara, S., Monticelli, D., Santos-Echeandía, J., Caprara, S., Monticelli, D., Santos-Echeandía, J., Caprara, S., Monticelli, D., 2013. Quantification of iron in seawater at the low picomolar range based on optimization of bromate/ammonia/dihydroxynaphthalene system by catalytic adsorptive cathodic stripping voltammetry. *Anal. Chem.* 85, 2486–2492.
- Laglera, L.M., Tovar-Sánchez, A., Iversen, M.H., González, H.E., Naik, H., Mangesh, G., Assmy, P., Klaas, C., Mazzocchi, M.G., Montresor, M., Naqvi, S.W.A., Smetacek, V., Wolf-Gladrow, D.A., 2017. Iron partitioning during LOHAFEX: Copepod grazing as a major driver for iron recycling in the Southern Ocean. *Mar. Chem.*
- Laglera, L.M., van den Berg, C.M.G., 2009. Evidence for geochemical control of iron by humic substances in seawater. *Limnol. Oceanogr.* 54, 610–619.
- Lam, P.J., Bishop, J.K.B., Henning, C.C., Marcus, M.A., Waychunas, G.A., Fung, I.Y., 2006. Wintertime phytoplankton bloom in the subarctic Pacific supported by continental margin iron. *Global Biogeochem. Cycles* 20, n/a-n/a.
- Langmuir, I., 1916. The Constitution and Fundamental Properties of Solids and Liquids. Part I. Solids. *J. Am. Chem. Soc.* 252, 2221–2295.
- Lannuzel, D., Bowie, A.R., Remenyi, T., Lam, P., Townsend, A., Ibisarri, E., Butler, E., Wagener, T., Schoemann, V., 2011. Distributions of dissolved and particulate iron in the sub-Antarctic and Polar Frontal Southern Ocean (Australian sector). *Deep Sea Res. Part II Top. Stud. Oceanogr.* 58, 2094–2112.
- Laukert, G., Frank, M., Bauch, D., Hathorne, E.C., Rabe, B., von Appen, W.-J., Wegner, C., Zieringer, M., Kassens, H., 2017. Ocean circulation and freshwater pathways in the Arctic Mediterranean based on a combined Nd isotope, REE and oxygen isotope section across Fram Strait. *Geochim. Cosmochim. Acta* 202, 285–309.
- Lazar, C.S., John Parkes, R., Cragg, B.A., L'Haridon, S., Toffin, L., 2012. Methanogenic activity and diversity in the centre of the Amsterdam Mud Volcano, Eastern Mediterranean Sea. *FEMS Microbiol. Ecol.* 81, 243–254.
- Liu, X., Millero, F.J., 2002. The solubility of iron in seawater. *Mar. Chem.* 77, 43–54.
- Lønborg, C., Middelboe, M., Brussaard, C.P.D., 2013. Viral lysis of *Micromonas pusilla*: Impacts on dissolved organic matter production and composition. *Biogeochemistry* 116, 231–240.
- Lupton, J., Ronde, C. de, Sprovieri, M., Baker, E.T., Bruno, P.P., Italiano, F., Walker, S., Faure, K., Leybourne, M., Britten, K., Greene, R., 2011. Active hydrothermal discharge on the submarine Aeolian Arc. *J. Geophys. Res. Solid Earth* 116.
- Maat, D.S., Brussaard, C.P.D., 2016. Both phosphorus and nitrogen limitation constrain viral proliferation in marine phytoplankton. *Aquat. Microb. Ecol.*
- Maat, D.S., Crawfurd, K.J., Timmermans, K.R., Brussaard, C.P.D., 2014. Elevated CO₂ and phosphate limitation favor *Micromonas pusilla* through stimulated growth and reduced viral impact. *Appl. Environ. Microbiol.* 80, 3119–27.
- Maat, D.S., Van Bleijswijk, J.D.L., Witte, H.J., Brussaard, C.P.D., 2016. Virus production in phosphorus limited *Micromonas pusilla* stimulated by a supply of naturally low concentrations of different phosphorus sources, far into the lytic cycle. *FEMS Microbiol. Ecol.* fiw136.
- Macdonald, R.W., Harner, T., Fyfe, J., 2005. Recent climate change in the Arctic and its impact on contaminant pathways and interpretation of temporal trend data. *Sci. Total Environ.* 342, 5–86.

- Mahmood, A., Abualhaija, M.M., van den Berg, C.M.G., Sander, S.G., 2015. Organic speciation of dissolved iron in estuarine and coastal waters at multiple analytical windows. *Mar. Chem.* 177, 706–719.
- Maldonado, M.T., Strzepek, R.F., Sander, S., Boyd, P.W., 2005. Acquisition of iron bound to strong organic complexes, with different Fe binding groups and photochemical reactivities, by plankton communities in Fe-limited subantarctic waters. *Global Biogeochem. Cycles* 19, n/a-n/a.
- Mari, X., Rassoulzadegan, F., Brussaard, C.P.D., Wassmann, P., 2005. Dynamics of transparent exopolymeric particles (TEP) production by *Phaeocystis globosa* under N- or P-limitation: A controlling factor of the retention/export balance. *Harmful Algae* 4, 895–914.
- Martin, J.H., Fitzwater, S.E., 1988. Iron deficiency limits phytoplankton growth in the north-east Pacific subarctic. *Nature* 331, 341–343.
- Martin, J.H., Fitzwater, S.E., Gordon, R.M., 1990. Iron deficiency limits phytoplankton growth in Antarctic waters. *Global Biogeochem. Cycles* 4, 5–12.
- Martínez Martínez, J., Boere, A., Gilg, I., van Lent, J.W.M., Witte, H.J., van Bleijswijk, J.D.L., Brussaard, C.P.D., 2015. New lipid envelope-containing dsDNA virus isolates infecting *Micromonas pusilla* reveal a separate phylogenetic group. *Aquat. Microb. Ecol.* 74, 17–28.
- Martínez Martínez, J., Swan, B.K., Wilson, W.H., 2014. Marine viruses, a genetic reservoir revealed by targeted viromics. *ISME J.* 8, 1079–1088.
- Mascle, J., Mary, F., Praeg, D., Brosolo, L., Camera, L., Ceramicola, S., Dupré, S., 2014. Distribution and geological control of mud volcanoes and other fluid/free gas features in the Mediterranean Sea and nearby Gulf of Cadiz. *Geo-Marine Lett.* 34, 89–110.
- Maslanik, J., Stroeve, J., Fowler, C., Emery, W., 2011. Distribution and trends in Arctic sea ice age through spring 2011. *Geophys. Res. Lett.* 38, 2–7.
- Mawji, E., Gledhill, M., Milton, J.A., Tarran, G.A., Ussher, S., Thompson, A., Wolff, G.A., Worsfold, P.J., Achterberg, E.P., 2008. Hydroxamate siderophores: occurrence and importance in the Atlantic Ocean. *Environ. Sci. Technol.* 42, 8675–8680.
- Mawji, E., Gledhill, M., Milton, J.A., Zubkov, M. V., Thompson, A., Wolff, G.A., Achterberg, E.P., 2011. Production of siderophore type chelates in Atlantic Ocean waters enriched with different carbon and nitrogen sources. *Mar. Chem.* 124, 90–99.
- Mawji, E., Gledhill, M., Worsfold, P.J., Achterberg, E.P., 2008. Collision-induced dissociation of three groups of hydroxamate siderophores: ferrioxamines, ferrichromes and coprogens/fusigens. *Rapid Commun. Mass Spectrom.* 22, 2195–2202.
- McCormack, P., Worsfold, P.J., Gledhill, M., 2003. Separation and Detection of Siderophores Produced by Marine Bacterioplankton Using High-Performance Liquid Chromatography with Electrospray Ionization Mass Spectrometry. *Anal Chem* 75, 2647–2652.
- Mengel, K., 1994. Iron availability in plant tissues - iron chlorosis on calcareous soils. *Plant Soil* 165, 275–283.
- Menzel Barraqueta, J., Klar, J.K., Gledhill, M., Schlosser, C., Shelley, R., Wenzel, B., Sarthou, G., Achterberg, E.P., 2018. Atmospheric aerosol deposition fluxes over the Atlantic Ocean : A GEOTRACES case study. *Biogeosciences Discuss.* 1–25.
- Middag, R., de Baar, H.J.W., Laan, P., Klunder, M.B., 2011. Fluvial and hydrothermal input of manganese into the Arctic Ocean. *Geochim. Cosmochim. Acta* 75, 2393–2408.
- Middag, R., Séférian, R., Conway, T.M., John, S.G., Bruland, K.W., de Baar, H.J.W., 2015. Intercomparison of dissolved trace elements at the Bermuda Atlantic Time Series

- station. Mar. Chem. 177, 476–489.
- Millero, F.J., 1998. Solubility of Fe (III) in seawater. Earth Planet. Sci. Lett. 154, 323–329.
- Millot, C., 1999. Circulation in the Western Mediterranean Sea. J. Mar. Syst. 20, 423–442.
- Millot, C., Taupier-Letage, I., 2005. Circulation in the Mediterranean Sea. pp. 29–66.
- Mills, M.M., Brown, Z.W., Lowry, K.E., van Dijken, G.L., Becker, S., Pal, S., Benitez-Nelson, C.R., Downer, M.M., Strong, A.L., Swift, J.H., Pickart, R.S., Arrigo, K.R., 2015. Impacts of low phytoplankton NO₃-: PO₄³⁻ utilization ratios over the Chukchi Shelf, Arctic Ocean. Deep. Res. Part II Top. Stud. Oceanogr. 118, 105–121.
- Mioni, C.E., Poorvin, L., Wilhelm, S.W., 2005. Virus and siderophore-mediated transfer of available Fe between heterotrophic bacteria: characterization using an Fe-specific bioreporter. Aquat. Microb. Ecol. 41, 233–245.
- Mojica, K.D.A., Brussaard, C.P.D., 2014. Factors affecting virus dynamics and microbial host-virus interactions in marine environments. FEMS Microbiol. Ecol. 89, 495–515.
- Mojica, K.D.A., Huisman, J., Wilhelm, S.W., Brussaard, C.P.D., 2016. Latitudinal variation in virus-induced mortality of phytoplankton across the North Atlantic Ocean. ISME J. 10, 500–513.
- Moore, J.K., Braucher, O., 2008. Sedimentary and mineral dust sources of dissolved iron to the World Ocean. Biogeosciences 5, 631–656.
- Mopper, K., Schultz, C.A., 1993. Fluorescence as a possible tool for studying the nature and water column distribution of DOC components. Mar. Chem. 41, 229–238.
- Mundy, C.J., Gosselin, M., Ehn, J., Gratton, Y., Rossnagel, A., Barber, D.G., Martin, J., Tremblay, J.É., Palmer, M., Arrigo, K.R., Darnis, G., Fortier, L., Else, B., Papakyriakou, T., 2009. Contribution of under-ice primary production to an ice-edge upwelling phytoplankton bloom in the Canadian Beaufort Sea. Geophys. Res. Lett. 36, 1–5.
- Murphy, J., Riley, J.P., 1962. A modified single solution method for the determination of phosphate in natural waters. Anal. Chim. Acta 27, 31–36.
- Nakayama, Y., Fujita, S., Kuma, K., Shimada, K., 2011. Iron and humic-type fluorescent dissolved organic matter in the Chukchi Sea and Canada Basin of the western Arctic Ocean. J. Geophys. Res. Ocean. 116, 1–16.
- National Snow and Ice Data Center, 2012. Arctic sea ice extent settles at record seasonal minimum [WWW Document]. Natl. Snow Ice Data Cent. URL <http://nsidc.org/arcticseainews/2012/09/arctic-sea-ice-extent-settles-at-record-seasonal-minimum/> (accessed 10.20.17).
- National Snow and Ice Data Center, 2015. Arctic sea ice reaches fourth lowest minimum [WWW Document]. Natl. Snow Ice Data Cent. URL http://nsidc.org/arcticseainews/2015/09/2015_arctic-minimum/ (accessed 10.20.17).
- Netz, D.J.A., Stith, C.M., Stümpfig, M., Köpf, G., Vogel, D., Genau, H.M., Stodola, J.L., Lill, R., Burgers, P.M.J., Pierik, A.J., 2012. Eukaryotic DNA polymerases require an iron-sulfur cluster for the formation of active complexes. Nat. Chem. Biol. 8, 125–132.
- Nishino, S., Kikuchi, T., Yamamoto-Kawai, M., Kawaguchi, Y., Hirawake, T., Itoh, M., 2011. Enhancement/reduction of biological pump depends on ocean circulation in the sea-ice reduction regions of the Arctic Ocean. J. Oceanogr. 67, 305–314.
- Nomikou, P., Papanikolaou, D., Alexandri, M., Sakellariou, D., Rousakis, G., 2013. Submarine volcanoes along the aegean volcanic arc. Tectonophysics 597–598, 123–146.
- Norman, L., Worms, I.A.M., Angles, E., Bowie, A.R., Nichols, C.M., Ninh Pham, A.,

- Slaveykova, V.I., Townsend, A.T., David Waite, T., Hassler, C.S., 2015. The role of bacterial and algal exopolymeric substances in iron chemistry. Mar. Chem. 173, 148–161.
- Not, F., Latasa, M., Marie, D., Cariou, T., Vaulot, D., Simon, N., 2004. A single species, *Micromonas pusilla* (Prasinophyceae), dominates the eukaryotic picoplankton in the Western English Channel. Appl. Environ. Microbiol. 70, 4064–4072.
- Not, F., Massana, R., Latasa, M., Marie, D., Colson, C., Eikrem, W., Pedrós-Alió, C., Vaulot, D., Simon, N., 2005. Late summer community composition and abundance of photosynthetic picoeukaryotes in Norwegian and Barents seas. Limnol. Oceanogr. 50, 1677–1686.
- Obata, H., Karatani, H., Nakayama, E., 1993. Automated Determination of Iron in Seawater by Chelating Resin Concentration and Chemiluminescence Detection. Anal. Chem. 65, 1524–1528.
- Obernosterer, I., Herndl, G.J., 2000. Differences in the optical and biological reactivity of the humic and nonhumic dissolved organic carbon component in two contrasting coastal marine environments 45, 1120–1129.
- Olli, K., Wassmann, P., Reigstad, M., Ratkova, T.N., Arashkevich, E., Pasternak, A., Matrai, P.A., Knulst, J., Tranvik, L., Klais, R., Jacobsen, A., 2007. The fate of production in the central Arctic Ocean - top-down regulation by zooplankton expatriates? Prog. Oceanogr. 72, 84–113.
- Omoregie, E.O., Mastalerz, V., De Lange, G., Straub, K.L., Kappler, A., Røy, H., Stadnitskaia, A., Foucher, J.P., Boetius, A., 2008. Biogeochemistry and community composition of iron- and sulfur-precipitating microbial mats at the Chefren mud volcano (Nile deep sea fan, eastern Mediterranean). Appl. Environ. Microbiol. 74, 3198–3215.
- Parekh, P., Follows, M.J., Boyle, E.A., 2005. Decoupling of iron and phosphate in the global ocean. Global Biogeochem. Cycles 19, 1–16.
- Paris, R., Desboeufs, K. V., Journet, E., 2011. Variability of dust iron solubility in atmospheric waters: Investigation of the role of oxalate organic complexation. Atmos. Environ. 45, 6510–6517.
- Passmore, R., Hsu, J., Liu, R.X., Tam, E., Cai, Y., Su, W., Frasca, J., Brigden, S.M., Comeau, A.M., Ortmann, A.C., Lavergne, A., Suttle, C.A., 2000. MPN Assay Analyzer.
- Passow, U., 2002. Transparent exopolymer particles (TEP) in aquatic environments. Prog. Oceanogr. 55, 287–333.
- Paucot, H., Wollast, R., 1997. Transport and transformation of trace metals in the Scheldt estuary. Mar. Chem. 58, 229–244.
- Pekey, H., 2006. The distribution and sources of heavy metals in Izmit Bay surface sediments affected by a polluted stream. Mar. Pollut. Bull. 52, 1197–1208.
- Pernet-Coudrier, B., Waeles, M., Filella, M., Quentel, F., Riso, R.D., 2013. Simple and simultaneous determination of glutathione, thioacetamide and refractory organic matter in natural waters by DP-CSV. Sci. Total Environ. 463–464, 997–1005.
- Peterson, B.J., Holmes, R.M., McClelland, J.W., Vörösmarty, C.J., Lammers, R.B., Shiklomanov, A.I., Shiklomanov, I.A., Rahmstorf, S., 2002. Increasing river discharge to the Arctic Ocean. Science 298, 2171–3.
- Peterson, B.J., McClelland, J.W., Curry, R., Holmes, R.M., Walsh, J.E., Aagaard, K., 2006. Trajectory Shifts in the Arctic and Subarctic Freshwater Cycle. Science (80-.). 313, 1061–1066.
- Pižeta, I., Sander, S.G., Hudson, R.J.M., Omanović, D., Baars, O., Barbeau, K.A., Buck, K.N., Bundy, R.M., Carrasco, G., Croot, P.L., Garnier, C., Gerringa, L.J.A., Gledhill, M., Hirose, K., Kondo, Y., Laglera, L.M., Nuester, J., Rijkenberg, M.J.A., Takeda, S.,

- Twining, B.S., Wells, M., 2015. Interpretation of complexometric titration data: An intercomparison of methods for estimating models of trace metal complexation by natural organic ligands. *Mar. Chem.* 173, 3–24.
- Pluskal, T., Castillo, S., Villar-Briones, A., Orešič, M., 2010. MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics* 11, 395.
- Pollak, M.J., 1951. The sources of the deep water of the eastern Mediterranean Sea. *J. Mar. Res.* 10, 128–152.
- Poorvin, L., Rinta-Kanto, J.M., Hutchins, D.A., Wilhelm, S.W., 2004. Viral release of iron and its bioavailability to marine plankton. *Limnol. Oceanogr.* 49, 1734–1741.
- Poorvin, L., Sander, S.G., Velasquez, I., Ibisanzmi, E., LeCleir, G.R., Wilhelm, S.W., 2011. A comparison of Fe bioavailability and binding of a catecholate siderophore with virus-mediated lysates from the marine bacterium *Vibrio alginolyticus* PWH3a. *J. Exp. Mar. Bio. Ecol.* 399, 43–47.
- Powell, R.T., Wilson-Finelli, A., 2003a. Importance of organic Fe complexing ligands in the Mississippi River plume. *Estuar. Coast. Shelf Sci.* 58, 757–763.
- Powell, R.T., Wilson-Finelli, A., 2003b. Photochemical degradation of organic iron complexing ligands in seawater. *Aquat. Sci. - Res. Across Boundaries* 65, 367–374.
- Press, W.H., Teukolsky, S.A., Vetterling, W.T., Flannery, B.P., 2007. *Numerical Recipes: The Art of Scientific Computing*, 3rd ed. Cambridge University Press, Cambridge, United Kingdom.
- Price, N.M., Ahner, B.A., Morel, F.M.M., 1994. The equatorial Pacific Ocean: grazer controlled phytoplankton populations in an iron-limited ecosystem. *Limnol. Oceanogr.* 39, 520–534.
- Puig, P., Madron, X.D. de, Salat, J., Schroeder, K., Martín, J., Karageorgis, A.P., Palanques, A., Roullier, F., Lopez-Jurado, J.L., Emelianov, M., Moutin, T., Houpert, L., 2013. Thick bottom nepheloid layers in the western Mediterranean generated by deep dense shelf water cascading. *Prog. Oceanogr.* 111, 1–23.
- Quentel, F., Filella, M., 2008. Quantification of refractory organic substances in freshwaters: Further insight into the response of the voltammetric method. *Anal. Bioanal. Chem.* 392, 1225–1230.
- R Development Core Team, 2008. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rabe, B., Karcher, M., Kauker, F., Schauer, U., Toole, J.M., Krishfield, R.A., Pisarev, S., Kikuchi, T., Su, J., 2014. Arctic Ocean basin liquid freshwater storage trend 1992–2012. *Geophys. Res. Lett.* 41, 961–968.
- Rabe, B., Schauer, U., Ober, S., Horn, M., Hoppmann, M., Korhonen, M., Pisarev, S., Hampe, H., Villacíeros, N., Savý, J.P., Wisotzki, A., 2016. Physical oceanography during POLARSTERN cruise PS94 (ARK-XXIX/3). Bremerhaven.
- Raiswell, R., Anderson, T.F., 2005. Reactive iron enrichment in sediments deposited beneath euxinic bottom waters: constraints on supply by shelf recycling. *Geol. Soc. London, Spec. Publ.* 248, 179–194.
- Rank, D., Özsoy, E., Salihoglu, I., 1999. Oxygen-18, deuterium and tritium in the Black Sea and the Sea of Marmara. *J. Environ. Radioact.* 43, 231–245.
- Reader, H.E., Stedmon, C.A., Nielsen, N.J., Kritzberg, E.S., 2015. Mass and UV-visible spectral fingerprints of dissolved organic matter: sources and reactivity. *Front. Mar. Sci.* 2, 1–10.
- Redfield, A., Ketchum, B., Richards, F., 1963. The influence of organisms on the composition of sea water. In: Hill, M. (Ed.), *The Sea*. Interscience, New York, pp. 26–77.
- Redfield, A.C., 1958. The biological control of chemical factors in the environment. *Am.*

Sci. 46, 230A–221.

- Reinthalter, T., Sintes, E., Herndl, G.J., 2008. Dissolved organic matter and bacterial production and respiration in the sea-surface microlayer of the open Atlantic and the western Mediterranean Sea. Limnol. Oceanogr. 53, 122–136.
- Rijkenberg, M.J.A., de Baar, H.J.W., Bakker, K., Gerringsa, L.J.A., Keijzer, E., Laan, M., Laan, P., Middag, R., Ober, S., van Ooijen, J., Ossebaar, S., van Weerlee, E.M., Smit, M.G., 2015. "PRISTINE", a new high volume sampler for ultraclean sampling of trace metals and isotopes. Mar. Chem. 177, 501–509.
- Rijkenberg, M.J.A., Gerringsa, L.J. a, Timmermans, K.R., Fischer, A.C., Kroon, K.J., Buma, A.G.J., Wolterbeek, B.T., de Baar, H.J.W., 2008a. Enhancement of the reactive iron pool by marine diatoms. Mar. Chem. 109, 29–44.
- Rijkenberg, M.J.A., Middag, R., Laan, P., Gerringsa, L.J.A., Van Aken, H.M., Schoemann, V., De Jong, J.T.M., De Baar, H.J.W., 2014. The distribution of dissolved iron in the West Atlantic Ocean. PLoS One 9.
- Rijkenberg, M.J.A., Powell, C.F., Dall'Osto, M., Nielsdottir, M.C., Patey, M.D., Hill, P.G., Baker, A.R., Jickells, T.D., Harrison, R.M., Achterberg, E.P., 2008b. Changes in iron speciation following a Saharan dust event in the tropical North Atlantic Ocean. Mar. Chem. 110, 56–67.
- Rijkenberg, M.J.A., Slagter, H.A., Rutgers van der Loeff, M., van Ooijen, J., Gerringsa, L.J.A., 2018a. Dissolved Fe in the Deep and Upper Arctic Ocean With a Focus on Fe Limitation in the Nansen Basin. Front. Mar. Sci. 5, 1–14.
- Rijkenberg, M.J.A., Slagter, H.A., Rutgers van der Loeff, M., Van Ooijen, J., Gerringsa, L.J.A., 2018b. Dissolved Fe in the Arctic Ocean shows Fe limitation in the Nansen Basin. Front. Mar. Sci. 1–14.
- Rijkenberg, M.J.A., Steigenberger, S., Powell, C.F., van Haren, H., Patey, M.D., Baker, A.R., Achterberg, E.P., 2012. Fluxes and distribution of dissolved iron in the eastern (sub-) tropical North Atlantic Ocean. Global Biogeochem. Cycles 26, GB3004.
- Ringbom, A., Still, E., 1972. The calculation and use of a coefficients. Anal. Chim. Acta 59, 143–146.
- Rodellas, V., Garcia-Orellana, J., Masqué, P., Feldman, M., Weinstein, Y., 2015. Submarine groundwater discharge as a major source of nutrients to the Mediterranean Sea. Proc. Natl. Acad. Sci. 112, 3926–3930.
- Roeske, T., Loeff, M.R. vd, Middag, R., Bakker, K., 2012. Deep water circulation and composition in the Arctic Ocean by dissolved barium, aluminium and silicate. Mar. Chem. 132–133, 56–67.
- Roether, W., Klein, B., Manca, B.B., Theocharis, A., Kioroglou, S., 2007. Transient Eastern Mediterranean deep waters in response to the massive dense-water output of the Aegean Sea in the 1990s. Prog. Oceanogr. 74, 540–571.
- Rolison, J.M., Middag, R., Stirling, C.H., Rijkenberg, M.J.A., de Baar, H.J.W., 2015. Zonal distribution of dissolved aluminium in the Mediterranean Sea. Mar. Chem. 177, 1–14.
- Romeo, A.M., Christen, L., Niles, E.G., Kosman, D.J., 2001. Intracellular chelation of iron by bipyridyl inhibits DNA virus replication: Ribonucleotide reductase maturation as a probe of intracellular iron pools. J. Biol. Chem. 276, 24301–24308.
- Rudels, B., 2008. Arctic Ocean Circulation. In: Steele, J., Turekian, K., Thorpe, S. (Eds.), Encyclopedia of Ocean Sciences. Academic Press, pp. 211–225.
- Rudels, B., 2012. Arctic Ocean circulation and variability - Advection and external forcing encounter constraints and local processes. Ocean Sci. 8, 261–286.
- Rudels, B., 2015. Arctic Ocean circulation, processes and water masses: A description of observations and ideas with focus on the period prior to the International Polar Year 2007–2009. Prog. Oceanogr. 132, 22–67.

- Rue, E., Bruland, K., 2001. Domoic acid binds iron and copper: A possible role for the toxin produced by the marine diatom *Pseudo-nitzschia*. Mar. Chem. 76, 127–134.
- Rue, E.L., Bruland, K.W., 1995. Complexation of iron(III) by natural organic ligands in the Central North Pacific as determined by a new competitive ligand equilibration/adsorptive cathodic stripping voltammetric method. Mar. Chem. 50, 117–138.
- Rutgers van der Loeff, M., Cai, P., Stimac, I., Bauch, D., Hanfland, C., Roeske, T., Moran, S.B., 2012. Shelf-basin exchange times of Arctic surface waters estimated from $^{228}\text{Th}/^{228}\text{Ra}$ disequilibrium. J. Geophys. Res. Ocean. 117.
- Rutgers van der Loeff, M.M., Key, R.M., Scholten, J., Bauch, D., Michel, A., 1995. ^{228}Ra as a tracer for shelf water in the arctic ocean. Deep Sea Res. Part II 42, 1533–1553.
- Sander, S.G., Koschinsky, A., 2011. Metal flux from hydrothermal vents increased by organic complexation. Nat. Geosci. 4, 145–150.
- Sarthou, G., Baker, A.R., Kramer, J., Laan, P., Laës, A., Ussher, S., Achterberg, E.P., de Baar, H.J.W., Timmermans, K.R., Blain, S., 2007. Influence of atmospheric inputs on the iron distribution in the subtropical North-East Atlantic Ocean. Mar. Chem. 104, 186–202.
- Sarthou, G., Jeandel, C., 2001. Seasonal variations of iron concentrations in the Ligurian Sea and iron budget in the Western Mediterranean Sea. Mar. Chem. 74, 115–129.
- Sarthou, G., Vincent, D., Christaki, U., Obernosterer, I., Timmermans, K.R., Brussaard, C.P.D., 2008. The fate of biogenic iron during a phytoplankton bloom induced by natural fertilisation: Impact of copepod grazing. Deep Sea Res. Part II Top. Stud. Oceanogr. 55, 734–751.
- Saydam, A.C., Senyuva, H.Z., 2002. Deserts: Can they be the potential suppliers of bioavailable iron? Geophys. Res. Lett. 29, 19-1–19-3.
- Schauer, U., 2016. The expedition PS94 of the Research Vessel POLARSTERN to the central Arctic Ocean in 2015. Berichte zur Polar- und Meeresforsch. - Reports Polar Mar. Res. 703, 170.
- Schlitzer, R., 2016. Ocean Data View.
- Schoemann, V., Becquevert, S., Stefels, J., Rousseau, V., Lancelot, C., 2005. Phaeocystis blooms in the global ocean and their controlling mechanisms: a review. J. Sea Res. 53, 43–66.
- Scholten, J.C., Rutgers van der Loeff, M.M., Michel, A., 1995. Distribution of ^{230}Th and ^{231}Pa in the water column in relation to the ventilation of the deep Arctic basins. Deep. Res. Part II 42, 1519–1531.
- Schroeder, K., Taillardier, V., Vetrano, A., Gasparini, G.P., 2008. The circulation of the western Mediterranean Sea in spring 2005 as inferred from observations and from model outputs. Deep Sea Res. Part I Oceanogr. Res. Pap. 55, 947–965.
- Schuur, E.A.G., Abbott, B.W., Bowden, W.B., Brovkin, V., Camill, P., Canadell, J.G., Chanton, J.P., Chapin, F.S., Christensen, T.R., Ciais, P., Crosby, B.T., Czimczik, C.I., Grosse, G., Harden, J., Hayes, D.J., Hugelius, G., Jastrow, J.D., Jones, J.B., Kleinen, T., Koven, C.D., Krinner, G., Kuhry, P., Lawrence, D.M., McGuire, A.D., Natali, S.M., O'Donnell, J.A., Ping, C.L., Riley, W.J., Rinke, A., Romanovsky, V.E., Sannel, A.B.K., Schädel, C., Schaefer, K., Sky, J., Subin, Z.M., Tarnocai, C., Turetsky, M.R., Waldrop, M.P., Walter Anthony, K.M., Wickland, K.P., Wilson, C.J., Zimov, S.A., 2013. Expert assessment of vulnerability of permafrost carbon to climate change. Clim. Change 119, 359–374.
- Schuur, E.A.G., McGuire, A.D., Grosse, G., Harden, J.W., Hayes, D.J., Hugelius, G., Koven, C.D., Kuhry, P., 2015. Climate change and the permafrost carbon feedback. Nature 520, 171–179.

- Sedwick, P.N., Church, T.M., Bowie, A.R., Marsay, C.M., Ussher, S.J., Achilles, K.M., Lethaby, P.J., Johnson, R.J., Sarin, M.M., McGillicuddy, D.J., 2005. Iron in the Sargasso Sea (Bermuda Atlantic Time-series Study region) during summer: Eolian imprint, spatiotemporal variability, and ecological implications. *Global Biogeochem. Cycles* 19, n/a-n/a.
- Sedwick, P.N., Sholkovitz, E.R., Church, T.M., 2007. Impact of anthropogenic combustion emissions on the fractional solubility of aerosol iron: Evidence from the Sargasso Sea. *Geochemistry, Geophys. Geosystems* 8, n/a-n/a.
- Sedwick, P.N., Sohst, B.M., Ussher, S.J., Bowie, A.R., 2015. A zonal picture of the water column distribution of dissolved iron(II) during the U.S. GEOTRACES North Atlantic transect cruise (GEOTRACES GA03). *Deep. Res. Part II Top. Stud. Oceanogr.* 116, 166–175.
- Serreze, M.C., Stroeve, J., 2015. Arctic sea ice trends, variability and implications for seasonal ice forecasting. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 373, 20140159.
- Serreze, M.C., Stroeve, J., Barrett, A.P., Boisvert, L.N., 2016. Summer atmospheric circulation anomalies over the Arctic Ocean and their influences on September sea ice extent: A cautionary tale. *J. Geophys. Res. Atmos.* 121, 11,463-11,485.
- Shaked, Y., Lis, H., 2012. Disassembling Iron Availability to Phytoplankton. *Front. Microbiol.* 3, 1–26.
- Sheik, A.R., Brussaard, C.P.D., Lavik, G., Lam, P., Musat, N., Krupke, A., Littmann, S., Strous, M., Kuypers, M.M.M., 2014. Responses of the coastal bacterial community to viral infection of the algae *Phaeocystis globosa*. *ISME J.* 8, 212–225.
- Sholkovitz, E.R., 1976. Flocculation of dissolved organic and inorganic matter during the mixing of river water and seawater. *Geochim. Cosmochim. Acta* 40, 831–845.
- Sholkovitz, E.R., 1993. The geochemistry of rare earth elements in the Amazon River estuary. *Geochim. Cosmochim. Acta* 57, 2181–2190.
- Slagter, H.A., Gerringa, L.J.A., Brussaard, C.P.D., 2016. Phytoplankton Virus Production Negatively Affected by Iron Limitation. *Front. Mar. Sci.* 3, 1–11.
- Slagter, H.A., Reader, H.E., Rijkenberg, M.J.A., Rutgers van der Loeff, M., de Baar, H.J.W., Gerringa, L.J.A., 2017. Organic Fe speciation in the Eurasian Basins of the Arctic Ocean and its relation to terrestrial DOM. *Mar. Chem.* 197, 11–25.
- Soria-Dengg, S., Reissbrodt, R., Horstmann, U., 2001. Siderophores in marine coastal waters and their relevance for iron uptake by phytoplankton: experiments with the diatom *Phaeodactylum tricornutum*. *Mar Ecol Prog Ser* 220, 73–82.
- Spokes, L.J., Jickells, T.D., 1995. Factors controlling the solubility of aerosol trace metals in the atmosphere and on mixing into seawater. *Aquat. Geochemistry* 1, 355–374.
- Stedmon, C.A., Amon, R.M.W., Rinehart, A.J., Walker, S.A., 2011. The supply and characteristics of colored dissolved organic matter (CDOM) in the Arctic Ocean: Pan Arctic trends and differences. *Mar. Chem.* 124, 108–118.
- Stedmon, C.A., Markager, S., Kaas, H., 2000. Optical Properties and Signatures of Chromophoric Dissolved Organic Matter (CDOM) in Danish Coastal Waters. *Estuar. Coast. Shelf Sci.* 51, 267–278.
- Stoderegger, K., Herndl, G.J., 1998. Stoderegger y Herndl, 1998 - Production and release of bacterial capsular material and its subsequent utilization by marine bacterioplankton. *Limnol. Oceanogr.* 43, 877–884.
- Strass, V.H., Nöthig, E.M., 1996. Seasonal shifts in ice edge phytoplankton blooms in the Barents Sea related to the water column stability. *Polar Biol.* 16, 409–422.
- Strickland, J., Parsons, T., 1972. A practical handbook of seawater analysis.
- Stroeve, J.C., Serreze, M.C., Holland, M.M., Kay, J.E., Malanik, J., Barrett, A.P., 2012. The Arctic's rapidly shrinking sea ice cover: A research synthesis. *Clim. Change*

- 110, 1005–1027.
- Strzepek, R.F., Maldonado, M.T., Hunter, K.A., Frew, R.D., Boyd, P.W., 2011. Adaptive strategies by Southern Ocean phytoplankton to lessen iron limitation: Uptake of organically complexed iron and reduced cellular iron requirements. Limnol. Oceanogr. 56, 1983–2002.
- Sukekava, C., Downes, J., Slagter, H.A., Gerringa, L.J.A., Laglera, L.M., 2018. Determination of the contribution of humic substances to iron complexation in seawater by catalytic cathodic stripping voltammetry. Talanta.
- Sunda, W.G., 1997. Control of dissolved iron concentrations in the world ocean, A comment. Mar. Chem. 57, 169–172.
- Sunda, W.G., Huntsman, S. a, 1997. Interrelated influence of iron, light and cell size on marine phytoplankton growth. Nature 390, 389–392.
- Suttle, C.A., 1993. Enumeration and isolation of viruses. In: Kemp, P.F., Sherr, B.F., Sherr, E.B., Cole, J.J. (Eds.), *Handbook of Methods in Aquatic Microbial Ecology*. CRC Press, pp. 121–134.
- Suttle, C.A., 2005. Viruses in the sea. Nature 437, 356–361.
- Tachikawa, K., Roy-Barman, M., Michard, A., Thouron, D., Yeghicheyan, D., Jeandel, C., 2004. Neodymium isotopes in the Mediterranean Sea: comparison between seawater and sediment signals. Geochim. Cosmochim. Acta 68, 3095–3106.
- Tagliabue, A., Aumont, O., DeAth, R., Dunne, J.P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J.K., Ridgwell, A., Sherman, E., Stock, C., Vichi, M., Völker, C., Yool, A., 2016. How well do global ocean biogeochemistry models simulate dissolved iron distributions? Global Biogeochem. Cycles 30, 149–174.
- Tagliabue, A., Bopp, L., Dutay, J.-C., Bowie, A.R., Chever, F., Jean-Baptiste, P., Bucciarelli, E., Lannuzel, D., Remenyi, T., Sarthou, G., Aumont, O., Gehlen, M., Jeandel, C., 2010. Hydrothermal contribution to the oceanic dissolved iron inventory. Nat. Geosci. 3, 252–256.
- Tagliabue, A., Bowie, A.R., Boyd, P.W., Buck, K.N., Johnson, K.S., Saito, M.A., 2017. The integral role of iron in ocean biogeochemistry. Nature 543, 51–59.
- Tanhua, T., Jones, E.P., Jeansson, E., Jutterström, S., Smethie, W.M., Wallace, D.W.R., Anderson, L.G., 2009. Ventilation of the arctic ocean: Mean ages and inventories of anthropogenic CO₂ and CFC-11. J. Geophys. Res. Ocean. 114, 1–11.
- Taylor, R.L., Semeniuk, D.M., Payne, C.D., Zhou, J., Tremblay, J.-É., Cullen, J.T., Maldonado, M.T., 2013. Colimitation by light, nitrate, and iron in the Beaufort Sea in late summer. J. Geophys. Res. Ocean. 118, 3260–3277.
- Testor, P., Gascard, J.C., 2003. Large-scale spreading of deep waters in the Western Mediterranean Sea by submesoscale coherent eddies. J. Phys. Oceanogr. 33, 75–87.
- Thuróczy, C.-E., Gerringa, L.J.A., Klunder, M., Laan, P., Le Guittion, M., de Baar, H.J.W., 2011. Distinct trends in the speciation of iron between the shallow shelf seas and the deep basins of the Arctic Ocean. J. Geophys. Res. 116, C10009.
- Thuróczy, C.-E., Gerringa, L.J.A., Klunder, M.B., Middag, R., Laan, P., Timmermans, K.R., de Baar, H.J.W., 2010. Speciation of Fe in the Eastern North Atlantic Ocean. Deep Sea Res. Part I 57, 1444–1453.
- Timmermans, K.R., Davey, M.S., van der Wagt, B., Snoek, J., Geider, R.J., Veldhuis, M.J.W., Gerringa, L.J.A., De Baar, H.J.W., 2001a. Co-limitation by iron and light of *Chaetoceros brevis*, *C. dichaeta* and *C. calcitrans* (Bacillariophyceae). Mar. Ecol. Prog. Ser. 217, 287–297.
- Timmermans, K.R., Gerringa, L.J.A., de Baar, H.J.W., van der Wagt, B., Veldhuis, M.J.W., de Jong, J.T.M., Croot, P.L., Boye, M., 2001b. Growth rates of large and small Southern Ocean diatoms in relation to availability of iron in natural seawater.

- Limnol. Oceanogr. 46, 260–266.
- Timmermans, K.R., Gledhill, M., Nolting, R.F., Veldhuis, M.J.W., de Baar, H.J.W., van den Berg, C.M.G., 1998. Responses of marine phytoplankton in iron enrichment experiments in the northern North Sea and northeast Atlantic Ocean. Mar. Chem. 61, 229–242.
- Timmermans, K.R., van der Wagt, B., de Baar, H.J.W., 2004. Growth rates, half saturation constants, and silicate, nitrate, and phosphate depletion in relation to iron availability of four large open-ocean diatoms from the Southern Ocean. Limnol. Oceanogr. 49, 2141–2151.
- Timmermans, K.R., van der Wagt, B., Veldhuis, M.J.W., Maatman, A., de Baar, H.J.W., 2005. Physiological responses of three species of marine pico-phytoplankton to ammonium, phosphate, iron and light limitation. J. Sea Res. 53, 109–120.
- Town, R.M., Filella, M., 2000. Dispelling the myths: Is the existence of L1 and L2 ligands necessary to explain metal ion speciation in natural waters? Limnol. Oceanogr. 45, 1341–1357.
- Tremblay, J.-É., Gagnon, J., 2009. The effects of irradiance and nutrient supply on the productivity of Arctic waters: a perspective on climate change. In: Nihoul, J.C.J., Kostianoy, A.G. (Eds.), *Influence of Climate Change on the Changing Arctic and Sub-Arctic Conditions*. Springer, Dordrecht, pp. 73–93.
- Tremblay, J.-É., Anderson, L.G., Matrai, P., Coupel, P., Bélanger, S., Michel, C., Reigstad, M., 2015. Global and regional drivers of nutrient supply, primary production and CO₂ drawdown in the changing Arctic Ocean. Prog. Oceanogr. 139, 171–196.
- Tremblay, J.-É., Simpson, K., Martin, J., Miller, L., Gratton, Y., Barber, D., Price, N.M., 2008. Vertical stability and the annual dynamics of nutrients and chlorophyll fluorescence in the coastal, southeast Beaufort Sea. J. Geophys. Res. Ocean. 113, 1–14.
- Trezzi, G., Garcia-Orellana, J., Rodellas, V., Santos-Echeandia, J., Tovar-Sánchez, A., Garcia-Solsona, E., Masqué, P., 2016. Submarine groundwater discharge: A significant source of dissolved trace metals to the North Western Mediterranean Sea. Mar. Chem. 186, 90–100.
- Twining, B.S., Baines, S.B., 2013. The trace metal composition of marine phytoplankton. Ann. Rev. Mar. Sci. 5.
- Twining, B.S., Rauschenberg, S., Morton, P.L., Vogt, S., 2015. Metal contents of phytoplankton and labile particulate material in the North Atlantic Ocean. Prog. Oceanogr. 137, 261–283.
- Ulfsbo, A., Cassar, N., Korhonen, M., Heuven, S. Van, Hoppema, M., Kattner, G., Anderson, L.G., 2014. Late summer net community production in the central Arctic Ocean using multiple approaches. Global Biogeochem. Cycles 28, 1129–1148.
- Ünlülata, Ü., Oğuz, T., Latif, M.A., Özsoy, E., 1990. On the physical oceanography of the Turkish Straits. In: *The Physical Oceanography of Sea Straits*. Springer, pp. 25–60.
- van de Poll, W.H., Janknegt, P.J., van Leeuwe, M.A., Visser, R.J.W., Buma, A.G.J., 2009. Excessive irradiance and antioxidant responses of an Antarctic marine diatom exposed to iron limitation and to dynamic irradiance. J. Photochem. Photobiol. B Biol. 94, 32–37.
- van de Poll, W.H.H., Boute, P.G.G., Rozema, P.D.D., Buma, A.G.J.G.J., Kulk, G., Rijkenberg, M.J.A.J.A., 2015. Sea surface temperature control of taxon specific phytoplankton production along an oligotrophic gradient in the Mediterranean Sea. Mar. Chem. 177, 536–544.
- van den Berg, C.M.G., 1995. Evidence for organic complexation of iron in seawater. Mar. Chem. 50, 139–157.
- van den Berg, C.M.G., 2006. Chemical speciation of iron in seawater by cathodic stripping

- voltammetry with dihydroxynaphthalene. *Anal. Chem.* 78, 156–163.
- van den Berg, C.M.G., Kramer, J.R., 1979. Determination of complexing capacities of ligands in natural waters and conditional stability constants of the copper complexes by means of manganese dioxide. *Anal. Chim. Acta* 106, 113–120.
- van den Berg, C.M.G., Nimmo, M., Daly, P., Turner, D.R., 1990. Effects of the detection window on the determination of organic copper speciation in estuarine waters. *Anal. Chim. Acta* 232, 149–159.
- van Haren, H., Millot, C., 2009. Slantwise convection: A candidate for homogenization of deep newly formed dense waters. *Geophys. Res. Lett.* 36.
- van Haren, H., Millot, C., Taupier-Letage, I., 2006. Fast deep sinking in Mediterranean eddies. *Geophys. Res. Lett.* 33, L04606.
- van Haren, H., The ANTARES Collaboration, 2014. High-frequency internal wave motions at the ANTARES site in the deep Western Mediterranean. *Ocean Dyn.* 64, 507–517.
- van Leeuwe, M.A., Stefels, J., 1998. Effects of iron and light stress on the biochemical composition of Antarctic *Phaeocystis* sp. (Prymnesiophyceae). II. Pigment composition. *J. Phycol.* 34, 496–503.
- van Leeuwe, M.A., Stefels, J., 2007. Photosynthetic responses in *Phaeocystis antarctica* towards varying light and iron conditions. *Phaeocystis, Major Link Biogeochem. Cycl. Clim. Elem.* 61–70.
- Vancoppenolle, M., Bopp, L., Madec, G., Dunne, J., Ilyina, T., Halloran, P.R., Steiner, N., 2013. Future arctic ocean primary productivity from CMIP5 simulations: Uncertain outcome, but consistent mechanisms. *Global Biogeochem. Cycles* 27, 605–619.
- Vaulot, D., 1989. CYTOPC: Processing software for flow cytometric data. *Signal and Noise* 2:8.
- Vaulot, D., Birrien, J.-L., Marie, D., Casotti, R., Veldhuis, M.J.W., Kraay, G.W., Chrétiennot-Dinet, M.-J., 1994. Morphology, ploidy, pigment composition, and genome size of cultured strains of *Phaeocystis* (Prymnesiophyceae). *J. Phycol.* 30, 1022–1035.
- Vaulot, D., Eikrem, W., Viprey, M., Moreau, H., 2008. The diversity of small eukaryotic phytoplankton ($\leq 3 \mu\text{m}$) in marine ecosystems. *FEMS Microbiol. Rev.* 32, 795–820.
- Velasquez, I., Nunn, B.L., Ibisani, E., Goodlett, D.R., Hunter, K.A., Sander, S.G., 2011. Detection of hydroxamate siderophores in coastal and Sub-Antarctic waters off the South Eastern Coast of New Zealand. *Mar. Chem.* 126, 97–107.
- Velasquez, I.B., Ibisani, E., Maas, E.W., Boyd, P.W., Nodder, S., Sander, S.G., 2016. Ferrioxamine Siderophores Detected amongst Iron Binding Ligands Produced during the Remineralization of Marine Particles. *Front. Mar. Sci.* 3, 1–14.
- Veldhuis, M.J.W., Timmermans, K.R., Croot, P., van der Wagt, B., 2005. Picophytoplankton; a comparative study of their biochemical composition and photosynthetic properties. *J. Sea Res.* 53, 7–24.
- Visser, F., Gerrings, L.J.A., Gaast, S.J. Van Der, Baar, H.J.W. De, Timmermans, K.R., Van Der Gaast, S.J., De Baar, H.J.W., Timmermans, K.R., 2003. The role of the reactivity and content of iron of aerosol dust on growth rates of two Antarctic diatom species. *J. Phycol.* 39, 1085–1094.
- von der Heyden, B.P., Roychoudhury, A.N., Mtshali, T.N., Tyliszczak, T., Myneni, S.C.B., 2012. Chemically and Geographically Distinct Solid-Phase Iron Pools in the Southern Ocean. *Science* (80-.). 338, 1199–1201.
- Vonk, J.E., Mann, P.J., Davydov, S., Davydova, A., Spencer, R.G.M., Schade, J., Sobczak, W. V., Zimov, N., Zimov, S., Bulygina, E., Eglington, T.I., Holmes, R.M., 2013. High biolability of ancient permafrost carbon upon thaw. *Geophys. Res. Lett.* 40, 2689–2693.
- Vonk, J.E., Sánchez-García, L., van Dongen, B.E., Alling, V., Kosmach, D., Charkin, A.,

- Semiletov, I.P., Dudarev, O. V., Shakhova, N., Roos, P., Eglinton, T.I., Andersson, A., Gustafsson, Ö., 2012. Activation of old carbon by erosion of coastal and subsea permafrost in Arctic Siberia. *Nature* 489, 137–140.
- Voorhis, A.D., Webb, D.C., 1970. Large vertical currents observed in a winter sinking region of the northwestern Mediterranean. Woods Hole Oceanographic Institution.
- Wagener, T., Guieu, C., Leblond, N., 2010. Effects of dust deposition on iron cycle in the surface Mediterranean Sea: results from a mesocosm seeding experiment. *Biogeosciences* 7, 3769–3781.
- Wagener, T., Pulido-Villena, E., Guieu, C., 2008. Dust iron dissolution in seawater: Results from a one-year time-series in the Mediterranean Sea. *Geophys. Res. Lett.* 35.
- Walker, S.A., Amon, R.M.W., Stedmon, C., Duan, S., Louchouarn, P., 2009. The use of PARAFAC modeling to trace terrestrial dissolved organic matter and fingerprint water masses in coastal Canadian Arctic surface waters. *J. Geophys. Res.* 114, 1–12.
- Weitz, J.S., Wilhelm, S.W., 2012. Ocean viruses and their effects on microbial communities and biogeochemical cycles. *F1000 Biol. Rep.* 8, 2–9.
- Wilhelm, C., Büchel, C., Fisahn, J., Goss, R., Jakob, T., LaRoche, J., Lavaud, J., Lohr, M., Riebesell, U., Stehfest, K., Valentin, K., Kroth, P.G., 2006. The Regulation of Carbon and Nutrient Assimilation in Diatoms is Significantly Different from Green Algae. *Protist* 157, 91–124.
- Wilhelm, S.W., King, A.L., Twining, B.S., LeCleir, G.R., DeBruyn, J.M., Strzepek, R.F., Breene, C.L., Pickmere, S., Ellwood, M.J., Boyd, P.W., Hutchins, D.A., 2013. Elemental quotas and physiology of a southwestern Pacific Ocean plankton community as a function of iron availability. *Aquat. Microb. Ecol.* 68, 185–194.
- Wilhelm, S.W., MacAuley, K., Trick, C.G., Nielsen, I., 1998. Evidence for the importance of catechol-type siderophores in the iron-limited growth of a cyanobacterium. *Limnol. Oceanogr.* 43, 992–997.
- Wilhelm, S.W., Suttle, C.A., 1999. Viruses and Nutrient Cycles in the Sea. *Bioscience* 49, 781.
- Wilhelm, S.W., Suttle, C.A., 2000. Viruses as regulators of nutrient cycles in aquatic environments. In: *Microbial Biosystems: New Frontiers. Proceedings of the VIII International Symposium on Microbial Ecology*. Atlantic Canada Society for Microbial Ecology, Halifax, Nova Scotia. pp. 551–556.
- Wilhelm, S.W., Trick, C.G., 1994. Iron-limited growth of cyanobacteria: Multiple siderophore production is a common response. *Limnol. Oceanogr.* 39, 1979–1984.
- Wozniak, A.S., Shelley, R.U., McElhenie, S.D., Landing, W.M., Hatcher, P.G., 2015. Aerosol water soluble organic matter characteristics over the North Atlantic Ocean: Implications for iron-binding ligands and iron solubility. *Mar. Chem.* 173, 162–172.
- Wu, J., Boyle, E., Sunda, W., Wen, L.S., 2001. Soluble and colloidal iron in the oligotrophic North Atlantic and North Pacific. *Science* 293, 847–849.
- Zapata, M., Rodríguez, F., Garrido, J.L., 2000. Separation of chlorophylls and carotenoids from marine phytoplankton: A new HPLC method using a reversed phase C8 column and pyridine-containing mobile phases. *Mar. Ecol. Prog. Ser.* 195, 29–45.
- Zhang, C., 2014. Essential functions of iron-requiring proteins in DNA replication, repair and cell cycle control. *Protein Cell* 5, 750–60.
- Zitter, T.A.C., 2004. Mud volcanism and fluid emissions in eastern Mediterranean neotectonic zones. *Applied geology*.