



University Medical Center Groningen

University of Groningen

Towards novel biomarkers and rational nutritional interventions in Inflammatory Bowel Disease

von Martels, Julius Zweder Hubertus

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

von Martels, J. Z. H. (2019). Towards novel biomarkers and rational nutritional interventions in Inflammatory Bowel Disease. [Groningen]: Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

4

INCREASED FECAL CALPROTECTIN LEVELS IN CROHN'S DISEASE CORRELATE WITH ELEVATED SERUM TH1- AND TH17-ASSOCIATED CYTOKINES

PLoS One 2018

Arno R. Bourgonje, Julius Z. H. von Martels, Paul de Vos, Klaas Nico Faber*,
Gerard Dijkstra*

** Shared last authors*

ABSTRACT

BACKGROUND

Patient-reported symptoms and endoscopic disease activity do not correlate well in Crohn's disease (CD). This warrants the need for reliable biomarkers to early detect active intestinal inflammation. Currently, the fecal calprotectin level is the most commonly used biomarker for inflammatory activity in CD. However, the diagnostic accuracy of the fecal calprotectin level is not fully efficacious and diagnosis may be further improved by the identification of other biomarkers for active CD. Here, we studied the association of pro-inflammatory serum cytokines with fecal calprotectin levels in CD patients.

METHODS

39 CD patients were included and subdivided into 'normal' (defined as < 200 mg/kg feces) and 'increased' (defined as > 200 mg/kg feces) fecal calprotectin level groups. Serum cytokine levels were quantified by an electrochemiluminescence multiplex assay (V-PLEX Human Biomarker 40-Plex Kit of Meso Scale Discovery[®]). Correlations between individual cytokines and the fecal calprotectin level were assessed using Spearman's correlation coefficient (ρ).

RESULTS

A highly significant positive correlation was observed between the pro-inflammatory serum cytokines IFN- γ and CRP and fecal calprotectin levels ($P < 0.01$). Moreover, fecal calprotectin levels showed a significant positive correlation with IL-6, TNF- β , SAA and IL-17A ($P < 0.05$).

CONCLUSION

We show that a positive correlation exists between multiple serum Th1- and Th17-associated cytokines and the fecal calprotectin level. These cytokines and CRP may serve as additional biomarkers for determining disease activity and evaluating treatment response in CD. Ultimately, this may result in more efficient treatment of active disease in CD patients and prevention of complications.

INTRODUCTION

Crohn's disease (CD) is a chronic inflammatory disease that mainly affects the gastro-intestinal tract and is characterized by an inappropriate and ongoing immune response.¹ Most patients suffer from a relapse-remitting disease course that is difficult to predict.^{2,3} Predicting the disease course is hampered by the poor availability of adequate disease biomarkers or symptoms that forecast a flare of inflammation. Longstanding sub-clinical disease activity increases the risk of various severe complications, such as stricturing or penetrating disease (i.e. fistula or abscess formation).⁴ Appropriate and prompt treatment of the inflammatory activity lowers the risk of these severe complications, and thus prevents future surgical interventions.

Currently, the only reliable approach for diagnosis of CD is an invasive ileo-colonoscopy. However, this procedure has several disadvantages, such as the risk of perforation, bleeding, relatively high costs and, most importantly, a high patient burden. Furthermore, a poor association exists between patient-reported symptoms and the observed endoscopic inflammatory activity. For instance, clinical risk scores, such as the Harvey Bradshaw Index (HBI) or the Crohn's Disease Activity Index (CDAI), cannot accurately predict active intestinal inflammation.^{5,6}

In clinical practice, fecal calprotectin levels are commonly used as non-invasive biomarker that significantly correlates with inflammatory disease activity and response to therapy.^{5,7-10} Calprotectin is a 36 kDa calcium- and zinc-binding protein dimer (consisting of S100A8 and S100A9) complex present in the cytosol of neutrophilic granulocytes.¹¹ Fecal calprotectin levels are representative of neutrophil migration into the intestinal mucosa that occurs in the process of intestinal inflammation. Despite its proven association with disease activity, the diagnostic accuracy may be further improved by inclusion of additional inflammatory biomarkers for active inflammation in CD.¹²⁻¹⁴ Many cytokines have been shown to be involved in disease pathogenesis and might give a more accurate representation of the inflammatory activity in CD relapses in combination with fecal calprotectin levels. Ultimately, this may aid in an improved detection of active disease and a more versatile and effective treatment.

A Th1-driven immune response with increased Th1-associated cytokines such as IFN- γ and TNF- α plays a pivotal role in the pathogenesis of CD.¹⁵ The importance of Th1-responses is also reflected by the clinical use of TNF- α antagonists, such as infliximab, adalimumab and certolizumab, which are effective treatment modalities in CD.¹⁶ Previously, quantification of cytokines for diagnosis of disease has been cumbersome due to low

serum concentrations of the relevant cytokines, but new techniques enable us to quantify most cytokines in a highly sensitive, reproducible and validated manner.¹⁷⁻¹⁹

The aim of this study is to identify potential serum cytokines that might serve as additional biomarkers for inflammatory disease activity in CD. A positive correlation between specific serum cytokines and fecal calprotectin levels might enhance the diagnostic potential for early recognition of disease exacerbations.

METHODS

STUDY POPULATION

Patients aged 19-67 years with an established diagnosis of Crohn's disease (CD) were included from March 2016 to April 2017 at the University Medical Center Groningen (UMCG), the Netherlands. In total, 39 CD patients were included and divided into two groups according to inflammatory disease activity, as determined by fecal calprotectin levels. Patients having a calprotectin level below 200 mg/kg were defined as the 'normal' calprotectin group (indicative of remissive disease) and patients with calprotectin levels above 200 mg/kg were defined as the 'increased' calprotectin group (indicative of inflammatory active CD). Clinically relevant data were obtained from medical records: age, gender, BMI, smoking history, maintenance medication, Montreal score, ileocecal resection and laboratory parameters. Fecal calprotectin levels were determined in the laboratory of the UMCG as a routine measurement.

CYTOKINE MEASUREMENT

Serum samples from all patients were collected and stored in 1 mL aliquots in the freezer (-20°C). After thawing, serum samples were centrifuged for 3 minutes at 2000 g to remove particulates prior to sample preparation and analysis. Measurement of serum cytokine levels was performed by the electrochemiluminescence (ECL) multiplex assay (Meso Scale Discovery (MSD[®]), Meso Scale Diagnostics, Rockville, MD). The MSD V-plex Pro-inflammatory panel 1 (IFN- γ , IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-13 and TNF- α), Cytokine panel 1 (GM-CSF, IL-5, IL-7, IL-12/23p40, IL-15, IL-16, IL-17A and TNF- β), Chemokine panel 1 (Eotaxin, MIP-1 β , Eotaxin-3, TARC, IP-10, MIP-1 α , MCP-1 and MDC), Angiogenesis panel 1 (VEGF, VEGF-C, VEGF-D, Tie-2, Flt-1, PlGF and bFGF) and Vascular injury panel 1 (SAA, CRP, VCAM-1 and ICAM-1) were used to detect a total of 37 cytokines. Calibration curves were created in order to calculate serum cytokine concentrations. Calibrator signals were fitted to a 4-parameter logistic model with weighting, providing the assay with a wide dynamic range of detection. Cytokine concentrations were calculated by back-fitting ECL signals to the calibration curves. Determination of final cytokine concentrations was performed using the MSD Discovery Workbench[®] analysis software. Concentrations of all cytokines were above the lower limit of detection (LLOD). For the V-plex Pro-inflammatory panel 1, Cytokine panel 1 and Angiogenesis panel 1 assays, standard volumes of 50 μ L of each serum sample were added and diluted 2 times. For the Chemokine panel 1, samples were diluted 4 times. For the Vascular injury panel 1, a standard volume of 25 μ L of each serum sample was added and 1000-fold diluted.

STATISTICS

Study population characteristics were presented as means \pm standard errors (SEM) or proportions (%), n). Serum cytokine distributions were presented as median \pm interquartile ranges (IQR) and shown in boxplots (10th-90th percentiles) grouped by inflammatory disease activity, as determined by the fecal calprotectin level. Correlations between fecal calprotectin levels and serum cytokine levels were established using the nonparametric Spearman's correlation coefficient (ρ). Data were analyzed using SPSS Statistics 23.0 software package for Windows. P -values ≤ 0.05 were considered as statistically significant.

RESULTS

Characteristics of the study population are presented in **Table 1**. Patients with a 'normal' fecal calprotectin level ($n = 22$) had a mean age of 40.5 ± 2.4 years and consisted of 5 males (22.7%) and 17 females (77.3%). Patients with an increased level of fecal calprotectin ($n = 17$) had a mean age of 39.7 ± 3.4 years and consisted of 5 males (29.4%) and 12 females (70.6%). Patients with increased fecal calprotectin levels had significantly higher C-reactive protein (CRP) levels ($P < 0.01$) and higher erythrocyte sedimentation rates (ESR) ($P < 0.05$) as compared to patients with fecal calprotectin levels in the normal range. No significant differences between groups were observed for disease location, medication use, smoking habits, surgery history and standard laboratory parameters.

TABLE 1: Study population characteristics ($n = 39$) of Crohn's disease patients with 'normal' (< 200 mg/kg) and increased (> 200 mg/kg) fecal calprotectin levels.

Characteristics	Calprotectin < 200 mg/kg (n = 22)	Calprotectin > 200 mg/kg (n = 17)	P value
Age (years)	40.5 ± 2.4	39.7 ± 3.4	0.788
Male gender	5 (22,7)	5 (29,4)	0.721
BMI (kg/m ²)	24.9 ± 1.4	25.8 ± 1.2	0.350
Active smoking	2 (9,1)	1 (5,9)	1.000
Maintenance medication			0.501
- None	7 (31,8)	3 (17,6)	
- Thiopurines	4 (18,2)	4 (23,5)	
- Mesalamine	4 (18,2)	2 (11,8)	
- TNF-antagonists	3 (13,6)	6 (35,3)	
- Combination therapy	4 (18,2)	2 (11,8)	
Montreal, Localization			0.194
- L1 (ileal)	11 (52,4)	4 (23,5)	
- L2 (colonic)	3 (14,3)	4 (23,5)	
- L3 (ileocolonic)	7 (33,3)	9 (52,9)	
Montreal, Behavior			0.322
- B1 (non stricturing, non penetrating)	14 (66,7)	8 (47,1)	
- B2 (stricturing)	5 (23,8)	8 (47,1)	
- B3 (penetrating)	2 (9,5)	1 (5,9)	
Ileocecal resection	8 (36,4)	5 (29,4)	0.740
Hemoglobin (mmol/l)	8.4 ± 0.2	8.4 ± 0.2	0.787
CRP (mg/l)	2.1 ± 0.6	8.2 ± 2.4	*0.002
ESR (mm/h)	12.5 ± 2.2	23.6 ± 4.2	*0.017
WBC ($\times 10^9/l$)	6.9 ± 0.5	6.9 ± 0.4	0.723
Thrombocytes ($\times 10^9/l$)	264 ± 13	308 ± 17	0.229
ASAT (U/l)	22 ± 1	23 ± 3	0.977
ALAT (U/l)	18 ± 2	23 ± 5	0.580
Creatinine ($\mu\text{mol/l}$)	68 ± 2	70 ± 3	0.681

BMI, body mass index; HBI, Harvey Bradshaw index; CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; WBC, white blood cell count; ASAT, aspartate aminotransferase; ALAT, alanine aminotransferase. Data are presented as numbers (proportions, n (%)) or mean \pm SE. Differences between groups were tested with Mann-Whitney U-test for continuous variables and Fisher's exact test for nominal variables. *P value < 0.05 was considered statistically significant.

In **Table 2**, serum cytokine levels (pg/mL) are shown as median values with interquartile ranges (IQR) for both groups. Cytokines are grouped in different experimental panels (Pro-inflammatory panel, Cytokine panel, Chemokine panel, Angiogenesis panel and Vascular injury panel).

TABLE 2: Median (IQR) of serum cytokine levels (pg/mL) in Crohn's disease patients with 'normal' (< 200 mg/kg) and increased (> 200 mg/kg) fecal calprotectin levels.

Cytokines	Calprotectin < 200 mg/kg	Calprotectin > 200 mg/kg	P-value
Pro-inflammatory panel			
IFN- γ	7.80 (4.09-19.72)	23.00 (10.13-38.72)	< 0.05*
IL-1 β	0.05 (0.04-0.08)	0.05 (0.03-0.13)	0.945
IL-2	0.18 (0.12-0.23)	0.17 (0.10-0.25)	0.932
IL-4	0.04 (0.02-0.07)	0.04 (0.03-0.06)	0.777
IL-6	0.78 (0.39-1.15)	1.29 (0.93-1.88)	< 0.01**
IL-8	11.91 (7.20-15.43)	12.63 (6.35-68.62)	0.955
IL-10	0.32 (0.19-0.44)	0.37 (0.24-0.74)	0.321
IL-12p70	0.09 (0.03-0.26)	0.07 (0.04-0.18)	0.624
IL-13	0.56 (0.42-0.91)	0.77 (0.55-1.79)	0.153
TNF- α	3.13 (2.50-3.67)	3.49 (2.82-4.79)	0.095
Cytokine panel			
GM-CSF	0.16 (0.07-0.29)	0.15 (0.07-0.24)	0.784
IL-5	0.26 (0.16-0.57)	0.26 (0.19-0.54)	0.883
IL-7	19.95 (13.57-28.08)	18.93 (10.18-27.09)	0.552
IL-12/23p40	118.18 (74.94-204.08)	158.38 (125.23-222.36)	0.157
IL-15	3.04 (2.31-5.66)	2.87 (2.44-5.74)	0.777
IL-16	284.06 (203.49-316.63)	285.97 (195.06-327.77)	0.755
IL-17A	5.25 (3.04-9.27)	8.22 (5.11-11.85)	0.058
TNF- β	0.49 (0.34-0.55)	0.55 (0.42-0.66)	0.051
Chemokine panel			
Eotaxin	266.32 (179.61-351.18)	240.14 (190.62-356.28)	0.955
MIP-1 β	90.12 (73.41-172.00)	80.40 (47.64-129.26)	0.322
Eotaxin-3	20.29 (15.72-30.32)	15.57 (11.79-22.39)	0.101
TARC	242.32 (138.99-512.79)	267.11 (156.86-359.57)	0.799
IP-10	2.99x10 ³ (2.40x10 ³ -4.62x10 ³)	3.60x10 ³ (2.25x10 ³ -4.66x10 ³)	0.590
MIP-1 α	16.27 (13.31-20.75)	16.75 (12.69-24.30)	0.966
MCP-1	217.13 (166.20-357.20)	261.95 (188.06-357.75)	0.630
MDC	1.33x10 ³ (1.02x10 ³ -1.69x10 ³)	1.28x10 ³ (1.07x10 ³ -1.66x10 ³)	0.977
Angiogenesis panel			
VEGF	469.52 (236.89-726.63)	382.16 (223.83-896.71)	0.755
VEGF-C	569.55 (442.96-737.47)	462.11 (375.45-628.34)	0.141
VEGF-D	1.07x10 ³ (770.27-1.47x10 ³)	901.29 (799.81-1.41x10 ³)	0.777
Tie-2	804.11 (652.49-969.74)	708.28 (626.90-845.61)	0.213
Flt-1	147.69 (126.14-167.65)	144.15 (129.16-171.30)	0.932
PIGF	6.28 (5.45-7.40)	5.80 (5.10-7.85)	0.876
bFGF	5.08 (2.05-21.97)	9.12 (2.29-27.11)	0.745
Vascular injury panel			
SAA	4.78x10 ⁶ (2.84x10 ⁶ - 8.87x10 ⁶)	9.02x10 ⁶ (4.64x10 ⁶ - 2.87x10 ⁷)	< 0.05*

TABLE 2 continued.

Cytokines	Calprotectin < 200 mg/kg	Calprotectin > 200 mg/kg	P-value
Vascular injury panel			
CRP	1.25x10 ⁶ (6.03x10 ⁵ - 4.67x10 ⁶)	6.79x10 ⁶ (3.61x10 ⁶ - 2.64x10 ⁷)	< 0.01**
VCAM-1	8.28x10 ⁵ (7.83x10 ⁵ -9.64x10 ⁵)	8.91x10 ⁵ (7.67x10 ⁵ -1.02x10 ⁶)	0.671
ICAM-1	4.97x10 ⁵ (4.27x10 ⁵ -6.00x10 ⁵)	5.63x10 ⁵ (4.99x10 ⁵ -7.06x10 ⁵)	0.062

In the Pro-inflammatory panel, concentrations of IFN- γ and IL-6 were significantly elevated in CD patients with 'increased' fecal calprotectin levels as compared to patients with 'normal' fecal calprotectin levels (IFN- γ : 23.00 pg/mL (IQR: 10.13-38.72) vs. 7.80 pg/mL (IQR: 4.09-19.72) ($P < 0.05$); IL-6: 1.29 pg/mL (0.93-1.88) vs. 0.78 pg/mL (0.39-1.15) ($P < 0.01$). In addition, in the Vascular injury panel, we found significantly higher concentrations of SAA and CRP in patients with increased fecal calprotectin levels as compared to the group with normal fecal calprotectin levels (SAA: 9.02x10⁶ pg/mL (IQR: 4.64x10⁶-2.87x10⁷) vs. 4.78x10⁶ pg/mL (2.84x10⁶-8.87x10⁶) ($P < 0.05$); CRP: 6.79x10⁶ pg/mL (IQR: 3.61x10⁶-2.64x10⁷) vs. 1.25x10⁶ pg/mL (IQR: 6.03x10⁵-4.67x10⁶) ($P < 0.01$). In the Cytokine panel, Chemokine panel and Angiogenesis panel, no significant differences in serum cytokine concentrations were observed between both groups. Interestingly, we also observed elevated concentrations of IL-17A and TNF- β in patients with increased fecal calprotectin levels, but these were borderline non-significant ($P = 0.058$ and $P = 0.051$, respectively).

Distributions of the significantly correlating serum cytokines between groups are shown in **Figure 1** (IFN- γ and CRP, $P < 0.01$) and **Figure 2** (IL-6, IL-17A, TNF- β and SAA, $P < 0.05$). A highly significant positive correlation was observed between the pro-inflammatory cytokines IFN- γ ($\rho = 0.523$, $P < 0.01$, **Figure 1A**) and CRP ($\rho = 0.511$, $P < 0.01$, **Figure 1B**) and fecal calprotectin levels. Significant, but less strong correlations were demonstrated between IL-6 ($\rho = 0.403$, $P < 0.05$, **Figure 2A**), IL-17A ($\rho = 0.352$, $P < 0.05$, **Figure 2B**), TNF- β ($\rho = 0.396$, $P < 0.05$, **Figure 2C**) and SAA ($\rho = 0.323$, $P < 0.05$, **Figure 2D**), and fecal calprotectin levels. No significant correlations were observed between the fecal calprotectin level and the other 31 investigated cytokines, chemokines and markers for angiogenesis and vascular injury. **Table 3** provides a complete overview of all (significant and non-significant) correlations between measured serum cytokine concentrations and fecal calprotectin levels.




FIGURE 1: Distribution of serum cytokine levels in patients with normal (blue, <200mg/kg) and increased (red, >200mg mg/kg) fecal calprotectin levels, shown in boxplots. **A** Serum CRP levels (pg/mL). **B** Serum IFN- γ levels (pg/mL). * $P < 0.01$.




FIGURE 2: Distribution of serum cytokine levels in patients with normal (blue, <200mg/kg) and increased (red, >200mg mg/kg) fecal calprotectin levels, shown in boxplots. **A** Serum IL-6 levels (pg/mL). **B** Serum IL-17A levels (pg/mL). **C** Serum TNF- β levels (pg/mL). **D** Serum SAA levels (pg/mL). * $P < 0.05$. ** $P < 0.01$.

TABLE 3: Correlations of fecal calprotectin levels (mg/kg) with serum cytokine levels (pg/mL) in patients with Crohn's disease.

Cytokines	Spearman's ρ	P-value
Pro-inflammatory panel		
IFN- γ	0.523	< 0.01**
IL-1 β	0.037	0.835
IL-2	-0.063	0.702
IL-4	0.111	0.526
IL-6	0.403	< 0.05*
IL-8	0.033	0.844
IL-10	0.135	0.411
IL-12p70	0.071	0.676
IL-13	0.101	0.647
TNF- α	0.255	0.117
Cytokine panel		
GM-CSF	0.004	0.981
IL-5	0.148	0.374
IL-7	-0.005	0.974
IL-12/23p40	0.214	0.191
IL-15	0.088	0.596
IL-16	-0.018	0.914
IL-17A	0.352	< 0.05*
TNF- β	0.396	< 0.05*
Chemokine panel		
Eotaxin	-0.003	0.986
MIP-1 β	-0.126	0.446
Eotaxin-3	-0.213	0.199
TARC	-0.112	0.499
IP-10	0.207	0.207
MIP-1 α	0.094	0.569
MCP-1	-0.025	0.878
MDC	0.088	0.593
Angiogenesis panel		
VEGF	0.019	0.907
VEGF-C	-0.256	0.116
VEGF-D	0.004	0.980
Tie-2	-0.205	0.211
Flt-1	0.087	0.599
PIGF	0.023	0.888
bFGF	-0.007	0.966
Vascular injury panel		
SAA	0.323	< 0.05*
CRP	0.511	< 0.01**
VCAM-1	0.148	0.367
ICAM-1	0.304	0.060

DISCUSSION

In this study, we show that increased fecal calprotectin levels – a broadly applied marker for the assessment of disease activity in CD patients – highly significantly correlates with elevated serum IFN- γ and CRP levels. Moreover, significant correlations were observed between serum levels of IL-6, IL-17A, TNF- β and SAA and fecal calprotectin levels. These findings indicate that identification of cytokine profiles might serve as an additional approach to determine inflammatory disease activity in CD patients.

Our data not only provide correlations between fecal calprotectin levels and inflammatory markers in the systemic circulation, but also confirm that Th1 responses are important in CD patients with active disease since IFN- γ levels were significantly elevated, as well as serum TNF- α levels, although the latter did not reach statistical significance. Furthermore, an interesting additional finding is the borderline non-significant enhancement of serum IL-17A levels ($P = 0.058$) in CD patients with increased fecal calprotectin levels. IL-17A has been shown to be a driving effector response for colitis in mice. However, transfer of T-cells from IL-17-deficient mice induced severe colitis in mice recipients and enhanced IFN- γ -producing T-cells.²⁰ Likewise, in humans, it is less clear whether IL-17 has a protective role or induces an effector response.^{21,22} In our study, we found elevated IL-17A levels specifically in CD patients with increased fecal calprotectin levels, suggesting that it is a marker for inflammatory disease activity instead of a protective marker. Interestingly, we also found that the Th17-stimulating cytokines IL-6 and TNF- β were enhanced in the group with increased fecal calprotectin levels, further supporting that an activated Th17 response is involved in these patients in addition to the Th1 response.

Our findings on the association between C-reactive protein (CRP) and fecal calprotectin levels corroborate the findings of other studies.^{23,24} CRP is an acute-phase protein, mainly produced by hepatocytes in response to systemic inflammation. Production occurs after stimulation by IL-6, IL-1 β and TNF- α . In clinical practice, CRP is used as a general biomarker for inflammation and is therefore commonly applied to monitor the disease course of CD. CRP rapidly increases after an acute-phase stimulus and has a short half-life, making CRP an useful marker for acute inflammatory events in CD. (25) In CD, active disease is significantly associated with both elevated CRP and fecal calprotectin levels.^{14,26} Elevated CRP levels at diagnosis are predictive for the requirement of future surgical interventions in CD and therefore for disease severity.²⁷ Also, increased CRP levels prior to treatment with the TNF- α -antagonist infliximab were predictive of a higher response rate and treatment success.^{25,28} Additionally, our data demonstrate that enhanced serum CRP levels correlate with inflammatory disease activity, as it was specifically enhanced in

the CD patient group with increased fecal calprotectin levels.

In the present study, serum IFN- γ levels were significantly elevated in CD patients with increased fecal calprotectin levels, as compared to patients with remissive disease. Likewise, it has previously been observed that circulating IFN- γ levels are strongly increased in IBD patients as opposed to healthy subjects.²⁹ IFN- γ is known for its central role in the Th1-driven immune response, which constitutes the major signaling pathway in the pathogenesis of CD.¹⁵ IFN- γ is predominantly produced by Th1-differentiated T-cells residing in the intestinal lamina propria upon stimulation by IL-12-producing macrophages in the nearby environment.³⁰ Therefore, since we observe a strong correlation between fecal calprotectin levels and serum IFN- γ , this cytokine may be considered as a representative marker of endoscopic disease activity driven by Th1-cell cytokine production (IL-6, IL-12, TNF- α and IFN- γ).^{15, 31} Given these findings, follow-up studies are warranted to determine whether therapeutic interference in patients with increased fecal calprotectin levels accompanied by high IFN- γ and CRP levels results in improved patient outcomes.

Finally, we observed a positive correlation between fecal calprotectin levels and serum amyloid A (SAA), an acute-phase protein implicated in multiple chronic inflammatory diseases and commonly found to be elevated in CD patients.³² SAA is an apolipoprotein of high-density lipoproteins (HDL) and plays a role in cholesterol transport to and from sites of inflammation.³³ It has previously been demonstrated that SAA correlates well with other acute-phase reactants, such as CRP and alpha-1-antichymotrypsin (alpha-1-ACT) and is suggested to be helpful in monitoring CD disease activity.^{24, 34}

In the present study, the concentrations of a large number of cytokines have been assessed using a highly-sensitive electrochemiluminescence (ECL) assay and were directly correlated to the fecal calprotectin level. To our best knowledge, no studies have focused on the direct correlations between fecal calprotectin levels and serum cytokine levels with such a wide dynamic range of detection, providing us with high sensitivity cytokine measurements. However, a limitation of this study is that no endoscopic results were obtained in this patient cohort, which is generally considered as gold standard for demonstrating disease activity in CD.^{35, 36}

In conclusion, we show that a positive correlation exist between multiple Th1- and Th17-associated cytokines and fecal calprotectin level, presenting additional non-invasive candidate biomarkers for clinical use in CD patients. Moreover, these candidate cytokine biomarkers could be of value in monitoring and treating disease activity in CD. Future studies should aim to further assess the diagnostic potential of a distinct cytokine profile

in relation to endoscopic activity measures.

ACKNOWLEDGEMENTS

We would like to thank research technicians Manon Buist-Homan and Janette Heegsma (Department of Gastroenterology and Hepatology) for their contributions to the experimental laboratory work.

REFERENCES

1. Sartor RB. Microbial influences in inflammatory bowel diseases. *Gastroenterology* 2008 Feb;134(2):577-594.
2. Cosnes J, Gower-Rousseau C, Seksik P, Cortot A. Epidemiology and natural history of inflammatory bowel diseases. *Gastroenterology* 2011 May 01;140(6):1785-1794.
3. Binder V, Hendriksen C, Kreiner S. Prognosis in Crohn's disease--based on results from a regional patient group from the county of Copenhagen. *Gut* 1985 February 01;26(2):146-150.
4. Colombel JF, Narula N, Peyrin-Biroulet L. Management Strategies to Improve Outcomes of Patients With Inflammatory Bowel Diseases. *Gastroenterology* 2017 February 01;152(2):361.e5.
5. Jones J, Loftus EV, Panaccione R, Chen LS, Peterson S, McConnell J, et al. Relationships between disease activity and serum and fecal biomarkers in patients with Crohn's disease. *Clin Gastroenterol Hepatol* 2008 November 01;6(11):1218-1224.
6. Sipponen T, Savilahti E, Kolho KL, Nuutinen H, Turunen U, Farkkila M. Crohn's disease activity assessed by fecal calprotectin and lactoferrin: correlation with Crohn's disease activity index and endoscopic findings. *Inflamm Bowel Dis* 2008 January 01;14(1):40-46.
7. Benitez JM, Meuwis MA, Reenaers C, Van Kemseke C, Meunier P, Louis E. Role of endoscopy, cross-sectional imaging and biomarkers in Crohn's disease monitoring. *Gut* 2013 December 01;62(12):1806-1816.
8. Boschetti G, Garnero P, Moussata D, Cuerq C, Preaudat C, Duclaux-Loras R, et al. Accuracies of serum and fecal S100 proteins (calprotectin and calgranulin C) to predict the response to TNF antagonists in patients with Crohn's disease. *Inflamm Bowel Dis* 2015 February 01;21(2):331-336.
9. Langhorst J, Elsenbruch S, Koelzer J, Rueffer A, Michalsen A, Dobos GJ. Noninvasive markers in the assessment of intestinal inflammation in inflammatory bowel diseases: performance of fecal lactoferrin, calprotectin, and PMN-elastase, CRP, and clinical indices. *Am J Gastroenterol* 2008 January 01;103(1):162-169.
10. Tibble J, Teahon K, Thjodleifsson B, Roseth A, Sigthorsson G, Bridger S, et al. A simple method for assessing intestinal inflammation in Crohn's disease. *Gut* 2000 October 01;47(4):506-513.
11. Foell D, Wittkowski H, Roth J. Monitoring disease activity by stool analyses: from occult blood to molecular markers of intestinal inflammation and damage. *Gut* 2009 June 01;58(6):859-868.
12. Lehmann FS, Burri E, Beglinger C. The role and utility of faecal markers in inflammatory bowel disease. *Therap Adv Gastroenterol* 2015 January 01;8(1):23-36.
13. Manz M, Burri E, Rothen C, Tchangui N, Niederberger C, Rossi L, et al. Value of fecal calprotectin in the evaluation of patients with abdominal discomfort: an observational study. *BMC Gastroenterol* 2012 January 10;12:5.
14. Sands BE. Biomarkers of Inflammation in Inflammatory Bowel Disease. *Gastroenterology* 2015 October 01;149(5):1285.e2.
15. Brand S. Crohn's disease: Th1, Th17 or both? The change of a paradigm: new immunological and genetic insights implicate Th17 cells in the pathogenesis of Crohn's disease. *Gut* 2009 August 01;58(8):1152-1167.
16. van Dullemen HM, van Deventer SJ, Hommes DW, Bijl HA, Jansen J, Tytgat GN, et al. Treatment of Crohn's disease with anti-tumor necrosis fac-

- tor chimeric monoclonal antibody (cA2). *Gastroenterology* 1995 July 01;109(1):129-135.
17. Park JH, Peyrin-Biroulet L, Eisenhut M, Shin JI. IBD immunopathogenesis: A comprehensive review of inflammatory molecules. *Autoimmun Rev* 2017 April 01;16(4):416-426.
 18. Neurath MF. Cytokines in inflammatory bowel disease. *Nat Rev Immunol* 2014 May 01;14(5):329-342.
 19.) Baumgart DC, Sandborn WJ. Crohn's disease. *Lancet* 2012 November 03;380(9853):1590-1605.
 20. O'Connor W, Kamanaka M, Booth CJ, Town T, Nakae S, Iwakura Y, et al. A protective function for interleukin 17A in T cell-mediated intestinal inflammation. *Nat Immunol* 2009 June 01;10(6):603-609.
 21. Sakuraba A, Sato T, Kamada N, Kitazume M, Sugita A, Hibi T. Th1/Th17 immune response is induced by mesenteric lymph node dendritic cells in Crohn's disease. *Gastroenterology* 2009 November 01;137(5):1736-1745.
 22. Fuss IJ. IL-17: intestinal effector or protector? *Mucosal Immunology* 2011;4:366-367.
 23. Saverymuttu SH, Hodgson HJ, Chadwick VS, Pepys MB. Differing acute phase responses in Crohn's disease and ulcerative colitis. *Gut* 1986 July 01;27(7):809-813.
 24. Niederau C, Backmerhoff F, Schumacher B, Niederau C. Inflammatory mediators and acute phase proteins in patients with Crohn's disease and ulcerative colitis. *Hepatogastroenterology* 1997 February 01;44(13):90-107.
 25. Vermeire S, Van Assche G, Rutgeerts P. C-reactive protein as a marker for inflammatory bowel disease. *Inflamm Bowel Dis* 2004 September 01;10(5):661-665.
 26. Solem CA, Loftus EV, Tremaine WJ, Harmsen WS, Zinsmeister AR, Sandborn WJ. Correlation of C-reactive protein with clinical, endoscopic, histologic, and radiographic activity in inflammatory bowel disease. *Inflamm Bowel Dis* 2005 August 01;11(8):707-712.
 27. Henriksen M, Jahnsen J, Lygren I, Stray N, Sauar J, Vatn MH, et al. C-reactive protein: a predictive factor and marker of inflammation in inflammatory bowel disease. Results from a prospective population-based study. *Gut* 2008 November 01;57(11):1518-1523.
 28. Louis E, Vermeire S, Rutgeerts P, De Vos M, Van Gossum A, Pescatore P, et al. A positive response to infliximab in Crohn disease: association with a higher systemic inflammation before treatment but not with -308 TNF gene polymorphism. *Scand J Gastroenterol* 2002 July 01;37(7):818-824.
 29. Singh UP, Singh NP, Murphy EA, Price RL, Fayad R, Nagarkatti M, et al. Chemokine and cytokine levels in inflammatory bowel disease patients. *Cytokine* 2016 January 01;77:44-49.
 30. Fuss IJ, Neurath M, Boirivant M, Klein JS, de la Motte C, Strong SA, et al. Disparate CD4+ lamina propria (LP) lymphokine secretion profiles in inflammatory bowel disease. Crohn's disease LP cells manifest increased secretion of IFN-gamma, whereas ulcerative colitis LP cells manifest increased secretion of IL-5. *J Immunol* 1996 August 01;157(3):1261-1270.
 31. Karczewski J, Swora-Cwynar E, Rzymiski P, Poniadzialek B, Adamski Z. Selected biologic markers of inflammation and activity of Crohn's disease. *Autoimmunity* 2015;48(5):318-327.

32. De Beer FC, Mallya RK, Fagan EA, Lanham JG, Hughes GR, Pepys MB. Serum amyloid-A protein concentration in inflammatory diseases and its relationship to the incidence of reactive systemic amyloidosis. *Lancet* 1982 July 31;2(8292):231-234.
33. Benditt EP, Hoffman JS, Eriksen N, Parmelee DC, Walsh KA. SAA, an apoprotein of HDL: its structure and function. *Ann N Y Acad Sci* 1982;389:183-189.
34. Chambers RE, Stross P, Barry RE, Whicher JT. Serum amyloid A protein compared with C-reactive protein, alpha 1-antichymotrypsin and alpha 1-acid glycoprotein as a monitor of inflammatory bowel disease. *Eur J Clin Invest* 1987 October 01;17(5):460-467.
35. Sipponen T. Diagnostics and prognostics of inflammatory bowel disease with fecal neutrophil-derived biomarkers calprotectin and lactoferrin. *Dig Dis* 2013;31(3-4):336-344.
36. Dignass A, Van Assche G, Lindsay JO, Lemann M, Soderholm J, Colombel JF, et al. The second European evidence-based Consensus on the diagnosis and management of Crohn's disease: Current management. *J Crohns Colitis* 2010 February 01;4(1):28-62.

